



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

aabc@abc.org.br

Academia Brasileira de Ciências

Brasil

Bezerra, Moisés F.; Lacerda, Luiz D.; Costa, Breno G.B.; Lima, Eduardo H.S.M.
Mercury in the sea turtle *Chelonia mydas* (Linnaeus, 1958) from Ceará coast, NE Brazil
Anais da Academia Brasileira de Ciências, vol. 84, núm. 1, 2012, pp. 123-128
Academia Brasileira de Ciências
Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32721622012>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Mercury in the sea turtle *Chelonia mydas* (Linnaeus, 1958) from Ceará coast, NE Brazil

MOISÉS F. BEZERRA¹, LUIZ D. LACERDA¹, BRENO G.B. COSTA¹ and EDUARDO H.S.M. LIMA²

¹Laboratório de Biogeoquímica Costeira, Instituto de Ciências do Mar, Universidade Federal do Ceará,
Av. Abolição 3207, 60165-081 Fortaleza, CE, Brasil.

²Fundação Centro Brasileiro de Proteção e Pesquisa das Tartarugas Marinhas. Acesso Projeto TAMAR 151,
Alto Alegre, 62592-000 Itarema, CE, Brasil.

Manuscript received on December 16, 2010; accepted for publication on April 18, 2011

ABSTRACT

Mercury concentrations in carapace fragments of the green turtle *Chelonia mydas* from the Ceará coast in NE Brazil are reported. Concentrations varied from <0.34 to 856.6 ng.g⁻¹ d.w., and were highest (average of 154.8 ng.g⁻¹ d.w.) in juveniles (n = 22), whereas lowest concentrations (average of 2.5 ng.g⁻¹ d.w.) were observed in adult/sub-adult animals (n = 3). There was a significant negative correlation between animal size and Hg concentration probably due to different diets between juveniles and sub-adults/adults. Carapace fragments, which are non-invasive, non-lethal substrates, may be of importance for monitoring purposes of these generally endangered species.

Key words: Pollution, metals, marine turtles, monitoring.

INTRODUCTION

Sea turtles are widely spread throughout the world's tropical and temperate oceans. They are long-living animals and forage on diverse marine food chains; therefore, they have been proposed as bio-indicators of anthropogenic impacts on oceanic systems (Kampalath et al. 2006, Day et al. 2005). Also, sea turtles have been the object of many studies on the effects of global scale, persistent contaminants, mostly metals and organic micro-pollutants, which

have been responsible for the decrease in their populations (Storelli and Marcotrigiano 2003, Lam et al. 2004). Such studies have been carried out among sea turtle populations in the Mediterranean (e.g. Storelli and Marcotrigiano 2003, Storelli et al. 2005) in the Pacific Ocean (e.g. Sakai et al. 2000a, b, Lam et al. 2004) and in the North Atlantic Ocean (e.g. Day et al. 2005). In Brazil, concentrations of metal pollutants, including Hg, are scarcely known, and the available results are based on a few individuals and restricted in geographical distribution along the country's coast (Soto et al. 2005, Barbieri 2009).

Correspondence to: Luiz Drude de Lacerda
E-mail: ldrude@pq.cnpq.br

Among the most ubiquitous global persistent pollutants, Hg is frequently found in relatively high concentrations in the open ocean biota due to worldwide atmospheric deposition, in particular in the northern hemisphere due to elevated anthropogenic emissions and its property of biomagnification through long food chains. Therefore, Hg is considered a key pollutant to which sea turtles are exposed (Kampalath et al. 2006, Day et al. 2005).

The coast of Ceará in NE Brazil is a known feeding, growing and migration corridor of 5 species of sea turtles. The green turtle, *Chelonia mydas* (Linnaeus, 1958), is the most frequent and abundant species that occurs in this littoral. The species is considered threatened of extinction in Brazil and at a global level (Lima 2001). *Chelonia mydas* is an omnivorous species. When juveniles, however, *C. mydas* mostly preys on small benthic and pelagic invertebrates, whereas as adults/sub-adults this turtle becomes almost strictly herbivorous, feeding on algae and sea grasses (Cardona et al. 2010).

The major pathway of Hg exposure and accumulation in long-living marine animals including birds, mammals and sea turtles is their diet (Kehrig et al. 2009), but little is known on the uptake, bioaccumulation and bioconcentration mechanisms that result in a given body or organ concentration mainly due to the difficulty in sampling and handling these animals under strict protocols based on wildlife protection standards. Most studies are based on results from dead or dying animals generally in small numbers (Storelli et al. 2005, Storelli and Marcotrigiano 2003, Lam et al. 2004). Therefore, any additional, reliable data on pollutant concentrations in these organisms are important to sum up a workable database to help the understanding of pollutant fate in this oceanic biota.

The major objective of this study is to provide additional Hg concentration data on the green turtle *C. mydas* from a region where no information is available. Also, it aims at testing the response of Hg concentration in carapace fragments, which

are non-invasive, non-destructive substrates, to animal size and diet, and finding out on their utility to improve future monitoring programs, respecting wildlife protection protocols, on Hg distribution in the marine biota.

MATERIALS AND METHODS

All samples collected for this study followed the accepted protocols for wildlife handling and sampling according to SISBIO License No. 21693-1 from October 2009. Ceará coastline extends through 573 km in the semi-arid region of NE Brazil. This western portion of this coast is an important feeding site and migration route of 5 from the 7 known species of sea turtles, which resulted in the establishment of a protection and conservation unit of the Brazilian Program on Sea Turtle Conservation (TAMAR/ICMBio) from the Ministry of the Environment. The unit is located in Almofala Beach, Itarema Municipality (02°56'18" S; 039°48'51" W) (Figure 1), where traditional fisheries use mainly fish weirs that frequently capture sea turtles (Lima 2001).

The sampling included 20 living individuals accidentally trapped in fish weirs and captured by the local TAMAR/ICMBio staff and 5 individuals that were found dead on the beach. The trapped turtles were immediately set free after the biometric data collection. The sampled animals varied in size from 27.4 to 103.1 cm (average 50.5 ± 17.0 cm) and weighted from 1.7 to 45.0 kg (average 11.4 ± 8.6 kg). The animals could be divided in roughly two groups based on size, one represented by 22 juvenil individuals with CCL varying from 36.4 and 55.3 cm and a second group of 3 sub-adult/adult individuals with CCL varying from 85.0 and 103.1 cm. Keratinized carapace fragments were collected from both groups. Only those fragments presenting signs of loss were collected in living animals, e.g. due to barnacle incrustation. Fragments were labeled and preserved frozen in plastic bags until analysis. Species identification and biometry, weight, curved carapace length (CCL) and carapace width (CW), were obtained during sampling.




Figure 1. Map showing the location of Almofala coast in Ceará State, Brazil, where samples were obtained.

In the laboratory keratin fragment samples were treated in an ultra-sonic bath to fully remove algae and other incrusting residues and sediment grains. Duplicate sub-samples of each individual weighting about 0.5 g were transferred to Teflon[®] tubes containing 10 mL of concentrated HNO₃ and left there for 1 hour. The tubes were then transferred to a microwave oven (MARS XPRESS – CEM), 400 W at 200° C for 30 min. After cooling, 1 mL of H₂O₂ was added and the solution diluted to 100 mL with ultra-pure (< 3μS) water. Mercury concentrations were quantified through cold vapor atomic absorption spectrophotometry (CV-AAS) in a Nippon Instrumentation Corporation (NIC RA-3) spectrophotometer. Simultaneously, Hg concentrations were measured in duplicate in two reference standards, and the recovery of referenced values varied from 105% for BCR 060 (aquatic plant tissue) with nominal Hg concentration of 340 ng.g⁻¹ to 106% for NIST 2976 (mussel tissue) with nominal Hg concentration of 64 ng.g⁻¹. The detection limit of the procedure was 0.59 ng (0.34 ng.g⁻¹) considering three times the standard deviation of reagent blanks (EPADYX) divided by the inclination of the calibration curve (2.5 – 20 ng).

RESULTS AND DISCUSSION

Mercury concentrations in carapace fragments among all sampled individuals varied from <0.34 to 856.6 ng.g⁻¹ d.w. The highest Hg concentrations, with an average of 154.8 ng.g⁻¹ d.w., were observed in the

juvenile group, whereas the lowest concentrations, with an average of 2.5 ng.g⁻¹ d.w., were observed in sub-adult/adult animals. There was a significant negative correlation ($P < 0.01$; $n = 25$) between animal size (CCL) and Hg concentration (Figure 2) when the entire range of size is included, but no significant correlation was found between Hg concentrations and weight, which was expected since individuals may be under different foraging state and degree of stress due to capture, which would affect their body weight. Even when only juveniles are plotted, the negative correlation between Hg concentrations and animal size can still be identified (Figure 3).




Figure 2. Significant negative correlation ($n = 25$; $p < 0.01$) between total Hg concentrations and curved carapace length (CCL) observed in the sea turtle *C. mydas* from the Ceará coast in NE Brazil.

Sexual maturation in *C. mydas* starts when individuals reach about 1 m in CCL (Heppell et al. 2003), and individuals smaller than 35 cm in straight carapace length (SCL) (corresponding to 40 to 50 cm in CCL) (Zug et al. 2002) and with about 6 kg are considered immature juveniles. The major group of animals analyzed in this study thus fell in this category. After hatching, juvenile *C. mydas* enter 3 to 5 years of oceanic stage and forage on invertebrates from the neuston, as well as small benthic animals (Senko et al. 2010). When they recruit to inshore foraging habitats their diet shifts from the omnivorous planktivory of a pelagic turtle to the herbivorous diet of an inshore immature turtle finally foraging exclusively on benthic algae and

sea grasses when reaching sexual maturity (Arthur and Balazs 2008, Bolten 2003, Cardona et al. 2010). This changing of the major diet items and the large range in size explain, in part, the high variability of Hg concentrations observed among individuals. Juvenile individuals showed a significant negative correlation between size and Hg concentrations, which is in accordance with a progressive acquiring of herbivory and can be interpreted as a response to higher organic Hg contents in the neuston and benthic animals, preyed by juveniles, relative to benthic plants, preyed by mature turtles. This may explain the higher concentrations found in the smaller animals, which should have just been recruited to the inshore foraging habitat. Also the preference of juveniles for coastal habitats can result in a higher exposure to Hg that is relatively abundant in these environments, as shown for the Ceará coast (Lacerda et al. 2007). Kampalath et al. (2006) also observed negative correlations between animal size and Hg content, but in the internal organs of *C. mydas*. These authors also associated this pattern with a change in diet between smaller and larger animals. It is important to consider that the pattern verified for the carapace fragments in this study is the same found for internal organs by these authors.




Figure 3. Significant negative correlation ($n=19$; $p<0.05$) between total Hg concentrations and curved carapace length (CCL) of living juvenile individuals of *C. mydas* from the Ceará coast in NE Brazil.

Few studies using the carapace fragments of *C. mydas* as an experimental subject have been published, making difficult the comparison of our results. Sakai et al. (2000a, b) found Hg concentrations varying from 2.09 to 2.79 ng.g^{-1}

in two adult individuals. These concentrations are in agreement with those found in the three adult individuals from the Ceará coast. They also found a good correlation between the Hg content in these carapaces and Hg concentrations in the whole body, confirming the possible use of carapace fragments as a desirable non-invasive sample for monitoring purposes. Lam et al. (2004) reported Hg concentrations in *C. mydas* varying from 200 ng.g^{-1} in lungs to 780 ng.g^{-1} in the liver of two juvenile individuals. In the same study they reported Hg concentrations in the muscles and liver of 3 adult individuals of 53 ng.g^{-1} and 124 ng.g^{-1} , respectively. Although the carapace was not analyzed by these authors, our average results, even for juveniles, are in the lower range of reported Hg concentrations for this species, which suggests a smaller Hg exposure.

The most available results are from other turtle species, which unfortunately hampers direct comparisons with our results, but may provide an overview of the Hg distribution in this group of animals. Day et al. (2005), for example, found comparable results, both regarding the concentration range and distribution among individuals of different size for *Caretta caretta*. A. Green, unpublished data observed a significant correlation between the Hg concentrations in the carapace with those present in other internal organs in freshwater turtle species from Asia. This author also found a significant negative correlation between Hg content and size of individuals, which is also associated with different diets. Unfortunately all these results are based on less than 10 animals, thus still avoiding the discussion of a general pattern of Hg fate in sea turtles. In Brazil, only two studies, to our knowledge, have quantified metal concentrations in sea turtles, and only one dealt with Hg (Soto et al. 2005). All of them have analyzed internal organs (Barbieri 2009). In another study, Schneider et al. (2009) published the only account of Hg concentrations in carapace fragments, of the freshwater turtle *Podocnemis erythrocephala* from

the Amazon region and found strong correlations between Hg and size of the individuals, but no significant correlations between Hg concentration in internal organs and size.

The Hg concentrations observed in the carapace fragments of the larger (adult/sub-adult) animals sampled in the present study are very low as a response to their herbivorous diet. However, those found in juveniles are much higher, which is also a response to the preferentially carnivorous diet. The concentrations are in general lower than the Hg content found in top marine carnivorous animals such as mammals and sea birds (Bond and Antony 2009, Kehrig et al. 2009).

CONCLUSION

The results presented in this study are the first ever published on the Hg concentrations in *C. mydas* in northeastern Brazil using keratinized carapace fragments. Concentrations are in the lower range of those observed in the carapaces of other species of sea turtles, and much lower than those observed in internal organs. On the other hand, carapace fragments proved to be a reliable, available and non-invasive, non-lethal subject for the Hg monitoring. Much more research, however, is needed to associate carapace concentrations with Hg body burden and environmental Hg levels, since carapace fragments could be the basis for a future monitoring program on the Hg contamination of pelagic environments along the Brazilian coast. Thus, it would avoid unnecessary deaths of animals and allow the work with living animals actually exposed to present Hg concentrations in the oceans.

ACKNOWLEDGMENTS

We are deeply indebted to Projeto TAMAR/ICMBio Regional Ceará for logistic support and sample collection. The Instituto Nacional de Ciência e Tecnologia de Transferência de Material Continente-

Oceano (INCT-TMCOcean), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq Proc. No. 573.601/2008-9) funded this research.

RESUMO

As concentrações de Hg em fragmentos de carapaça de *Chelonia mydas* no litoral do Ceará, nordeste do Brasil, são reportadas. Concentrações variaram de <0,34 a 856,6 ng.g⁻¹ em peso seco, e foram maiores (média de 154,8 ng.g⁻¹ em peso seco) em indivíduos juvenis (n = 22), enquanto que as menores concentrações (média de 2,5 ng.g⁻¹ em peso seco) foram observadas em indivíduos adultos/sub-adultos (n = 3). Houve uma correlação negativa significativa entre tamanho do animal e concentração de Hg provavelmente devido a diferença de dieta entre juvenis e sub-adultos/adultos. Fragmentos de carapaça, que constituem substratos não-invasivos e não letais, podem ser importantes para fins de monitoramento ambiental dessas espécies ameaçadas de extinção.

Palavras-chave: Poluição, metais, tartarugas marinhas, monitoramento.

REFERENCES

- ARTHUR KE AND BALAZS GH. 2008. Comparison of immature green turtle (*Chelonia mydas*) diets among seven sites in the main Hawaiian Islands. *Pacific Sci* 62: 205-217.
- BARBIERI E. 2009. Concentration of heavy metals in tissues of green turtles (*Chelonia mydas*) sampled in the Cananéia estuary, Brazil. *Braz J Oceanogr* 57: 243-248.
- BOLTEN AB. 2003. Variation in sea turtle life history patterns: Neritic vs. oceanic development stages. In: Lutz PL, Musick JA and Wyneken J (Eds), *THE BIOLOGY OF SEA TURTLES*. CRC Press, Washington, DC 2: 243-257.
- BOND AL AND ANTONY WD. 2009. Total and methyl mercury concentrations in seabird feathers and eggs. *Environ Contam Toxicol* 56: 286-291.
- CARDONA L, CAMPOS P, LEVY Y, DEMETROPOULOS A AND MARGARITOULIS D. 2010. Asynchrony between dietary and nutritional shifts during the ontogeny of green turtles (*Chelonia mydas*) in the Mediterranean. *J Exper Mar Biol Ecol* 393: 83-89.

- DAY RD, CHRISTOPHER SJ, BECKER PR AND WHITAKER DW. 2005. Monitoring mercury in the loggerhead sea turtle, *Caretta caretta*. *Environ Sci Technol* 39: 437-446.
- HEPPELL SS, SNOVER ML AND CROWDER LB. 2003. Sea turtle population ecology. In: Lutz PL, Musick JA and Wyneken J (Eds), THE BIOLOGY OF SEA TURTLES. CRC Press, Washington, DC 2: 275-299.
- KAMPALATH R, GARDNER SC, MÉNDEZ-RODRÍGUEZ L AND JAY JA. 2006. Total and methylmercury in three species of sea turtles of Baja California Sur. *Mar Pollut Bull* 52: 1784-1832.
- KEHRIG HA, FERNANDES KWG AND MALM O. 2009. Transferência trófica de mercúrio e selênio na costa norte do Rio de Janeiro. *Quim Nov* 32: 1822-1828.
- LACERDA LD, SANTOS JA, CAMPOS RC, GONÇALVES RA AND SALLES R. 2007. Total-Hg and organic-Hg in *Cephalopholis fulva* (Linnaeus, 1758) from inshore and offshore waters of NE Brazil. *Braz J Biol* 67: 493-498.
- LAM JCW, TANABE S, CHAN SAKF, YUEN EKW, LAM MHW AND LAM PKS. 2004. Trace element residues in tissues of green turtles (*Chelonia mydas*) from South China Waters. *Mar Pollut Bull* 48: 164-192.
- LIMA EHS. 2001. Helping the people help the turtles: The work of Projeto TAMAR-IBAMA in Almofala, Brazil. *Mar Turtles Newslet* 91: 7-9.
- SAKAI H, SAEKI K, HICHIHASHI H, KAMIZAKI N, TANABE S AND TATSUKAWA R. 2000a. Growth-related changes in heavy metal accumulation in green turtle (*Chelonia mydas*) from Yaeyama Islands, Okinawa, Japan. *Archiv Environ Contam Toxicol* 39: 378-385.
- SAKAI H, SAEKI K, HICHIHASHI H, SUGANUMA H, TANABE S AND TATSUKAWA R. 2000b. Species-specific distribution of heavy metals in tissues and organs of loggerhead turtle (*Caretta caretta*) and green turtle (*Chelonia mydas*) from Japanese coastal waters. *Mar Pollut Bull* 40: 701-709.
- SCHNEIDER L, BELGER L, BURGER J AND VOGT RC. 2009. Mercury bioaccumulation in four tissues of *Podocnemis erythrocephala* (Podocnemididae: Testudines) as a function of water parameters. *Sci Tot Environ* 407: 1048-1054.
- SENKO J, LÓPEZ-CASTRO MC, KOCH V AND NICHOLS WF. 2010. Immature East Pacific green turtle (*Chelonia mydas*) use multiple foraging areas off the Pacific Coast of Baja California Sur, Mexico: First evidence from mark-recapture data. *Pacific Sci* 64: 125-130.
- SOTO JMR, SOARES JT, CELINI AAOS AND SANTOS RCA. 2005. Concentração de mercúrio total em tecidos de *Caretta caretta* (Linnaeus, 1758) (Reptilia, Cheloniidae) encalhadas na costa sul do Rio Grande do Sul, Brasil. In: 2ª Jornada de Conservação e Pesquisa de Tartarugas Marinhas no Atlântico Sul Ocidental, Rio Grande, p. 25-27.
- STORELLI MM AND MARCOTRIGIANO GO. 2003. Heavy metal residues in tissues of marine turtles. *Mar Pollut Bull* 46: 397-400.
- STORELLI MM, STORELLI A, D'ADDABBO R, MARANO C, BRUNO R AND MARCOTRIGIANO GO. 2005. Trace elements in loggerhead turtles (*Caretta caretta*) from the eastern Mediterranean Sea: overview and evaluation. *Environ Pollut* 135: 163-170.
- ZUG GR, BALAZS GH, WETHERALL JA, PARKER DM AND MURAKAWA SKK. 2002. Age and growth of Hawaiian green turtle (*Chelonia mydas*): An analysis based on skeletochronology. *Fish Bull* 100: 117-127.