



Avances en Biomedicina  
E-ISSN: 2244-7881  
avanbiomed.idic@ula.ve  
Universidad de los Andes  
Venezuela

Vielma, Silvana  
Inflammatory mediators in patients with Dengue infection: role in early endothelial damage  
Avances en Biomedicina, vol. 2, núm. 1, octubre, 2013, pp. 48-48  
Universidad de los Andes  
Mérida, Venezuela

Available in: <http://www.redalyc.org/articulo.oa?id=331330397009>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System  
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal  
Non-profit academic project, developed under the open access initiative

## Inflammatory mediators in patients with Dengue infection: role in early endothelial damage

Silvana Vielma, MD, PhD.

Departamento de microbiología e inmunología, Universidad de Los Andes. Mérida. Venezuela.

### [CONFERENCIA]

#### Abstract

Several mechanisms have been proposed to explain the pathogenesis of dengue virus (DENV) infection. One of the most important finding is the production of pro-inflammatory cytokines as responsible of endothelial cells activation, plasma leakage and subsequently, disease severity.

Methods: Our aim was to determined levels of inflammatory mediators (IL-8, TNF- $\alpha$ ), soluble cell adhesion molecules (sICAM-1 and sVCAM-1) and soluble lymphocyte activation markers (sIL-2R $\alpha$ , sTNF-RII/p75) during the course of DENV infection. Serum samples from fifty-two patients with confirmed DENV infection were collected during the acute (0-3 days after fever onset) and critical/recovery (4-7 days after onset) phases of the diseases. As controls, sera from non-febrile individuals were included. Patients were classified as dengue without warning signs (DwoWS), dengue with warning signs (DwWS) and severe dengue (SD).

Results: Levels of sIL-8 and sTNF- $\alpha$  were significant higher in cases of DwWS during the critical phase of the disease. A significant increase of sICAM-1 and sVCAM-1 were detected during the critical phase of the disease compare with controls, however, the highest levels of sVCAM-1 were observed during the acute phase of SD. Finally, sIL2-R showed a significantly increased during the acute phases of SD, while sTNF-RII/p75 was elevated in DwWS during the critical stage of infection.

Conclusion: sIL2-R and sVCAM-1 may be early markers of lymphocyte and endothelial damage in the acute phases of DENV infection in patients with SD.

**Keywords:** Pro-inflammatory cytokines, Endothelial damage, sICAM-1, sVCAM-1, sIL-2R $\alpha$ , sTNF-RII/p75, dengue without warning signs (DwoWS), dengue with warning signs (DwWS) and severe dengue (SD).