

**Revista Mexicana de
Astronomía y Astrofísica**

Revista Mexicana de Astronomía y Astrofísica

ISSN: 0185-1101

rmaa@astroscu.unam.mx

Instituto de Astronomía

México

Ayala, S.; Raga, A. C.; Curiel, S.

An Optical Spectroscopic Study of HH 110: a Turbulent Mixing Layer?

Revista Mexicana de Astronomía y Astrofísica, vol. 15, septiembre, 2003, p. 134

Instituto de Astronomía

Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=57115050>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

AN OPTICAL SPECTROSCOPIC STUDY OF HH 110: A TURBULENT MIXING LAYER?

S. Ayala,¹ A. C. Raga,² and S. Curiel³

The HH 110 jet extends $\sim 3'$ in length and consists of numerous knots forming the flow. Noriega-Crespo et al. (1996) found that the turbulent optical and near-infrared morphology of the HH 110 jet is consistent with that of a boundary layer. In this work, we have analyzed some line ratios along and across the jet in order to make a quantitative comparison with the line ratios predicted by the current mixing layer models.

We present calibrated intermediate resolution long-slit spectra along and across the jet. We calculate different relative line ratios and determine the electronic density (N_e) for all the sub-spectra. Then, we have constructed diagrams where the intensity distribution, different line ratios, and N_e are shown as a function of spatial position. All positions across the jet clearly show a monotonous decrement in the excitation degree from the east edge to the west edge. Along the jet there is no specific tendency for physical parameters.

The mean value for E_{B-V} along the jet is $E_{B-V} = 0.5 \pm 0.4$. Electronic densities derived from the [S II] 6717/31 ratio are within $50 \geq N_e \geq 1500 \text{ cm}^{-3}$. The observed line ratios in the HH 110 jet are similar to HH objects with high excitation, although the [O III] emission lines are not detected.

We compare three line ratios in HH 110 and several mixing layer models (Binette et al. 1999) and plane-parallel shock models (Hartigan, Morse, & Raymond 1994) and line ratios for HH objects with low excitation. The HH 110 line ratios do not fit with plotted mixing layer models. Line ratios predicted by mixing layer models would be classified as low excitation, while we have classified the HH 110 jet as a high-excitation object.

Using the [N II] 6584/H α and [O I] 6300/H α ratios and N_e we have estimated a T_e and X (model independent parameter, as in Bacciotti & Eislöffel




Fig. 1. Spatial distribution of the derived electronic temperature (T_e) and ionization fraction (X) along slits I, II, III and IV. We show the spatial intensity distribution of H α and [S II] 6717/31 lines.

1999), assuming that O and N dominate the charge-exchange reactions in an optically thin medium. Figure 1 presents the results for all analyzed slits. Values obtained for T_e in HH 110 are similar and slightly increase from the east edge to the west edge. The average temperature in the jet is $T_e = 4700 \pm 500 \text{ K}$. In the case of the ionization fraction estimated across the jet (slits II, III and IV), we can observe how these values decrease from the east edge to the west edge. These results are clearly consistent with the expected tendency of these physical parameters in a turbulent mixing layer. The values for T_e and X (~ 0.1) are similar to the same parameters estimated in other HH objects (HH 46/47, HH 24E, HL Tau and the Th 28 jet).

REFERENCES

- Bacciotti, F., & Eislöffel, J. 1999, A&A, 342, 717
 Binette, L., Cabrit, S., Raga, A., & Cantó, J. 1999, A&A, 346, 260
 Hartigan, P., Morse, J. A., & Raymond, J. 1994, ApJ, 436, 125
 Noriega-Crespo, A., Garnavich, P. M., Raga, A. C., Cantó, J., & Böhm, K.-H. 1996, ApJ, 462, 804

¹Instituto de Astronomía, Observatorio Astronómico Nacional, UNAM, Apartado Postal 877, 22800 Ensenada, B.C., México (sayala@astro.unam.mx).

²Instituto de Ciencias Nucleares, UNAM, Apartado Postal 70-543, 04510 México, D. F., México.

³Instituto de Astronomía, UNAM, Apartado Postal 70-264, 04510 México, D. F., México.