

**Revista Mexicana de
Astronomía y Astrofísica**

Revista Mexicana de Astronomía y Astrofísica

ISSN: 0185-1101

rmaa@astroscu.unam.mx

Instituto de Astronomía

México

de Ugarte Postigo, A.; Gorosabel, J.; Thöne, C. C.; Sánchez-Ramírez, R.

THE ROLE OF GTC IN GAMMA-RAY BURST SCIENCE

Revista Mexicana de Astronomía y Astrofísica, vol. 42, 2013, pp. 73-74

Instituto de Astronomía

Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=57127862035>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

THE ROLE OF GTC IN GAMMA-RAY BURST SCIENCE

A. de Ugarte Postigo,^{1,2} J. Gorosabel,¹ C. C. Thöne,¹ R. Sánchez-Ramírez¹ on behalf of a larger collaboration

RESUMEN

Durante el último año, el telescopio GTC de 10.4 m ha reivindicado su lugar como una de las instalaciones punteras a nivel mundial en el estudio de las explosiones de rayos gamma (GRB). GTC es el único telescopio de clase 8–10 m en terreno europeo, lo que le coloca en una situación privilegiada para las observaciones de oportunidad. El hecho de que se opera fundamentalmente en modo de servicio lo hace óptimo para este tipo de observaciones. En este artículo presentamos una visión general de los estudios de GRBs que se han realizado desde el comienzo de las operaciones de GTC.

ABSTRACT

Over the last year the 10.4 m GTC telescope has claimed its role as one of the leading facilities for gamma-ray burst (GRB) research world wide. GTC is the only large telescope within European soil, giving it an outstanding position for fast target of opportunity observations. The predominant service mode operation, makes it specially well suited for these kind of observations. We give an overview of the results obtained in GRB studies since the beginning of scientific operations.

Key Words: gamma-ray burst: general

1. GAMMA-RAY BURSTS

Gamma-ray bursts (GRBs) are the most luminous explosions in the Universe, releasing isotropic equivalent energies in the range of 10^{51} – 10^{54} erg in gamma-rays. In optical, they can reach, for a short period of time, luminosities 10^4 times brighter than typical quasars. Thanks to this, they have been observed at almost any redshift, from our nearby environment (GRB 980425 was the nearest one, at $z=0.08$, Galama et al. 1998) to the very distant Universe (GRB 090429 is the current record holder with a photometric redshift of $z=9.4$, Cucchiara et al. 2011, being GRB 090423 the furthest spectroscopically confirmed at $z=8.2$, Tanvir et al. 2009; Salvaterra et al. 2009). Their optical spectra is normally well reproduced by a clean, simple power law, making them not only great laboratories for extreme physics, but also ideal light houses to probe the material within the line of sight between the GRB and the observer. Here we give some examples of the ongoing work in the GRB field using the 10.4 m GTC.

2. FAST TARGET OF OPPORTUNITY OBSERVATIONS AT GTC

GTC has a very important role to play in target of opportunity observations, and in particular in

the ones that require fast response, due to its privileged location. Almost all the 8–10 m class telescopes that are currently in operation are located between longitudes of 70 and 155 deg West. Only the SALT telescope is located at an eastern longitude from GTC. However, SALT can only observe objects on transit through a restricted elevation range. This leaves GTC with 3.5 hours advantage (and after a 15 hour gap) over any other 8–10 m class telescope, to rapidly observe any event that occurs during the night of the eastern Hemisphere. Furthermore the service mode, dominant at GTC, is optimal for ToO observations. As an example we show, in Figure 1 the spectrum of GRB 110503A, which was observed with GTC. Thanks to the rapid delivery of the pre-reduced data by the GTC staff, the redshift of the afterglow was published in an astronomical circular just 42 minutes after observing (de Ugarte Postigo et al. 2011a).

3. CURRENT WORK WITH GTC

The large aperture of GTC is well suited for obtaining spectroscopy of the afterglows and host galaxies of GRBs, as well as very deep imaging. Furthermore, the tuneable filters available in OSIRIS, allow us to observe the spatially resolved hosts of nearby GRBs. Here we give some examples of our work:

Spectroscopy of GRB afterglows is the most reliable way to determine the distance at which they happen, which is essential to convert observations

¹Instituto de Astrofísica de Andalucía, Glorieta de la Astronomía, s/n, 18008, Granada, Spain (deugarte@iaa.es).

²Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, 2100, Copenhagen Ø, Denmark).

