



Acta Zoológica Mexicana (nueva serie)

ISSN: 0065-1737

azm@ecologia.edu.mx

Instituto de Ecología, A.C.

México

Servín, Jorge

Duration and frequency of chorus howling of the mexican wolf (*Canis lupus baileyi*)

Acta Zoológica Mexicana (nueva serie), núm. 80, 2000, pp. 223-231

Instituto de Ecología, A.C.

Xalapa, México

Available in: <http://www.redalyc.org/articulo.oa?id=57508013>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

DURATION AND FREQUENCY OF CHORUS HOWLING OF THE MEXICAN WOLF (*CANIS LUPUS BAILEYI*)

Jorge SERVÍN

Instituto de Ecología, A. C., Centro Regional Durango, Departamento de Fauna Silvestre,
Apdo. Postal 632, C.P. 34100, Durango, Dgo., MEXICO
e-mail: loboservin@omanet.com.mx

RESUMEN

El lobo mexicano (*Canis lupus baileyi*) es una especie amenazada de extinción y su biología básica ha sido poco estudiada. En este artículo se aporta información sobre la duración en tiempo y la frecuencia diaria de los aullidos en grupo del lobo mexicano a lo largo de un año. Los datos fueron obtenidos de una manada de cinco individuos adultos (tres machos y dos hembras), mantenidos en cautiverio en la Reserva de la Biosfera "La Michilía", en la Sierra Madre Occidental en el Estado de Durango, México. Doscientos ochenta y seis aullidos fueron registrados para la manada durante un año. Los aullidos en grupo fueron más comunes durante el período de apareamiento (Enero/Febrero) y tendieron a escucharse en el amanecer y anochecer, como en las poblaciones norteñas. La frecuencia diarias de aullidos en grupo, presentó una variación significativa a lo largo del año ($p < 0.005$); durante la temporada de apareamiento, se encontró un promedio de 2.3 aullidos por día y con una duración de 59.2 ± 6.6 seg; mientras que en Agosto sólo se registraron 0.9 aullidos al día y con menor duración (26.7 ± 3.5 seg). La duración promedio del coro de aullidos de este grupo fue de 41 ± 9 seg, tiempo más reducido con respecto a lo encontrado para las manadas de lobos en E.U.A. y Canadá.

Palabras clave: Aullidos, *Canis lupus baileyi*, duración, frecuencia, grupo, lobo mexicano.

ABSTRACT

The Mexican wolf (*Canis lupus baileyi*) is an endangered species but remains poorly studied. I report the Mexican wolf chorus howling frequency through the year, from a pack with five adults (three males and two females) in captivity in "La Michilía" Biosphere Reserve at the Western Sierra Madre in Durango, Mexico. These is the first quantitative howling data reported for the Mexican wolf. Two hundred and eighty six chorus howls were recorded between January and December. Chorus howling was most common during the breeding season (January/February), and trends to listen at the dusk and sunrise, likely in wolves from northern populations. Daily howling chorus varied significantly in frequency between months ($p < 0.005$); in February, chorus occurred 2.3 per day and duration of 59.2 ± 6.6 sec, whereas in August 0.9 chorus per day and duration of 26.7 ± 3.5 sec was recorded. The mean duration of chorus were 41 ± 9 sec for this wolf pack, were shorter than those reported from wolves packs of the northern United States and Canada.

Key words: *Canis lupus baileyi*, duration, fequency, howling, Mexican wolf.

INTRODUCTION

The Mexican wolf (*Canis lupus baileyi*) has been described as the smallest in size of the 24 American subspecies of *Canis lupus* (Young and Goldman 1944). Two of which *C.l. monstrabilis* and *C.l. baileyi* were recognized for Mexico (Hall 1981). Although now restricted to the West Sierra Madre and adjoining tableland region of Western Mexico (McBride 1980, Servín 1986, 1996), *C. l. baileyi* formerly inhabited portions of southeastern Arizona, southern New Mexico, and western Texas, as well as reaching into the southern Valley of México (Young and Goldman 1944). Currently its population is fragmented into small areas in the States of Durango and probably Chihuahua and Sonora (Servín 1996). This endangered subspecies is protected by Mexican law throughout its historic range (Mech, 1982). Biological knowledge of this species in Mexico is scarce, although conservation and restoration plans have been started (Aldama-Garisonain 1996, García-Moreno *et al.* 1996, Parson 1998, Siminski 1998). Its social behavior has been studied (Servín 1991), and growth rates (Servín 1997) and count of the remaining wild populations were undertaken in Durango State (Servín 1986, 1996).

The wolf is a strict carnivore with a complex spatial and social organization that is mediated by visual, olfactory and acoustic communication (Mech, 1970). Wolf howling has crucial social roles both within the pack as well as among packs (Mech 1970, Harrington and Mech 1979). Within a pack, howling helps to bring together its members and coordinate social activities and movements. Among packs, howling conveys information about the locations of neighbors, thereby helping to lower aggressive encounters (Joslin 1967, Harrington and Mech 1979, 1983, Mech 1970). All data previously reported on howling has come from North American wolf populations (Harrington and Mech 1983, Joslin 1976, Mech 1970). On the other hand, no data on howls have been obtained on the Mexican wolf.

Little is known of the natural history of the Mexican wolves. Some data obtained from old trappers in the mountains, suggested that wolves were rare, that packs were uncommon, and that individual wolves are more likely to hunt their small prey alone rather than in groups. Recently, it has been suggested that social behavior and groups sizes have evolved differently in southern, as compared to northern ecosystems (Bernardz 1988, Servín 1991). Thus, it is possible that sound communication in southern wolves has different qualities, such as duration and frequency howling, compared to those in northern wolves. *C. l. baileyi* has special scientific interest, because as a group, they are the most genetically distinct population of North American gray wolf, and our finding would support hypothesis of subtle adaptation to the

ecological conditions at the extreme southern limits of the species' range. If this is the case, then it could be expected that howling could be different in the Mexican wolf. This study describes chorus howling, including mean daily frequencies and average duration, throughout a year. Despite the limitations inherent in a captive study, this information can be used in the field to look for wild Mexican wolves in remote areas. It is important to know, when and how often Mexican wolves howl, to be able to use howling as a censusing tool.

MATERIALS AND METHODS

Howling data were obtained from a pack of five adult wolves (three males and two females) housed in a 1.5 ha. enclosure at the Biosphere Reserve (BR) "La Michilia", Durango; 80 km south of Durango city and 55 km west of Vicente Guerrero, in "La Sierra de Michis", at an altitude of 2450 meters. Annual precipitation is 65-80 cm, 70% of which falls from July to September; the dry season is from January to May. Annual average temperature is 11°C (range -14°C until 38°C). The dominant vegetation is oak-pine forest (*Quercus* spp. and *Pinus* spp.). The enclosure is situated within the original historic range of the Mexican wolf. Social structure in the pack was stable during this study, with a dominant pair, a subordinate pair, a low ranking male, and without pups (Servín 1991).

All instances of chorus howling were recorded around the enclosure (<1000 m of distance), usually from the Biological Station "Piedra Herrada" located 500 m from the wolves enclosure, thus allowing almost continuous monitoring of wolf howls. All howling records were spontaneous; wolves were not stimulated by any mechanism like sirens that could have provided stimulation for their howling.

I considered a chorus howl, as an event during which two or more members of the pack howled at the same time. A single howl by one wolf (solo howl) was omitted from the chorus howl count. Duration of howls, was the length of time that the chorus howls lasted and was measured in seconds; daily frequency of howls was the number of howls in one sample day. Monthly averages were obtained from these data.

Howling was sampled during 20 days every month, from 06:00 to 10:00 in the morning and 18:00 to 24:00 h at night, for a total of 10 h of howling data collection each day, 200 h each month and 2400 h per year. Monthly data based on the frequency and duration of howlings were analyzed using descriptive statistics. Monthly frequencies and duration of howling were compared with "goodness of fit" χ^2 test, under the null hypothesis

of no monthly variations. Monthly mean variation in howling duration was analyzed by one way analysis of variance (ANOVA) (Sokal & Rohlf 1981).

RESULTS

A total of two hundred and eighty six chorus howls were recorded for the wolf pack at "La Michilia" BR between January and December 1986. Data suggest that chorus howling tends to occur more frequently at dusk and sunrise. A representative sample for a statistical analysis was not obtained for comparing nighttime and daytime data, since samples were unequal and insufficient. However some trends were observed in howling schedule.

Daily frequency of chorus howls

Daily frequency of chorus howling showed significant differences during different times of the year ($\chi^2=40.29$; $df=11$; $p<0.005$). During February, when wolves had their breeding season, they displayed the highest daily mean frequency of howls ($\bar{x}=2.3$ howls/day); after the breeding season, the mean frequency of howls dropped, so that in March there was a mean of 1.2 howls each day, and the frequency remained low until November ($\bar{x}\leq 1.2$ howls/day). Finally, in December the mean increased again ($\bar{x}=1.5$ howls/day) (Fig. 1).




Figure 1

Mean of the chorus howling per day during the year for a Mexican wolf pack in Durango, México.

Duration of chorus howling

Mean duration of chorus howling of "La Michilia" pack was 41 ± 9 sec (range 20 to 79 sec). Mean monthly duration of chorus howling varied significantly throughout the year ($F=124$; $df=11$; $p<0.001$) (Figure 2). During February, the pack had the highest mean duration of chorus howling (59.2 ± 6.6 sec); the mean duration of howls abruptly decreased in March (39.8 ± 4.7 sec), and remained low until December ($\bar{x} \leq 49$ sec). In January the mean increased again (51.7 ± 3.5 sec) (Fig. 2).




Figure 2

Mean and standard deviation of the chorus howling durations throughout the year for a Mexican wolf pack in Durango, Mexico.

A comparison of Chorus Howling

An important datum obtained in this research, is the seasonal change in the chorus duration of Mexican wolves and its increase during breeding season.

The duration of chorus howling emitted by the Mexican wolf (41 ± 9 sec) is shorter than that reported from wolves in the northern United States, for which the mean was 60 sec (range 30 to 130 sec) (Harrington 1989); for Canadian wolves the mean was 85 sec, but the range was not reported (Joslin 1967). Hence, the mean duration of chorus howling varies among wolves from Canada, the U.S.A. and Mexico (Fig. 3).




Figure 3
Comparison between gray wolf chorus howling duration in Canada, United States of America and Mexico.

DISCUSSION

Zimen (1982) reported an increase of howls/day and its duration during the breeding season; also a preference for howling between 20:00-08:00 h is also reported by Joslin (1967) and Harrington and Mech (1978). The following characteristics were present in "La Michilia" wolves: howling occurred between sunset and sunrise, seasonal variation in the duration of chorus howling, and an increase during breeding season. Chorus howling usually occurs when wolves are resting, or in rally, a behavior called "greeting ceremony" by some authors (Mech 1970, Zimen 1982).

Daily frequency of chorus howls

Howling increased before the breeding season (December) and reached its peak in the last two weeks of February, when the reproductive season and successful mating occurred (Servín 1991,1997). These data suggest a close association between high daily frequency of howling and the breeding season, and the social behaviors (aggressive, sexual and friendly behaviors) related to both (Klinghammer 1978, Servín 1991, Zimen 1982). These recorded howls had similar seasonality to those reported

from northern wolves in Canada (Joslin 1967) and elsewhere in the United States (Harrington and Mech 1979, Klinghammer 1978) and Europe (Zimen 1982).

Some authors believe that the low frequency of daily howls was due to the reluctance of adults to howl when their pups were small. These authors reported that adults would not howl until the pups were 6 to 9 weeks old; this lack of vocalization would help to protect the vulnerable pups from possible predators (Joslin 1967, Harrington and Mech 1978, Rabb *et al.* 1967, Zimen 1982). In "La Michilia" pack, pups were born at the end of April (Servín 1997), and the adults stopped their howls until late July, more than 10 weeks later.

Duration of chorus

It is interesting that there were significant differences in the duration of chorus howling throughout the year, particularly in the breeding season. These findings strengthen the hypothesis that howling in wolves has crucial social roles, both within the pack and among packs (Mech and Harrington 1978).

These data suggest that howls could be an interesting parameter to be studied in order to understand communications among packs throughout a latitudinal gradient. Since in temperate areas the duration of howling is shorter than in the Arctic zone. Harrington (1989) reported that chorus in very small packs (i.e., just two adults) had shorter duration than those from larger packs.

Regional comparison of Chorus Howling

The average of the chorus howling in "La Michilia" pack is shorter than those reported from wolves of the northern United States (Harrington 1989), and Canadian wolves (Joslin 1967). Hence, the howling range in group varies among wolves from Canada, U.S.A. and Mexico (Fig. 3).

These data support the hypothesized relationship between latitude and average pack size, with small group size expected from the southern regions in Mexico (Servín 1991). Data from Canada reported the duration of individual wolf howls as lasting from five to eleven seconds; hence, packs with ten or more wolves will howl in chorus for more than one minute. Considering this Northern howling pattern, in Mexico the howling chorus would be shorter than one minute, as I found and report in this paper. However, Harrington (1989) has reported that the relationship between pack size and duration of chorus howling was not a linear one. Packs of two or three adults were different, but there were not noticeable differences in packs of five or six adults. However, does not take into consideration the significant variation during the year.

ACKNOWLEDGEMENTS

This paper is dedicated to Jaime Gonzalez (†) for his important field work developed in surveys wolves in the mountains of Mexico. The draft was kindly revised by F.H. Harrington, L.D. Mech, P. Joslin, J. Packard, S. Lindaker, Z. Tang-Martínez, I. A. Rosen and two anonymous referees. Thanks are due to Pedro Reyes-Castillo, Gonzálo Halffter, Alberto González and Sonia Gallina, for their encouragement for the Mexican Wolf Project; to Martha Vences, Macamen Huxley, Elias Chacón, Federico Alvarado, Jose Medina and José Angel Moreno for their field work assistance. This project was funded by Departamento de Fauna Silvestre (902-06) and is a contribution to the MAB-UNESCO Conservation Program.

LITERATURE CITED

- Aldama-Garisonain, A.** 1996. Proyecto de conservación y recuperación del lobo gris mexicano (*Canis lupus baileyi*). Informe Técnico. SEMARNAP-INE, 30 pp.
- Bernardz, J.A.** 1988. *The Mexican wolf: biology, history, and prospects for reestablishment in New Mexico*. U. S. Fish & Wildlife Service, Albuquerque, New Mexico, Endangered Species Report 18.
- García-Moreno, J., M.D. Matocq, M.S. Roy, E. Geffen & R.K. Wayne.** 1996. Relationships and genetic purity of the endangered Mexican wolf based on analysis of microsatellite loci. *Conserv. Biol.* 10:376-387.
- Hall, E.R.** 1981. *The mammals of North America*. 2nd. Ed: Wiley & Sons. 1181 pp.
- Harrington, F.H.** 1989. Chorus howling by wolves: Acoustic structure, pack size and the beaugeste effect. *Bioacoust.* 2:117-136.
- Harrington, F.H. & L.D. Mech.** 1978. Howling at two Minnesota wolf pack summer homesites. *Can. J. Zool.* 56:2024-2028.
- _____. 1979. Wolf howling and its role in a territory maintenance. *Behav.* 58:207-249.
- _____. 1983. Wolf pack spacing: howling as a territory-independent spacing mechanism in a territorial population. *Behav. Ecol. & Sociobiol.* 12:161-168.
- Joslin, P.** 1967. Movements and homesites of timber wolves in Algonquin Park. *Amer. Zool.* 7:279-288.
- Klinghammer, E.** 1978. Analysis of 14 months of daily howl records in a captive wolf pack. Pp. 153-181 In: E. Klinghammer (Ed). *Proceedings of Symposium On the Behavior and ecology of wolves*. Garland Press, U.S.A.
- McBride, R.T.** 1980. The Mexican wolf (*Canis lupus baileyi*): a historical review and observations on its status and distribution. U.S. Fish and Wildlife Service, Albuquerque, New Mexico, Endangered Species Report 8.
- Mech, L.D.** 1970. *The wolf: the ecology and behavior of an endangered species*. Doubleday, N. Y.
- _____. 1982. The IUCN-SSC wolf specialist group. Pp. 327-333. In: F.H. Harrington & P.C. Paquet (Eds). *Wolves of the World*. Noyes Publications, Park Ridge, New Jersey.

- Parson, D.R.** 1998. "Green fire" returns to the Southwest: reintroduction of the Mexican wolf. *Wildl. Soc. Bull.* 26:799-807.
- Rabb, G.L., J.H. Woolpy & B.E. Ginsburg.** 1967. Social relations in a group of captive wolves. *Amer. Zool.* 7:305-311.
- Servín, J.** 1986. Estudio para la recuperación del lobo mexicano (*Canis lupus baileyi*) en el Estado de Durango, II Etapa. Informe Técnico, Instituto de Ecología-SEDUE, México, 66 pp.
- _____. 1991. Algunos aspectos de la conducta social del lobo mexicano (*Canis lupus baileyi*) en cautiverio. *Acta Zool. Mex. (n.s.)*, 45:1-41.
- _____. 1996. Prospección y búsqueda del lobo mexicano (*Canis lupus baileyi*) en el estado de Durango, México. Informe Técnico. Instituto de Ecología-CONABIO, México, 31 pp.
- _____. 1997. El período de apareamiento, nacimiento y crecimiento del lobo mexicano (*Canis lupus baileyi*). *Acta Zool. Mex., (n.s.)*, 71:45-56.
- Siminski, P.** 1998. *Mexican gray wolf international studbook*. Arizona-Sonora Desert Museum. Tucson, Arizona, 104 pp.
- Sokal, R.R. & F.J. Rohlf.** 1981. *Biometry*. W. H. Freeman and Co., San Francisco, Calif.
- Young, S.P. & E.A. Goldman.** 1944. *The wolves of North America. Part II, Classification of wolves*. The American Wildlife Institute, Washington, D.C.
- Zimen, E.** 1982. A wolf pack sociogram. Pp. 282-322. *In: F.H. Harrington and P. Paquet (Eds). Wolves of the World. Perspectives of behavior, ecology and management*. Noyes Publications Park Ridge, New Jersey.

Recibido: 9 de febrero 1999
Aceptado: 28 de marzo 2000