



Boletín Latinoamericano y del Caribe de
Plantas Medicinales y Aromáticas

ISSN: 0717-7917

editor.blacpma@usach.cl

Universidad de Santiago de Chile
Chile

MUÑOZ-RAMIREZ, Alethia; PEREZ-GUTIERREZ, Rosa Martha; GARCIA-BAEZ, Efren; MOTA-FLORES, Jose Maria

Antimicrobial activities of diterpene labdane from seeds of *Byrsonima crassifolia*

Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas, vol. 13, núm. 1, 2014, pp.
31-37

Universidad de Santiago de Chile
Santiago, Chile

Available in: <http://www.redalyc.org/articulo.oa?id=85629766004>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Artículo Original | Original Article

Antimicrobial activities of diterpene labdane from seeds of *Byrsonima crassifolia*

[Actividad antimicrobial de diterpenos labdanos de la semilla de *Byrsonima crassifolia*]

Alethia MUÑIZ-RAMIREZ¹, Rosa Martha PEREZ-GUTIERREZ²,
Efren GARCIA-BAEZ³ & Jose Maria MOTA-FLORES²

¹Departamento de Biotecnología y Bioingeniería, Cinvestav-IPN, Av. IPN 2508, Col. San Pedro Zacatenco, Mexico D.F., CP 07360.

²Laboratorio de Investigación de Productos Naturales, Escuela Superior de Ingeniería Química e Industrias Extractivas, IPN. Av. Instituto Politécnico Nacional s/n, Unidad Profesional Adolfo López Mateos, Col. Zacatenco, CP 07758, Mexico D.F.

³Departamento de Biotecnología, UPIBI-IPN, Col. San Pedro Zacatenco, Mexico D.F., CP 07360.

Contactos / Contacts: Rosa Martha PEREZ-GUTIERREZ - E-mail address: rmpg@prodigy.net.mx

Abstract: Two compounds from the hexane extract of seeds of *Byrsonima crassifolia* were isolated and their structures elucidated using extensive spectroscopic analyses. These compounds are derived from the new labdane diterpene Labda-17-(1) and the known antimicrobial Labda-8 (17)-(2). The present study was aimed to study the effect antimicrobial of novel diterpene 1 against bacterial pathogens showed a moderate activity with MIC values 18.79-70.12 µg/ml and a MBC ranging between 250-1000 µg/ml against all assayed microorganisms.

Keywords: *Byrsonima crassifolia*, diterpene, labdane, antimicrobial

Resumen: Se aislaron dos compuestos del extracto de hexano de semillas de *Byrsonima crassifolia* y sus estructuras se elucidaron por medio de extensos análisis espectroscópicos. Estos compuestos derivados del labdano corresponden al nuevo diterpeno Labda-17- (1) y el conocido antimicrobiano Labda-8(17)-(2). En el presente estudio se estudió el efecto antimicrobiano del nuevo diterpeno 1 sobre algunas bacterias patógenas mostrando sobre de estas una actividad moderada, con valores de MIC de 18.79-70.12 µg/ml y un rango de MBC que oscila entre 250-1000 µg/ml frente a todos los microorganismos ensayados.

Palabras Clave: *Byrsonima crassifolia*, diterpeno, labdano, antimicrobiano

Recibido | Received: February 24, 2013

Aceptado en versión corregida | Accepted in revised form: May 8, 2013

Publicado en línea | Published online: January 30, 2014

Declaración de intereses | Declaration of interests: The authors are immensely thankful to the Instituto de Ciencia y Tecnología del Distrito Federal for their support.

Este artículo puede ser citado como / This article must be cited as: A Muñoz-Ramírez, RM Pérez-Gutiérrez, E García-Baez, JM Mota-Flores. 2014. Antimicrobial activities of diterpene labdane from seeds of *Byrsonima crassifolia*. *Bol Latinoam Caribe Plant Med Aromat* 13(1): 31 – 37.

INTRODUCTION

Byrsonima crassifolia is a tropical tree, commonly known as “nanche”, distributed widely in México, Central and South America (Holmgren *et al.*, 1990). Nanche fruit is edible and bright yellow when after ripened, it has sweet taste and slightly bitter taste. In México, nanche is consumed as juice, liquor, jelly and candy. Since prehispanic times it has been used as medicine, reports on ethnobotanical uses include: bark, to promote bleeding in females, to facilitate childbirth and snakebite treatment; aerial parts, used as diuretic, febrifuge, to expel placenta and for diarrhea treatment; fruit, used for fever treatment and to induce a pleasant dizziness; bark and branches, assist in tightening loose teeth and fish poison; and finally, seeds for dysentery, infections, wound healing and anti-inflammatory (Bejar and Malone, 1993). Phytochemical studies indicate that nanche plant contains esters (Alves and Franco, 2003), epicatechins (Geiss *et al.*, 1995), glycolipids (Rastrelli *et al.*, 1997). A pharmacological study of leaf and bark extracts displayed spasmogenic effects (Bejar and Malone, 1993). In another study, a chloroformic extract from bark showed anti-inflammatory activity when evaluated by the Croton oil model (Silva *et al.*, 2007). Furthermore, the antioxidant activity of extracts from leaves, fruits and bark has been determinate (Maldini *et al.*, 2009). The ethyl-acetate extract of roots exhibited antibacterial activity (Martinez-Vazquez *et al.*, 1999). Also, the aqueous extract of leaves inhibited some dermatophytes (Caceres *et al.*, 1993). The ethanol extract of leaves showed trypanocidal activity against *Leishmania mexicana* promastigotes (Berger *et al.*, 1998). The hypoglycemic activity of hexane extract have previously been reported (Perez-Gutierrez *et al.*, 2010). Based on traditional uses of seeds from *B. crassifolia* and given the lack of scientific studies on its potential pharmacological properties, the aim of this study was to investigate the potential antibacterial activity of a new diterpene labdane from seed of *B. crassifolia*.

MATERIALS AND METHODS

General experimental procedures

IR spectra were recorded on a Perkin-Elmer FTRI 1720X. Optical rotations were measured by a Perkin-Elmer 192 polarimeter equipped with a sodium lamp (589 nm). ¹H-NMR and ¹³CNMR spectra were taken at a Bruker DRX-300 NMR spectrometer, with UXNMR software package, was used for NMR experiments;

chemical shifts are reported in δ (ppm), downfield relative to TMS as an internal standard. The NMR experiments were carried out using the conventional pulse sequences as described in the literature. HREIMS were measured on a JEOL HX 110 mass spectrometer. Precoated TLC silica gel 60 F254 aluminum sheets and Sephadex LH-20 from Sigma-Aldrich (St. Louis, USA) were used. Column chromatography was carried on Silica gel 60 (230-400 mesh, Merck Co. New Jersey (USA), solvents used as eluents from Fermont (California, USA).

Plant material

B. crassifolia L. belong to the Malpighiaceae family, fruit were collected in Morelos state and were taxonomically authenticated in the Herbarium of Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional. A voucher specimen of the plant is stored for reference (No. 8976).

Extraction, isolation and characterization of the *B. crassifolia* diterpene derived

Seeds was air dried and the ground (4 kg) was extracted twice with hexane for 3 h. The seeds extracts were combined and evaporated *in vacuo* to generate a residue (225 g). The resulting extract was loaded onto a silica gel column chromatography and eluted with petroleum ether-acetone-hexane 2:1:0.5 and 7 fractions (F1-F-7) have been obtained. These fractions were then tested for antimicrobial activity. Active fractions were pooled together according to their similarities provides by thin layer chromatography analysis. The fraction F3 was the fraction that showed antimicrobial properties. F3 was subjected to silica gel column chromatography eluted with CHCl₃-EtOAc (11:2) to produce five fractions (F3-1 to F3-5). The active fraction F3-4 was subjected to chromatographed over silica gel column using CH₂Cl₂-CHCl₃ 3:1.5 to yield four subfractions (F34-1 to F34-4). The F34-1 fraction was further purified by preparative plate using petroleum ether-EtOAc 1:4 and visualized with UV at 254 nm. Fraction F341-2 was separated by Sephadex LH-20 using a gradient of CHCl₃-MeOH (from 10:1 to 5:1) to yield **1** (80 mg) and **2** (67 mg).

Compound 1:

Yellowish oil; $[\alpha]_D^{24}$ -21.4° (CHCl₃, c 0.68); IR ν_{\max} 2924, 2854, 1710, 1645, 1460, 1375, 1163, 1096, 895 cm⁻¹; m/z 588.9254 [M+H]⁺, (calc, for C₃₈H₆₈O₄, 588.9217); ¹H NMR (CDCl₃, 300 MHz) δ : 1.55 (m, H-1a), 1.76

(dd, $J = 13.1, 3.4$ Hz, H-1b), 1.46 (dd, $J = 13.1, 3.4$ Hz, H-2a), 1.66 (m, H-2b), 1.22 (m, H-3a), 1.58 (m, H-3b), 1.34 (m, H-5), 1.70 (m, H-6), 1.33 (1H, m, 7a), 1.39 (m, H-7b), 1.84 (m, H-8), 1.96 (m, H-9), 1.39 (m, H-11a), 1.59 (m, H-11b), 2.16 (m, H-12a), 1.76 (m, H-12b), 5.3 (t, $J = 6.4$ Hz, H-14), 3.74 (2H, d, $J = 7.2$ Hz, H-15), 0.96 (3H, s, H-16), 4.08 (2H, d, $J = 2.8$ Hz, H-17), 1.17 (s, H-18), 1.18 (s, H-19), 0.75 (s, H-20), 2.02 (s, COOMe), 2.04, (s, COOMe), 1.18-1.25 (CH_2); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 33.21 (C-1), 18.23 (C-2), 34.65 (C-3), 41.64 (C-4), 52.24 (C-5), 25.71 (C-6), 39.89 (C-7), 37.82 (C-8), 54.32 (C-9), 36.81 (C-10), 24.16 (C-11), 43.21 (C-12), 129.89 (C-13), 131.18 (C-14), 68.72 (C-15), 16.52 (C-16), 62.54 (C-17), 33.71 (C-18), 22.35 (C-19), 14.23 (C-20), 172.13 (COOMe), 20.98 (COOMe), 173.07 ($\text{CH}_2\text{-COO-CH}_2$).

Screening for antibacterial activity

Compound **1** was dissolved in DMSO at a final concentration of 10 mg/ml and serial dilutions were prepared using the same solvent which were aseptically dispensed onto sterilized 6 mm filter paper discs and dried. The disk-diffusion method (Gaydos, 1982) was used to measure the antimicrobial activity of the compounds. The different organisms were seeded over previously sterilized Mueller-Hinton agar for bacteria. The zones of growth inhibition were observed around dried disks (Whatman 6 mm diameter). Each plate was inoculated with 0.1 ml of a bacterial culture directly from the 24 h broth culture diluted to match 0.5 MacFarland standard (108 Colony Forming Units (CFU)/ml). Inoculums on the plates were aseptically spread using a sterile glass rod to avoid contamination and allowed to stand for 2 - 3 minutes in a refrigerator at 4° C. The discs loaded with **1** were then placed onto the seeded plates. The bacterial cultures were incubated at 37° C for 24hr. After the incubation period, the zones of inhibition were measured and recorded in mm as described by Elgayyar *et al.*, (2000). Negative controls were done by using sterile 6mm discs soaked in 10 ml of DMSO and dried.

Minimum inhibitory concentrations (MIC) and minimum bactericidal concentrations (MBC)

Compound **1** from the antibacterial screening was tested for Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC). The MIC was determined on 96-well microdilution plates according the published protocols (Micheal *et*

al., 2003). The MBC were determined by sub-culturing 0.1 ml of all the tubes showing no growth on Nutrient Agar (NA). After 24 h incubation at 37° C, the first plate showing no growth was recorded as the MBC. The MICs accuracy was checked against standard antibiotics. Sterility of the media was controlled by use of broth only in a negative control tube, and growth of the organism was checked by broth plus microorganism in question in a positive control tube. All the controls were subjected to the same conditions as the tests. Antimicrobial sensitivity and resistance were confirmed by use of standard discs containing ciprofloxacin (10 µg/ml).

RESULTS AND DISCUSSION

Two compounds were isolated from hexane extract of seeds of *B. crassifolia* after purification through silica gel column chromatography. Compound **1** showed a molecular ion at m/z 588.9254 in its HREIMS and the presence of 38 carbons on the ^{13}C NMR spectrum suggested a molecular formula $\text{C}_{38}\text{H}_{68}\text{O}_4$. DEPT experiments classified the protonated carbon signals to six methyls, twenty three methylenes, four methines and five quaternary carbons. The IR spectrum of **1** showed absorption bands at ν_{max} 1645 cm^{-1} to double bond and 1710 and 1163 cm^{-1} corresponding to an ester group, it was supported by the presence of signal of quaternary carbons at δ_{C} 172.13 and δ_{C} 173.07 in its ^{13}C NMR spectrum. Its ^1H NMR spectrum showed signals for four tertiary methyl groups (δ_{H} 0.96, s; 1.17, s; 1.18, s; 0.75, s), one trisubstituted double bonds (δ_{H} 5.3, t, $J = 6.4$ Hz) and two acetoxyl groups (δ_{H} 2.02, s and 2.04, s). In the ^{13}C NMR spectrum, two acetoxycarbonyl carbons (δ_{C} 172.13, and 173.07), and the one 129.89, trisubstituted double bonds characterized by the presence of the two olefinic carbons (δ_{C} and 131.18). Inspection of the 2D NMR data (^1H - ^1H COSY, ^1H - ^1H TOCSY, HSQC and HMBC) allowed the assignments of all the signals observed in the 1D NMR spectra and revealed the labdane skeleton of the diterpene **1** (Bohlmann and Czerson, 1979; Timmermann *et al.*, 1986). Connectivities in the HMBC spectrum from H-17 to C-7, C-8 and C-9 (δ_{C} 54.32) corroborated the location of the side chain to be at C-9, whose methine proton H-9 (δ_{H} 1.96) formed an isolated proton spin system with H-11 (δ_{H} 1.39 and δ_{H} 1.59) and H-12 (δ_{H} 1.76 and 2.16) as deduced from the COSY and TOCSY spectra (Itokawa *et al.*, 1988). A trisubstituted double bond was located at C-13 (δ_{C} 129.89) and C-14 (δ_{C} 131.18) on the basis of $^2\text{J}_{\text{C-H}}$ and $^3\text{J}_{\text{C-H}}$ HMBC correlations from

Me-16 (δ_H 0.96) to C-12 (δ_C 43.21), C-13 and C-14, and from the olefinic proton H-14 (δ_H 5.3) to C-12 (Figure 1). The double bond at C-13 was assigned as the E configuration because of the ROESY correlation observed between H-15 and Me-16, and by comparison of NMR data with those of related compounds (Urones *et al.*, 1995; Su *et al.*, 1996). Large number of methylene groups (22.7-34.4), suggesting the presence of fatty acid esterified with the labdane derivative at C-15. It is supported by the HMBC correlation between methylene protons H₂-15 (δ_H 4.08, 2H, d, J = 7.2 Hz) and methylene chain and the quaternary carbon δ_C 173.07. Analysis of 1H - 1H COSY showed clear correlation between H5/H6, H6/H7, H11/H12 (Figure 2). One methyl group (δ_C 16.52), and large number of methylene groups (δ_C

22.6-34.5) (DEPT) suggesting the presence of monoenic fatty acid esterified with a labdane derivative at C-15. The NMR data of the long-chain moiety were in good accordance with those for Δ^9 -18:1 *cis* fatty acid esters (Mannina *et al.*, 1999). Therefore, compound **1** was identified as labda-17-acetoxi-13E-en-15-palmitate. All data of 1H NMR and ^{13}C NMR spectra, 1H - 1H , COSY and HMBC connectivities of **2** allowed the identification of labda-8(17),13E-dien-19-carboxy-15-yl palmitate which was previously isolated from *Cretan propolis* (Peopova *et al.*, 2009) exhibited a weak antimicrobial activity against *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Klebsiella pneumonia*, *Escherichia coli*, *Candida tropicalis* and *Candida glabrata*.




Figure 1
Compounds **1** and **2** isolated from *B. crassifolia*




Figure 2
Selected 1H - 1H COSY (bold) and 1H - ^{13}C HMBC (arrow) correlations of **1**

Screening of the *in vitro* antimicrobial activity of diterpenoid **1** was studied against Gram positive microorganisms *Bacillus cereus*, *Bacillus megaterium*, *Bacillus subtilis*, *Staphylococcus aureus*, and Gram negative microorganisms as *Escherichia coli*, *Salmonella paratyphi*, *Pseudomonas aeruginosa*, *Salmonella typhi*, *Shigella boydii*, *Salmonella dysenteriae*, *Vibrio mimicus* and *Vibrio parahemolyticus* of significant importance using the filter paper disc agar diffusion technique. All the tested bacteria were standard strains from ATCC (American type culture collection). The results showed variations in the antimicrobial properties of diterpene labdane **1** (Table 1). Compound **1** had strong activity (inhibition zone ≥ 20 mm), moderate activity (inhibition zone < 20 -12 mm) and no inhibition (zone

< 12 mm). The most effective **1** was that of, *S. paratyphi* (23.87 ± 0.65 mm, zdi), followed by *P. aeruginosa* (21.45 ± 0.53 mm, zdi), *S. typhi* (20.72 ± 0.56 mm, zdi), *B. subtilis* (20.89 ± 2.90 mm, zdi). In contrast, showed a moderate activity against *B. megaterium* (19.20 ± 3.42 mm, zdi), *B. cereus* (18.76 ± 0.68), *Staphylococcus aureus* (17.95 ± 1.78), *Escherichia coli* (13.35 ± 1.43), and *S. dysenteriae* (16.52 ± 1.88). However, no significant antimicrobial effect against *Vibrio mimicus* (12.08 ± 4.12), *Sarcinia lutea* (12.17 ± 1.23), *V. parahemolyticus* (11.87 ± 4.20) and *Shigella boydii* (0.0 mm) was observed. Gram-positive bacterial cells were, in general, more resistant to the presence of antimicrobial than Gram-negative.

Table 1
Antimicrobial activity of the compound 1 obtained from seeds of *Byrsonima crassifolia*

Test microorganisms	Inhibition zone diameter (mm)	MIC ($\mu\text{g/ml}$)	MBC ($\mu\text{g/ml}$)
Gram positive bacteria			
<i>Bacillus cereus</i> ATCC 11778	18.76 ± 0.68	38.34 ± 1.87	500
<i>B. megaterium</i> ATCC 12872	19.20 ± 3.42	43.26 ± 2.76	500
<i>B. subtilis</i> ATCC 6633	20.89 ± 2.90	41.28 ± 3.29	500
<i>Staphylococcus aureus</i> ATCC 6538	17.95 ± 1.78	36.79 ± 1.87	500
<i>Sarcinia lutea</i> ATCC 9341	12.17 ± 1.23	58.84 ± 4.76	1000
Gram negative bacteria			
<i>Escherichia coli</i> ATCC 25922	13.35 ± 1.43	69.20 ± 2.76	1000
<i>Pseudomonas aeruginosa</i> ATCC 9027	21.45 ± 0.53	32.41 ± 4.11	500
<i>Salmonella paratyphi</i> ATCC 6539	23.87 ± 0.65	20.27 ± 1.04	250
<i>S. typhi</i> ATCC 43971	20.72 ± 0.56	18.79 ± 2.54	250
<i>Shigella boydii</i> ATCC 9207	0.0	-	-
<i>S. dysenteriae</i> ATCC 13313	16.52 ± 1.88	45.67 ± 3.87	500
<i>Vibrio mimicus</i> ATCC 33654	12.08 ± 4.12	66.78 ± 5.14	1000
<i>V. parahemolyticus</i> ATCC 43996	11.87 ± 4.20	70.12 ± 4.98	1000

Zone diameter inhibition (ZDI) of compound 1 (20 μg) per disk

However, more precise data on antimicrobial properties were obtained through determinations of bacteriostatic and bactericidal concentrations. The minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) the results are shown in Table 1. The most bacteriostatic properties was against all Gram-negative strains assayed. Martinez-Vazques *et al.*, (1999) studied ethyl acetate, methanol extracts of roots and stems of *B. crassifolia* against *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Salmonella typhi*, *Shigella flexneri*, *Staphylococcus aureus*, *Staphylococcus*

epididymis, *Streptococcus pneumoniae* and *Micrococcus luteus*. The ethyl acetate of roots was the most active. In another research had been previously studied methanol extract from *B. crassifolia* against *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Candida albicans* (Navarro *et al.*, 1996). The results obtained by us indicate the existence of antimicrobial compounds in this plant, and show a good correlation between the reported studies of *B. crassifolia* uses in traditional medicine against infectious diseases and the experimental data of such extracts toward the most

common pathogens. Also results indicate the existence of antimicrobial compounds in the hexane extract of *B. crassifolia*, the phytochemical isolation and identification of two active compounds indicates that the antimicrobial activity of the seed of *B. crassifolia* is due mainly to the presence of compounds labdane diterpene.

In conclusion the present study shows that antimicrobial activity is present in the seeds of *B. crassifolia*. Bioassay with the purified diterpene 1 indicated that exhibited strong activity against Gram negative bacterias. The results suggest that further investigations against other enteric Gram-positive and Gram-negative bacteria, which are frequently resistant to drugs commonly used in therapy, may be carried out.

ACKNOWLEDGEMENT

The authors are immensely thankful to the Instituto de Ciencia y Tecnologia del Distrito Federal for their support.

REFERENCES

- Alves GL, Franco MRB. 2003. Headspace gas chromatography-mass spectrometry of volatile compounds in murici (*Byrsonima crassifolia*). **J Chromatogr A** 985: 297 - 301.
- Bejar E, Malone MH. 1993. Pharmacological and chemical screening of *Byrsonima crassifolia* a medicinal tree from México. Part I. **J Ethnopharmacol** 39: 141 - 158.
- Berger I, Barrientos AC, Cáceres A, Hernández M, Rastrelli L, Passreiter CM, Kubelka W. 1998. Plants used in Guatemala for the treatment of protozoal infections: Activity of extracts and fractions of five Guatemalan plants against *Trypanosoma cruzi*. **J Ethnopharmacol** 62: 107 - 115.
- Bohlmann F, Czerson H. 1979. Neue labdan- und pimaren-derivate aus *Palafoxia rosea*. **Phytochemistry** 18: 115 - 118.
- Caceres A, Brenda B, Lopez R, Giron MA, Logemann H. 1993. Plants used in Guatemala for the treatment of dermatophytic infections. I. Screening for antimycotic activity of 44 plant extracts. **J Ethnopharmacol** 40: 207 - 213.
- Elgayyar M, Draughon FA, Golden DA, Mount JN. 2000. Antimicrobial activity of essential oils from plants against selected pathogenic and saprophytic microorganisms. **J Food Prot** 64: 1019 - 1024.
- Gaydos JM, Harrington BJ. 1982. Agar disk diffusion for the quality control testing of Autobac elution disks. **Antimicrob Agents Chemother** 21: 516 - 518.
- Geiss F, Heinrich M, Hunkler D, Rimplerl H. 1995. Proanthocyanidins with (+)-epicatechin units from *Byrsonima crassifolia* bark. **Phytochemistry** 39: 635 - 643.
- Holmgren KH, Holmgren NH, Barnett LC. 1990. **Index Herbariorum. Part 1: The herbaria of the world**. 8a ed. New York Botanical Garden. Nueva York, USA.
- Itokawa H, Yoshimoto S, Morita H. 1988. Diterpenes from the rhizomes of *Alpinia formosana*. **Phytochemistry** 27: 435 - 438.
- Maldini M, Sosa S, Montoro P, Giangaspero A, Balick MJ, Pizza CD, Loggia R. 2009. Screening of the topical anti-inflammatory activity of the bark of *Acacia cornigera* Willdenow, *Byrsonima crassifolia* Kunth, *Sweetia panamensis* Yakovlev and the leaves of *Sphagneticola trilobata* Hitchcock. **J Ethnopharmacol** 122: 430 - 433.
- Mannina L, Luchinat C, Emanuele MC, Segre A. 1999. Acyl positional distribution of glycerol tri-esters in vegetable oils: a ¹³C-NMR study. **Chem Phys Lipids** 103: 47 - 55.
- Martinez-Vazquez M, Gonzalez-Esquinca AR, Cazares LL, Moreno GMN, Garcia-Argaer AN. 1999. Antimicrobial activity of *Byrsonima crassifolia* (L.). H.B.K. **J Ethnopharmacol** 66: 79 - 82.
- Micheal JP, Chan ECS, Krieg NR. 2003. Microbiology. Tata McGraw-Hill Publishing Company Limited, New Delhi, India.
- Navarro V, Villarreal ML, Rojas G, Lozoya X. 1996. Antimicrobial evaluation of some plants used in Mexican traditional medicine for the treatment of infectious diseases. **J Ethnopharmacol** 53: 143 - 147.
- Rastrelli L, De Tommasi N, Berger I, Caceres A, Saravia A, De Simone F. 1997. Glycolipids from *Byrsonima crassifolia*. **Phytochemistry** 5: 647 - 650.
- Silva EM, Souza JNS, Rogez H, Rees JF, Larondelle Y. 2007. Antioxidant activities and polyphenolic contents of fifteen selected plant species from the Amazonian region. **Food Chem** 101: 1012 - 1018.

- Su W, Fang J, Cheng Y. 1996. Diterpenoids from leaves of *Cryptomeria japonica*. **Phytochemistry** 41: 255 - 261.
- Peopova MP, Chinou IB, Marekov IN, Bankova VS. 2009. Terpenes with antimicrobial activity from *Cretan propolis*. **Phytochemistry** 70: 1262 - 1271.
- Perez-Gutierrez RM, Muñiz-Ramirez A, Gomez Gomez Y, Bautista-Ramirez E. 2010. Antihyperglycemic, antihyperlipidemic and antiglycation of *Byrsonima crassifolia* fruits. **Plant Foods Hum Nutr** 20717728.
- Timmermann BN, Hoffmann JJ, Jolad SD, Bates RB, Stahaan TJ. 1986. Labdane diterpenoid from *Grindella discoidea* (Asteraceae). **Phytochemistry** 25: 1389 - 1392.
- Urones JG, Isidro S, Marcos IM, Oliva NM, Garrido JH, Vanessa MH. 1995. Labdane diterpenes from *Halimium viscosum*. **Phytochemistry** 38: 663-666.