



Investigación y Educación en Enfermería

ISSN: 0120-5307

ISSN: 2216-0280

Imprenta Universidad de Antioquia

de Castro Figueiredo Pereira Coelho, Raquel; Nunes Garcia, Sabrina; Marcondes, Larissa; Jacinto da Silva, Flaviane Andriele; de Paula, Aline; Puchalski Kalinke, Luciana

**Impact on the quality of life of women with breast cancer
undergoing chemotherapy in public and private care¹**

Investigación y Educación en Enfermería, vol. 36, no. 1, e4, 2018, January-April

Imprenta Universidad de Antioquia

DOI: 10.17533/udea.iee.v36n1e04

Available in: <http://www.redalyc.org/articulo.oa?id=105255813004>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Impact on the quality of life of women with breast cancer undergoing chemotherapy in public and private care

Raquel de Castro Figueiredo Pereira Coelho¹
Sabrina Nunes Garcia²
Larissa Marcondes³
Flaviane Andreele Jacinto da Silva⁴
Aline de Paula⁵
Luciana Puchalski Kalinke⁶

Impact on the quality of life of women with breast cancer undergoing chemotherapy in public and private care

Objective. To compare the general quality of life of women with breast cancer undergoing chemotherapy in a public and a private institution. **Methods.** Longitudinal observational study including 115 women with breast cancer in an outpatient chemotherapy treatment at two health institutions, one public and one private, in the city of Curitiba (Brazil). The Quality of Life Questionnaire-C30 and the Quality of Life Questionnaire - Breast Cancer Module instruments were applied at three moments of treatment (onset, 40-50 days, and 85-95 days after initiation). **Results.** Women's global health status was affected in both groups since the first phase of chemotherapy (76.2 points in the private institution and 74.6 points in the public institution, considering 100 points = maximum health). In both groups there was worsening of quality of life over time, which was lower in women in the private



Original article



UNIVERSIDAD
DE ANTIOQUIA
1803

- 1 Nurse, Masters. Hospital de Clínicas da Universidade Federal do Paraná. Curitiba-Pr, Brasil.
email: raque.gabi@gmail.com
- 2 Nurse, Masters. Instituto de Oncologia do Paraná – IOP. Curitiba-Pr, Brasil.
email: sgarcia@iop.com.br
- 3 Nurse, Masters. student. Universidade Federal do Paraná. Curitiba-Pr, Brasil.
email: marcondes.lari@gmail.com
- 4 Nursing undergraduate student. Universidade Federal do Paraná. Curitiba-Pr, Brasil.
email: fla.andreele@gmail.com
- 5 Nursing undergraduate student. Universidade Federal do Paraná. Curitiba-Pr, Brasil.
email: aline.paulla94@gmail.com
- 6 Nurse, Ph.D. Adjunt Professor, Universidade Federal do Paraná. Curitiba-Pr, Brasil.
email: lucianakalinke@yahoo.com.br

Conflicts of Interest: None.

Received: July 24th, 2017.

Accepted: November 20th, 2017.

How to cite this article: Coelho RCFP, Garcia SN, Marcondes L, Silva FAJ, de Paula A, Kalinke LP. Impact on the quality of life of women with breast cancer undergoing chemotherapy in public and private care. Invest. Educ. Enferm. 2018; 36(1):e04.

DOI: 10.17533/udea.iee.v36n1e04



Investigación y Educación en

Enfermería

—Nursing Research and Education—

Vol. 36 No1, • February 15th 2018 • ISSN: 2216-0280

institution (second phase = 75.0 and third phase = 74; $p=0.47$), compared to those in the public institution (second phase = 71.5 and third phase 69.1; $p=0.02$). Regarding the type of institution, the most committed functions were the emotional and social in the private, and physical and pain in the public. **Conclusion.** The global quality of life of women with breast cancer deteriorated as a result of chemotherapy in both institutions, and it was lower in those receiving private care. Nursing should consider the type of institution responsible for women's care in order to provide comprehensive care that considers the functions affected at each phase of treatment.

Descriptors: quality of life; breast neoplasms; private sector; public sector; ambulatory care facilities.

Compromiso de la calidad de vida de mujeres con cáncer de mama sometidas a quimioterapia en la atención pública y privada

Objetivo. Comparar la calidad de vida general de mujeres con cáncer de mama sometidas a quimioterapia en una institución pública y otra privada. **Métodos.** Estudio observacional longitudinal en el que participaron 115 mujeres con cáncer de mama en tratamiento ambulatorio de quimioterapia en dos instituciones de salud, una pública y una privada, de la ciudad de Curitiba (Brasil). Los instrumentos *Quality of Life Questionnaire-C30* y *Quality of Life Questionnaire - Breast Cancer Module* se aplicaron en tres momentos del tratamiento (inicio, 40-50 días y 85-95 días post-inicio). **Resultados.** La calidad de vida global de las mujeres ya estaba afectada en ambos grupos desde la primera etapa de la quimioterapia (76.2 puntos en la privada y 74.6 puntos en la pública, siendo 100 puntos el equivalente de máxima salud). También se pudo apreciar que, aunque en ambos grupos hubo empeoramiento de la calidad de vida con el tiempo, este fue menor en las mujeres en la atención privada (segunda etapa = 75.0 y tercera etapa = 74; $p=0.47$), comparadas con las que recibieron atención pública (segunda etapa = 71.5 y tercera etapa 69.1; $p=0.02$). Según tipo de institución, las funciones más comprometidas fueron la emocional y la social en la privada, y la física y el dolor en la pública. **Conclusión.** La calidad de vida global de las mujeres con cáncer de mama se deterioró como consecuencia de la quimioterapia en las dos instituciones, siendo menor en las que recibieron atención

privada. Enfermería debe también considerar el tipo de institución responsable de la atención de la mujer, con el fin de brindar un cuidado integral que tenga en cuenta las funciones afectadas en cada fase da tratamiento.

Descritores: calidad de vida; neoplasias de la mama; sector privado; sector público; instituciones de atención ambulatoria.

Comprometimento da qualidade de vida de mulheres com câncer de mama submetidas a quimioterapia no atendimento público e privado

Objetivo. Comparar a qualidade de vida geral de mulheres com câncer de mama submetidas a quimioterapia em uma instituição pública e outra privada. **Métodos.** Estudo observacional, longitudinal, em que participaram 115 mulheres com câncer de mama em tratamento ambulatorial de quimioterapia em duas instituições de saúde, uma pública e uma privada, na cidade de Curitiba (Brasil). Os instrumentos *Quality of Life Questionnaire-C30* e *Quality of Life Questionnaire – Breast Cancer Module* foram aplicados em três momentos do tratamento (início, 40-50 dias e 85-95 dias após o início). **Resultados.** A qualidade de vida global das mulheres foi afetada em ambos os grupos desde a primeira fase da quimioterapia (76.2 pontos na privada e 74.6 pontos na pública, sendo 100 pontos o equivalente ao máximo de saúde). Também foi possível observar em ambos os grupos que houve piora na qualidade de vida ao longo do tempo, esta foi menor nas mulheres em atendimento privado (segunda etapa = 75.0 e terceira etapa = 74; $p=0.47$), comparadas com aquelas que receberam atenção pública (segunda etapa = 71.5 e terceira etapa 69.1; $p=0.02$). De acordo com o tipo de instituição, as funções mais comprometidas foram a emocional e a social na privada, e física e a dor na pública. **Conclusão.** A qualidade de vida global das mulheres com câncer de mama deteriorou-se em consequência da quimioterapia nas duas instituições, sendo esta menor naquelas que recebem atenção privada. A enfermagem deve considerar o tipo de instituição responsável pelo atendimento das mulheres, a fim de fornecer um atendimento integral que leve em consideração as funções afetadas em cada fase do tratamento.

Descritores: qualidade de vida; neoplasias da mama; setor privado; setor público; instituições de assistência ambulatorial.

Introduction

Breast cancer is one of the most frequent and feared diseases among women. It has negative impact given its severity, unpredictable evolution, mutilation and self-image alterations that compromise physical, psychological and social aspects and affect women's quality of life (QoL). Antineoplastic chemotherapy is one of the most important forms of breast cancer control and represents an advance in its cure.⁽¹⁾ However, despite prolonging life and improving prognosis, chemotherapy is associated with a lower global quality of life (GQoL) score, possibly related to its toxicity and serious side effects.⁽²⁾ Both the diagnosis and chemotherapy change patients' routine and QoL. They lead to negative feelings due to the uncertainty of prognosis, possible surgical processes, and coping with the possibility of death.

The search for improved care and QoL of women with breast cancer undergoing chemotherapy is an important aspect to be measured because this assesses the disease dimensions and creates parameters for daily healthcare practices in health services.⁽³⁾ Thus, by assessing the QoL it is possible to verify the treatment and impact, and plan nursing actions for patients' adherence and rehabilitation, and better QoL conditions during treatment and survival. Knowing the sociodemographic profile of women with breast cancer is key to provide targeted and comprehensive care according to each moment of the therapeutic treatment. Highlighting possible profile differences according to the care institution may alert the health team to perform actions, guidelines and care adjusted to women's needs and reality. The physical, emotional, social and cognitive impacts must be constantly evaluated to provide accurate care to women experiencing breast cancer, from diagnosis to the end of life.

Thus, the objective of the present study was to evaluate and compare the general quality of life and sociodemographic and clinical profile of women with breast cancer undergoing chemotherapy treatment at a public and a private institution.

Methods

This is an observational, longitudinal and analytical study. It is part of the thematic project entitled 'Quality of life of cancer patients undergoing chemotherapy'. The study was conducted between October 2012 and May 2015 in chemotherapy outpatient clinics of a private and a public institution considered reference in the care for this patient profile. Both clinics are located in the city of Curitiba, state of Paraná, southern Brazil. The non-probabilistic sample consisted of all eligible women during the study period. Inclusion criteria were: female gender with proven diagnosis of breast cancer, aged over 18 years, no previous chemotherapy treatment for cancer and indication for initiation of chemotherapy. Women on palliative therapy and had lost follow-up were excluded. A total of 131 women were selected, of which 16 were ineligible because they had not been approached before chemotherapy.

In the first stage of the study, 115 women participated in the study, of which 48 were from the private institution and 67 from the public institution. In the second stage, which occurred between 40 and 50 days after the first administration of chemotherapy (period of adverse effects occurrence),⁽⁴⁾ a woman was discontinued in the public institution. In the third stage, 85 to 95 days after the first, (period of control of adverse events),⁽⁴⁾ five women were discontinued in the public institution. In the private institution there was no discontinuity in any of the stages. Discontinuations occurred because of chemotherapy toxicity or loss of follow-up between the second and third stages. A total of 791 questionnaires were collected in the three stages of the study.

Three questionnaires were used for data collection. Questionnaire 1 had sociodemographic and clinical data, and was applied only in the first stage. Questionnaire 2 was the Quality of Life Questionnaire-C30 (QLQ-C30), developed by the European Organization for Research and Treatment of Cancer (EORTC), version 3.0, and was used to assess the QoL of cancer patients. Questionnaire 3 was the Quality of Life Questionnaire - Breast

Cancer Module (QLQ-BR23), and applied in the three stages. It is directed to the basic disease with issues related to breast cancer, version 1.0. The QLQ-C30 and QLQ-BR23 questionnaires used were the translated and validated versions for Brazil.⁽⁵⁾

The analysis of sociodemographic and clinical data was by absolute and relative frequency. Data from the QLQ-C30 and QLQ-BR23 questionnaires had a 0-100 score according to the Scoring Manual of European Organization for Research and Treatment of Cancer.⁽⁶⁾ For QoL-related scores such as emotional functioning, the closer to 100 represents greater functionality and corresponds to a better QoL. For the comparison of data between institutions, was applied the non-parametric Mann Whitney test, and for the comparison of stages, was applied the Friedman non-parametric test complemented by the Least Significant Difference Test (LSD), considering $p \leq 0.05$ as significant. All ethical precepts were respected in the study, and both questionnaires were authorized upon registration of the project with the EORTC. It was approved by the Ethics and Research Committee under protocol number 5301 for the private institution, and number 518.067 for the public institution.

Results

In the present study, participants' mean age in the private institution was 46 years (min = 24, max = 69), and in the public it was 51.3 years (min = 30, max = 77), which was a significant difference ($p=0.02$). Table 1 shows sociodemographic data of the 115 participants. Regarding women's educational level, in the private institution 54% ($n = 26$) had complete higher education, whereas in the public institution 43.3% ($n = 29$) had less than nine years of study, which includes incomplete or complete primary education. The family income in minimum wages was another highlight; in the private institution, the average income was 12 salaries (US\$ 3,069.76), while in the public institution the average was 2.8 salaries (US\$ 716.27).

Table 1. Sociodemographic characteristics of women with breast cancer undergoing chemotherapy according to type of institution. Curitiba, PR, Brazil, 2015

Variables	Private n=48		Public n=67		p-value
	Number	%	Number	%	
Marital status					0.57
Single	4	8.5	8	11.9	
Married or living together	32	66.5	41	61.2	
Separated or divorced	9	18.8	8	11.9	
Widows	3	6.3	10	14.9	
Number of children					0.50
None	3	6.3	10	14.9	
Up to 2	43	89.6	43	64.2	
3 or more	2	4.2	14	20.8	
Educational level (years of study)					0.74
< 9	4	8.5	29	43.3	
9 to 12	17	35.5	19	28.4	
> 12	26	54.2	17	25.4	
No answer	1	2.1	2	2.9	
Occupation					0.41
Active	27	56.5	38	56.7	
Retired	2	4.2	9	13.4	
Housework	14	29.2	17	25.4	
Unemployed	5	10.5	3	4.5	

In relation to clinical data (Table 2), in the private institution 68.5% (n = 33) reported having no comorbidity. In the public institution, hypertension stood out in 35.8% of women, and 49.3% reported not having any secondary diseases. Clinical stage III was frequent in 82% and 39.5% of women in public and private institutions, respectively, and AC-T (Adriamycin® and Cyclophosphamide® followed by Taxol®) was the most used protocol.

The QoL results were evaluated with the QLQ-C30 questionnaire, the Global health status in the private institution was considered satisfactory with a value of 76.2%, but it declined in the second and third phases. In the public institution, there was also a decrease between the first and third phases, but in this institution the change was greater. As for significant differences between phases, the p-value of the Global health status in the

private institution was 0.47, and in the public institution was 0.02 (Table 3).

Still in the comparison between treatment phases with the QLQ C-30 questionnaire, in the public institution were observed significant changes in values of physical and emotional functioning, fatigue, nausea and vomiting, and in the private institution, in physical and social functioning, fatigue, nausea and vomiting. When using the QLQ BR23 questionnaire, in the private institution, the systemic effects and body image stood out, and in the public institution, systemic effects and breast symptoms had significance (Table 4).

When comparing institutions (Table 5) using the QLQ-C30 questionnaire, the significant domains in the private were emotional functioning (first phase) and social functioning (third phase). In the public institution, physical functioning and pain (both in the first phase) were significant.

Table 2. Clinical characteristics of women with breast cancer undergoing chemotherapy according to type of institution. Curitiba, PR, Brazil, 2015

Variables	Private n=48		Public n=67		p-value
	Number	%	Number	%	
Smoking habits					0.17
Smoker	5	10.5	13	19.4	
Non smoker	36	75	43	64.2	
Former smoker	7	14.5	11	16.4	
Alcohol intake					0.28
Does not drink alcohol	40	83.5	51	76.1	
Moderate intake	6	12.5	10	15	
Heavy drinker	2	4	6	8.9	
Physical activity practice					0.04
No practice	27	56.5	50	74.6	
Eventual practice	14	29	15	18	
Regular practice	7	14.5	5	7.4	

Table 3. Global health status scores of women with breast cancer undergoing chemotherapy according to stage of treatment and type of institution. Curitiba, PR, Brazil, 2012-2015

Institution	First phase	Second phase	Third phase	p-value
Private	76.2	75.2	74	0.47
Public	74.6	71.5	69.1	0.02

Table 4. Significant domains between chemotherapy treatment phases in women with breast cancer by type of institution. Curitiba, PR, Brazil, 2012-2015

Scale and domain	<i>p</i> -value	Private			<i>p</i> -value	Public		
		1 st x2 nd stages	1 st x3 rd stages	2 nd x3 rd stages		1 ^a x2 ^a stages	1 st x3 rd stages	2 nd x3 rd stages
QLQ C-30								
Global health status	0.47	ns*	ns	ns	0.02	ns	ns	ns
Physical functioning	<0.01	ns	<0.01	ns	<0.01	ns	<0.05	ns
Emotional functioning	<0.01	ns	<0.05	ns	0.01	<0.05	<0.05	ns
Fatigue	<0.01	0.01	<0.01	ns	<0.01	<0.01	<0.01	ns
Nausea and vomiting	<0.01	<0.05	0.05	ns	<0.01	<0.01	<0.01	ns
QLQ-BR23				ns				
Systemic effects	<0.01	<0.01	<0.01	ns	<0.01	<0.01	<0.01	ns
Breast symptoms	<0.01	<0.01	<0.01	ns	<0.01	<0.05	<0.05	ns

(*) ns: not significant

Table 5. Significant domains in the comparison between private and public institutions. Curitiba, PR, Brazil, 2012-2015

Questionnaire	Function	Phase	Private institution		Public institution		U	p-value
			n	Sum of ranks	n	Sum of ranks		
C30	Physical functioning	1st	48	3151	67	3519	1241	0.037
C30	Emotional functioning	1st	48	3205	67	3465	1187	0.017
C30	Pain	1st	48	2361	67	4309	1185	0.016
C30	Social functioning	3rd	48	2272.5	61	3722.5	1096.5	0.024
BR23	Body image	1st	47	3050	67	3505	1227	0.045
BR23	Systemic effects	1st	48	2245	67	4425	1069	0.002
BR23	Systemic effects	2nd	48	2347	66	4208	1171	0.017

Discussion

QoL studies in women with breast cancer aim to identify the impact of diagnosis, treatment, adverse effects and biopsychosocial needs throughout the course, allowing that nurses plan actions to improve these women's living conditions. In this study, the public institution reveals a more heterogeneous population in relation to age, educational level and professional occupation. Socioeconomic conditions are less favorable with less access to health services for periodic follow-up, and when it is done, greater time is required for diagnosis given the available resources in gratuitous care.

The private institution characteristic is also a component to be considered as it can show specific results for this scenario because of the population's social characteristics. In private institutions, women often have more favorable socioeconomic conditions and access to health services because of the possibility of payment, and this aspect facilitates routine examinations and frequent consultations hence allowing an early diagnosis.

In the private institution, the mean age was lower than the national. This is an uncommon fact in the literature that constitutes 5 to 7% of breast cancer cases. In contrast, in the public institution, women had an average age similar to the national of around 50 years.^(7,8) The explanation

for this condition can be that younger women at reproductive ages are more likely to seek medical service, therefore are more examined, which can provide an early diagnosis of the disease.⁽⁹⁾ However, statistics predicting breast cancer in relation to age are changing, and the results of the research in the private institution corroborate with this information. These studies^(10,11) show the predominance of age groups below the national average, with increased number of cases as women's age increased. International studies are in contrast to the findings of the present study, and indicate the occurrence in the age group over 50 years, women's mean age of 61.8 years in Sweden⁽¹²⁾ and 58 years in the United States.⁽¹³⁾

In addition to the age group, other factors deserve health professionals' attention, such as educational level and socioeconomic status. In the private institution, several women have studied more than 12 years, but in the public institution, many women studied less than nine years, which demonstrates the educational level difference between women with breast cancer in chemotherapy treatment at the two institutions. This corroborates the fact that the higher the educational level the greater the insertion of these women in the labor market, and consequently, the greater knowledge about prevention methods.⁽¹⁴⁾

According to Rosa and Radünz,⁽¹⁵⁾ low educational level may be tied to low socioeconomic status. That was also a finding in the present study,

in which around 56% of women in the private and public sectors were economically active. However, the average family wage in the private institution was high. In the public institution, in turn, it was characterized as low socioeconomic status. Simeão et al.⁽¹⁶⁾ emphasize that similar socioeconomic and cultural level characteristics allow a better understanding and planning of the needs for the physical and psychological benefit of affected women. In contrast, economically active and productive women suffer a greater impact on QoL after diagnosis and treatment, triggered by the feeling of impotence in the face of illness, of their social and family life, and fearing the future for the uncertainty of the present, because they conciliate job activities with the treatment.⁽¹⁷⁾ In counterpoint, authors⁽²⁾ of a Chinese study highlight the association of a high family income with better QoL of patients with breast cancer in each measured domain. Guidelines and approaches to this group need to be conducive of effective communication throughout treatment by alleviating stressful effects resulting from changes in their emotional, social and family life.

The comparison of clinical staging between institutions showed the public institution had 42.5% more women with stage III than the private institution. This demonstrates women start treatment in more aggressive stages of breast cancer in public institutions, and the access of women from a private institution is more precocious. These data corroborate with the study in Africa. The objective was to understand the late diagnosis and impact of the stage of breast cancer among Libyan women. Stage III was the most expressive, with a diagnosis time of more than six months, which potentially results in high mortality rates.⁽¹⁸⁾ The late diagnosis is focused on several factors related to the population's characteristics, low educational and socioeconomic level, and tracking policies, which make the access to early diagnosis difficult, worsens the prognosis, and leads to more aggressive treatments with impact on women's survival and QoL. Health professionals' interventions should be performed at all levels of care to provide agile assistance and according to their reality. QoL is a multidimensional

construct that allows identifying the psychosocial aspects altered by cancer and its treatments. It is a support tool for nurses and other professionals in the management of alterations in women with breast cancer.

Data from QLQ-C30 questionnaires at the private institution show a higher QoL score compared to the public institution. However, the same domains are scored in both institutions: physical functioning, emotional functioning, fatigue, nausea and vomiting. These are common effects in patients receiving chemotherapy for breast cancer. In both institutions, most women with breast cancer arrived for treatment already in stage III, leading to the choice for the most aggressive therapy that causes greater decline in physical functioning. A Brazilian study with the same population profile is in agreement with these data.⁽¹⁾ It shows physical functioning with lower means, 60.5% and 64.43% respectively, which characterizes impairment in this domain with a decrease in QoL.

Accordingly, a study in Turkey that evaluated QoL and nursing care satisfaction during chemotherapy using the FACT-G questionnaire demonstrated the domain of physical well-being was compromised, with the score of 14.85%, which evidenced that more advanced stages experience more physical symptoms than initial stages.⁽¹⁹⁾ In contrast, Hoyer et al.⁽¹²⁾ observed scores greater than 80% in physical functioning among Swedish women on chemotherapy, and mean values above 75% also for social functioning, indicating no impairment of QoL in relation to these functions. Analogous results were found in a similar study conducted in Malaysia, with mean values above 75% for physical and social functioning, which positively characterizes the maintenance of living conditions, and social and leisure activities among affected women.⁽²⁰⁾ The impairment of physical and social functioning reflects a reduction in the socio-occupational participation of these women regarding their involvement in daily situations often impaired by decreased mobility in the homolateral arm of breast cancer⁽¹⁾ that results in worsening QoL concerning these domains.

Emotional functioning was compromised in the three phases in both institutions, showing that patients were worried, fearful and emotionally fragile throughout treatment, compromising QoL in this domain. In their studies, authors⁽¹⁾ emphasized marked changes in emotional functioning, with mean values of 48.4%, which indicate the QoL impairment of patients undergoing chemotherapy. Women with breast cancer live with emotional disorders daily, such as feelings of imminent death and fear, and behaviors caused by the threat of cancer given the actual or idealized proximity of incapacity or risk of life that lead to fear, anguish, shame, and feelings of discrimination that are difficult situations to be embraced emotionally by patients.^(1,3)

A study conducted in China,⁽²¹⁾ assessed the relationship between depression, anxiety and QoL of 269 women during treatment, and 148 women who had completed treatment one year before by using the Hospital Anxiety and Depression Scale-Cantonese/Chinese version and FACT-G questionnaires. It showed an association between high levels of anxiety with worse physical and functional functioning, and impairment in emotional functioning during and after therapy. This indicates an impairment of QoL regardless of social status, staging and modality of treatment. The QLQ-BR23 questionnaire scores show there is a singularity in relation to systemic effects, but with significance for the public institution. Patients of the public institution had significant scores when evaluated in relation to systemic effects of chemotherapy. This may occur because of the treatment aggressiveness at more advanced stages of the disease. Another aspect to be considered is that women attended at private institutions often have access and better conditions to obtain medications that minimize side effects.⁽¹⁴⁾

Other results presented with the QLQ-BR23 questionnaire showed a significantly higher score for the private institution related to body image, and in the public institution for breast symptoms. It is noteworthy that breast cancer therapy can affect several dimensions of women's lives. It is directly related to the symptomatology and women's perception of their body image, besides

compromising the QoL.⁽¹⁴⁾ The altered body image is an uninterrupted sign of the presence of cancer that maintains and aggravates feelings of anxiety, fear, revolt, and psychologically affects women's interpersonal relationships. Therefore, cancer nurses must develop actions to favor and stimulate social, family, affective, and sexual life, and the QoL in relation to self-image.

The emotional and social reactions caused by breast cancer diagnosis in the life of a woman are singular. She receives the news of cancer as something unexpected, threatening, capable to reach the integrity and proof of female existence, since breasts are the symbolism and concept that women make of themselves. Feelings of death, stigma of mutilation and feelings that may alter self-esteem can cause change and devaluation of these patients' social aspect.⁽²²⁾ In a comparison between institutions regarding altered domains in quality of life, there was a higher score for social functioning in women under care in the private institution. It is noteworthy that these women have higher educational level, high family income, and are inserted in the labor market. As treatment is long and mutilating, it takes them away from their routine, which possibly led to the direct alteration in this domain.

In the public institution, pain was a significant factor in the change in QoL. This fact can be justified by the high number of late diagnoses that result in more discomfort to women, and more aggressive treatments. Like fatigue, it is a frequent, persistent symptom that causes physical and emotional exhaustion to patients and their families because of difficulties in daily activities and self-care, changes in mood and concentration.⁽²³⁾ In the study by Guimarães and Anjos,⁽²⁴⁾ pain was identified before chemotherapy in 20% of the women, with increased mean values, intensification of the symptom during therapy, and losses in the QoL. It reverberates in daily life with emotional, social and functional impairment, and prevents women from performing their routine activities. Measuring pain is one of nurses' actions and it requires knowledge of evaluation methods and factors that trigger and overcome this symptom by considering its subjectivity.

In nursing care for women with breast cancer, the prediction of coping strategies that preserve self-esteem, social and emotional functioning, and physical effects of cancer and chemotherapy treatment should be part of the daily life of these professionals. The bonding between women and nurses during this period may help women to better understand their body changes, making the process of coping with the disease less stressful and exhausting⁽²⁵⁾ and consequently, resulting in a better QoL. A limiting factor in this study was the size and lack of randomness of the sample. The reduced number allows to consider the findings only for the population in question. It was also difficult to find studies comprising results in mixed institutions for comparison with our data.

Conclusion

In this study, were investigated the possible impacts of breast cancer and chemotherapy treatment on women's QoL in public and private institutions. Chemotherapy deteriorated women's QoL in both institutions, and it was lower in patients at the public institution. Differences between public and private institutions may impact the QoL differently. In the study, the only difference found in women's characteristics was their age at the time of diagnosis, which was lower in the private institution. Nursing should have a better understanding about the type of institution responsible for women's care in order to identify the domains and affected symptoms, which will allow the provision of comprehensive care to these women at each phase of treatment hence improving their QoL.

References

1. Fangel LMV, Panobianco MS, Kebbe LM, Almeida AM, Gozzo T O. Qualidade de vida e desempenho de atividades cotidianas após tratamento das neoplasias mamárias. *Acta Paul. de Enferm.* 2013; 26(1):93-100.
2. Yan B, Yang L, Hao L, Yang C, Quan L, Wang L, et al. Determinants of Quality of Life for Breast Cancer Patients in Shanghai, China. *PLoS One.* 2016; 11(4):e0153714.
3. Terra FS, Costa AMDD, Damasceno LL, Lima TS, Filipini CB, Leite MAC. Avaliação da qualidade de vida de pacientes oncológicos submetidos à quimioterapia. *Rev. Bras. Clin. Med.* 2013; 11(2):112-7.
4. Gozzo TO, Nascimento TG, Panobianco MS, Almeida AM. Ocorrência de neutropenia em mulheres com câncer de mama durante tratamento quimioterápico. *Acta Paul. Enferm.* 2011; 6:810-4.
5. Michels FAZ, Latorre MRDO, Maciel MS. Validity, reliability and understanding of the EORTC-C30 and EORTC- BR23, quality of life questionnaires specific for breast cancer. *Rev. Bras. Epidemiol.* 2013; 16(2):352-63.
6. European Organization Research and Treatment of Cancer. (EORTC). Brussels: EORTC; 2014.
7. Ministério da Saúde (BR), Instituto Nacional de Câncer José Alencar Gomes da Silva (INCA). Estimativa 2016: Incidência de câncer no Brasil. Rio de Janeiro: Inca; 2017 [cited 15 Nov 2017]. Available from: <http://www.inca.gov.br/estimativa/2016/>
8. Renck DV, Barros F, Domingues MR, Gonzales MC, Sclowitz ML, Caputo ED, et al. Equidade do acesso ao rastreamento mamográfico do câncer de mama com intervenção de mamógrafo móvel no Sul do Rio Grande do Sul, Brasil. *Cad. Saúde Pública.* 2014; 30(1):88-96.
9. Lima ALP, Rolim NCOP, Gama MEA, Pestana AL, Silva EL, Cunha CLF. Rastreamento oportunístico do câncer de mama entre mulheres jovens no Estado do Maranhão, Brasil. *Cad. Saúde Pública.* 2011; 27(7):1433-9.
10. Keegan TH, DeRouen MC, Press DJ, Kurian AW, Clarke CA. Occurrence of breast cancer subtypes in adolescent and young adult women. *Breast Cancer Res.* 2012; 14(2):R55.
11. Pinheiro AB, Lauter DS, Medeiros GC, Cardoso IR, Menezes LM, Barreto RM, et al. Câncer de mama em mulheres jovens: análise de 12.689 casos. *Rev bras cancerol.* 2013; 59(3):351-9.
12. Hoyer M, Johansson B, Nordin K, Bergkvist L, Ahlgren J, Lidin-Lindqvist A, et al. Health-related quality of life among women with breast cancer – a population-based study. *Acta Oncol.* 2011; 50:1015-26.

13. Sio TT, Chang K, Jayakrishnan R, Wu D, Politi M, Malacame D, et al. Patient age is related to decision-making, treatment selection and perceived quality of life in breast cancer survivors. *World J. Surg. Oncol.* 2014; 12:230.
14. Rodrigues CS, Givisiez JHN. Estudo da acessibilidade ao serviço de saúde de pacientes com câncer de mama no município de campos dos Goytacazes. In: XVII Encontro Nacional de Estudos Populacionais; 2010 set. 20-4. Available from: http://www.alapop.org/Congreso2012/DOCSFINAIS_PDF/ALAP_2012_FINAL811.pdf
15. Rosa LM, Radunz V. Do sintoma ao tratamento adjuvante da mulher com câncer de mama. *Texto Contexto Enferm.* 2013; 22(3):713-21.
16. Simeão SFAP, Landro ICR, De Conti MHS, Gatti MAN, Delgallo WD, De Vitta A. Qualidade de vida em grupos de mulheres acometidas de câncer de mama. *Ciênc. Saúde Coletiva.* 2013; 18(3):779-88.
17. Silva MM, Silva JÁ, Esteves LO, Mesquita MGR, Stipp MAC, Duarte SCM. Perfil sociodemográfico e clínico de pessoas em tratamento quimioterápico: subsídios para o gerenciamento em enfermagem. *Rev. Eletr. Enf.* 2013; 15(3):704-12.
18. Eramah E, Abdalla F, Buhmeida A, Larbesj E, Pyrhonen S, Collan Y. Diagnosis delay in Libyan female breast cancer. *BMC Res. Notes.* 2012; 5(452):1-8.
19. Bayran Z, Durna Z, Akin S. Quality of life during chemotherapy and satisfaction with nursing care in Turkish breast cancer patients. *Eur. J. Cancer Care.* 2014; 23:675-84.
20. Yusuf A, Ahmad Z, Keng SL. Quality of life in Malay and Chinese women newly diagnosed with breast cancer in Kelantan, Malaysia. *Asian Pac. J. Cancer Prev.* 2013; 14:435-40.
21. Ho SSM, So WK, Leung DY, Lai ET, Chan CW. Anxiety, depression and quality of life in Chinese women with breast cancer during and after treatment: A comparative evaluation. *Eur. J. Oncol. Nurs.* 2013; 17:877-82.
22. Anjos JC, Alayala A, Hofelmann DA. Fatores associados ao câncer de mama. *Cad. Saúde Coletiva.* 2012; 20(3).
23. Lamino DA, Mota DDCF, Pimenta CAM. Prevalência e comorbidade de dor e fadiga em mulheres com câncer de mama. *Rev. Esc. Enferm. USP.* 2011; 45(2):508-14.
24. Guimarães AGC, Anjos ACY. Caracterização sociodemográfica e avaliação da qualidade de vida em mulheres com câncer de mama em tratamento quimioterápico adjuvante. *Rev. Bras. Cancerol.* 2012; 58(4):581-92.
25. Gomes NS, Silva SR. Avaliação da autoestima de mulheres submetidas à cirurgia oncológica mamária. *Texto Contexto Enferm.* 2013; 22(2):509-16.