



Investigación y Educación en Enfermería

ISSN: 0120-5307

ISSN: 2216-0280

Imprenta Universidad de Antioquia

Ortega Barco, María Alejandra; Muñoz de Rodríguez, Lucy
Evaluation of the Nursing Care Offered during the Parturition Process. Controlled Clinical Trial of an Intervention based on Swanson's Theory of Caring versus Conventional Care¹

Investigación y Educación en Enfermería, vol. 36, no. 1, e5, 2018, January-April

Imprenta Universidad de Antioquia

DOI: 10.17533/udea.iee.v36n1e05

Available in: <http://www.redalyc.org/articulo.oa?id=105255813005>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UAEH
redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Evaluation of the Nursing Care Offered during the Parturition Process. Controlled Clinical Trial of an Intervention based on Swanson's Theory of Caring versus Conventional Care

María Alejandra Ortega Barco¹
Lucy Muñoz de Rodríguez²



Original article



UNIVERSIDAD
DE ANTIOQUIA
1803

Evaluation of the Nursing Care Offered during the Parturition Process. Controlled Clinical Trial of an Intervention based on Swanson's Theory of Caring versus Conventional Care

Objective. This work sought to compare the evaluation of the nursing care provided during the parturition process in the intervention group based on Swanson's theory of caring versus that of the control group that received conventional care. **Methods.** Preventive-type controlled clinical trial conducted in a tier II hospital in San Gil, Santander (Colombia). During the parturition process, the intervention group received care based on Swanson's theory of caring ($n=20$) and the control group received conventional care ($n=23$). During immediate postpartum, the mothers were applied the Professional care rating scale by Swanson, validated into Spanish in Colombia. **Results.** Assessment of

1 Nurse, Masters. Universidad Nacional de Colombia, at Bogotá (Colombia).

email: maortegab@unal.edu.co

2 Nurse, Masters. Professor emeritus, Universidad Nacional de Colombia, at Bogotá (Colombia).

email: lmunozdr@unal.edu.co

Conflicts of Interest: None.

Received: July 28th, 2017

Accepted: November 14th, 2017

How to cite this article: Ortega MA, Muñoz L. Evaluation of the Nursing Care Offered during the Parturition Process. Controlled Clinical Trial of an Intervention based on Swanson's Theory of Caring versus Conventional Care. Invest. Educ. Enferm. 2018; 36(1):e05.

DOI: 10.17533/udea.iee.v36n1e05



Investigación y Educación en

Enfermería

—Nursing Research and Education—

Vol. 36 No1, • February 15th 2018 • ISSN: 2216-0280

professional care in the intervention group was Excellent with an average for the scale total of 59.8 points of a possible maximum of 60 points, while in the control group it was Good with 50.2 ($p<0.0001$). In the 15 items that make up the scale and in both subscales (Compassionate Healer and Competent Healer) higher scores were also observed in the intervention group compared to those of the control group. Conclusion. The intervention derived from Swanson's theory of caring was associated to a better evaluation of professional nursing care by women during the parturition process.

Descriptors: nursing care; behavior rating scale; labor, obstetric; parturition; controlled clinical trial.

Valoración del cuidado de Enfermería brindado durante el proceso de parto. Ensayo clínico controlado de una intervención basada en la Teoría de Swanson versus cuidado convencional

Objetivo. Comparar la valoración del cuidado de enfermería brindado durante el proceso de parto en el grupo de Intervención basada en la Teoría de Swanson versus la del grupo de Control que recibió cuidado convencional. **Métodos.** Ensayo clínico controlado de tipo preventivo realizado en un hospital de segundo nivel de atención en San Gil, Santander (Colombia). Durante el proceso de parto el grupo de *Intervención* recibió cuidado basado en la Teoría de Swanson ($n=20$) y al grupo de *Control* se le practicó el cuidado convencional ($n=23$). En el postparto inmediato se aplicó a las madres la *Escala de Valoración del cuidado profesional* de Swanson, validada al español en Colombia. **Resultados.** La valoración del cuidado profesional en el grupo de *Intervención* fue *Excelente* con un promedio para el total de la escala de 59.8 puntos de un máximo posible de 60, mientras que en el grupo de *Control* fue *Bueno* con 50.2 ($p<0.0001$). En los 15 ítems que conforman la escala y en las dos subescalas (Sanador Compasivo y Sanador Competente) también se observaron mayores puntajes en el grupo de *Intervención* comparados con los del grupo de

Control. **Conclusión.** La intervención derivada de la teoría de Swanson se asoció a una mejor valoración del cuidado profesional de enfermería por parte de las mujeres durante el proceso de parto.

Descriptores: atención de enfermería; escala de evaluación de la conducta; trabajo de parto; parto; ensayo clínico controlado.

Valorização do cuidado de Enfermagem brindado durante o processo de parto. Ensaio clínico controlado de uma intervenção baseada na Teoria de Swanson versus cuidado convencional

Objetivo. Comparar a valorização do cuidado de enfermagem brindado durante o processo de parto no grupo de Intervenção baseada na Teoria de Swanson versus a do grupo de Controle que recebeu cuidado convencional. **Métodos.** Ensaio clínico controlado de tipo preventivo realizado num hospital de segundo nível de atenção em San Gil, Santander (Colômbia). Durante o processo de parto o grupo de *Intervenção* recebeu cuidado baseado na Teoria de Swanson ($n=20$) e ao grupo de *Controle* se lhe praticou o cuidado convencional ($n=23$). No pós-parto imediato se aplicou às mães a *Escala de Valorização do cuidado profissional* de Swanson, validada ao espanhol na Colômbia. **Resultados.** A valorização do cuidado profissional no grupo de *Intervenção* foi *Excelente* com uma média para o total da escala de 59.8 pontos de um máximo possível de 60, enquanto que no grupo de *Controle* foi *Bom* com 50.2 ($p<0.0001$). Nos 15 itens que conformam a escala e nas duas sub-escalas (Curador Compassivo e Curador Competente) também se observaram maiores pontuações no grupo de *Intervenção* comparados com os do grupo de *Controle*. **Conclusão.** A intervenção derivada da teoria de Swanson se associou a uma melhor valorização do cuidado profissional de enfermagem por parte das mulheres durante o processo de parto.

Descritores: cuidados de enfermagem; escala de avaliação comportamental; trabalho de parto; parto; ensaio clínico controlado.

Introduction

The parturition process is a complex phenomenon due to different factors that interact amongst themselves,⁽¹⁾ which can increase the risk of complications and death, and which additionally affect the mother's wellbeing. Likewise, women during the parturition process report feeling mainly fear,⁽¹⁻³⁾ which alters their performance and comfort, especially in primiparous.⁽¹⁾ These negative emotions,⁽¹⁾ constant pain, affectation in the capacity to make decisions,⁽⁴⁾ communication difficulties with the health professionals,⁽¹⁾ and care guided with biological approach translate into negative perception toward the care givers and the parturition.⁽⁴⁾

For women to have a positive experience,⁽⁴⁾ participation is recommended from nurses trained and experts in obstetrics during parturition care,⁽⁵⁾ given that this favors the naturalness of the event and diminishes rates of caesareans and complications. Through humane care, they can reduce potential problems that arise;⁽⁶⁾ however in practice, nursing care is provided routinely,^(7,8) with knowledge of the task involved, but without disciplinary theoretical orientation, and although they are suitable professionals to achieve broad benefits and satisfaction of the woman during the parturition process, little time is assigned for this purpose, dedicating over 70% of the time to objectives different from specific care⁽⁸⁾ due to different barriers⁽⁹⁾ identified by the nurses and by the women.

Even so, nurses have been seen as leaders^(10,11) to favor empowerment and control during the parturition process;⁽¹⁰⁾ women value individualized information, explanations, advice, and nursing care⁽²⁾ and identify the professionals as humanized, respectful, and sensitive to the women's expressions of pain and joy;⁽²⁾ however, these perceptions are sporadic.-

In light of this panorama, it is necessary to provide care based on the theory of the discipline that permits appreciating benefits and wellbeing during the parturition process. Swanson's theory of nursing as informed caring for the well-being of others⁽¹²⁾ is the support of the intervention proposed in this study, which guides nurses through the five care processes proposed: Maintaining belief in the woman's capacities to confront the situation, approaching with affection, offering hope and favoring autonomy;⁽¹⁾ Knowing the meaning of the event, without judgments and understanding specific manifestations; Being with - listening to concerns, being available, offering emotional support, and understanding feelings; Doing for - offering physical measures of comfort and information, controlling fear and pain;⁽²⁾ Enabling – protecting the woman from any type of harm and favoring self-care, guiding in the process. A theoretical analysis was performed and, supported by existing evidence, the specific activities were constructed for nurses to apply care processes during the parturition process.

Seeking to compare the assessment of nursing care offered during the parturition process in the intervention group based on Swanson's theory of

caring versus that of the control group, which received conventional care, the intervention proposed was applied and nurses were evaluated, in both groups, with the Professional care rating scale by the same author.⁽¹³⁾

Methods

This was a preventive-type controlled clinical trial. A minimum sample size was calculated of 20 women for each of the study groups, bearing in mind an alpha value of 0.05, a beta value of 0.8, and a minimum difference between groups of 8 points for the scale total. The participants were women over 14 years of age who were hospitalized to care for the parturition process in the maternity service at the Regional Hospital in San Gil, Santander (Colombia) during the period from September to November 2016, and who signed the informed consent.

Assignment of the women to the study groups was done according to the day they were hospitalized during labor. The control group was comprised by the mothers who received conventional care from the maternity service staff plus the skin-to-skin contact during the immediate postpartum, while the intervention group was provided care based on the care processes proposed by Swanson in the Theory of informed caring for the well-being of others.⁽¹³⁾ For this intervention, a last-year Nursing student was trained through 6 h of theory, 5 h of supervised practical hours, and 40 h of independent practice on the specific activities of the intervention proposed to be applied during labor and expulsion.

During the immediate postpartum, another last-year Nursing student who ignored to what group the mothers had been assigned, applied the Professional care rating scale based on Swanson's theory of caring⁽¹³⁾ to the participants from the intervention and control groups. The rating scale has 15 items with four Likert-type response options: 1 = never, 2 = sometimes yes, sometimes no, 3 = most of the time, and 4 = always. The general classification of

professional care is considered according to the total score, thus: <23 = Deficient; 23 to 37 = Regular; 38 to 51 = Good; and 52 to 60 = Excellent⁽¹⁴⁾. The scale applied received semantic validation for Colombia with excellent reliability (Cronbach's alpha: 0.90).⁽¹⁴⁾ To store the information, the study used an electronic device with the scale aggregated, connected to an immediate response to the principal researcher's e-mail and a meticulous record of the women's answers was kept.

This research was approved by the Ethics Committee of the Faculty of Nursing at Universidad Nacional de Colombia and by the Ethics Committee at the Regional Hospital in San Gil. Respect for the dignity of the women prevailed, along with the protection of their rights, and their wellbeing. The study kept in mind the ethical principles contemplated in Legislation 911 of 2004: Beneficence, Non-maleficence, Justice, Autonomy, Fidelity, and Reciprocity. The study was classified as minimum risk and participant information confidentiality was protected.

Results

During the study period, 107 women in labor were admitted to the Hospital's maternity service, 43 of them fulfilled the inclusion criteria; 20 were assigned to the intervention group and 23 to the control group.

Sociodemographic characteristics

The mean age in the intervention group was 25.15 ± 7.91 years versus 24.35 ± 5.77 years in the control group ($p=0.893$). Table 1 shows that the groups are similar regarding the sociodemographic variables and obstetric history. Most of the women in both groups are between 20 and 35 years of age, are registered in the subsidized regime and report their level of studies to complete secondary. Other common characteristics in these women is that they come from municipalities near San Gil to have their first parturition, with gestations at full term, having

as antecedents compliance of prenatal controls and attendance to the motherhood and fatherhood preparation course. Principally, the women in both study groups had no complications during gestation

or labor. The parturition lasted 14.85 ± 12.11 h in the intervention group versus 17.65 ± 10.45 h in the control group ($p=0.420$), with this difference not being statistically significant.

Table 1. General characteristics of the study groups

Variable	Group				p value*
	Intervention (n=20)		Control (n=23)		
	Number	%	Number	%	
Sociodemographic Characteristics					
Age group in years					0.622
15-17	4	20	4	17.39	
18-19	3	15	2	8.70	
20-25	4	20	8	34.78	
26-35	6	30	8	34.78	
36 and more	3	15	1	4.35	
Marital status					0.408
Married	5	25	4	17.39	
Single	5	25	3	13.04	
Common law	10	50	16	69.57	
Healthcare Regime					0.266
Contributive	4	20	2	8.70	
Subsidized	16	80	21	91.30	
Schooling					0.225
Up to primary	9	45	7	30.43	
Up to secondary	9	45	13	56.52	
Technical or university	2	10	0	0.00	
Origin					0.498
San Gil	6	30	8	34.78	
Other municipalities	14	70	15	65.22	
Obstetric history					
Parity					0.867
0	11	55	14	60.87	
1	4	20	5	21.74	
2	3	15	3	13.04	
3	1	5	1	4.35	
4	1	5	0	0.00	
Gestation to term					0.342
Early (37-38.6 weeks)	10	50	9	39.13	
Full (39-40.6 weeks)	10	50	14	60.87	
Planned gestation					0.447
Yes	10	50	14	60.87	
No	10	50	9	39.13	

Table 1. General characteristics of the study groups. (Cont.)

Variable	Group				p value*
	Intervention (n=20)		Control (n=23)		
	Number	%	Number	%	
Number of prenatal controls					0.994
0	1	5	1	4.35	
1-3	1	5	1	4.35	
4-8	14	70	17	73.91	
9 and more	4	20	4	17.39	
Motherhood and fatherhood preparation course					0.865
Yes	17	85	20	86.96	
No	3	15	3	13.04	
Complications during gestation					0.405
Yes	5	25	4	17.39	
No	15	75	19	82.61	
Complications during parturition					0.431
Yes	3	15	2	8.70	
No	17	85	21	91.30	
Use of oxytocin					0.261
Induction	6	30	7	30.43	
Conduction	10	50	15	65.22	
No	4	20	1	4.35	

* Chi square test probability value

Results in the Professional care rating scale

Assessment of professional care in the intervention group was *Excellent* (average 59.8 ± 0.6 , minimum = 58, maximum = 60), while the control group was *Good* (average 50.2 ± 7.8 , minimum = 39, maximum = 60). Table 2 reviews the average values obtained in each of the items, which per subscale and scale total were higher in the intervention group compared to those of the control group, and all these differences were statistically significant. Specifically, in the intervention group the scores of the 15 items were between 4 and very close to 4 (3.95-3.97), indicating that the questions were mostly answered with the option "always".

It was not found that the scale's total score was affected by possible interactions between the study group with the variables of use of oxytocin ($p = 0.85$) or time of labor duration ($p = 0.57$).

Discussion

Most of the women in parturition process for both groups are within the age group 20 to 35 years, corresponding to the ages of highest rates of fertility for Colombia,⁽⁶⁾ an important variable given that the two extreme groups, adolescents and women over 35 years of age are of obstetric risk. These data reiterate that found in Europe, Latin America, and Colombia where similar ages are reported.^(1,15-17)

Women hospitalized for the birth of their children came from municipalities close to San Gil and many had basic primary and secondary studies, differing from data reported in other Colombian studies^(15,18) in which between 50% and 60% of the women had higher levels of schooling; in this research, <5% had this level of schooling and it may be because most of the women come from small towns near San Gil, while the

Table 2. Averages per item, subscale, and scale of the Professional Care Rating Scale according to study group

Subscale	Item	Group		p value*
		Intervention	Control	
Compassionate Healer	1. Made you feel well?	3.95±0.22	3.48±0.59	0.002
	2. Showed a positive attitude toward you and their work?	4±0	3.34±0.71	<0.001
	3. Listened attentively?	4±0	3.17±0.88	<0.001
	4. Let you express your feelings?	3.95±0.22	2.74±0.96	<0.001
	5. Showed interest for what happened to you?	4±0	3.04±0.87	<0.001
	6. Understood your symptoms and concerns?	4±0	3.26±0.62	<0.001
	7. Showed they were prepared to do their work?	3.95±0.22	3.52±0.51	0.002
	<i>Average of the Compassionate Healer scale</i>	3.97±0.07	3.22±0.55	<0.001
Competent Healer	8. Encouraged you to take care of yourself during labor?	4±0	3.56±0.51	0.001
	9. Was respectful with you?	4±0	3.65±0.58	0.004
	10. Offered help and collaboration?	4±0	3.17±0.88	<0.001
	11. Performed labor control correctly?	3.95±0.22	3.47±0.59	0.002
	12. Did you feel cared for during your labor?	4±0	3.39±0.66	<0.001
	13. Clearly explained the indications to follow?	4±0	3.26±0.86	<0.001
	14. Was kind with you?	4±0	3.61±0.49	0.002
	15. Treated you like a person?	4±0	3.52±0.51	<0.001
	<i>Average of the Competent Healer scale</i>	3.99±0.03	3.5±0.5	<0.001
	Average of the scale total	59.8±0.6	50.2±7.8	<0.001

* Mann-Whitney U test probability value

Colombian studies were held in Bogotá in sectors with prevalence of high socioeconomic level.^(15,18) This reveals the need for intervention to women in parturition process in this province or in regions with similar characteristics. More than half the women were in the parturition of their first child; research in different continents report predominance of primiparous women,^(1,15-20) which shows that, in general, women are in a group that requires greater care because they are upon the birth of their first child. Changes of different intensities occur during the parturition process, which behave as a complex phenomenon, especially for the primiparous women, who feel more fear than multiparous women;⁽¹⁾ thus, the interrelationship with nurses improves the woman's performance and wellbeing during the parturition process. It must be highlighted that the parturitions occurred in pregnancies with gestational ages to term, which require principally of care and of the nurse's continuous presence^(7,21)

to contribute to the naturalness of the process.

Most of the women reported that their gestation was unplanned and this has been associated to the development of post-traumatic stress;⁽⁷⁾ additionally, they comply with prenatal controls, but do not attend the motherhood and fatherhood preparation course – although they required it because most were primiparous. This corresponds to a wake-up call for care to be offered from the initial stages of gestation to the moment of parturition; the nurses' presence and the support they provide during labor can diminish the onset of postpartum stress.⁽²²⁾

Regarding the results of applying the *Professional care rating scale*, it was noted that the score was higher in the intervention group compared to the control group, showing the benefits of applying this care intervention during the mothers' parturition process. These data are a reminder that women request and grant importance to the relational

component of nursing care, which influences upon the care assessment, corresponding to that found in Latin American studies.^(15,18)

Given that the participants from the intervention group scored the 15 items from the scale with values around 4, it is important to analyze some items in the control group that deserve attention: the reactive with the lowest score was # 4- Let you express your feelings? (2.74), which beckons nurses to keep in mind activities from the process “Being with”, broadening care toward feelings, remembering that women during parturition process have mixed feelings.⁽¹⁾ Item 1- Made you feel well? Obtained an average score in the control group of 3.48. By using the linear interpolation of the averages of each item from the scale with its total score, it was found that this score corresponds to the lower limit for professional care considered *Excellent*, hence, the care process “Maintaining belief” requires being delved into by nurses who apply the intervention.

In the *Competent* Healer subscale for the control group, the lowest average scores were: 13- Clearly explained the indications to follow?; and 10- Offered help and collaboration?, which calls on nurses in general to improve in the education and direct care of women during parturition process; these scores obtained a classification of *Excellent* in the intervention group. It should be remembered that women who do not feel cared for by nurses report development of postpartum post-traumatic stress⁽⁷⁾, which drives to continue delving into this care intervention and its benefits during the parturition process, expulsion and in the immediate, middle and late postpartum.

The scale’s total score for the women who received intervention was in the category of

Excellent, while being classified as *Good* for the control group, which agrees with the data found in other research,⁽²³⁾ which also applied during prenatal control an intervention derived from Swanson’s theory of caring, with before and after measurements, and finding that women value nurses better after the intervention.

Finally, it was observed that in this study’s care intervention, the scores were equal in both subscales, that is, they keep in mind the care aspects of Compassionate Healer and Competent Healer, while for the control group the score of the Compassionate Healer subscale was lower. Studies related to the nursing care offered to women during parturition process demonstrate that its evaluation is conceived through two components: one technical – less strong, but not less important – and another relational, considered the most important,^(8,15) hence, the interrelation between the nurses and women influences directly on the perception of the care assessment.⁽¹⁵⁾

The conclusion herein is that the intervention derived from Swanson’s theory of caring was associated to a better assessment of the professional nursing care by women during the parturition process. Research should be conducted to continue with the assessment of this study’s intervention, as well as explore the different variables related to the process that can modify the scores manifested by the women in the subscales during the evaluation of the professional care and the variables susceptible to modification by the care intervention. Likewise, future research should review the impact of nursing care with evaluations at different times, like during the middle and late postpartum, as well as in women who have been programmed for caesarean.

References

1. Dodou H, Paiva D, Marculino E, Cavalcante M, Nery P, Sousa N. A contribuição do acompanhante para a humanização do parto e nascimento: percepções de puérperas. *Esc. Anna Nery Rev. Enferm.* 2014; 18(2):262-9.
2. Hodnett E, Gates S, Hofmeyr G, Sakala C. Continuous support for women during childbirth. *Cochrane Database of Systematic Reviews*. 2012; 10:CD003766.
3. Lally J, Thomson R, McPhail S, Exley C. Pain relief in labour: a qualitative study to determine how to support women to make decisions about pain relief in labour. *BMC Pregnancy childbirth*. 2014; 14:6.
4. Palinski J, Kissula S, Pessoa J, Rejane N, Gualda D. Women's perception of the process of labor coaching: study descriptive. *Online Braz. J. Nurs.* 2012; 11(2):274 -88.
5. Organización Mundial de la Salud OMS. Mortalidad materna, Nota descriptiva N° 348 [Internet]. 2016 [cited 21 nov 2016]. Available from: <http://www.who.int/mediacentre/factsheets/fs348/es/>
6. Ministerio de Salud y de la Protección Social. Análisis de Situación de Salud Colombia, 2015 [cited 21 nov 2016]. Bogotá: El Ministerio. Available from: <https://www.minsalud.gov.co/sites/rid/Lists/BibliotecaDigital/RIDE/VS/ED/PSP/asis-2015.pdf>
7. Andersen L, Melvaer L, Videbech P, Lamont R, Joergensen J. Risk factors for developing post-traumatic stress disorder following childbirth: a systematic review. *Acta Obstet. Gynecol. Scand.* 2012; 91:1261-72.
8. De Fátima C, Pereira N, Odeh L, Goulart B. Parto de adolescentes: elementos qualitativos da assistência. *Rev. Esc. Enferm. USP.* 2012; 46(2):287-94.
9. Souza TG, Gaíva MAM, Modes PSSA. A humanização do nascimento: percepção dos profissionais de saúde que atuam na atenção ao parto. *Rev. Gaúcha Enferm.* 2011; 32(3):479-86.
10. Brasil EGM, Queiroz MVO, Fernandes AFC, Costa RF, Xavier EO. Perception of women on the care in the childbirth: contributions to nursing. *Acta Scientiarum Health Sciences*. 2013; 35(2):195-200.
11. Hanson L, VandeVusse L. Supporting labor progress toward physiologic birth. *J. Perinatal Neonatal Nurs.* 2014; 28(2):101-7.
12. Swanson K. Nursing as informed caring for the well-being of others. *Image J. Nur. Sch.* 1993; 25(4):352-7.
13. Watson J. Assessing and measuring caring in nursing and health science. New York: Springer Pub.; 2002.
14. Vesga L, Ruíz C. Validez y confiabilidad de una escala de cuidado profesional en español. *Av. Enferm.* 2016; 34(1):69-78.
15. Saavedra D. Comportamientos de cuidado del profesional de enfermería que percibe la gestante en trabajo de parto y parto de una clínica privada [Internet]. Bogotá: Universidad Nacional de Colombia; 2013 [cited 7 Dec 2016]. Available from: <http://www.bdigital.unal.edu.co/11168/1/deisylorenasaavedrainfante.2013.pdf>
16. Silva FMB, Paixao TCR, Oliveira SMJV, Leite JS, Riesco MLGonzalez, Osava RH. Care in a birth center according to the recommendations or the World Health Organization. *Rev. Esc. Enferm. USP.* 2013; 47(5):1031-8.
17. Giraldo D, González E, Henao C. Experiencias de las mujeres durante el trabajo de parto y parto. *Av. Enferm.* 2015; 33(2):271-81.
18. Díaz E, Mendoza E. Percepción que tienen las puérperas en relación con el cuidado brindado por el personal de enfermería durante la atención del parto en una institución de cuarto nivel durante el primer semestre del año 2013 [Internet]. Bogotá: Pontificia Universidad Javeriana; 2013. Available from: <https://repository.javeriana.edu.co/handle/10554/13773>
19. Grilo C, d'Orsi E, Soares R, Alves J, Bastos M, Schneck C, et al. Implementação da presença de acompanhantes durante a internação para o parto: dados da pesquisa nacional Nascer no Brasil. *Cad. Saúde Publica.* 2014; 30:140-53.
20. Whitburn L, Jones L, Davey M. Small, R. Women's experiences of labour pain and the role of the mind: an exploratory study. *Midwifery.* 2014; 30:1029-35.
21. Organización Mundial de la Salud. Recomendaciones de la OMS para la conducción del trabajo de parto [Internet]. 2015 [cited Dec 1 2016]. Available from: http://www.who.int/topics/maternal_health/directrices_OMS_parto_es.pdf

22. Harris R, Ayers S. What makes labour and birth traumatic?: A survey of intrapartum 'hospots'. *Psychol.Health*. 2012; 27(10):1166-77.
23. Reyes D. Valoración del cuidado profesional de Enfermería que realizan las adolescentes gestantes antes y después de una intervención de cuidado [Internet]. Bogotá: Universidad Nacional de Colombia; 2016 [cited Nov 28 2016]. Available from: <http://www.bdigital.unal.edu.co/52921/1/63533904.2016.pdf>.