



Investigación y Educación en Enfermería

ISSN: 0120-5307

ISSN: 2216-0280

Imprenta Universidad de Antioquia

Gandhi, Sailaxmi; Poreddi, Vijayalakshmi; Nagarajaiah;
Palaniappan, Marimutthu; Nikhil Reddy, S. Sai; BadaMath, Suresh
Indian nurses' Knowledge, Attitude and Practice towards use of physical restraints in psychiatric patients
Investigación y Educación en Enfermería, vol. 36, no. 1, e10, 2018, January-April
Imprenta Universidad de Antioquia

DOI: 10.17533/udea.iee.v36n1e10

Available in: <http://www.redalyc.org/articulo.oa?id=105255813010>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UDEM  redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Indian nurses' Knowledge, Attitude and Practice towards use of physical restraints in psychiatric patients

Sailaxmi Gandhi¹
Vijayalakshmi Poreddi²
Nagarajaiah³
Marimutthu Palaniappan⁴
S. Sai Nikhil Reddy⁵
Suresh BadaMath⁶

Indian nurses' Knowledge, Attitude and Practice towards use of physical restraints in psychiatric patients

Objective. To assess nurses' knowledge, attitude and practice towards using physical restraints among psychiatric patients. **Methods.** A descriptive cross sectional survey was carried out among conveniently selected sample of nurses working in psychiatry departments at a tertiary care center. The data was collected using self reported questionnaires of Suen. **Results.** The findings revealed that nurses had good knowledge (7.2 ± 1.7 , maximum possible=11), favorable attitudes 30.8 ± 3.3 (maximum possible=48) and good practice 31.2 ± 6.2 (maximum possible=42) about use of physical restraints in psychiatric patients. Females had better knowledge ($p < 0.001$), attitudes ($p < 0.05$) than males towards use of physical restraints. Nurses those had more than ten years of experience found to have more favorable attitudes towards using physical restraints than nurses with less experience ($p < 0.05$) and nurses with higher education



Original article



UNIVERSIDAD
DE ANTIOQUIA
1803

- 1 Ph.D. Additional Professor, Department of Nursing National Institute of Mental Health and Neuro Sciences, (Institute of National Importance), Bangalore (India).
email: sailaxmi63@yahoo.com
- 2 RN, RM, BSN, MSN. College of Nursing, National Institute of Mental Health and Neuro Sciences, (Institute of National Importance), Bangalore (India).
email: pvijayalakshmi63@gmail.com
- 3 PhD, Former Additional Professor, Department of Nursing National Institute of Mental Health and Neuro Sciences, (Institute of National Importance), Bangalore (India).
email: dr.nagarajaiah@gmail.com
- 4 PhD, Additional Professor, Department of Bio-Statistics, National Institute of Mental Health and Neuro Sciences, (Institute of National Importance), Bangalore (India).
email: p_marimutthu@hotmail.com
- 5 Second year MBBS. Bangalore Medical College and Research Institute, Bangalore (India).
email: saithereddy@gmail.com
- 6 MD, Psychiatry. Professor, Department of Psychiatry National Institute of Mental Health and Neuro Sciences, (Institute of National Importance), Bangalore (India).
email: nimhans@gmail.com

Conflicts of interest: None.

Received: June 8th, 2017.

Accepted: November 24th, 2017.

How to cite this article: Gandhi S, Vijayalakshmi P, Nagarajaiah, Marimutthu P, Reddy SSN, Suresh BM. Indian nurses' Knowledge, Attitude and Practice towards use of physical restraints in psychiatric patients. Invest. Educ. Enferm. 2018; 36(1):e10.

DOI: 10.17533/udea.iee.v36n1e10



Investigación y Educación en

Enfermería

—Nursing Research and Education—

Vol. 36 No1, • February 15th 2018 • ISSN: 2216-0280

differed significantly on practice score than nurses with basic education in nursing ($p<0.05$). **Conclusion.** This study revealed good knowledge, positive attitudes and good practices among nurses about using physical restraints in mental health services. However there is need to improve even more nurses practice through continuing education programs on this topic.

Descriptors: cross-sectional studies; health knowledge, attitudes, practice; restraint, physical; psychiatric nursing.

Conocimientos, actitudes y prácticas de las enfermeras hindúes sobre el uso de restricciones físicas en pacientes psiquiátricos

Objetivo. Evaluar los conocimientos, las actitudes y las prácticas de las enfermeras hacia el uso de restricciones físicas en los pacientes psiquiátricos. **Métodos.** Se realizó un estudio descriptivo de corte transversal en una muestra por conveniencia de enfermeras que laboraban en departamentos de psiquiatría de un hospital de atención terciaria al sur de la India. Los datos se recolectaron utilizando el cuestionario de Suen, el cual fue contestado por autorreporte. **Resultados.** Los hallazgos revelaron que en cuanto al uso de restricciones físicas en pacientes psiquiátricos las enfermeras tenían buen conocimiento (media = 7.2 ± 1.7 , máximo posible=11), actitudes favorables (media = 30.8 ± 3.3 , máximo posible=48) y buenas prácticas (media = 31.2 ± 6.2 , máximo posible=42). Las mujeres tenían mejores conocimientos ($p<0.001$) y actitudes ($p<0.05$) que los varones hacia el uso de restricciones físicas. Las enfermeras con más de diez años de experiencia tuvieron actitudes más favorables hacia el uso de estas restricciones que las de menor experiencia ($p<0.05$) y las enfermeras con educación superior difirieron significativamente en el puntaje de la práctica comparadas con aquellas con educación básica en enfermería ($p<0.05$). **Conclusión.** Este estudio reveló buenos conocimientos, actitudes positivas y buenas prácticas de las enfermeras sobre el uso de restricciones físicas en los servicios de salud mental. Sin embargo, es necesario

mejorar aún más las prácticas de las enfermeras a partir de programas de educación continua sobre este tema

Descritores: estudios transversales; conocimientos, actitudes y práctica en salud; restricción física; enfermería psiquiátrica.

Conhecimentos, atitudes e práticas das enfermeiras hindus sobre o uso de restrições físicas em pacientes psiquiátricos

Objetivo. Avaliar os conhecimentos, as atitudes e as práticas das enfermeiras para o uso de restrições físicas nos pacientes psiquiátricos. **Métodos.** Se realizou um estudo descritivo de corte transversal numa amostra por conveniência de enfermeiras que trabalhavam em departamentos de psiquiatria de um hospital de atenção terciária no sul da Índia. Os dados foram recolhidos utilizando o questionário de Suen, o qual foi contestado por auto-reporte. **Resultados.** As descobertas revelaram que em quanto ao uso de restrições físicas em pacientes psiquiátricos as enfermeiras tinham bom conhecimento (media = 7.2 ± 1.7 , máximo possível=11), atitudes favoráveis (media= 30.8 ± 3.3 , máximo possível=48) e boas práticas (média= 31.2 ± 6.2 , máximo possíveis=42). As mulheres tinham melhores conhecimentos ($p < 0.001$) e atitudes ($p < 0.05$) que os homens para o uso de restrições físicas. As enfermeiras com mais de dez anos de experiência tiveram atitudes mais favoráveis para o uso destas restrições que as de menor experiência ($p < 0.05$) e as enfermeiras com educação superior deferiram significativamente na pontuação da prática comparadas com aquelas com educação básica na enfermagem ($p < 0.05$). **Conclusão.** Este estudo revelou bons conhecimentos, atitudes positivas e boas práticas das enfermeiras sobre o uso de restrições físicas nos serviços de saúde mental. Embora, seja necessário melhorar ainda mais as práticas das enfermeiras através de programas de educação continua sobre este assunto.

Descritores: estudos transversais; conhecimentos, atitudes e prática em saúde; restrição física; enfermagem psiquiátrica.

Introduction

The use of Physical Restraints (PR) within the psychiatric services is a common practice worldwide. Physical restraints refer to any physical methods of restricting a person's freedom of movement, physical activity or normal access to his or her body.⁽¹⁾ Use of restraints in mental health care setting perceived differently as some people view restraint of psychiatric patients as a violation of basic human rights, others as a necessity to reduce the risk of harm among psychiatric patients whether it is directed toward self or toward others⁽²⁾ and others as a therapeutic modality. However, throughout history the practice of restraint has been associated with punishment, custodial care, institutional abuse and neglect.⁽³⁾ On the other hand, using of physical restraints among psychiatric patients raises a range of ethical, legal, and clinical questions.^(4,5) Psychiatric management in India often includes the practice of physical restraints to control aggressive patients not only in psychiatric hospitals and also in places of traditional healing.⁽⁶⁾ For example, the Erwadi tragedy, which was a fire accident, occurred on August 06, 2001 and 28 psychiatric patients died at Moideen Badhusha Mental Home, because they had been fettered and could not escape. Today, nine years later, and despite the existence of laws against such practices, nothing has changed.⁽⁷⁾ Drew et al also reported that mental health policies and laws in low- and middle-income countries failed to incorporate international human rights and best practice standards to prevent violation of human rights. India was not an exception. The new Mental Health Care Bill 2016 focused on a rights-based protection of persons with mental illness and guidelines for using seclusion and restraints.⁽⁸⁾

Earlier studies report that elderly and persons with mental illness have a higher rate of physical restraint use worldwide.⁽⁹⁾ Nurses are ethically obligated to ensure the patient's basic rights and restraints must not be used for coercion, punishment, discipline, or staff convenience.⁽¹⁰⁾ Further, nurses are the health care providers who initiate restraint use and attribute its use to ensuring the safety of the restrained and the others.⁽¹¹⁾ Their role starts with the selection of appropriate restraint device to ending with modifying the patient care plan based on an hourly assessment of the patient's response and physical condition.⁽¹²⁾ The nurses' positive attitudes towards physical restraints may influence their practice in general.⁽¹³⁾ A recent study among psychiatric nurses showed ambivalent attitudes towards use of physical restraints among mental health consumers.⁽¹⁴⁾ Further, it was argued that nurses' views and attitudes toward the use of physical restraints may create conflicts with patients' rights, including their autonomy in making decisions for their own care.⁽¹⁵⁾ In a recent cross sectional surveys from India and Australia, most psychiatrists opined that coercion violates the patients' integrity and may harm therapeutic relationships.⁽¹⁶⁾ Since nurses' knowledge and attitude towards physical restraints is closely related to their practice, it is critical to assess nurses' knowledge, attitude and practice towards use of physical restraints to develop

standardized guidelines in mental health care settings. Further, there were no formal studies on this issue from India. Hence, the present study was aimed to assess knowledge, attitude and practice towards use of physical restraints among nurses those working in psychiatric health care settings.

Methods

The present study was carried out among nurses working at a tertiary care center in South India in the month of November 2015. Descriptive design was adopted for the present study. The study sample was selected through convenient sampling technique. The inclusion criteria for this study as follows; a) both gender, b) nurses those were working in different psychiatric units with minimum one year of experience. A total of 140 nurses were eligible to participate in the study. However, six of the participants were absent during data collection, four of them refused to participate and two of the incomplete questionnaires were discarded. Hence, the final sample consisted of 128 nurses with 86% response rate.

Data were collected through self administered questionnaire which comprised of 4 parts: 1- *Socio demographic profile*: included background information of the participants such as age, gender, experience and education; 2- *Knowledge questionnaire about the use of physical restraints*: used to assess nurses knowledge, towards using the Physical Restraints in psychiatric patients⁽¹⁷⁾ and consisted of 11 items with closed ended responses such as 'true' or false. The correct responses were given a score of '1' and incorrect responses a score of '0'. Thus a score of 11 represented 100% correct responses; 3- *Attitudes toward the use of physical restraints*: This was a four point Likert scale with 12 items (4 negatively worded items) used to assess nurses' attitudes towards use of physical restraints.⁽¹⁷⁾ The responses were scored with a four for "strongly agree", to a one for "strongly disagree". Thus high scores reflected positive attitudes and low scores reflected negative

attitudes. Items 4, 9, 10 and 12 were negative and their scores were reversed. Thus, high scores with cutoff point >24 reflected positive attitudes and low scores with cutoff ≤ 24 reflected negative attitudes (potential range: 12–48); 4- *Nurses' practice towards use of physical restraints*: This part addresses nurses' practice about use of physical restraints in mental health care settings. The participants were asked to respond to a three-point Likert scale ranged from 'always' ⁽³⁾ to never⁽¹⁾. Item no 10 was a negative item and the responses need to be reversed. Thus a score of 14 indicates the most undesirable practice and a score of 42 the best practice in terms of restraint use. The reliability coefficients for the knowledge, attitudes and practice scales were 0.65, 0.61 and 0.94 respectively.⁽¹⁷⁾

The questionnaire was piloted among 15 of participants and found that the study was feasible. Hence no necessary modifications were made. The lead researcher approached the nurses various departments of psychiatry units and the objective and procedure of the study were explained to potential participants. After obtaining permission from the administrators of the hospital, researchers met the participants individually at their workplaces and explained the purpose of the research data that would be collected from the nurses who were willing to participate in the study. Participants were reminded that they could withdraw from the study at any time and confidentiality of the data was assured. After obtaining written informed consent, the self-reported questionnaires were distributed to participants. For the Statistical analysis, negatively worded items were reverse coded and the data were analyzed using 'R' statistical software. The results were presented in narratives and tables. The t-test was used to determine whether significant differences existed between the participants regarding mean knowledge, attitude and practice scores. Statistical significance was assumed at $p < 0.05$.

Results

This study represents 79% of the nurses working at a Psychiatric hospital in South India. Among 128

respondents, a majority were females (77.3%). The mean age of the nurses was 36.9 ± 8.66 (range 24-59 yrs, $M \pm SD$) and 50% of them were

below 35 years of age. More number of them were experienced more than 11yrs (55.5%) and Diploma (GNM) holders (57.8%).

Table 1. Characteristics of the participants (n=128)

Variable	Group	Frequency	Percentage
Age	<25	3	2.3
	26-35	64	50
	36-45	38	29.7
	46-55	19	14.8
	>55	4	3.1
Gender	Male	29	22.7
	Female	99	77.3
Experience	≤10yrs	57	44.5
	≥11yrs	71	55.5
Education	GNM	74	57.8
	BSN	43	33.6
	Others	11	8.6

With regard to participants responses to knowledge about use of restraints, a majority of them agreed that 'physical restraints are safety garments designed to prevent injury' (78.1%), 'restraints should be released every 2 hours' (79.7%), and 'when patient is restrained skin breakdown may increase' (79.7%). While 39.1% agreed, a majority (60.9%) disapproved to the item 'restraints should be used when one cannot watch the patient closely'.

Most of the participants answered correctly to the items related to the rights of the patients i.e. 'patients are allowed to refuse to be placed in a restraint' (61.7%) and 'physical restraints require a consent form from the family member' (74.2%). However, the mean score on knowledge scale was above average of 7.2 ± 1.7 (maximum possible= 11), that indicates good knowledge about use of restraints among the participants (Table 2).

Table 2. Participants' knowledge towards use of physical restraints

Item	Agree		Disagree	
	f	%	f	%
1. Physical restraints are safety, garments designed to prevent injury.	100*	78.1	28	21.9
2. Restraints should be used when one cannot watch the patient closely	50	39.1	78*	60.9
3. Patients are allowed to refuse to be placed in a restraint	79*	61.7	49	38.3
4. A physical restraint (safety, vest, garment) requires a consent form from the family member	95*	74.2	32	25
5. A restraint should be released every 2 hours, if the patient is awake	102*	79.7	26	20.3
6 Restraints should be put on snugly so that there is no space between the restraint and the patients skin	42	32.8	86*	67.2
7. When a patient is restrained skin breakdown may increase.	102*	79.7	26	20.3
8. When a patient is restrained in bed, the restraint should not be attached to the side rail	79*	61.7	49	38.3
9. A patient should never be restrained while lying flat in bed because of the danger of choking	78*	60.9	50	39.1
10. Good alternatives to restraints do not exist	40	31.3	88*	68.8
11. Deaths have been linked to the use of vest restraints	32*	25	96	75

(*) Correct responses

Table 3 depicts the participants' attitudes towards use of physical restraints among psychiatric patients. A majority of the participants hold positive attitudes towards rights of the participants as they agreed with 1 and 2 items i.e. 'I feel that family members have the right to refuse the use of restraints' (78.9%) and 'If I were the patient, I feel I should have the right to refuse or resist when restraints are placed on me' (75.8%). Unfortunately, nearly half of the participants ($n=61$, 47.6%) disapproved that 'patient suffers a loss of dignity when placed in restraints'. While 40.6% of the participants felt guilty, majority of them disagreed for placing a patient in restraints.

Similarly, nearly three fourth of the participants did not feel embarrassed when the family enters the room of a patient who is restrained. However, more than 56% of the participants felt bad if the patients got upset or become more disoriented after the restraints have been applied. It was disappointing to note that 45.3% of the participants opined that 'applying restraints assures legal protection to his/her self and his/her hospital'. The mean score on this section was 30.8 ± 3.3 (ranged from 24 to 45) of a maximum possible of 48 and suggest favorable attitudes among the participants towards use of physical restraints among persons with mental illness.

Table 3. Participants' responses to Attitude towards physical restraints questionnaire

Item	A/SA		DA/SDA	
	f	%	f	%
1. I feel that family members have the right to refuse the use of restraints	101	78.9	27	21.1
2. If I were the patient, I feel I should have the right to refuse or resist when restraints are placed on me	97	75.8	31	24.2
3. I feel guilty placing a patient in restraints	52	40.6	76	59.4
4. I feel that the main reason that restraints are used is that the hospital is short staffed	27	21.1	101	78.9
5. I feel embarrassed when the family enters the room of a patient who is restrained.	33	25.8	95	74.2
6. It makes me feel bad if the patients get more upset after restraints are applied.	73	57.1	55	43
7. It makes me feel bad when patients become more disoriented after the restraints have been applied.	72	56.2	56	43.8
8. A patient suffers a loss of dignity when placed in restraints	67	52.4	61	47.6
9. It is important to apply restraints to assure legal Protection for myself and my hospital	58	45.3	70	54.7
10. I feel that placing a patient in restraints can decrease nursing care time	29	22.6	99	77.4
11. I believe that restraints increase the risk of strangulation	51	39.9	77	60.1
12. I believe that restraints decrease the number of patients to fall	82	64.1	46	35.9

Table 4 addresses the nurses practices with regard to use of physical restraints in psychiatric patients. A majority (93%) of the nurses agreed that they always/sometimes try alternative nursing measures before restraining the patient. More number of the participants stated that they sometimes than always restraints the patients only with physicians' order and they suggest the doctor when they feel that patient doesn't need to be restrained. Majority of the participants expressed

that they check the restraint every two hours to make sure they are in position and inspect the skin of the patient for abrasions or skin tears. Unfortunately, 79.7% opined that more patients are restrained when there is short of staff. Similarly, more number of the nurses felt sometimes than always work together to discover ways to control patients' behavior than use of restraints and assesses if the restraint to be removed. The mean score of nurses' practice related to use of

restraints in psychiatric patients was 31.2 ± 6.2 (range 15-42) of maximum possible of 42. These

findings indicate good practice related to use of physical restraints among nurses.

Table 4. Nurses' practice towards using of physical restraints in Psychiatric patients

Items	Always		Sometimes		Never	
	f	%	f	%	f	%
1. I try alternative nursing measures before restraining the patient	58	45.3	61	47.7	9	7
2. When I restrain a patient, I make this decision only with a physician's order	50	39.1	61	47.7	17	13.3
3. When I feel that the patient does not need to be restrained. I make this suggestion to the doctor	41	32	66	51.6	21	16.4
4. I answer the call for the patient who is restrained as soon as possible	57	44.5	56	43.8	15	11.7
5. I check the restraint every two hours to make sure they are in the proper position	71	55.5	44	34.4	13	10.2
6. I inspect the skin of the patient for abrasions or skin tears if I bath the Patient who is restrained	66	51.6	45	35.2	17	13.3
7. I tell family members why the patient is being restrained	63	49.2	45	35.2	20	15.6
8. I explain to the patient why the restraint is being applied	74	57.8	42	32.8	12	9.4
9. I tell the patient when the restraints will be removed	54	42.2	52	40.6	22	17.2
10. More patients are restrained when we are short of staff than when we are fully staffed	102	79.7	18	14.1	8	6.3
11. In our hospital staff members work together to discover ways to control patients behavior other than use of physical restraints	33	25.8	74	57.8	21	16.4
12. I frequently assess if the restraint should be removed	34	26.6	77	60.2	17	13.3
13. When physical restraint is applied. I record on the nursing charts the type of restraint used, the reason for adopting it, the time when the application commences, and the related nursing care required	62	48.4	54	42.2	12	9.4
14. I frequently evaluate and record the effect of physical restraint when it is applied to a patient	56	43.8	57	44.5	15	11.7

On correlation analysis, it was revealed statistically significant association between gender with knowledge ($r=0.30$, $p<0.001$) and attitude ($r=0.21$, $p<0.05$) of the participants. Females had better knowledge, attitudes than males towards use of physical restraints. However there was no significant difference between gender and practice of the nurses with related use of physical restraints. Nurses those had more than ten years of experience found to be favorable attitudes than nurses with less experience ($r=0.17$, $p<0.05$). On one way analysis, nurses with higher education differed significantly on practice score than nurses with basic education in nursing ($p<0.05$). A significant correlation was found between nurses'

total scores of knowledge ($p<0.01$), attitude ($p<0.001$) with their practice of using physical restraints.

Discussion

This was a first cross sectional survey that assessed nurses' knowledge, attitude and practice towards physical restraints in psychiatric patients. The findings reveled that nurses had better knowledge and favorable attitudes as reflected in good practice of using physical restraints in psychiatric patients. There were significant association between gender with knowledge and attitude of the participants

towards use of the physical restraints. However, a positive significant correlation was found between knowledge, attitude and practice scores of the participants.

In the present study, majority (79%) of the nurses agreed that skin breakdown may increase when patients are restrained. These findings were in line with earlier studies that found 'dehydration, choking, circulatory and skin problems, loss of muscle strength and mobility, pressure sores, incontinence and injury from associated physical/mechanical restraint, injury from other patients, increased psychological distress and, in rare circumstances, death were the adverse events associated with use of physical restraints among patients.⁽¹⁸⁾ Although, nurses in the current study, endorsed good knowledge and positive attitudes towards rights of patients, nearly one quarter of nurses disagreed that 'family members have the right to refuse the use of Physical restraints'. These findings were in consistent with earlier studies conducted among Psychiatric nurses⁽¹⁴⁾ and ICU nurses.⁽¹⁹⁾ These findings suggest need to improve nurses' knowledge on patients' rights and ethical issues related to use of physical restraints.⁽¹⁴⁾

In supporting the previous studies,⁽¹⁹⁾ nearly half the participants in this study disapproved with the statement "A patient suffers a loss of dignity when placed in restraints". It is unfortunate to note the nurses' unawareness about the feelings of the psychiatric patients when they were restrained and this negative attitude also increases the use of physical restraints. Published evidence report that patients as a result of being restrained reported that they felt angry, helpless, sad, and powerless, punished, embarrassed, and that their right to autonomy and privacy has been violated, in addition to a feeling of loss of self worth, degradation, demoralization and humiliation while they are restrained.⁽²⁰⁾ Further, recent studies in India report that most of the psychiatrists' had negative attitudes towards coercion among psychiatric patients as it violates the integrity of the patients.⁽²¹⁾ Earlier research report array of emotional reactions among nurses while restraining the patients such as anxiety, anger, feeling bored

or distressed, crying, inadequacy, hopelessness, frustration, fear, guilt, dissatisfaction, isolation, being overwhelmed, feeling drained, vengeance and repugnance.⁽²²⁾ In line with these findings, the present study also showed guilty and disagreement among the participants to place patients on restraints.

In the current study, nearly three fourths of the participants did not feel embarrassed when the family enters the room of a patient who is restrained and support the earlier evidence.⁽¹⁹⁾ Further nearly 43% of the participants did not feel bad if the patients got upset or become more disoriented after the restraints have been applied. Earlier studies also indicate no emotional reaction among nurses.⁽²²⁾ These findings could be attributed to the fact the practice of physical restraints had become so ritualized that it does not provoke any reaction among nurses.⁽²³⁾

A vast majority of the nurses in this study agreed that they would try alternative nursing measures before restraining the patients. These results were in concordance with previous research among nurses that found several alternative methods that could be used before applying Physical Restraints such as providing companionship and supervision, offering physical and diversional activities, playing soft music, manipulating environments.⁽¹⁴⁾ Interestingly, merely 9.4% of the nurses in this study agreed that they never record on the nursing charts about the type and the reason for applying restraints. These findings were inconsistent with earlier studies that indicate less than half of the nurses do not record data about PR use in patient's chart.⁽¹⁴⁾ However, majority (79.7%) of the nurses felt that more patients are restrained when they are short of staff. These findings were inconsistent with other studies⁽¹⁹⁾ and suggest the need for creating awareness among nurses regarding the reasons for applying the restraints.

The result of the current study is consistent with earlier studies⁽¹⁹⁾ found that more than 50% of the patients were restrained physically and removed without written medical orders. However, this

number is lesser than the present study where 39% of the nurses restrain the patients with doctors' order. In line with previous findings,⁽¹⁴⁾ in the present study nurses those had more than ten years of experience found to be favorable attitudes than nurses with less experience. These findings suggest that better experience improves nurses' decision making about proper use of physical restraints. In the current study, female nurses with higher education had better knowledge and positive attitudes towards use of physical restraints and these findings were in support of the earlier evidence.⁽²⁴⁾ In addition, nurses' knowledge, attitude and practice scores towards use of physical restraints were positively correlated. These findings were in agreement with other studies.⁽¹⁹⁾

The present study has certain limitations such as cross sectional survey, small and convenient sampling and the data was collected from a single setting. Hence, it is difficult to generalise

the findings. However, the study contributes to the literature by assessing nurses' knowledge, attitudes and practices towards use of physical restraints in mental health care settings which is a common practice in India. The findings may be helpful to nursing administrators to develop standardized protocols about using of physical restraints among psychiatric patients.

Conclusion

This study revealed good knowledge, positive attitudes and good practices among nurses about using physical restraints in mental health services. Further, in line with earlier findings,⁽⁶⁾ our study also showed that nurses with higher education had good practice related to use of physical restraints. These findings indicate the need for continuing education about use of physical restraints among nurses may help to protect human rights of patients with mental illness.

References

1. Martin B. Restraint use in acute and critical care settings: changing practice. *AACN clinical issues*. 2002; 13(2):294-306.
2. Gelkopf M, Roffe Z, Behrbalk P, Melamed Y, Werbluff N, Bleich A. Attitudes, opinions, behaviors, and emotions of the nursing staff toward patient restraint. *Issues Ment. Health Nurs*. 2009; 30(12):758-63.
3. Moosa MYH, Jeenah FY. The Use of Restraints in Psychiatric Patients. *S. Afr. J. Psychiatr*. 2009; 15(3):72-5.
4. Mohr WK. Restraints and the code of ethics: An uneasy fit. *Arch. Psychiatr. Nurs*. 2010; 24(1):3-14.
5. Steinert T, Lepping P, Bernhardsgrütter R, Conca A, Hatling T, Janssen W, et al. Incidence of seclusion and restraint in psychiatric hospitals: a literature review and survey of international trends. *Soc. Psychiatry Psychiatr. Epidemiol*. 2010; 45(9):889-97.
6. Khandelwal S, Deb K, Krishnan V. Restraint and seclusion in India. *Indian J. Soc. Psychiatry*. 2015; 31(2):141-7.
7. Ghanashyam B, Nagarathinam S. India is failing the mentally ill as abuses continue. *Lancet*. 2010 13; 376(9753):1633-4.
8. Antony JT. The mental health care bill 2013: A disaster in the offing? *Indian J. Psychiatry*. 2014; 56(1):3-7.
9. Lai CK, Chow SK, Suen LK, Wong IY. The effect of a restraint reduction program on physical restraint rates in rehabilitation settings in Hong Kong. *Rehabil. Res. Pract*. 2011; 2011:284604.
10. Springer G. When and how to use restraints. *Am. Nurse Today*. 2015; 10:1. Available from: <https://www.americannursetoday.com/use-restraints/>
11. Lai CKY. Nurses using physical restraints: Are the accused also the victims? – A study using focus group interviews. *BMC Nurs*. 2007; 6:5.
12. De Jonghe B, Constantin J-M, Chanques G, Capdevila X, Lefrant J-Y, Outin H, et al. Physical restraint in mechanically ventilated ICU patients: a survey of French practice. *Intensive Care Med*. 2013; 39(1):31-7.
13. Wynn R. Staff's attitudes to the use of restraint and seclusion in a Norwegian university psychiatric hospital. *Nord J. Psychiatry*. 2003; 57(6):453-9.
14. Mahmoud AS. Psychiatric Nurses' Attitude and Practice toward Physical Restraint. *Arch. Psychiatr. Nurs*. 2017; 31(1):2-7.
15. Al-Khaled T, Zahran E, El-Soussi A. Nurses' related factors influencing the use of physical restraint in critical care units. *J. Am. Sci*. 2011; 7(8):13-22.
16. Kinner SA, Harvey C, Hamilton B, Brophy L, Roper C, McSherry B, et al. Attitudes towards seclusion and restraint in mental health settings: findings from a large, community-based survey of consumers, carers and mental health professionals. *Epidemiol. Psychiatr. Sci*. 2016; 12:1-10.
17. Suen K. Knowledge, attitude and practice of nursing home staff towards physical restraints in Hong Kong nursing homes. *Asian J. Nurs. Stud*. 1999; 5(2):73-86.
18. Barnett R, Stirling C, Pandyan AD. A review of the scientific literature related to the adverse impact of physical restraint: gaining a clearer understanding of the physiological factors involved in cases of restraint-related death. *Med. Sci. Law*. 2012; 52(3):137-42.
19. Azab S, Abu Negm L. Use of physical restraint in intensive care units (ICUs) at Ain Shams University hospitals, Cairo. *J. Am. Sci*. 2013; 9(4):230-40.
20. Elgamal M. Patients and staff attitudes toward physical restraint. *Current Psychiatry*. 2006; 13(3):474.
21. Gowda GS, Noorthoorn EO, Kumar CN, Nanjegowda RB, Math SB. Clinical correlates and predictors of perceived coercion among psychiatric inpatients: A prospective pilot study. *Asian J. Psychiatry*. 2016; 22:34-40.
22. Kamel NMF, Maximus MHF, Gaafar MM. Reactions of Patients and Psychiatric Hospital Staff About Physical Restraint. *Alexandria Sci. Nurs. J*. 2007; 6(2):1-22.
23. Sequeira H, Halstead S. The psychological effects on nursing staff of administering physical restraint in a secure psychiatric hospital: 'When I go home, it's then that I think about it'. *Br. J. Forensic Pract*. 2004; 6(1):3-15.
24. Chien W-T, Lee IY. Psychiatric nurses' knowledge and attitudes toward the use of physical restraint on older patients in psychiatric wards. *Int. J. Mult. Res. Approaches*. 2007; 1(1):52-71.