



Suma Psicológica

ISSN: 0121-4381

ISSN: 2145-9797

Fundación Universitaria Konrad Lorenz

Seco Ferreira, Diogo Conque; Oliveira, Walter Lisboa; Costa
Delabrida, Zenith Nara; Faro, André; Cerqueira-Santos, Elder
Intolerance of uncertainty and mental health in Brazil during the Covid-19 pandemic
Suma Psicológica, vol. 27, no. 1, 2020, January-June, pp. 62-69
Fundación Universitaria Konrad Lorenz

DOI: <https://doi.org/10.14349/sumapsi.2020.v27.n1.8>

Available in: <https://www.redalyc.org/articulo.oa?id=134265182008>

- ▶ [How to cite](#)
- ▶ [Complete issue](#)
- ▶ [More information about this article](#)
- ▶ [Journal's webpage in redalyc.org](#)

The logo for Redalyc.org, featuring the text 'redalyc.org' in a stylized font with a red dot above the 'y'.

Scientific Information System Redalyc


Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative



SUMA PSICOLÓGICA

<http://sumapsicologica.konradlorenz.edu.co>



Intolerance of uncertainty and mental health in Brazil during the Covid-19 pandemic

Diogo Conque Seco Ferreira*, Walter Lisboa Oliveira, Zenith Nara Costa Delabrida, André Faro, Elder Cerqueira-Santos

Universidade Federal de Sergipe, Brazil

Received 30 June 2020; accepted 7 July 2020

KEYWORDS

Intolerance of uncertainty, depression, anxiety, stress, COVID-19, pandemic

Abstract The first Brazilian diagnosed with COVID-19 was identified on February 25th, resulting in a series of governmental actions to prepare the population for the effects of the pandemic. Nevertheless, geographic and socioeconomic characteristics of the country, and the strategies adopted may have contributed to generating a widespread feeling of uncertainty in the population; uncertainty about the disease, how to prevent it, its severity, and its impact on political and economic issues. This study was designed to investigate the relationship between intolerance of uncertainty and common mental health disorders. Participants were contacted via social media messages and encouraged to fill out an online questionnaire with socioeconomic questions, a short measure of intolerance of uncertainty (IUS-12), and a measure of mental health indicator (DASS-21). With a sample of 924 participants from Sergipe, Brazil, three binomial logistic regressions were performed, one for each DASS-21 outcome, categorized by a median. The women in the sample showed a higher probability of having stress, anxiety and depression scores above the median. Over the entire sample, both subscales of IUS-12 were significantly and positively related to all three DASS-21 subscales. These findings are discussed in the context of mental health during a pandemic, and future directions for research are also presented.

© 2020 Fundación Universitaria Konrad Lorenz. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Intolerância a incerteza, depressão, ansiedade, estresse, COVID-19, pandemia

Intolerância a incerteza e saúde mental no Brasil durante a pandemia de Covid-19

Resumo O primeiro brasileiro diagnosticado com COVID-19 foi identificado dia 25 de fevereiro, resultando em uma série de ações governamentais para preparar a população para os efeitos da pandemia. No entanto, características geográficas e socioeconômicas do país, e as estratégias adotadas, podem ter contribuído para a criação de uma sensação generalizada de incerteza na população; incerteza a respeito da doença, como preveni-la, sua gravidade, e seu impacto em questões políticas e econômicas. Este estudo foi elaborado para investigar a relação entre Intolerância à Incerteza e transtornos mentais comuns. Os participantes foram contactados por mensagens em redes sociais e convidados a preencher um questionário online

* Autor para correspondencia.

Correo electrónico: diogoseco@ufs.br

com questões socioeconômicas, um instrumento breve de Intolerância a Incerteza (IUS-12) e um indicador de saúde mental (DASS-12). Com uma amostra de 924 participantes de Sergipe, Brasil, três regressões logísticas binomiais foram realizadas, uma para cada desfecho do DASS-21, categorizados pela mediana. As mulheres da amostra apresentaram maior probabilidade de escores estresse, ansiedade e depressão acima da mediana. Em toda a amostra, ambas subescalas da IUS-12 foram significativa e positivamente relacionadas às três subescalas da DASS-12. Estes achados são discutidos no contexto de saúde mental durante uma pandemia, e futuras direções de pesquisa também são apresentadas.

© 2020 Fundación Universitaria Konrad Lorenz. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/bync-nd/4.0/>).

Despite the first cases of COVID-19 having been reported in December 2019, in Wuhan, China, it would take two months for the first Brazilian patient to be identified - a man living in São Paulo, on February 25th (Croda et al., 2020). The Brazilian government's response was swift, with both international and local health organizations coordinating efforts to harmonize, plan and organize activities with the stakeholders involved (Croda & Garcia, 2020). Channels for direct contact with the population were established early on, through instant messaging apps and cell phone apps, to provide reliable information and prevent the propagation of fake news and/or distorted facts (Oliveira, Duarte, França, & Garcia, 2020). Guidance for prevention behaviours were also provided, which included recommendations for social distancing, hand washing procedures and, as of April 22nd, the use of cloth face masks in public venues (Oliveira et al., 2020)

In sharp contrast to the rapid governmental response, public opinion has been confused due to different measures and proposals from different official agencies (State-level vs. Central government) and the inevitable rise of fake news, along with the propagation of distorted facts. The Brazilian Ministry of Health itself acknowledges the necessity for heterogeneous strategies across States and Regions, considering the continental size of the country (Oliveira et al., 2020) and this may have contributed to the uncertainty faced by the population regarding the procedures to be adopted. If there is something certain about the first trimester of the year 2020, it is that people are uncertain about what is going to happen in the immediate and distant future. In commenting on the situation in Brazil, Gomes Mello and Rodrigues (2020) state that there is pervasive uncertainty about the disease, how to tackle it, and its economic, commercial, and political toll.

Although seldom confused in laymen terms (De Groot & Thurik, 2018), uncertainty and risk appear to describe two different scenarios. If you are capable of describing, not only the outcomes of a situation, but the distribution of probabilities of ex-ante outcomes, you are dealing with a risk scenario; on the other hand, if you only know the outcomes, but have little information regarding their probability distribution, then you are facing an uncertain scenario (see Knight, 1921, for the canonical definition). While betting on a single outcome from a die toss exemplifies a risky situation, betting on a horse in a horse race seems to involve an even greater deal of uncertainty. There are several findings from behavioural economics (Baillon, Cabantous, & Wakker, 2012) to psychology (Slovic, 2010) that corroborate the difference between these phenomena and the

extent to which they are related to other psychological events (such as Depression, Anxiety and Stress).

Those findings also point to the fact that sensitivity to uncertain events is heterogeneously distributed in a population and there has been considerable efforts to devise and sharpen measurement tools to detect it (Gosselin et al., 2008; Greco & Roger, 2001; Roma & Hope, 2017). One aspect of our reactions to uncertainty frequently studied has been our intolerance to it. Intolerance of uncertainty (IU) has been defined as "the tendency of an individual to consider the possibility of a negative event occurring as unacceptable, irrespective of the probability of occurrence" (Carleton, Norton & Asmundson, 2007, p. 105). Although the concept was initially proposed to be associated with generalized anxiety disorder (GAD) (Dugas, Gosselin & Ladouceur, 2001; Dugas, Gagnon, Ladouceur & Freeston, 1998), it has recently been discovered as a contributing factor to a variety of mental disorders, from obsessive compulsive disorder (OCD) (Holaway, Heimberg & Coles, 2006), to social anxiety (Carleton, Collimore & Asmundson, 2010), depression (McEvoy & Mahoney, 2011) and panic disorder (Tolin, Abramowitz, Brigidi, & Foa, 2003). McEvoy and Erceg-Hurn (2016) go as far as to consider it a trans-diagnostic and trans-therapy factor, which means that treatment for several distinct mental disorders, from different theoretical backgrounds, should benefit from tackling intolerance of uncertainty. In a study following standardized treatment of 256 participants diagnosed with either GAD, social anxiety depression or depression, the authors found consistent evidence that changes in IU were closely related to changes in different symptoms related to the disorders.

In a scenario involving viral illness in a pandemic situation, Taha Matheson, Cronin, and Anisman, (2014) addressed the sequential interplay between IU, coping strategies, and anxiety based on 1,027 self-reported questionnaires during the 2009 H1N1 pandemic. The authors warn that, should the government officials fail to prevent the worry and hysteria related to viral contraction, the uncertain estimates related to the threat may increase perception of uncertainty and thus increase anxiety. Elledge, Brand, Regens and Boatright, (2008) investigated the public understanding of the Avian Flu, a threatening viral illness, around 2006, in a North American city, and found little evidence of solid knowledge in the population researched. Results indicate that participants were ill-informed about the severity of the illness, about the United States' coping potential and about the proximity of a vaccine or effective treatment. Although authors consider it safe to assume that participants were exposed to information about the avian flu, it

seems clear that “accurate communication of risk [and uncertainty] was faulty”. Satici, Saricali, Satici and Griffiths (2020) have studied a large sample of Turkish nationals and established an inverse relation between IU and Mental Wellbeing, measured by the Warwick-Edinburgh Mental Well-Being Scale (WEMWBS; Tennant et al. 2007). Rumination and a Fear of COVID-19 (measured by the FCV-19S Scale; Ahorsu et al. (2020) partially mediated the relation, elucidating possible psychological mechanisms responsible for the decrease of mental wellbeing in high IU participants facing the pandemic. Bakioglu, Korkmaz, and Ercan (2020) found similar results in another Turkish sample, presenting evidence for the relation of Fear of Covid-19, IU, mental health indicators (depression, anxiety and stress) and positivity. The authors’ proposed model indicates the role of IU and mental health indicators as mediators of the relation between fear of COVID-19 and positivity, again designating a relevant role played by IU.

As of May 25th, 2020, the COVID-19 pandemic has escalated into a major world-wide health crisis, with 5,304,772 confirmed cases and 342,029 deaths (BRASIL, 2018). In Brazil, there have been 22,013 deaths and 347,398 confirmed cases, even though lack of effective testing makes this estimate less reliable. Lipsitch, Phil, Swerdlow and Finelli, (2020) question the use of confirmed cases as good indicators to model the current epidemic’s dynamics, partially because of the relevant role of asymptomatic infected people in spreading the virus, so the correct number of infections will remain uncertain for some time. An additional problem is the lack of criteria for differential clinical diagnosis, since clinical features of COVID-19 are non-specific and may be seen in several different respiratory illnesses (Phua et al., 2020). This situation may characterize the COVID-19 pandemic as a particularly challenging situation to cope with. Uncertainty about the disease’s transmission and about the effectiveness of prevention procedures may be related to severe psychological distress, including anxiety, depression, and acute stress disorder (Gardner & Moalef, 2015; Jiang et al., 2020; Tavares Lima et al., 2020). These factors warrant research on mental health assessment, support, treatment, and services related to COVID-19 (Xiang et al., 2020).

The current study investigates how differential responses to the uncertainty of events (measured by IUS-12) may be related to depression, anxiety and stress symptomatology (measured by DASS-21) during the early days of the COVID-19 pandemic in Sergipe, a north eastern state in Brazil. The research presented here is a part of the first of a three-wave monitoring project aimed at describing mental health indicators during the initial, intermediate, and final moments of the COVID-19 pandemic in Brazil.

Method

Participants

The sample comprised of 924 participants, from both genders, from 18 to 72 years old [mean (*M*) = 36.8; standard deviation (*SD*) = 11.70], residing in Sergipe, a state in the north eastern region of Brazil. The majority reported being in the capital (69.7%; *n* = 613) during the survey, and the others in different cities of the state (30.3%, *n* = 267). Most of the sample (95.5%; *n* = 882) reported having complete or

incomplete higher education and the rest (4.5%; *n* = 42) said they had complete or incomplete high school education. Regarding skin colour, people who declared themselves to be Brazilian Asians, indigenous or other, were excluded from the sample, due to the small amount. More than half declared themselves to be “pardo” - Brazilian mixed ethnicity (54.3%, *n* = 502), followed by white (34.2%, *n* = 316) and black (11.5%; *n* = 106). Most (61.7%, *n* = 570) said they had a fixed income and the rest said they were unemployed or did not have a fixed income (38.3%; *n* = 354).

Instruments

Sociodemographic Questionnaire. Used to collect information such as self-identified gender (male, female), age (in years) and education (complete or incomplete high school, and complete or incomplete higher education).

Reduced Intolerance Uncertainty Scale (IUS-12) (Carleton et al., 2007; Kretzmann, 2018). Composed of 12 items on the scale of the type of agreement (01 to 07), distributed into two dimensions: (1) Prospective Anxiety and (2) Inhibitory Anxiety. Cronbach’s alpha was .79; .86 and .88 for Prospective and Inhibitory Anxiety subscales and total scores, respectively.

Depression, Anxiety and Stress Scale - Short Form (DASS-21) (Apóstolo, Mendes & Azeredo, 2006; Vignola & Tucci, 2014). Likert scale of occurrence of symptomatic changes (4 points), distributed into 21 items, which generates independent scores on anxiety symptoms, depression and stress over the last week. Cronbach’s data in this research were .83, .88, .89 and .94 for the factors of anxiety, depression, stress and the whole scale, respectively.

Procedures and Ethical Aspects

A survey was conducted from April 3 to April 16, 2020, which corresponded to a period of quarantine and social isolation imposed by the government of Sergipe (Decreto 40560 de 16 de Março, 2020). The sampling was made by convenience and snowballing methods. Participants received a public invitation on social media to participate in an online survey regarding mental health in the confinement period. The first research screen presented a free and informed Consent Form online. The project was approved by the National Council for Ethics in Research [Conselho Nacional de Ética em Pesquisa (CONEP); register n. 3.954.144].

Data Analysis

Data were analysed using SPSS (v. 24) for descriptive statistics (absolute and relative frequencies, media, media and standard deviations), and inferential statistics (binomial logistic regressions and a multinomial). For binomial logistic regressions, the dependent variables were scores for anxiety, depression and stress, all dichotomized by the median, considering the non-parametric distribution of the total sample. In the binomial models, the factors were Prospective and Inhibitory Anxiety, gender (male or female) and age (divided into four groups: up to 30 years old, between 30 to 40 years old, between 40 to 50 years old, and more than 50 years old).

All regressions were performed through the Backward LR methods (binomial logistic regression), which do not require the selection of variables before the final modelling (Field, 2009). For the evaluation of binomial models, the following indicators were observed: Omnibus test (expected to be statistically significant), Nagelkerke's R^2 (expected the higher the better, corresponding to the explained variance of the final model), Hosmer and Lemeshow test (expected not to be statistically significant) and correct predictive capacity of the model (expected to be around 70%). The multicollinearity assessment was carried out for all models and no problems were found in the composition or final solution of the models. It is worth noting that all Odds Ratio (OR) values below 1 were converted by the formula $1 / OR$ to standardize the description of the findings. The p-value adopted was less than .05 for all steps of binomial regressions.

Results

Sample Profile

The sample's mean age was 36.8 years old ($SD = 11.70$). After categorization, the proportion for each age group was 35.3% ($n = 326$) up to 30 years of age, 26.7% ($n = 247$) between 30 and 40 years of age, 23.8% ($n = 220$) between 40 and 50 years old and 14.2% ($n = 131$) over 50 years old. Most of the participants were female (79.0%; $n = 730$).

Due to the non-normal distribution of anxiety, depression and stress scores in the present sample, its median was obtained as the most reliable indicator of the sample midpoint. The median was 2, 4 and 6 points for anxiety, depression and stress, respectively. After categorizing the outcomes according to the midpoint, 48.8% ($n = 451$) of the participants were above the median of the anxiety score, 41.9% ($n = 387$) above the median depression, and 44.8% ($n = 414$) above the median in the stress variable. On IUS-12, the factors Prospective Anxiety and Inhibitory Anxiety averaged 21.3 ($SD = 6.31$) and 13.0 ($SD = 5.63$) points, respectively. When classified into three strata of scores, the distribution in the Prospective Anxiety factor was as follows: low (up to 18 points) with 34.5% ($n = 319$), moderate (between 18 and 24 points) with 32, 6% ($n = 301$) and high (greater than 24 points) with 33.1% ($n = 306$). For the Inhibitory Anxiety factor, the low group (up to 9 points) was 33.4% ($n = 309$), moderate (between 9 and 15 points) with 33.4% ($n = 309$) and high (greater than 15 points) with 33.1% ($n = 306$) individuals.

Logistic regressions

Three binomial logistic regressions were performed, one for each DASS-21 outcome, categorized by a median. Table 1 shows the results of each model, as well as their adjustment indicators. In stress outcome, all independent variables were included in the final model. The female gender had three times more chances to be in the group above the stress median ($OR = 2.7$) and the age showed a relatively linear relationship. People up to 30 years old had a higher presence among those above the median of anxiety compared to the other age groups, namely: almost twice as many chances than those between 30 and 40 years old ($OR = 0.5$ and $1/OR = 1.9$), two and a half times more chances

than those between 40 and 50 years old ($OR = 0.4$ and $1/OR = 2.5$) and also twice as many chances than those over 50 years old ($OR = 0.5$ and $1/OR = 2.0$). On IUS-12, Prospective Anxiety factor, individuals in the high group (greater than 24 points) exhibited chances about twice as high ($OR = 1.9$) as those in the low group (up to 18 points). There was no significance for the moderate group (greater than 18 and up to 24 points) in relation to the low BP group. In the factor Inhibitory Uncertainty, the participants classified as moderate (between 9 and 15 points) had twice as many chances to score higher on the stress scale ($OR = 2.1$). The high group (more than 15 points) had almost four times more chances of being above the stress median ($OR = 3.9$).

On the anxiety scale, it was seen that the final model was composed by the variables gender, age and Inhibitory Anxiety, a subscale of the Intolerance of Uncertainty Scale. The female model showed higher chances to be in the group above the anxiety median compared to the male model ($OR = 2.4$). Regarding age, the proportion was similar to that found for stress: the youngest had higher chances. Those up to 30 years old had higher chances for high anxiety scores compared to those between 30 and 40 years old ($OR = 0.6$; $1/OR = 1.7$), those between 40 and 50 years old ($OR = 0.4$; $1/OR = 2.5$) and those over 50 years old ($OR = 0.4$; $1/OR = 2.5$). The factor Inhibitory Anxiety of IUS-12 also exhibited a relative progression of risk. Individuals who scored moderate (between 9 and 15 points) and high (greater than 15 points) had higher chances of composing the stratum of people with scores above the anxiety median ($OR = 2.2$ and $OR = 5.9$, respectively).

Finally, for the depression outcome, the same variables independent of the anxiety model remained in the final solution. The female gender with high odds ($OR = 1.5$) and age followed a similar pattern: younger (up to 30 years old) were more exposed, and in all comparisons the OR values were 2.5, i.e. two and a half times more likely to be in the group with scores above the median depression. The factor Inhibitory Anxiety was also significant, with high Odds ratio. The group with a moderate score showed 3.6 more chances and those with a high score had 9.9 times more chances of being in the group with depression scores above the median.

Discussion

Our results corroborate existing findings in regards to the higher propensities of depression, anxiety and stress in women (Novais, Monteiro, Roque, Correia-Nevez, & Sousa, 2016; Rincón-Cortés, Herman, Lupien, Maguire, & Shansky, 2019; Salk, Hyde, & Abramson, 2017) and young age groups (e.g., Flint et al., 2010; Novais et al., 2016). Stress and Anxiety, in particular, were found to be up to three times more prominent in women, a result in line with those obtained by Bakioğlu, et al., (2020) and Wang et al., (2020). Several factors have been considered as causes for this robust finding, including physiological and socio-economical differences (Kuehner, 2017). A recent rise in the role of female headed households (BRASIL, 2018) may add to the explanation of these results, considering the heavy economic impact of the COVID-19 pandemic. Younger participants in our research also demonstrated a higher chance of manifesting Depression, Anxiety and Stress, consistently displaying 2 to 2.5 times the odds of being over the median in regards

Table 1 Binomial Logistic Regression Indicators for Stress, Anxiety and Depression in DASS-21

Binomial regressions		Odds Ratio (OR)	1/OR	p-value
Stress³				
Age	Up to 30 years old	-	-	-
	Between 30 and 40 years old	0.5	1.9	.001
	Between 40 and 50 years old	0.4	2.5	< .001
	Over 50 years old	0.5	2.0	.002
Prospective Anxiety (IUS-12)	Low (up to 18 points)	-	-	-
	Moderate (between 18 and 24 points)	1.0	-	.756
	High (more than 24 points)	1.9	-	.002
Inhibitory Anxiety (IUS-12)	Low (up to 9 points)	-	-	-
	Moderate (between 9 and 15 points)	2.1	-	< .001
	High (more than 15 points)	3.9	-	< .001
Gender	Male	-	-	-
	Female	2.7	-	< .001
Anxiety¹				
Age	Up to 30 years old	-	-	-
	Between 30 and 40 years old	0.6	1.7	.004
	Between 40 and 50 years old	0.4	2.5	< .001
	Over 50 years old	0.4	2.5	.002
Inhibitory Anxiety (IUS-12)	Low (up to 9 points)	-	-	-
	Moderate (between 9 and 15 points)	2.2	-	< .001
	High (more than 15 points)	5.9	-	< .001
Gender	Male	-	-	-
	Female	2.4	-	< .001
Depression²				
Age	Up to 30 years old	-	-	-
	Between 30 and 40 years old	0.4	2.5	< .001
	Between 40 and 50 years old	0.4	2.5	< .001
	Over 50 years old	0.4	2.5	< .001
Inhibitory Anxiety (IUS-12)	Low (up to 9 points)	-	-	-
	Moderate (between 9 and 15 points)	3.6	-	< .001
	High (more than 15 points)	9.9	-	< .001
Gender	Male	-	-	-
	Female	1.5	-	.031

Notes.

* Variables with no statistical significance in each model were excluded from the table.

1. Stress outcome: Omnibus test = 166.952 ($p < .001$). Hosmer and Lemeshow X^2 test = 4.672 ($p = .792$). Nagelkerke $R^2 = .221$ (22.1%). Percentage of correctly predicted cases = 68%.

2. Anxiety outcome: Omnibus test = 160.686 ($p < .001$). Hosmer and Lemeshow X^2 test = 3,484 ($p = 0.900$). Nagelkerke $R^2 = .213$ (21.3%). Percentage of correctly predicted cases = 69%.

3. Depression outcome: Omnibus test = 205,600 ($p < .001$). Hosmer and Lemeshow X^2 test = 5,513 ($p = .598$). Nagelkerke $R^2 = .268$ (26.8%). Percentage of correctly predicted cases = 72%.

to all of these measures. These results are not spurious and appear to show a trend in recent COVID-19 mental health monitoring studies. Huang and Zhao (2020) have observed a similar trend in a Chinese sample, while Liu et al., (2020) found the same result with an American sample and attributes this to the distress in managing work and school responsibilities, the latter being one of the first areas hit by social isolation and quarantine protocols. Although not directly investigated, it is feasible that underdeveloped coping skills and other psychological adaptive factors (such as Self-Esteem, Orth, & Robins, 2014) in younger participants play a relevant role throughout this public health crisis.

The association between the scores of intolerance of uncertainty and the results of the DASS-21 scale are also evident in our results, and these too corroborate the existing literature (Bottesi, Noventa, Freeston, & Ghisi, 2019). Participants over the 3rd tertile on the Prospective Anxiety subscale of IUS-12 were almost twice as likely to have higher than median scores when it comes to Stress. Greco and Rogers (2003) had similar results, although with a physiological measure of stress and a different instrument to measure intolerance of uncertainty. It is worth noting that the Prospective Anxiety was only significantly associated with Stress. As defined by Hong and Lee (2015, p. 606), Prospective Anxiety “seems to represent a desire for predictability of future events, triggered by anxious apprehension about uncertainty, and prompting engagement in strategies (e.g., seeking more information) to reduce uncertainty” and is closely related to worrying and symptoms of Obsessive-Compulsive Disorder.

The Inhibitory Anxiety subscale was significantly related to all three DASS-21 subscales, and participants in the 3rd tertile on this scale were 4, 6 and 10 times more likely to have higher than median scores of Stress, Anxiety and Depression, respectively. This concept measures paralysis and the compromised functioning in reaction to uncertainty (Berenbaum, Bredemeier & Thompson, 2008), so its extreme impact on depression measures should be expected, but also the expressive effects on Stress and Anxiety can be highlighted in the current results. Hong and Lee (2015) obtained results similar to ours, with Depression and Anxiety being the most impacted by higher scores on the Inhibitory subscale of IUS-12.

In sum, our results describe a scenario in which intolerance of uncertainty plays a relevant role in the mental health of participants during the initial phase of the COVID-19 pandemic. All measures employed were significantly and positively related to uncertainty intolerance, so our research is in line with the evidence suggesting that Intolerance of Uncertainty may be regarded as a transdiagnostic factor (McEvoy & Erceg-Hurn, 2016) related to Depression, Anxiety and Stress, even though we did not investigate any behavioural or psychological mechanism that could be responsible for the observed results. Some features of our study must be considered before extrapolating our findings. First, our participant sample was big ($n > 1000$), but recruited in a non-systematic fashion, through social media apps, rendering it unrepresentative of Sergipe's population. Second, since this is the first wave of a three-part monitoring research project, it remains to be evaluated how possible changes in the public perception of the threats posed by the pandemic will affect the

inhabitants of Sergipe, and how these changes will affect mental health, and behavioural and psychological measures. For future research, it is suggested that the link between mental health measures and specific behavioural indicators (such as propensity to remain socially isolated or to adopt prevention strategies) be directly investigated. Additionally, the impact of the pandemic on the development or the severity of specific anxiety disorders should be investigated (e.g. social phobia, health anxiety, generalized anxiety disorder, and agoraphobia), as well as the role of different coping strategies and positive emotions (e.g. hope) to deal with the daily adversities related to COVID-19. And finally, following the results obtained by Bakioğlu et al., (2020) and Satici et al. (2020), more work investigating possible variables mediating the effects of IU on mental health, such as optimism, worry and hope, should be conducted. This would help to elucidate whether IU plays a role as a predictor, as suggested by Carleton (2016) or as a mediating variable, as proposed by Bakioğlu et al., (2020).

References

- Apóstolo, J. L. A., Mendes, A. C., & Azeredo, Z. A. (2006). Adaptação para a língua portuguesa da Depression, Anxiety and Stress Scale (DASS). *Revista Latino Americana de Enfermagem*, 14(6), 863-871.
- Ahorsu, D. K., Lin, C. Y., Imani, V., Saffari, M., Griffiths, M. D., & Pakpour, A. H. (2020). The Fear of COVID-19 Scale: Development and Initial Validation. *International Journal of Mental Health and Addiction*, 1-9. Advance online publication. <https://doi.org/10.1007/s11469-020-00270-8>
- Baillon, A., Cabantous, L., & Wakker, P. (2012). Aggregating imprecise or conflicting beliefs: An experimental investigation using modern ambiguity theories. *Journal of Risk and Uncertainty*, 44, 115-147. <https://doi.org/10.1007/s11166-012-9140-x>
- Bakioğlu, F., Korkmaz, O., Ercan, H. (2020). Fear of COVID-19 and positivity: Mediating role of intolerance of uncertainty, depression, anxiety, and stress. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00331-y>
- Berenbaum, H., Bredemeier, K., & Thompson, R. J. (2008). Intolerance of uncertainty: Exploring its dimensionality and associations with need for cognitive closure, psychopathology, and personality. *Journal of Anxiety Disorders*, 22(1), 117-125. <https://doi.org/10.1016/j.janxdis.2007.01.004>
- Bottesi, G., Noventa, S., Freeston, M. H., & Ghisi, M. (2019). Seeking certainty about intolerance of uncertainty: Addressing old and new issues through the Intolerance of Uncertainty Scale - revised. *PLOS ONE*, 14(2), e0211929. <https://doi.org/10.1371/journal.pone.0211929>
- BRASIL (2018). *Estatísticas de gênero: Indicadores sociais das mulheres no Brasil*. (Estudos e pesquisa, informação demográfica e socioeconômica, n.38). Brasília: Instituto Brasileiro de Geografia e Estatística. Retrieved from https://biblioteca.ibge.gov.br/visualizacao/livros/liv101551_informativo.pdf in June, 15.
- Carleton, R. N., Collimore, K. C., & Asmundson, G. J. G. (2010). “It’s not just the judgements—It’s that I don’t know”: Intolerance of uncertainty as a predictor of social anxiety. *Journal of Anxiety Disorders*, 24(2), 189-195. <https://doi.org/10.1016/j.janxdis.2009.10.007>
- Carleton, R. N., Norton, M. A. P. J., & Asmundson, G. J. G. (2007). Fearing the unknown: A short version of the Intolerance of Uncertainty Scale. *Journal of Anxiety Disorders*, 21(1), 105-117. <https://doi.org/10.1016/j.janxdis.2006.03.014>
- Carleton, R. N. (2016). Into the unknown: A review and synthesis of contemporary models involving uncertainty. *Journal of Anxiety Disorders*, 39, 30-43. <https://doi.org/10.1016/j.janxdis.2016.02.007>

- Croda, J. H. R., & Garcia, L. P. (2020). Resposta imediata da vigilância em saúde à epidemia da COVID-19. *Epidemiologia e Serviços de Saúde*, 29(1), e2020002. <https://doi.org/10.5123/S1679-49742020000100021>
- Croda, J. O., Wanderson K., Frutuoso, R. L., Mandetta, L. H., Baia-da-Silva, D. C., Brito-Sousa, J. D., Monteiro, W. M., & Lacerda, M. V. G. (2020). COVID-19 in Brazil: advantages of a socialized unified health system and preparation to contain cases. *Revista da Sociedade Brasileira de Medicina Tropical*, 53, e20200167. Epub April 17, 2020. <https://dx.doi.org/10.1590/0037-8682-0167-2020>
- De Groot, K., & Thurik, R. (2018). Disentangling risk and uncertainty: When risk-taking measures are not about risk. *Frontiers in Psychology*, 9(Article 2194). <https://doi.org/10.3389/fpsyg.2018.02194>
- Decreto 40560 de 16 de março, Pub. L. No. Diário Oficial do Estado de Sergipe (2020). <https://segrase.se.gov.br/ver-flip/3350/#/e:3350/p:1?find=coronavirus>
- Dugas, M. J., Gosselin, P., & Ladouceur, R. (2001). Intolerance of uncertainty and worry: Investigating specificity in a nonclinical sample. *Cognitive Therapy and Research*, 25(5), 551-558. <https://doi.org/10.1023/A:1005553414688>
- Dugas, M. J., Gagnon, F., Ladouceur, R., & Freeston, M. H. (1998). Generalized anxiety disorder: A preliminary test of a conceptual model. *Behaviour Research and Therapy*, 36(2), 215-226. [https://doi.org/10.1016/S0005-7967\(97\)00070-3](https://doi.org/10.1016/S0005-7967(97)00070-3)
- Elledge, B. L., Brand, M., Regens, J. L., & Boatright, D. T. (2008). Implications of public understanding of avian influenza for fostering effective risk communication. *Health Promotion Practice*, 9(4_suppl), 54S-59S. <https://doi.org/10.1177/1524839908319089>
- Field, A. (2009). *Discovering Statistics Using SPSS*. 3rd Edition, Sage Publications Ltd., London.
- Flint, A. J., Peasley-Miklus, C., Papademetriou, E., Meyers, B. S., Mulsant, B. H., Rothschild, A. J., Whyte, E. M., & STOP-PD Study Group (2010). Effect of age on the frequency of anxiety disorders in major depression with psychotic features. *The American Journal of Geriatric Psychiatry: Official Journal of the American Association for Geriatric Psychiatry*, 18(5), 404-412. <https://doi.org/10.1097/jgp.0b013e3181c294ac>
- Gardner, P. J., & Moallem, P. (2015). Psychological impact on SARS survivors: Critical review of the English language literature. *Canadian Psychology/Psychologie canadienne*, 56(1), 123-135. <http://dx.doi.org/10.1037/a0037973>
- Gomes, L. A. de V., Mello, A. M. de, & Rodrigues, A. C. de A. (2020). *Vencendo as incertezas para superar a pandemia do vírus*. *Jornal Da USP*. <https://jornal.usp.br/?p=312817>
- Gosselin, P., Ladouceur, R., Evers, A., Laverdière, A., Routhier, S., & Tremblay-Picard, M. (2008). Evaluation of intolerance of uncertainty: Development and validation of a new self-report measure. *Journal of Anxiety Disorders*, 22(8), 1427-1439. <https://doi.org/10.1016/j.janxdis.2008.02.005>
- Greco, V., & Roger, D. (2001). Coping with uncertainty: The construction and validation of a new measure. *Personality and Individual Differences*, 31(4), 519-534. [https://doi.org/10.1016/S0191-8869\(00\)00156-2](https://doi.org/10.1016/S0191-8869(00)00156-2)
- Greco, V., & Roger, D. (2003). Uncertainty, stress, and health. *Personality and Individual Differences*, 34(6), 1057-1068. [https://doi.org/10.1016/S0191-8869\(02\)00091-0](https://doi.org/10.1016/S0191-8869(02)00091-0)
- Hong, R. Y., & Lee, S. S. M. (2015). Further clarifying prospective and inhibitory intolerance of uncertainty: Factorial and construct validity of test scores from the Intolerance of Uncertainty Scale. *Psychological Assessment*, 27(2), 605-620. <https://doi.org/10.1037/pas0000074>
- Holaway, R. M., Heimberg, R. G., & Coles, M. E. (2006). A comparison of intolerance of uncertainty in analogue obsessive-compulsive disorder and generalized anxiety disorder. *Journal of Anxiety Disorders*, 20(2), 158-174. <https://doi.org/10.1016/j.janxdis.2005.01.002>
- Huang, Y., & Zhao, N. (2020). Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: A web-based cross-sectional survey. *Psychiatry Research*, 288, 112954. <https://doi.org/10.1016/j.psychres.2020.112954>
- Jiang, F., Deng, L., Liangqing, Z., Yin, C., Chi Wai, C., & Zhengyuan, X. (2020). Review of the clinical characteristics of coronavirus disease 2019 (COVID-19). *Journal of General Internal Medicine*, 35, 1545-1549. <https://doi.org/10.1007/s11606-020-05762-w>
- Knight, F. H. (1921). *Risk, Uncertainty and Profit*. Sentry Press.
- Kretzmann, R. P. (2018). *Intolerance of Uncertainty Scale (IUS-12) para uso no Brasil: Adaptação Transcultural e Propriedades Psicométricas* (Issue 2) [(dissertação de mestrado) Universidade Federal do Rio Grande do Sul].
- Kuehner, C. (2017). Why is depression more common among women than among men? *The Lancet Psychiatry*, 4(2), 146-158. [https://doi.org/10.1016/S2215-0366\(16\)30263-2](https://doi.org/10.1016/S2215-0366(16)30263-2)
- Lipsitch, M., Phil, D., Swerdlow, D. L., & Finelli, L. (2020). Defining the epidemiology of Covid-19 - studies needed. *The New England Journal of Medicine*, 1-3.
- Liu, C. H., Zhang, E., Wong, G. T. F., Hyun, S., & Hahm, H. "Chris." (2020). Factors associated with depression, anxiety, and PTSD symptomatology during the COVID-19 pandemic: Clinical implications for U.S. young adult mental health. *Psychiatry Research*, 290(May). <https://doi.org/10.1016/j.psychres.2020.113172>
- McEvoy, P. M., & Erceg-Hurn, D. M. (2016). The search for universal transdiagnostic and trans-therapy change processes: Evidence for intolerance of uncertainty. *Journal of Anxiety Disorders*, 41, 96-107. <https://doi.org/10.1016/j.janxdis.2016.02.002>
- McEvoy, P. M., & Mahoney, A. E. J. (2011). Achieving certainty about the structure of intolerance of uncertainty in a treatment-seeking sample with anxiety and depression. *Journal of Anxiety Disorders*, 25, 112-122. <https://doi.org/10.1016/j.janxdis.2010.08.010>
- Novais, A., Monteiro, S., Roque, S., Correia-Neves, M., & Sousa, N. (2016). How age, sex and genotype shape the stress response. *Neurobiology of Stress*, 6, 44-56. <https://doi.org/10.1016/j.ynstr.2016.11.004>
- Oliveira, W. K. de, Duarte, E., França, G. V. A. de, & Garcia, L. P. (2020). Como o Brasil pode deter a COVID-19. *Epidemiologia e Serviços de Saúde*, 29(2), e2020044. <https://doi.org/10.5123/S1679-49742020000200023>
- Orth, U., & Robins, R. W. (2014). The development of self-esteem. *Current Directions in Psychological Science*, 23(5), 381-387. <https://doi.org/10.1177/0963721414547414>
- Phua, J., Weng, L., Ling, L., Egi, M., Lim, C.-M., Divatia, J. V.,... Du, B. (2020). Intensive care management of coronavirus disease 2019 (COVID-19): Challenges and recommendations. *The Lancet Respiratory Medicine*, 8(5), 506-517. [https://doi.org/10.1016/S2213-2600\(20\)30161-2](https://doi.org/10.1016/S2213-2600(20)30161-2)
- Rincón-Cortés, M., Herman, J. P., Lupien, S., Maguire, J., & Shansky, R. M. (2019). Stress: Influence of sex, reproductive status and gender. *Neurobiology of Stress*, 10, 100155. <https://doi.org/10.1016/j.ynstr.2019.100155>
- Roma, V. G., & Hope, D. A. (2017). Are we certain about which measure of intolerance of uncertainty to use yet? *Cognitive Behaviour Therapy*, 46(4), 327-338. <https://doi.org/10.1080/16506073.2016.1252792>
- Salk, R. H., Hyde, J. S., & Abramson, L. Y. (2017). Gender differences in depression in representative national samples: Meta-analyses of diagnoses and symptoms. *Psychological Bulletin*, 143(8), 783-822. <https://doi.org/10.1037/bul0000102>

- Satici, B., Saricali, M., Satici, S.A., Griffiths, M.D. (2020). Intolerance of uncertainty and mental wellbeing: Serial mediation by rumination and fear of COVID-19. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00305-0>
- Slovic, P. (2010). The psychology of risk. *Saúde e Sociedade*, 19(4), 731-747. <https://dx.doi.org/10.1590/S0104-12902010000400002>
- Taha, S., Matheson, K., Cronin, T., & Anisman, H. (2014). Intolerance of Uncertainty, appraisals, coping, and anxiety: The case of the 2009 H1N1 pandemic. *British Journal of Health Psychology*, 19(3), 592-605. <https://doi.org/10.1111/bjhp.12058>
- Tavares Lima, C. K., de Medeiros Carvalho, P. M., Silva Lima, I. A. A., de Oliveira Nunes, J. V. A., Saraiva, J. S., de Souza, R. I.,...Rolim Neto, M. L. (2020). The emotional impact of coronavirus 2019-nCoV (new coronavirus disease). *Psychiatry Research*, 287, 112915. <https://doi.org/10.1016/j.psychres.2020.112915>
- Tennant, R., Hiller, L., Fishwick, R., Platt, S., Joseph S., Weich, S.,...Stewart-Brown, S. (2007). The Warwick-Edinburgh Mental Well-being Scale (WEMWBS): development and UK validation. *Health and Quality of Life Outcomes* 5, 63. <https://doi.org/10.1186/1477-7525-5-63>
- Tolin, D. F., Abramowitz, J. S., Brigidi, B. D., & Foa, E. B. (2003). Intolerance of Uncertainty in obsessive-compulsive disorder. *Journal of Anxiety Disorders*, 17, 233-242.
- Vignola, R. C. B., & Tucci, A. M. (2014). Adaptation and validation of the depression, anxiety and stress scale (DASS) to Brazilian Portuguese. *Journal of Affective Disorders*, 155(1), 104-109. <https://doi.org/10.1016/j.jad.2013.10.031>
- Wang, C., Pan, R., Wan, X., Tan, Y., Xu, L., Ho, C. S.,...Ho, R. C. (2020). Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. *International Journal of Environmental Research and Public Health*, 17, 17- 29. <https://doi.org/10.3390/ijerph17051729>
- World Health Organization. Coronavirus disease (COVID-19): Situation report—126. Accessed May 30, 2020. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200525-covid-19-sitrep-126.pdf?sfvrsn=887dbd66_2
- Xiang, Y., Yang, Y., Li, W., Zhang, L., Zhang, Q., Cheung, T., & Ng, C. H. (2020). Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. *The Lancet Psychiatry*, 7(3), 228-229. [https://doi.org/10.1016/S2215-0366\(20\)30046-8](https://doi.org/10.1016/S2215-0366(20)30046-8)