



Revista EIA
ISSN: 1794-1237
revista@eia.edu.co
Escuela de Ingeniería de Antioquia
Colombia

Environmental Risk assessment of Chlorpyrifos and TCP in Aquatic Ecosystems

Gomez Arguello, Mayra Alejandra; Narvaez Valderrama, Jhon Fredy; Amaringo Villa, Fredy Alberto; Molina Pérez, Francisco José

Environmental Risk assessment of Chlorpyrifos and TCP in Aquatic Ecosystems

Revista EIA, vol. 17, núm. 34, 2020

Escuela de Ingeniería de Antioquia, Colombia

Disponible en: <https://www.redalyc.org/articulo.oa?id=149264860010>

DOI: <https://doi.org/10.24050/reia.v17i34.1313>

Environmental Risk assessment of Chlorpyrifos and TCP in Aquatic Ecosystems

Evaluación de Riesgo ambiental de Clorpirifos y TCP en Ecosistemas Acuáticos

Avaliação de Risco Ambiental do Clorpirifos e TCP em Ecosistemas Aquáticos

Mayra Alejandra Gomez Arguello
mayra.gomeza@udea.edu.co

Universidad de Antioquia, Colombia

Jhon Fredy Narvaez Valderrama
jhon.narvaez@uniremington.edu.co

Universidad de Antioquia, Colombia

Fredy Alberto Amaringo Villa fredy.amaringo@udea.edu.co
Universidad de Antioquia, Colombia

Francisco José Molina Pérez francisco.molina@udea.edu.co
Universidad de Antioquia, Colombia

Revista EIA, vol. 17, núm. 34, 2020

Escuela de Ingeniería de Antioquia,
Colombia

Recepción: 23 Febrero 2019

Aprobación: 18 Junio 2020

DOI: <https://doi.org/10.24050/reia.v17i34.1313>

Redalyc: <https://www.redalyc.org/articulo.oa?id=149264860010>

Abstract: Chlorpyrifos is a commonly used pesticide that belong to the organophosphorus pesticides (OPPs) group. Chlorpyrifos is widely used in residential environments and agriculture for pest control. As many OPPs, Chlorpyrifos degrades faster to more complex and toxic compounds under natural conditions, thus diverse effects of these compounds over aquatic species are unknown. Due to the risk that pesticides as Chlorpyrifos, which mainly breaks down to 3,5,6- trichloro-2-pyridinol (TCP) on the ecosystems, there is an increasing need to intensify the environmental monitoring and ecotoxicological risk assessment for both substances. Risk assessment provides a systematic approach for characterizing the nature and magnitude of the risks associated with environmental health hazards. However, in countries like Colombia, where the use of Chlorpyrifos is widespread, the number of investigations on the dynamics and risk involved in the presence of this class of substances in water bodies is limited.

Keywords: Chlorpyrifos, Risk Assessment, Aquatic ecosystems, Degradation products, Organophosphorus pesticides, Colombia.

Resumen: El clorpirifos es un plaguicida de uso común que pertenece al grupo de compuestos organofosforados (OPP). El clorpirifos se usa ampliamente en entornos residenciales y agrícolas para el control de plagas de insectos. Como muchos de los OPP, El clorpirifos se degrada rápidamente a compuestos más complejos y tóxicos en condiciones naturales, por lo que se desconocen los diversos efectos de estos compuestos sobre las especies acuáticas. Debido al riesgo de que los plaguicidas como el Clorpirifos, que se descompone principalmente en 3,5,6-tricloro-2-piridinol (TCP) en los ecosistemas, existe una necesidad de intensificar y ampliar los datos de monitoreo ambiental y la evaluación de riesgo ecotoxicológico para ambas sustancias. La evaluación de riesgos proporciona un enfoque sistemático para caracterizar la naturaleza y la magnitud de los riesgos asociados con los peligros para la salud ambiental. Sin embargo, en países como Colombia, donde el uso del Clorpirifos está muy extendido, el número de investigaciones sobre la dinámica y el riesgo que implica la presencia de esta clase de sustancias en cuerpos de agua es limitado.

Palabras clave: Clorpirifos, evaluación de riesgos, ecosistemas acuáticos, productos de degradación, pesticidas organofosforados, Colombia..

Resumo: O clorpirifos é um pesticida comumente usado para o grupo de compostos organofosforados (OPP). O clorpirifos é amplamente utilizado em ambientes residenciais e agrícolas para o controle de pragas de insetos. Como muitas OPPs, o clorpirifos é rapidamente degradado em compostos mais complexos e tóxicos em condições naturais, portanto os vários efeitos desses compostos sobre as espécies aquáticas são desconhecidos. Devido ao risco de pesticidas, tais como clorpirifos, decompõe-se principalmente em 3,5,6-tricloro-2-piridinol (TCP) em ecossistemas, existe uma necessidade crescente de melhorar e estender os dados de monitorização e ambientais a avaliação do risco ecotoxicológico para ambas as substâncias. A avaliação de risco fornece uma abordagem sistemática para caracterizar a natureza e a magnitude dos riscos associados aos perigos para a saúde ambiental. No entanto, em países como a Colômbia, onde o uso do clorpirifos é generalizado, o número de investigações sobre a dinâmica e o risco envolvido na presença dessa classe de substâncias em corpos d'água é limitado.

Palavras-chave: Clorpirifos, avaliação de risco, ecossistemas aquáticos, produtos de degradação, pesticidas organofosforados, Colômbia.

INTRODUCTION

Since the prohibition in the production and use of organochloride pesticides in the 1970s, due to their high bioaccumulation and toxic biological effects, organophosphorus pesticides have increased, especially in developing countries, where agriculture is the main economic activity Moussavi, et al. (2014). This is due to the proven high-efficiency of these substances to control pests at competitive costs. Even though pesticides offer economic benefits to the agricultural sector, the relationship between agricultural production and its environmental impact could be described as negative since the interaction between pesticides components also affects other populations in addition to the target population. Furthermore, due to their persistent nature, pesticides with high half-life ($t_{1/2}$) remain in the ecosystem longer periods and enter into the food chain Chawla, et al. (2018), affecting the vegetation cover, soil quality and aquatic systems Burgues, et al. (2012).

Factors such as land use and pesticide application pattern, rainfall intensity and irrigation strategy, soil type, landscape and field slope, physicochemical properties such as adsorption, absorption, solubility and Log Kow, and natural degradation processes are crucial to determine the environmental fate and persistence of pesticides, in surface water bodies, sediments and their own biota Sabatier, et al. (2014).

In Colombia, by the first quarter of 2018 there were 444 companies registered as manufacturers, formulators, importers, exporters, packagers and distributors of chemical pesticides for agricultural use Instituto Colombiano Agropecuario. (2018). In 2016, about 6,871.88 m³ and 6,831.43 Tn of Chlorpyrifos were commercialized. This data suggests a significant consumption of agrochemicals in the country, where Chlorpyrifos was the third most commercialized pesticide behind glyphosate and Mancozeb Instituto Colombiano Agropecuario ICA. (2017).

Taking into account the lack of appropriate technology, situations arise where use and misuse of pesticides produce considerable waste, adding to the cost and contributing to the adverse environmental and health consequences. Inappropriate application of pesticides increase the available fractions which may be accumulated in the food chain and may be transferred to soil, air, ground, and surface water Abhilash, Singh. (2009).

Due to the broad-spectrum insecticidal activity, OPPs are widely used in residential environment and agriculture for pest control Jain. (2017). However, those substances from polluted environments lead to acute toxicity on aquatic organisms (especially fish) while excessive exposure makes humans prone to acute phosphorus poisoning through phosphorylation of acetylcholinesterase (AChE) Dahiya, et al. (2017), and cause cholinergic effects over different species exposed Wu, et al. (2017). Furthermore, more chronic effects at low levels may arise on biota. Therefore, it is important to determine the effects that these substances generate in the ecosystems, which may be measured through different tools. Among these are Risk Assessments (ERA), which is a process to collect, organize, analyze and present scientific information to improve decision making US EPA. (2018b), that use a formal process of problem formulation (PF) to narrow the focus of the assessment to address key questions and, from these, develop the risk hypotheses US EPA. (1998). This methodology has been applied widely for a variety of substances such as Persistent Organic Compounds (POPPs) because it allows to determine levels of harm, prioritize issues, and inform policy for contaminated land management Andres, et al. (2018).

Chlorpyrifos

Chlorpyrifos belong to the organophosphorus compounds which are part of a large group of synthetic chemicals that can be broadly classified as esters, amides or thioester derivatives from phosphoric, phosphonic, phosphorothioic or phosphonothioic acids. Hundreds of those chemicals have been synthesized and studied for biological activity and many have been used commercially. As OPPs Chlorpyrifos is characterized by its low solubility in water, a high water-oil partition coefficient (LogKow), and a low vapor pressure Riches. (2014). Chlorpyrifos (Figure 1) is a widely commercialized insecticide used in rice, coffee, banana, among other crops; has a broad spectrum and moderate toxicity with a half-life (persistence) of 10–120 days in the environment Maya, et al. (2011), while its main degradation product is the 3,5,6- trichloro-2-pyridinol (TCP), which is a broad-spectrum antimicrobial and toxic metabolite Mauriz, et al. (2007), that is subsequently degraded to organochloride compounds and carbon dioxide Testai, Buratti, Di Consiglio. (2010). Chlorpyrifos mechanism of action is to interfere with (AChE), that is essential for the proper working of the nervous system of humans among others Xu, et al. (2008). The excessive use of chlorpyrifos and its

inappropriate application significantly increase the risks to human health, animals and the environment Agudelo, Jaramillo, Peñuela. (2012).




Figure 1.

A) Chemical structure of Chlorpyrifos, B) Chemical structure of TCP United States Environmental Protection Agency. (2018)

TCP is considered as one of the major degradation product of Chlorpyrifos Yang, et al. (2005) and exhibits a high degradation capacity, high water solubility and high migration capacity Lei, Huo, Zhou. (2017), It is moderately mobile due to its greater water solubility, which causes the widespread contamination in soils and in the aquatic environment Yang, et al. (2005) . Also TCP has antimicrobial activity, and it has been classified as toxic, persistent and mobile by the US EPA with a half-life ranging from 65 to 360 days in soil Maya, et al. (2011).

Physical and chemical characteristics of Chlorpyrifos and TCP (Table 1) are the principal factors that govern the fate in the different environmental matrices Solomon, Giesy. (2014), towards the aquatic systems, as a result of transport phenomena as well as volatilization Singare. (2016), diffusion Giesy, et al. (2014), advection Thibodeaux, Mackay. (2011), dispersion Hemond, Fechner-Levy. (2015) and sorption Gebremariam, et al. (2012). There, the rate of Chlorpyrifos degradation depends on environmental conditions, such as pH, temperature, UV radiation, and microbiota, therefore as being a xenobiotic that alters the quality of water and as a result of Chlorpyrifos presence and its degradation products, which may result even more hazardous than the parental compounds, affect the ecosystem Ríos-González. (2010).

Table 1

Physicochemical properties of Chlorpyrifos and its degradation product TCP Cheng, et al. (2007); National Center for Biotechnology Information. (2018); US EPA. (2018a).

| | CHLORPYRIFOS | TCP |
|---|---|--|
| Chemical name | O,O-diethyl-O-(3,5,6-trichloro-2-pyridyl)phosphorothioate | 3,5,6-Trichloro-2-pyridinol |
| Molecular Weight (g/mol) | 350.6 | 198.44 |
| Empirical and Structural Formula | C ₉ H ₁₁ Cl ₃ NO ₃ PS | C ₅ H ₂ Cl ₃ NO |
| CAS Registry Number | 2921-88-2 | 6515-38-4 |
| Melting point (°C) | 41.5 – 42.5 | 208 - 209 |
| Vapor pressure (mmHg) | 2.03E-05 at 25°C | 1.03E-05 at 25°C |
| Density (g/mL) | 1.51 at 21°C | 1.67 at 26 °C |
| Solubility in water (g/L) at 25°C | 1.12 | 89 |
| Partition coefficient (n-octanol and water) | log K _{ow} = 5.1 | log K _{ow} = 3.21 |

Chlorpyrifos in the aquatic environments and Its Risk Assessment

Acute and chronic exposure to common-use pesticides remains a considerable threat to non-target species, despite the continued effort to synthesize compounds with high target specificity and low environmental persistence Simpson, Jeyasingh and Belden. (2017), in the case of Chlorpyrifos its behavior in surface water may be given by complex interactions of factors related to its application, agronomic practices, climatological conditions during and after the application, soil pedology and chemistry, hydrologic responses of drainage systems, and its physicochemical properties that affect mobility and persistence under those environmental settings Williams, et al. (2014). Chlorpyrifos may cause acute toxicity effects by inhibition of AChE occurring at low-level exposure in organisms that lack the target enzyme Giddings, et al. (2014); besides its long half-life in water, the effects of chlorpyrifos on aquatic ecosystems at different trophic levels are attracting more interest Zhao, Chen. (2016). Chlorpyrifos is potentially toxic to most organisms that show differences in susceptibility, this is because of the differences in rates of adsorption, distribution, metabolism, and excretion among species Solomon, Giesy. (2014).

Due to the threat that pesticides as Chlorpyrifos pose to the ecosystems, and owing to the growing public awareness on the need of protecting both ecosystems and human health from the risks related to chemical pollution Kuzmanović, et al. (2015). There is an increasing need to intensify the environmental monitoring data and Ecotoxicological Risk Assessment(ERA) Carazo-Rojas, et al. (2018). Recently, many monitoring programs and scientific studies have put more focus on

the occurrence, distribution and fate of pesticides and their potential environmental effects in individual rivers, reservoirs and lakes Chen, et al. (2018)

The ERA provide a systematic approach for characterizing the nature and magnitude of the risks associated with environmental health hazards Department of Health and Ageing of Australia. (2004), and allow to estimate the likelihood that undesired effects might occur or are occurring as a result of exposure to one or more stressors Guo, et al. (2013), in view of ecological risk assessments involving multiple chemical, physical and biological stressors are, by their very nature, complex Moore. (2001).

Researchers have been developing different approaches for ERA in order to make more holistic assessments due to the complexity of the ecosystems and the substances in the aquatic systems Tsaboula, et al. (2016); using tools such as multimedia fate, exposure and effects models Guillén, et al. (2012); Houbraeken, et al. (2017); the influence of physicochemical factors such as temperature, volatilization, precipitation Delnat, et al. (2019) ,Beeck, Verheyen, Stoks. (2018), Potter, Coffin. (2017), Pereira, Cerejeira, Daam. (2017); ERAs based on the Ecosystem services Raimondo, et al. (2019), and ecotoxicity assays Janssens, Op De Beeck, Stoks. (2017).

Table 2 shows a summary of ERA conducted for some organophosphorus compounds including Chlorpyrifos in different environmental matrices and sample techniques, which found that Chlorpyrifos was systematically associated with hazard to the studied system.

Table 2
Risk Assessments related with Chlorpyrifos and other organophosphorus compounds

| Assessment | Environmental Matrix | Country | Source |
|---|--|-----------------|-------------------------------------|
| Framework For Ecotoxicological Risk Assessment, EPA method | Aquatic organisms | United States | (Giddings <i>et al.</i> , 2014) |
| Joint Probability Curve | surface waters | United States | (Wang, Singhasemanon and Goh, 2016) |
| Biological Response Indicator Devices Gauging Environmental Stressors (BRIDGES) | Passive Samplers in surface waters | United States | (Hillwalker <i>et al.</i> , 2010) |
| PRISW-1: Short- Term Pesticide Risk Index for Surface Water System | Polar Organic Chemical Integrative Sampler in surface waters | Lebanon | (Aisha <i>et al.</i> , 2017) |
| PRIMET model | Aquatic Ecosystems | Mexico | (Ríos-González, 2010) |
| Prospective Aquatic Environmental Risk Assessment | Aquatic Ecosystems | The Netherlands | (Rico <i>et al.</i> , 2016) |
| Ecotoxicological essays | Aquatic Ecosystems | Spain | (Rivetti <i>et al.</i> , 2017) |
| The probabilistic cumulative risk assessment | Soil | United States | (Li, 2018) |
| First-Tier Risk Quotient (RQ) Approach | Aquatic Ecosystems | Bangladesh | (Sumon <i>et al.</i> , 2018) |
| Toxicity Units | Aquatic Ecosystems | Ghana | (Affum <i>et al.</i> , 2018) |
| ChimERA fate | Shallow Aquatic Ecosystems | Italy | (Morselli <i>et al.</i> , 2018) |

n spite of the increasing number of ERAs made around the world, there is a limited number of investigations about OPPs pesticides in equatorial zones such as Colombia. In 2015 Narvaez, (2015) conducted research about the occurrence and preliminary risk assessment of Chlorpyrifos and its principal degradation product TCP in a reservoir in Colombia, where the Risk Quotient was less than 0.1, which indicate a presumption of no danger to the population. Nevertheless, the preliminary results indicated that the TCP as a partial transformation of Chlorpyrifos leads to a product with possible impact on the population.

Although there have been studies on the presence and monitoring of organophosphorus pesticides in aquatic ecosystems in latitudes close to the tropics Carro, et al. (2012); Singare. (2016); Huang, Tornero-Velez, Barzyk. (2017); Bedoya-Ríos, et al. (2018); Moncaleano-Niño, et al. (2018)., it is necessary to do more research about the dynamic, and impact of the OPPs, because in equatorial regions like Colombia, due

to its distinctive climatological conditions processes such as occurrence, distribution and fate of organophosphorus compounds as Chlorpyrifos and its degradation products are still poorly known (61).

Quantitative Structure Activity Relationship (QSARs) models of Chlorpyrifos and TCP for Ecological Risk Assessment

Although *Daphnia magna*, an important freshwater invertebrate species in aquatic food webs, has been used worldwide for many years as a representative test species for the ecotoxicological evaluation of chemicals, this approach is expensive and time-consuming Toropova, et al. (2016). Also, based on the reduction in the use of living organisms to carry on test, QSAR models appear as an alternative, since for ethical reasons, such as avoiding animal testing, QSAR methods may be also preferable Guillén, et al. (2012). Another factor that explain why QSAR models are popular is because they can be developed using experimental data of a small data set, which can be used to get predictions for a large dataset provided they fall inside the applicability domain of the models Khan, Roy, Benfenati. (2019).

QSAR, sometimes defined as “in silico” computational models, enable the prediction of physicochemical or biological properties of the compound by comparing them to other known molecules. In QSAR, different types of descriptors associated with chemical structures are quantitatively correlated with their physicochemical properties as melting point, water solubility, etc., environmental fate, ecotoxicity and other effect end points related to human health Guillén, et al. (2012), Hamadache, et al. (2014). These QSAR models have proven to be very valuable as prioritization tools to classify compounds according to their toxicity and, as more information becomes available, to predict toxicity Benfenati, et al. (2017), and because of that utility the use of QSAR is recommended in early detection of environmental hazards by several regulatory agencies like European Centre for the validation of Alternative Methods (ECVAM) of the European Union, United States Environmental Protection Agency (US EPA), European Union Commission's Scientific Committee on Toxicity, and the Agency for Toxic Substances and Disease Registry (ATSDR) Khan, Roy, Benfenati. (2019).

In the specific case of Chlorpyrifos and its principal degradation product, preliminary QSAR models indicate that the anaerobic conditions in reservoir sediments favor the reductive dehalogenation mechanisms of CPF and TCP, which could carry out the formation of pyridine, a compound with high carcinogenic activity Narvaez, et al. (2014).

Conclusion

The environmental risk assessment has proven to be an appropriate tool for the estimation of environmental health hazards and therefore the well-being of the human being. This is why ERA it's widely used by government regulatory agencies around the world as the basis for decision making for the management of resources. Risk assessments associated with the exposure of organophosphorus pesticides such as chlorpyrifos have helped the authorities to restrict the use of these substances in their territories, however, these investigations have been carried out in latitudes with climatic and geomorphological characteristics different from those of Colombia, where the use of this pesticides is common in agriculture and livestock.

Due to the limited number of investigations on the dynamics of organophosphorus pesticides, such as chlorpyrifos and its degradation products in Colombian ecosystems, the number of associated ecological risk assessments has also been limited. That is why it is necessary to conduct future researches using tools such as QSAR models that are low cost and with easy access to information in order to predict the toxic effects of these substances widely used in the agricultural sector of Colombia and with this, present scientific bases that allow Colombian governmental entities to manage policies for the regulation of the use of these chemical compounds.

Acknowledgments

The authors thank the Department of Science and Technology of Colombia (COLCIENCIAS) for its financial support within the framework of the 727 of 2015 national scholarships.

References

- Abhilash, P. C.; Singh, N. (2009). 'Pesticide use and application: An Indian scenario', *Journal of Hazardous Materials*, 165(1–3), pp. 1–12. doi: 10.1016/j.jhazmat.2008.10.061.
- Affum, A. O.; Acquah, S.O.; Osa, S.D.; Kwaansa-Ansah, E.B. (2018). 'Distribution and risk assessment of banned and other current-use pesticides in surface and groundwaters consumed in an agricultural catchment dominated by cocoa crops in the Ankobra Basin, Ghana', *Science of the Total Environment*. Elsevier B.V., 633, pp. 630–640. doi: 10.1016/j.scitotenv.2018.03.129.
- Agudelo C, R. M.; Jaramillo, M. L.; Peñuela, G. (2012). 'Comparison of the removal of chlorpyrifos and dissolved organic carbon in horizontal sub-surface and surface flow wetlands', *Science of the Total Environment*. Elsevier B.V., 431, pp. 271–277. doi: 10.1016/j.scitotenv.2012.05.045.
- Aisha, A. A.; Hneine, W.; Mokh, S.; Devier, M.H.; Budzinski, H.; Jaber, F. (2017). 'Monitoring of 45 pesticides in Lebanese surface water using polar

- organic chemical integrative sampler (POCIS)', *Ocean Science Journal*, pp. 1–12. doi: 10.1007/s12601-017-0041-4.
- Arias Espana, V.A.; Rodriguez Pinilla, A.R.; Bardos, P.; Naidu, R. (2018). 'Contaminated land in Colombia#: A critical review of current status and future approach for the management of contaminated sites', *Science of the Total Environment*. Elsevier B.V., 618, pp. 199–209. doi: 10.1016/j.scitotenv.2017.10.245.
- Bedoya-Ríos, D. F. Lara-Borrero, J.A.; Duque-Pardo, V.; Madera-Parra, C.A.; Jimenez, E.M.; Toro, A.F. (2018). 'Study of the occurrence and ecosystem danger of selected endocrine disruptors in the urban water cycle of the city of Bogotá, Colombia', *Journal of Environmental Science and Health, Part A*. Taylor & Francis, 53(4), pp. 317–325. doi: 10.1080/10934529.2017.1401372.
- Op de Beeck, L.; Verheyen, J.; Stoks, R. (2018). 'Competition magnifies the impact of a pesticide in a warming world by reducing heat tolerance and increasing autotomy', *Environmental Pollution*. Elsevier Ltd, 233, pp. 226–234. doi: 10.1016/j.envpol.2017.10.071.
- Benfenati, E. Toropov, A.A.; Toropova, A.P.; Marzo, M.; Dorne, J.L.; Georgiadis, N. (2017). 'QSAR models for predicting acute toxicity of pesticides in rainbow trout using the CORAL software and EFSA's OpenFoodTox database', *Environmental Toxicology and Pharmacology*, 53(May), pp. 158–163. doi: 10.1016/j.etap.2017.05.011.
- Burgues, M.; Golin, A.; Maydana, G.; Portapila, M. (2012). 'Análisis preliminar de contaminación en aguas superficiales provenientes de fertilizantes y pesticidas utilizados en las actuales prácticas agrícolas', in *1er Encuentro De Investigadores En Formación En Recursos Hídricos*. Buenos Aires, Argentina. Available at: http://www.ina.gov.ar/pdf/ifrrhh/02_003_Burgues.pdf.
- Carazo-Rojas, E.; Pérez-Rojas, G.; Pérez-Villanueva, M.; Chinchilla-Soto, C.; Chin-Pampillo, J.S.; Aguilar-Mora, P.; Alpízar-Marína, M.; Masís-Mora, M.; Rodríguez-Rodríguez, C.E.; Vryzas, Z. (2018). 'Pesticide monitoring and ecotoxicological risk assessment in surface water bodies and sediments of a tropical agro-ecosystem', *Environmental Pollution*, 241, pp. 800–809. doi: 10.1016/j.envpol.2018.06.020.
- Carro, A. M.; Fernández, S.; Racamonde, I.; García-Rodríguez, D.; González, P.; Lorenzo, R.A. (2012). 'Dispersive liquid-liquid microextraction coupled with programmed temperature vaporization-large volume injection-gas chromatography-tandem mass spectrometry for multiclass pesticides in water', *Journal of Chromatography A*. Elsevier B.V., 1253, pp. 134–143. doi: 10.1016/j.chroma.2012.06.089.
- Chawla, P.; Kaushik, R.; Shiva Swaraj, V.J.; Kumar, N. (2018). 'Organophosphorus pesticides residues in food and their colorimetric detection', *Environmental Nanotechnology, Monitoring and Management*. Elsevier, 10(July), pp. 292–307. doi: 10.1016/j.enmm.2018.07.013.
- Chen, Y.; Yu, K.; Hassan, M.; Xu, C.; Zhang, B.; Gin, K.Y.H.; He, Y. (2018). 'Occurrence, distribution and risk assessment of pesticides in a river-reservoir system', *Ecotoxicology and Environmental Safety*. Elsevier Inc., 166(July), pp. 320–327. doi: 10.1016/j.ecoenv.2018.09.107.
- Cheng, Y.; Sang, S.X.; Huang, H.Z.; Liu, X.J.; Ouyang, J.B. (2007). 'Variation of Coenzyme F420 Activity and Methane Yield in Landfill Simulation of

- Organic Waste', *Journal of China University of Mining and Technology*, 17(3), pp. 403–408. doi: 10.1016/S1006-1266(07)60114-X.
- Dahiya, V.; Chaubey, B.; Dhaharwal A.K.; Pal, S. (2017) 'Solvent-dependent binding interactions of the organophosphate pesticide, chlorpyrifos (CPF), and its metabolite, 3,5,6-trichloro-2-pyridinol (TCPy), with Bovine Serum Albumin (BSA): A comparative fluorescence quenching analysis', *Pesticide Biochemistry and Physiology*. Elsevier Inc., 139, pp. 92–100. doi: 10.1016/j.pestbp.2017.04.011.
- Delnat, V.; Tran, T.T.; Janssens, L.; Stoks, R. (2019). 'Daily temperature variation magnifies the toxicity of a mixture consisting of a chemical pesticide and a biopesticide in a vector mosquito', *Science of the Total Environment*. Elsevier B.V., 659, pp. 33–40. doi: 10.1016/j.scitotenv.2018.12.332.
- Department of Health and Ageing of Australia. (2004). 'ENVIRONMENTAL HEALTH RISK ASSESSMENT, Guidelines for assessing human health risks from environmental hazards'. Canberra, Australia, p. 258. Available at: <http://enhealth.nphp.gov.au/council/pubs/pubs.htm>.
- Gebremariam, S. Y.; Beutel, M.W.; Yonge, D.R.; Flury, M.; Harsh, J.B. (2012). 'Adsorption and Desorption of Chlorpyrifos to Soils and Sediments', *Reviews of Environmental Contamination and Toxicology, Volume*, 215, pp. 123–155. DOI: 10.1007/978-1-4614-1463-6_3.
- Giddings, J. M.; Williams, W.M.; Solomon, K.R.; Giesy, J.P. (2014). 'Risks to Aquatic Organisms from Use of Chlorpyrifos in the United States', *Reviews of Environmental Contamination and Toxicology*. Edited by P. J. Giesy and R. K. Solomon. Cham: Springer International Publishing, (231), pp. 119–162. doi: 10.1007/978-3-319-03865-0_5.
- Giesy, J. P.; Solomon, K.R.; Cutler, G.C.; Giddings, J.M.; Mackay, D.; Moore, D.R.J.; Purdy, J.; Williams, W.M. (2014). 'Ecological risk assessment of the uses of the organophosphorus insecticide chlorpyrifos, in the United States', *Reviews of Environmental Contamination and Toxicology*. Saskatoon, Canada, pp. 1–11. doi: 10.1007/978-3-319-03865-0_1.
- Guillén, D.; Ginebreda, A.; Farré, M.; Darbra, R.M.; Petrovic, M.; Gros, M.; Barceló, D. (2012). 'Prioritization of chemicals in the aquatic environment based on risk assessment: Analytical, modeling and regulatory perspective', *Science of the Total Environment*. Elsevier B.V., 440, pp. 236–252. doi: 10.1016/j.scitotenv.2012.06.064.
- Guo, G.; Wu, F.; He, H.; Zhang, R.; Li, H. (2013). 'Ecological Risk Assessment of Organochlorine Pesticides in Surface Waters of Lake Taihu, China', *Human and Ecological Risk Assessment*, 19(4), pp. 840–856. doi: 10.1080/10807039.2012.691811.
- Hamadache, M.; Khaouane, L.; Benkortbi, O.; Si Moussa, C.; Hanini, S.; Amrane, A. (2014). 'Prediction of Acute Herbicide Toxicity in Rats from Quantitative Structure–Activity Relationship Modeling', *Environmental Engineering Science*, 31(5), pp. 243–252. doi: 10.1089/ees.2013.0466.
- Hemond, H. F.; Fechner-Levy, E. (2015). *CHEMICAL FATE AND TRANSPORT IN THE ENVIRONMENT*. third edit. Elsevier.
- Hillwalker, W. E.; Allan, S.E.; Tanguay, R.L.; Anderson, K.A. (2010). 'Exploiting lipid-free tubing passive samplers and embryonic zebrafish to link site specific contaminant mixtures to biological

- responses', *Chemosphere*. Elsevier Ltd, 79(1), pp. 1–7. doi: 10.1016/j.chemosphere.2010.02.001.
- Houbraken, M. *et al.* (2017). 'Science of the Total Environment Multi-residue determination and ecological risk assessment of pesticides in the lakes of Rwanda', *Science of the Total Environment*. Elsevier B.V., 576, pp. 888–894. doi: 10.1016/j.scitotenv.2016.10.127.
- Huang, H.; Tornero-Velez, R.; Barzyk, T. M. (2017). 'Multi-class chemical exposure in rural Peru using silicone wristbands', *Journal of Exposure Science and Environmental Epidemiology*, 27(6), pp. 544–550. doi: 10.1038/jes.2017.15.
- Instituto Colombiano Agropecuario, I. (2018). *Empresas Titulares de Registros de Plaguicidas-abril 2018*. Available at: <https://www.ica.gov.co/Areas/Agricola/Servicios/Regulacion-y-Control-de-Plaguicidas-Quimicos/Listados/2009/EMPRESAS-PLAGUICIDAS-PQUA-15-04-09.aspx>.
- Instituto Colombiano Agropecuario ICA. (2017). *ESTADÍSTICAS DE COMERCIALIZACIÓN DE PLAGUICIDAS QUÍMICOS DE USO AGRÍCOLA 2016*. Bogotá, Colombia. Available at: https://www.ica.gov.co/Areas/Agricola/Servicios/Regulacion-y-Control-de-Plaguicidas-Quimicos/Estadisticas/Cartilla-Plaguicidas-2016_22-01-18.aspx.
- Jain, R. B. (2017). 'Association between thyroid function and urinary levels of 3,5,6-trichloro-2-pyridinol: data from NHANES 2007–2008', *Environmental Science and Pollution Research*. Environmental Science and Pollution Research, 24(3), pp. 2820–2826. doi: 10.1007/s11356-016-8007-0.
- Janssens, L.; Op De Beeck, L.; Stoks, R. (2017). 'Stoichiometric Responses to an Agricultural Pesticide Are Modified by Predator Cues', *Environmental Science and Technology*, 51(1), pp. 581–588. doi: 10.1021/acs.est.6b03381.
- Khan, K.; Roy, K.; Benfenati, E. (2019). 'Ecotoxicological QSAR modeling of endocrine disruptor chemicals', 369(February), pp. 707–718.
- Kuzmanović, M.; Ginebreda, A.; Petrović, M.; Barceló, D. (2015). 'Risk assessment based prioritization of 200 organic micropollutants in 4 Iberian rivers', *Science of the Total Environment*, 503–504, pp. 289–299. doi: 10.1016/j.scitotenv.2014.06.056.
- Lei, W.; Huo, X.; Zhou, X. (2017). 'Adsorption Characteristics and Its Parameters Estimation of 3,5,6-trichloro-2-pyridinol in Purple Soil', *Transactions of the Chinese Society for Agricultural Machinery*, 48(5), pp. 267–274. doi: 10.6041/j.issn.1000-1298.2017.05.033.
- Li, Z. (2018). 'Introducing relative potency quotient approach associated with probabilistic cumulative risk assessment to derive soil standards for pesticide mixtures', *Environmental Pollution*. Elsevier Ltd, 242, pp. 198–208. doi: 10.1016/j.envpol.2018.06.076.
- Mauriz, E.; Calle, A.; Manclús J. J.; Montoya, A.; Lechuga, L. M. (2007). 'On-line determination of 3,5,6-trichloro-2-pyridinol in human urine samples by surface plasmon resonance immunosensing', *Analytical and Bioanalytical Chemistry*, 387(8), pp. 2757–2765. doi: 10.1007/s00216-007-1175-5.
- Maya, K.; Singh, R.S.; Upadhyay, S.N.; Dubey, S. K. (2011). 'Kinetic analysis reveals bacterial efficacy for biodegradation of chlorpyrifos and its

- hydrolyzing metabolite TCP', *Process Biochemistry*, 46(11), pp. 2130–2136. doi: 10.1016/j.procbio.2011.08.012.
- Moncaleano-Niño, A. M.; Acosta, A.L.; Gómez-Cubillos, M.C.; Villamil, L.; Abrens, M.J. (2018). 'Cholinesterase activity in the cup oyster *Saccostrea* sp. exposed to chlorpyrifos, imidacloprid, cadmium and copper', *Ecotoxicology and Environmental Safety*. Elsevier Inc., 151(January), pp. 242–254. doi: 10.1016/j.ecoenv.2017.12.057.
- Moore, D. R. J. (2001). 'The *Anna Karenina* Principle Applied to Ecological Risk Assessments of Multiple Stressors', *Human and Ecological Risk Assessment: An International Journal*, 7(2), pp. 231–237. doi: 10.1080/20018091094349.
- Morselli, M.; Terzaghi, E.; Galimberti, F.; Di Guardo, A. (2018) 'Pesticide fate in cultivated mountain basins: The improved DynAPlus model for predicting peak exposure and directing sustainable monitoring campaigns to protect aquatic ecosystems', *Chemosphere*. Elsevier Ltd, 210, pp. 204–214. doi: 10.1016/j.chemosphere.2018.06.181.
- Moussavi, G.; Hossaini, H.; Jafari, S. J.; Farokhi, M. (2014). 'Comparing the efficacy of UVC, UVC/ZnO and VUV processes for oxidation of organophosphate pesticides in water', *Journal of Photochemistry and Photobiology A: Chemistry*, 290, pp. 86–93. doi: 10.1016/j.jphotochem.2014.06.010.
- Narvaez, J. F. (2015). *DINÁMICA Y EVALUACIÓN PRELIMINAR DE RIESGO AMBIENTAL DE PLAGUICIDAS Y PRODUCTOS DE DEGRADACIÓN EN LOS EMBALSES LA FE Y RIOGRANDE II – COLOMBIA, POR MEDIO DE MUESTREADORES PASIVOS*. Universidad de Antioquia.
- Narvaez Valderrama, J. F.; Berrio Puerta, J.A.; Correa Zuluaga, S.; Palacio Baena, J. A.; Molina Pérez, F. J. (2014). 'DEGRADACIÓN HIDROLÍTICA DE CLORPIRIFOS Y EVALUACIÓN DE LA TOXICIDAD DEL EXTRACTO HIDROLIZADO CON *Daphnia pulex*', *Revista Politécnica*. Medellín, Antioquia: Fondo Editorial POLI, pp. 9–15. National Center for Biotechnology Information (2018) 3,5,6-Trichloro-2-pyridinol, PubChem Compound Database.
- Pereira, A. S.; Cerejeira, M. J.; Daam, M. A. (2017). 'Toxicity of environmentally realistic concentrations of chlorpyrifos and terbuthylazine in indoor microcosms', *Chemosphere*, 182, pp. 348–355. doi: 10.1016/j.chemosphere.2017.05.032.
- Potter, T. L.; Coffin, A. W. (2017). 'Assessing pesticide wet deposition risk within a small agricultural watershed in the Southeastern Coastal Plain (USA)', *Science of the Total Environment*, 580, pp. 158–167. doi: 10.1016/j.scitotenv.2016.11.020.
- Raimondo, S.; Sharpe, L.; Oliver, L.; McCaffrey, K. R.; Purucker, S. T.; Sinnathamby, S.; Minucci J. M. (2019). 'A unified approach for protecting listed species and ecosystem services in isolated wetlands using community-level protection goals', *Science of the Total Environment*. Elsevier B.V., 663, pp. 465–478. doi: 10.1016/j.scitotenv.2019.01.153.
- Riches, J. (2014). *Analysis of Organophosphorus Chemicals, Best Synthetic Methods: Organophosphorus (V) Chemistry*. Elsevier Ltd. doi: 10.1016/B978-0-08-098212-0.00007-8.

- Rico, A.; Van den Brink, P. J.; Gylstra, R.; Focks, A.; Brock, T. C. M. (2016). 'Developing ecological scenarios for the prospective aquatic risk assessment of pesticides', *Integrated environmental assessment and management*, 12(3), pp. 510–521. doi: 10.1002/ieam.1718.
- Ríos-González, A. (2010). *Uso de modelos predictivos y conceptuales para la evaluación ambiental y el análisis de la percepción de riesgo por uso de plaguicidas: Una opción para el manejo de riesgos en Chiapas. TESIS, Ecosur*. El Colegio de la Frontera Sur.
- Rivetti, C.; López-Perea, J.J.; Laguna, C.; Piña, B.; Mateo, R.; Eljarrat, E.; Barceló, D.; Barata, C. (2017). 'Integrated environmental risk assessment of chemical pollution in a Mediterranean floodplain by combining chemical and biological methods', *Science of the Total Environment*. Elsevier B.V., 583, pp. 248–256. doi: 10.1016/j.scitotenv.2017.01.061.
- Sabatier, P.; Poulenard, J.; Fanget, B.; Reyss, J. L.; Develle, A. L.; Wilhelm, B.; Ployon, E.; Pignol, C.; Naffrechoux, E.; Dorioz, J. M.; Montuelle, B.; Arnaud, F. (2014). 'Long-term relationships among pesticide applications, mobility, and soil erosion in a vineyard watershed', *Proceedings of the National Academy of Sciences*, 111(44), pp. 15647–15652. doi: 10.1073/pnas.1411512111.
- Simpson, A. M.; Jeyasingh, P. D.; Belden, J. B. (2017). 'Assessment of biochemical mechanisms of tolerance to chlorpyrifos in ancient and contemporary *Daphnia pulex* genotypes', *Aquatic Toxicology*. Elsevier, 193(October), pp. 122–127. doi: 10.1016/j.aquatox.2017.10.012.
- Singare, P. U. (2016). 'Distribution and risk assessment of suspected endocrine-disrupting pesticides in creek water of Mumbai, India', *Marine Pollution Bulletin*. Elsevier Ltd, 102(1), pp. 72–83. doi: 10.1016/j.marpolbul.2015.11.055.
- Solomon, K.; Giesy, J. P. (2014). 'Ecological Risk Assessment for Chlorpyrifos in Terrestrial and Aquatic Systems in the United States', *Reviews of Environmental Contamination and Toxicology*, p. 282 pp. doi: 10.1007/978-3-319-03865-0.
- Sumon, K. A.; Rashid, H.; Peeters, E.; Bosma, R. H.; Van den Brink, P. J. (2018). 'Environmental monitoring and risk assessment of organophosphate pesticides in aquatic ecosystems of north-west Bangladesh', *Chemosphere*. Elsevier Ltd, 206, pp. 92–100. doi: 10.1016/j.chemosphere.2018.04.167.
- Testai, E.; Buratti, F. M.; Di Consiglio, E. (2010). 'Chlorpyrifos', *Hayes' Handbook of Pesticide Toxicology*, pp. 1505–1526. doi: 10.1016/B978-0-12-374367-1.00070-7.
- Thibodeaux, L.; Mackay, D. (2011). *Handbook of chemical mass transport in the environment*. Edited by L. J. Thibodeaux and D. Mackay. Boca Raton, USA: CRC Press.
- Toropova, A. P.; Toropov, A. A.; Veselinović, A. M.; Veselinović, J. B.; Leszczynska, D.; Leszczynski, J. (2016). 'Monte Carlo-based quantitative structure-activity relationship models for toxicity of organic chemicals to *Daphnia magna*', *Environmental Toxicology and Chemistry*, 35(11), pp. 2691–2697. doi: 10.1002/etc.3466.
- Tsaboula, A.; Papadakis, E. N.; Vryzas, Z.; Kotopoulou, A.; Kintzikoglou, K.; Papadopoulou-Mourkidou, E. (2016). 'Environmental and human risk hierarchy of pesticides: A prioritization method, based on monitoring,

- hazard assessment and environmental fate', *Environment International*. Elsevier Ltd, 91, pp. 78–93. doi: 10.1016/j.envint.2016.02.008.
- United States Environmental Protection Agency, U. S. E. (2018). 'Ecological Estructure Activity Relationships (ECOSAR) Predictive Model'. Available at: <https://www.epa.gov/tsca-screening-tools/ecological-structure-activity-relationships-ecosar-predictive-model>.
- US EPA. (1998). 'Guidelines for Ecological Risk Assessment'. Washington, DC, p. 188. Available at: https://www.epa.gov/sites/production/files/2014-11/documents/eco_risk_assessment1998.pdf.
- US EPA .(2018a). *Ecological Structure Activity Relationships (ECOSAR) Predictive Model*. Available at: <https://www.epa.gov/tsca-screening-tools/ecological-structure-activity-relationships-ecosar-predictive-model>.
- US EPA. (2018b). 'Watershed Ecological Risk Assessment', pp. 1–32. Available at: <http://www.epa.gov/watertrain>.
- Wang, D.; Singhasemanon, N.; Goh, K. S. (2016). 'A statistical assessment of pesticide pollution in surface waters using environmental monitoring data: Chlorpyrifos in Central Valley, California', *Science of The Total Environment*, 571, pp. 332–341. doi: 10.1016/j.scitotenv.2016.07.159.
- Williams, W. M.; et al. (2014) 'Ecological Risk Assessment for Chlorpyrifos in Terrestrial and Aquatic Systems in the United States', *Reviews of Environmental Contamination and Toxicology*, 231, pp. 77–117. doi: 10.1007/978-3-319-03865-0.
- Wu, S.; Li, D.; Wang, J.; Zhao, Y.; Dong, S.; Wang, X. (2017). 'Gold nanoparticles dissolution based colorimetric method for highly sensitive detection of organophosphate pesticides', *Sensors and Actuators, B: Chemical*. Elsevier B.V., 238, pp. 427–433. doi: 10.1016/j.snb.2016.07.067.
- Xu, G.; Zheng, W.; Li, Y.; Wang, S.; Zhang, J.; Yan, Y. (2008). 'Biodegradation of chlorpyrifos and 3,5,6-trichloro-2-pyridinol by a newly isolated Paracoccus sp. strain TRP', *International Biodeterioration and Biodegradation*, 62(1), pp. 51–56. doi: 10.1016/j.ibiod.2007.12.001.
- Yang, L.; Zhao, Y. H.; Zhang, B. X.; Yang, C. H.; Zhang, X. (2005). 'Isolation and characterization of a chlorpyrifos and 3,5,6-trichloro-2- pyridinol degrading bacterium', *FEMS Microbiology Letters*, 251(1), pp. 67–73. doi: 10.1016/j.femsle.2005.07.031.
- Zhao, J.; Chen, B. (2016) 'Species sensitivity distribution for chlorpyrifos to aquatic organisms: Model choice and sample size.', *Ecotoxicology and environmental safety*, 125, pp. 161–9. doi: 10.1016/j.ecoenv.2015.11.039.

Notas de autor

mayra.gomeza@udea.edu.co