



Acta Medica Colombiana

ISSN: 0120-2448

Asociacion Colombiana de Medicina Interna

Hernández-Sómerson, Mario Andrés; Ortiz-Arroyo, Jaime  
Bullous hemorrhagic dermatosis at distant sites  
Acta Medica Colombiana, vol. 45, no. 4, 2020, October-December, p. 67  
Asociacion Colombiana de Medicina Interna

DOI: <https://doi.org/10.36104/amc.2020.1738>

Available in: <https://www.redalyc.org/articulo.oa?id=163168638013>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UAEH  redalyc.org

Scientific Information System Redalyc  
Network of Scientific Journals from Latin America and the Caribbean, Spain and  
Portugal

Project academic non-profit, developed under the open access initiative

## Bullous hemorrhagic dermatosis at distant sites

MARIO ANDRÉS HERNÁNDEZ-SÓMERSON, JAIME ORTIZ-ARROYO • BOGOTÁ, D.C. (COLOMBIA)

DOI: <https://doi.org/10.36104/amc.2020.1738>



Figure 1. Lesions on the right arm.



Figure 2. Lesions on the left foot.

A 72-year-old man was hospitalized for acute heart failure. He had a history of arterial hypertension being treated with enalapril and hydrochlorothiazide. Thromboprophylaxis was begun at admission with 40 mg of subcutaneous enoxaparin daily, in the abdomen. Three days later, tense, blood-filled bullous skin lesions with a 5-15 mm diameter appeared. The lesions were on the right arm (Figure 1) and the left foot (Figure 2); they had an erythematous halo and were not painful.

Clotting, C-reactive protein, ESR and complete blood count were normal. A skin biopsy was performed, which showed a subepidermal lesion with hemorrhagic material in the papillary and reticular dermis with scant neutrophils and no vasculitis or thrombosis. These findings confirm the diagnosis of bullous hemorrhagic dermatosis at distant sites related to enoxaparin. This is an uncommon and self-resolving adverse effect of heparins, unrelated to the dose and at sites distant from their application. This patient was treated by discontinuing the medication, and the lesions resolved seven days later with no recurrence after three months of follow-up.

Dr. Mario Andrés Hernández-Sómerson: Especialista en Medicina Interna, Servicio de Clínicas Médicas Hospital Universitario Mayor – MEDERI, Docente Facultad de Ciencias de la Salud Universidad del Rosario; Dr. Jaime Ortiz-Arroyo: Servicio de Clínicas Médicas, Hospital Universitario Mayor – MEDERI, Bogotá, D.C. (Colombia). Correspondencia: Mario Andrés Hernández-Sómerson. Bogotá, D.C. (Colombia). E-mail: somerson18@hotmail.com

Received: 26/III/2019 Accepted: 12/VIII/2020

