



Acta Medica Colombiana

ISSN: 0120-2448

Asociacion Colombiana de Medicina Interna

MUÑOZ-LOMBO, JENNY PATRICIA; TABARES-BURBANO, ADRIÁN; OCAMPO-CHAPARRO, JOSÉ MAURICIO; CARVAJAL-ORTIZ, REYNALDO; CASANOVA-VALDERRAMA, MARÍA EUGENIA; REYES-ORTIZ, CARLOS ALFONSO

Multimorbidity and geriatric syndromes. Their effect on mortality in older adults with sepsis

Acta Medica Colombiana, vol. 47, no. 1, 2022, January-March, pp. 7-14

Asociacion Colombiana de Medicina Interna

DOI: <https://doi.org/10.36104/amc.2022.2125>

Available in: <https://www.redalyc.org/articulo.oa?id=163172407002>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Multimorbidity and geriatric syndromes

Their effect on mortality in older adults with sepsis

JENNY PATRICIA MUÑOZ-LOMBO, ADRIÁN TABARES-BURBANO, JOSÉ MAURICIO OCAMPO-CHAPARRO, REYNALDO CARVAJAL-ORTIZ, MARÍA EUGENIA CASANOVA-VALDERRAMA. CALI (COLOMBIA)
CARLOS ALFONSO REYES-ORTIZ • TALLAHASSEE (USA)

DOI: <https://doi.org/10.36104/amc.2022.2125>

Abstract

Introduction: sepsis is diagnosed in more than 60% of older adults (OAs) worldwide. These OAs often have multimorbidity and one of the geriatric syndromes, leading to physical, cognitive and psychosocial disability with consequently high healthcare costs, resulting in a serious public health problem.

Objective: to determine the impact of multimorbidity and geriatric syndromes on the 30-day mortality rate of OAs hospitalized for sepsis in an acute geriatric unit

Materials and methods: an observational, analytical, nested case-control study.

Results: 238 patients with a mean age of 83.15 ± 7.12 were analyzed; 52.1% were women and 99% had at least one comorbidity; the 30-day mortality was 34%. Urinary tract infection was the main cause of hospitalization (42.9%), with microbiological isolation achieved in 43.3% of cases and *Escherichia coli* being the most common causal agent (46.6%). Multiple logistic regression showed that chronic kidney disease (OR 2.1 95% CI 1.1-4.8; $p=0.037$), delirium (OR 3.1 95% CI 1.6-5.8; $p=0.001$) and disability (Barthel index <60 ; OR 3.4 95% CI 1.5-7.5; $p=0.002$) were significantly related to 30-day mortality in patients with sepsis admitted to an acute geriatric unit.

Conclusion: in OAs hospitalized for sepsis, multimorbidity, chronic kidney disease and geriatric syndromes (represented by delirium and disability) were the predictors of 30-day mortality. (*Acta Med Colomb* 2022; 47. DOI: <https://doi.org/10.36104/amc.2022.2125>).

Key words: *hospitalization, sepsis, geriatric syndrome, multimorbidity, comprehensive geriatric assessment, mortality.*

Dres. Jenny Patricia Muñoz-Lombo, Adrián Tabares-Burbano, Reynaldo Carvajal-Ortiz, María Eugenia Casanova-Valderrama: Universidad Libre, Facultad Ciencias de la Salud, Departamento de Medicina Interna, Grupo Interinstitucional de Medicina Interna (GIMI 1); Dr. José Mauricio Ocampo-Chaparro: Universidad Libre, Facultad Ciencias de la Salud, Departamento de Medicina Interna, Grupo Interinstitucional de Medicina Interna (GIMI 1). Universidad del Valle, Facultad de Salud, Departamento Medicina Familiar. Cali (Colombia); Dr. Carlos Alfonso Reyes-Ortiz: PhD. *Institute of Public Health, Behavioral Sciences & Health Education, College of Pharmacy and Pharmaceutical Sciences, Florida A & M University*. Tallahassee, Florida (USA). Correspondencia: Dr. Carlos A. Reyes-Ortiz. Tallahassee, Florida (USA). E-Mail: carlosareyesort@gmail.com, carlos.reyesortiz@fam.u.edu
Received: 26/II/2021 Accepted: 09/IX/2021

Introduction

Sepsis is defined as a potentially fatal organ dysfunction caused by the host's response to infection, whether in healthy or injured tissue, causing pathophysiological and biochemical disorders that cover a broad spectrum of clinical and laboratory manifestations ranging from bacteremia to septic shock (1). Early diagnosis and prompt, appropriate treatment are key for limiting damage and controlling the disease (2).

Considered to have one of the highest burdens of morbidity and mortality in the twenty-first century, sepsis very often affects older adults (OAs) in whom, despite advances in diagnosis and treatment, mortality continues to be high, causing more than 60% of hospital deaths in those over the age of 65 and up to 80% in those 80 years old or older (3-5).

Several factors predispose OAs to sepsis, resulting from typical aging changes such as oxidative stress, mitochondrial suppression, telomere shortening and DNA damage, which cause cellular disorders in cytokine regulation

and the neuroendocrine response. These modify protein metabolism which, coupled with low food intake, leads to malnutrition and changes in the intestinal microbiota, potentiating a proinflammatory state and immunosenescence which may cause or worsen neurological conditions, triggering an impaired functional status with disuse atrophy, sarcopenia secondary to accelerated muscle loss, sluggishness and fatigue (6).

Furthermore, OAs often have multimorbidity, defined as the conjunction of several chronic conditions which require rehabilitation or a long period of health care (7), which is frequently advanced and a significant risk factor for developing infectious diseases, especially pneumonia, urinary tract infections and skin and soft tissue infections (8). These infections may begin with atypical symptoms such as delirium, functional decline, weight loss, fatigue, dizziness and falls, symptoms which are part of geriatric syndromes and cause special challenges in diagnosis and treatment. This population should be cared for by an

interdisciplinary team (7, 9), using the interdisciplinary team work model, caring for patients at various care levels (acute units, mid-stay units, day hospitals, and home care).

According to epidemiological reports in the United States, despite medical advances, greater access to health care and an improved quality of life, sepsis continues to cause up to 30% mortality in hospitalized patients (3), which may reach 50% if there is septic shock. These percentages are comparable to the mortality caused by cardiovascular disease (10). The vast majority of survivors develop medical, mental and functional complications which lead to physical, cognitive and psychosocial disability (11, 12), with high costs for the healthcare system (13), thus becoming a public health issue (4). In 2017, Colombia reported 224,187 deaths, 62% of which were in people over the age of 65, with sepsis being the fifth cause of death, nationwide (14). Thus, the objective of this study was to determine the effect of multimorbidity and geriatric syndromes on mortality in OAs hospitalized for sepsis in an acute geriatric unit.

Materials and methods

Design

An observational, analytical nested case-control study was carried out in a cohort of patients diagnosed with sepsis and cared for in an acute geriatric unit.

The study protocol was approved as a no-risk study by the Technical-Scientific and Bioethical Committee of Universidad Libre and Clínica Universitaria Rafael Uribe Uribe. The patients or some caregivers were interviewed to obtain information during hospitalization. There were no conflicts of interest

Study population

Of the cohort of 2,054 OA patients 60 years old or older seen at Corporación Universitaria Rafael Uribe Uribe (CU-RUU) in Cali (Colombia) from January 2012 to September 2015, all met at least one of the following inclusion criteria for treatment in the acute geriatric unit (AGU): taking eight or more medications, a subacute cerebrovascular event, a disabling illness or prior severe or total functional dependence, frequent hospital readmissions (two or more per month), prior mental conditions (delirium on admission or baseline dementia), the presence of multiple geriatric syndromes (frailty, pressure ulcers, falls), body mass index less than 20 kg/m², an inadequate support network, living in a geriatric institution, or patients over the age of 80 with an acute medical disease. From this cohort of 2,054 patients, the records of those diagnosed with sepsis were reviewed, with cases being defined as all the records in which death occurred within 30 days of admission, and controls being the survivors. Pairing was done with one case for every two controls.

Informed consent was obtained from every patient or from a relative or caregiver, in cases where the OA was

unable to fill it out. Subsequently, the questionnaire was applied, which was completed on admission by one of the physicians on the interdisciplinary team trained to collect the standardized data from the comprehensive geriatric assessment (CGA) scales.

The data obtained was extracted from the geriatric interdisciplinary team's database, who assessed and treated the patient, family and caregivers, based on the biopsychosocial model, during their hospital stay. Thus, each professional involved applied geriatric clinical measurement scales from his/her discipline, as follows: geriatrician: information related to the biological domain (cause of hospitalization, comorbidities, death, medications); physical therapist: information related to the functional domain (activities of daily living); psychologist: information related to the mental domain (depression, delirium, cognitive impairment); and social worker: information related to the social and family domain (sociodemographic variables, the presence of a caregiver, relationship with the caregiver, and social/familial condition).

Outcome

Mortality within 30 days of admission to the AGU. The information on mortality was provided by the Valle del Cauca Departmental Secretariat of Health through death certificates up to October 2015.

Main independent variable

Sepsis was diagnosed based on clinical signs and symptoms of a systemic inflammatory response as a host response to infection associated with biological laboratory markers such as leukocytosis, neutrophilia, C-reactive protein and microbiological isolation, when possible, along with the support of imaging tools showing findings compatible with an infectious process (1, 15). It is important to keep in mind that the study was carried out in a hospital unit, and not in intermediate or intensive care, where variables such as SOFA or APACHE are not routinely used to determine the patients' severity of illness.

Covariables

The analysis included age, sex and known variables which could affect the outcome; laboratory values were used as numeric variables. The multimorbidity model was defined as the presence of two or more chronic conditions which cannot be cured but can be controlled using medications or other treatments; in other words, chronic diseases (16). Functional status was assessed using the Barthel Index (BI) which evaluates 10 basic activities of daily living, assigning a predetermined value for autonomy/independence on admission; the score ranges from 0 to 100, with 0 being maximum dependence and 100 total independence. The BI has been recommended as the instrument of choice for measuring physical disability and was used as a numeric and categorical variable. A BI score less than 60 was

considered to be disability (17,18), which was defined as a difficulty or limitation in carrying out basic activities of daily living (19).

For evaluating delirium, the Confusion Assessment Method (CAM) was used, which evaluates four characteristics: acute onset and fluctuating course, inattention, disordered thinking and a change in level of consciousness, with the presence of characteristics 1 and 2 and 3 or 4 being considered positive for delirium (20). Kidney function was assessed through the estimated glomerular filtration rate (GFR) using the equation derived from Levey et al.'s study (*Modification of Diet in Renal Disease Study equation [MDRD-4] or MDRD-IDMS*) (21), and chronic kidney disease (CKD) was defined as a GFR less than 60 ml/min/1.73 m² or kidney injury present for three months or more, based on the 2013 KDIGO definition. Kidney injury markers were not considered for the definition of CKD (22).

Analysis plan

Descriptive statistics were used to summarize the characteristics of the study population. Quantitative variables were presented using their measures of central tendency and dispersion. The Kolmogorov-Smirnov test was used to compare the distribution of the data. Qualitative variables were summarized as percentages and presented in frequency tables. The statistical significance of the bivariate associations with the outcome variable (30-day mortality) was analyzed using the Mann-Whitney U test for numerical variables and Chi² or Fisher's exact test for dichotomous variables, as applicable, taking a p value ≤ 0.05 as significant for rejecting the null hypotheses.

For exploring the multivariate association between the exposure variables and the outcome variable, the odds ratio (OR) was used as the measure of association, with its respective 95% confidence intervals (95%CI).

The morbidity and mortality and geriatric syndrome models were defined as theoretical constructs which encompass a set of interrelated aspects. The variables which showed a statistical significance <0.5 in the bivariate analysis were used to create these models. All the data was recorded in Excel and analyzed with IBMSPSS version 20.0 software.

Results

Of the 2,054 records of patients cared for in the AGU at Corporación Universitaria Rafael Uribe Uribe (CRUU), 238 met the inclusion criteria for analysis in this study and were selected. The reasons for exclusion are shown in Figure 1.

The average age was 83.2±7 years, and 52.1% were women. Ninety-nine percent of the patients had some chronic disease, with hypertension (73.9%) and diabetes mellitus (31.1%) being the most frequent. Likewise, cerebrovascular disease (CVD) and CKD had statistically significant differences between the groups in favor of cases (deceased patients), which was borderline for coronary disease (Table

1). The bivariate analysis for the geriatric syndromes model showed statistically significant differences for 30-day mortality after admission to the AGU: 48% for cases with delirium and 87% for cases with disability (Table 1).

Regarding the site of infection, urinary tract infection was found to be the most frequent cause of hospitalization, followed by pneumonia and soft tissue infections with 42.9, 38.7 and 16.0%, respectively. Microbiological isolation was achieved in 43.2% of the cases, 100% of which belonged to the Enterobacteriaceae family, with *Escherichia coli* being the most common causal agent in 46.6%, with a natural resistance pattern in 11.8% (Table 2).

In the multimorbidity model, the bivariate analysis showed that CKD (OR 2.5, 95% CI 1.2-5.5; p=0.017) and CVD (OR 2.1, 95% CI 1.1-4.0; p=0.023) were statistically significant variables (Table 3). For the geriatric syndromes model, delirium (OR 4.5, 95% CI 2.5-8.2; p<0.001) and disability (OR 5.6, 95% CI 2.7-11.9; p<0.001) (Table 4) were statistically significant.

The multiple regression analysis, which included all the significant variables from the multimorbidity and geriatric syndrome models, showed that CKD (OR 2.1, 95% CI 1.1-4.8, p=0.037), *delirium* (OR 3.1 95% CI 1.6-5.8; p=0.001) and disability (OR 3.4 95% CI 1.5-7.5; p=0.002) were statistically significantly associated with 30-day mortality in OAs hospitalized for sepsis in an AGU (Table 5).

Discussion

In this study, we evaluated the presence of multimorbidity and geriatric syndromes as risk factors for mortality in a Colombian OA population hospitalized for sepsis in an AGU. As seen in the results, CKD, delirium and disability were the main factors associated with 30-day mortality in OAs hospitalized for sepsis.

Studies have shown that age is an independent risk factor for mortality both for all causes as well as for sepsis (5, 12), also suggesting that multimorbidity plays an important role in this aspect (4, 23), as seen in a retrospective study evaluating the outcomes of adults with severe sepsis admitted to 171 ICUs in Australia and New Zealand. This study reported a 14% mortality rate in patients without comorbidities vs. 26.4% in patients with multimorbidity, which was statistically significant (p <0.001). In this same cohort, adults who were 44 years old or younger had a 7.3% mortality rate vs. 30.4% in adults 85 years old or older (p <0.001) (4). This agrees with the findings of our study, in which 30-day mortality was 34% and more than 99% of the patients had multimorbidity.

Furthermore, an increased risk of death has been found in patients with a history of CKD and lung disease (24-28), justifying admission to the ICU when the Charlson index is ≥ 3 (29). This may be due to an immunological disorder in patients with CKD, as they tend to develop leukocyte abnormalities, coagulation disorders and hypercoagulability states which lead to greater mortality in patients with sepsis

(30), similar to our study in which there were statistically significant differences between the mortality of people without CKD (8.9%) and those with CKD (19.8%), with a 110% greater chance of dying in OAs with sepsis and CKD compared with OAs with sepsis but without CKD.

Comprehensive geriatric assessment is considered to be the cornerstone of geriatric practice, an essential tool for approaching geriatric syndromes which could be the first sign of an infectious process in OAs (9), and an interdisciplinary teamwork model, caring for patients at different care levels (acute units, mid-stay units, day hospitals, home care). Within this group of syndromes, the role of delirium as a predictor of mortality stands out. Pisani et al. reported that for OAs 60 years old or older admitted to the ICU, the greater the number of days with delirium, the greater the risk of mortality in the year following admission (HR 1.10; 95% CI, 1.02–1.18) (31). Likewise, the meta-analysis published by Salluh et al. showed a greater mortality (RR 2.2, 95% CI from 1.78–2.70; $p < 0.001$), longer duration of mechanical ventilation ($p < 0.001$) and longer hospital stay both in the ICU ($p < 0.001$) as well as the general ward ($p < 0.001$) in patients who developed delirium during their hospital stay (32), along with greater mortality. This finding was seen in

our study, in which patients who developed delirium had a 2.1 times greater chance of dying than OAs without delirium.

Another condition included in the analyses was the presence of disability assessed using the BI. As indicated by Wu L W et al. in their 2016 cohort of 1,834 patients from 60 to 84 years old, disability doubles the risk of death from any cause (HR 2.3 [95% CI 1.3–3.9], $p < 0.005$) (33). Likewise, other studies have shown an association between disability and long-term mortality in patients hospitalized for infectious processes. In a retrospective study of Spanish patients between 75 and 84 years old hospitalized for pneumonia, Núria Torner et al. found that the factors significantly associated with 30-day mortality were: having any degree of disability (OR 3.7, 95% CI, 2.3–5.8; $p < 0.001$), age ≥ 85 years (OR 3.0, 95% CI, 1.7–5.3; $p < 0.001$), impaired cognitive function (OR 1.9, 95% CI, 1.2–3.1; $p = 0.005$) and ICU admission (OR 2.6, 95% CI, 1.7–5.2; $p = 0.009$) (34). Cillóniz C et al. also carried out a prospective observational study of 2,149 adult patients with pneumonia over a period of 12 years, whose results showed that mortality increased with age (65–74 years, 6.9%; 75–84 years, 8.9%; > 85 years, 17.1%; $P < 0.001$) and was associated with an increase in chronic conditions (neurological conditions causing dis-




Figure 1. Study flow chart.

ability; OR, 2.1; 95% CI; 1.5-2.1), pneumonia severity index IV or V (OR, 3.2; 95% CI, 1.8-6.0), bacteremia (OR, 1.7; 95% CI, 1.1-2.7) and the presence of multidrug-resistant germs (OR, 2.4; 95% CI, 1.3-4.3) (35). On the other hand,

Quintero et al., in their study of OA patients hospitalized for pneumonia in an AGU recorded a 30-day fatality of 26.3%; 93% of those who died of pneumonia were dependent prior to hospitalization, compared with OAs without dependence, with a statistically significant difference, indicating that the

Table 1. Sociodemographic, mental and functional characteristics of the study population.

Characteristics	General	Status after sepsis diagnosis		P value
	n (%) (n=238)	Deceased n (%) (n=81)	Alive n (%) n=157	
Age*	83.15±7.12	83.02±7.31	83.22±7.05	0.84 ⁱ
Male	114 (47.90)	42 (51.85)	72 (45.86)	0.38
HTN (n=234)	173 (73.93)	61 (75.31)	112 (73.20)	0.73
Hypothyroidism	35 (14.71)	10 (12.35)	25 (15.92)	0.46
Diabetes mellitus	74 (31.09)	25 (30.86)	49 (31.21)	0.96
Congestive heart failure	27 (11.34)	10 (12.35)	17 (10.83)	0.73
Coronary disease	16 (6.72)	9 (11.11)	7 (4.46)	0.05
Cerebrovascular disease	48 (20.17)	23 (28.40)	25 (15.92)	0.02
Atrial fibrillation	16 (6.72)	7 (8.64)	9 (5.3)	0.40
Chronic kidney disease	30 (12.61)	16 (19.75)	14 (8.92)	0.02
Arterial occlusive disease	11 (4.62)	5 (6.17)	6 (3.82)	0.52 ^f
Multimorbidity	236 (99.16)	80 (98.77)	156 (99.36)	1.00
Anemia	9 (3.78)	4 (4.94)	5 (3.18)	0.49 ^f
Dementia	43 (18.07)	17 (20.99)	26 (16.56)	0.40
Neoplasms	22 (9.24)	9 (11.11)	13 (8.28)	0.48
Chronic obstructive pulmonary disease	49 (20.59)	16 (19.75)	33 (21.02)	0.82
Polypharmacy	134 (56.30)	50 (61.73)	84 (53.50)	0.22
Delirium	66 (27.73)	39 (48.15)	27 (17.20)	0.00
Disability	158 (66.39)	71 (87.65)	87 (55.41)	0.00
Hospital stay (days)**	10 (6 – 15)	11 (5 – 21)	10 (7 – 14)	0.31 ^m

*Mean and standard deviation; **Median and interquartile range; ⁱStudent's T; ^mMann – Whitney – Wilcoxon; ^fFisher's exact test.

Table 2. Microbiological characteristics of the study population..

Characteristics	General	Status after the sepsis diagnosis		P value
	n (%) (n=238)	Deceased n (%) (n=81)	Living n (%) n=157	
Microbiological isolation	103 (43.28)	29 (35.80)	74 (47.13)	0.10
<i>Escherichia coli</i> : natural pattern	28 (11.76)	1 (1.23)	27 (17.20)	0.00
<i>Escherichia coli</i> : ESBL	20 (8.40)	6 (7.41)	14 (8.92)	0.69
<i>Klebsiella pneumoniae</i> : natural pattern	7 (2.94)	0 (0.00)	7 (4.46)	0.10 ^f
<i>Klebsiella pneumoniae</i> : ESBL	11 (4.62)	6 (7.41)	5 (3.18)	0.19 ^f
<i>Klebsiella pneumoniae</i> : KPC	3 (1.26)	3 (3.70)	0 (0.00)	0.04 ^f

^fFisher's exact test.
ESBL: Extended spectrum β-lactamases, KPC: *Klebsiella pneumoniae* carbapenemase

Table 3. Risk factors for 30-day mortality from sepsis. Results of the logistic regression of the model consisting of multimorbidity variables.

Multimorbidity	Cases (n=81) (%)	Controls (n= 157) (%)	OR	CI (95%)	P value
Cerebrovascular disease	28.4	15.9	2.1	1.1 – 4.0	0.023
Chronic kidney disease	19.8	8.9	2.5	1.2 – 5.5	0.017
Coronary disease	11.1	4.5	2.7	0.9 – 7.5	0.052
Diabetes mellitus	30.9	31.2	1.0	0.6 – 1.8	0.956
Hypertension	75.3	73.2	1.1	0.6 – 2.1	0.727
Arterial occlusive disease	6.2	3.8	1.7	0.5 – 5.6	0.671

Table 4. Risk factors for 30-day mortality from sepsis. Results of the logistic regression according to the model consisting of geriatric syndrome variables.

Geriatric syndrome	Cases (n=81) (%)	Controls (n= 157) (%)	OR	CI (95%)	P value
Delirium	48.1	17.2	4.5	2.5 – 8.2	<.0001
Dementia	21.0	16.6	1.3	0.7 – 2.6	0.400
Disability	87.7	55.4	5.7	2.7 – 11.9	<.0001
Polypharmacy	75.3	73.2	1.1	0.6 – 2.1	0.727

Table 5. Risk factors for 30-day mortality from sepsis. Results of the logistic regression according to the final model consisting of significant variables.

Variables	OR	CI (95%)	P value
Cerebrovascular disease	1.5	0.7 – 3.0	0.274
Chronic kidney disease	2.1	1.1 – 4.8	0.037
Delirium	3.1	1.6 – 5.8	0.001
Disability	3.4	1.5 – 7.5	0.002

independent predictors of mortality were: age 90 or above (RR=1.6; 95% CI: 1.1-2.7; p=0.04), multilobar involvement (RR=1.9; 95% CI: 1.1-3.3; p=0.02), elevated blood urea nitrogen (median ≥ 22.5 ; RR=3.9; 95% CI: 1.7-9.3; p<0.01), and a Lawton scale score of zero on admission (RR=3.2; 95% CI 1.1-9.8; p=0.04) (36).

With regard to the infectious process, 65-year-old OAs are more prone than young adults to have gram negative bacterial infections, as they have a 1.3 greater chance of being infected by these microorganisms (12). Within the Enterobacteriaceae family, *Escherichia coli* is the most commonly identified organism in urine cultures obtained from patients with urinary sepsis, occurring in 50% of cases, but other gram-negative bacteria may be present, such as *Proteus spp*, *Klebsiella spp* and *Pseudomonas spp* (37). These findings are similar to those of our study in which the most common causal agent was *Escherichia coli* in 46.6%.

Furthermore, a prospective cohort study of OAs found a 24% mortality rate for patients with negative blood cultures and 31% for those with positive blood cultures (38),

similar to our study, with the main source of infection being intra-abdominal (18.6%), followed by healthcare-associated pneumonia and community-acquired pneumonia with 17 and 12.4%, respectively (39), with the latter being the second cause of death in our study.

Despite the available reports, the data on the epidemiology and prognosis of sepsis in OAs in our setting is scant. This is partially explained by the fact that this population is excluded from most research studies (3, 12), despite age being a known risk factor which independently predicts mortality, early death after hospital admission, and institutionalization (5, 12, 40), all of which lends greater relevance to our study.

This study's limitations include its retrospective nature which prevents the analysis of other variables which could affect the outcome, leading to information bias.

In conclusion, in our study of OAs hospitalized for sepsis, the main factors associated with mortality within 30 days of admission to an AGU were delirium, functional decline and CKD. Therefore, it is essential to identify and intervene in these factors to impact on the mortality of OA patients hospitalized for sepsis.

Sources of financing

Financed by Universidad Libre, Cali branch.

References

- Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2016 [Internet]. Vol. 43, Intensive Care Medicine. 2017 [cited 2019 Oct 8]. 304–377 p. Available from: <http://link.springer.com/10.1007/s00134-017-4683-6>
- Rivers E, Nguyen B, Havstad S, Ressler J, Muzzin A, Knoblich B, et al. Early

- goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med*. 2001 Nov 8;345(19):1368–77.
3. Pinsky; DAL-ZLCC. Epidemiology of severe sepsis in the United States: Analysis of incidence, outcome, and associated costs of care. *Crit Care Med*. 2001;29(7):1303–10.
 4. Kaukonen KM, Bailey M, Suzuki S, Pilcher D, Bellomo R. Mortality related to severe sepsis and septic shock among critically ill patients in Australia and New Zealand, 2000–2012. *JAMA - J Am Med Assoc*. 2014;311(13):1308–16.
 5. Nasa P, Juneja D, Singh O, Dang R, Arora V. Severe sepsis and its impact on outcome in elderly and very elderly patients admitted in intensive care unit. Vol. 27, *Journal of Intensive Care Medicine*. 2012. p. 179–83.
 6. Destarac LA, Ely W. Sepsis in Older Patients: An Emerging Concern in Critical Care. *Adv SEPSIS*. 2002;2.
 7. Nicholson K, Makovski TT, Griffith LE, Raina P, Stranges S, van den Akker M. Multimorbidity and comorbidity revisited: refining the concepts for international health research [Internet]. Vol. 105, *Journal of Clinical Epidemiology*. Elsevier USA; 2019 [cited 2020 Sep 9]. p. 142–6. Available from: <http://www.jclinepi.com/article/S0895435618305432/fulltext>
 8. Martín S, Pérez A, Aldecoa C. Sepsis and immunosenescence in the elderly patient: A review. Vol. 4, *Frontiers in Medicine*. Frontiers Media S.A.; 2017.
 9. González Montalvo JI, Alarcón Alarcón T. Grandes síndromes geriátricos. Concepto y prevención de los más importantes. *Med - Programa Form Médica Contin Acreditado*. 2003 Jan;8(108):5778–85.
 10. Mayr FB, Yende S, Angus DC. Epidemiology of severe sepsis. Vol. 5, *Virulence*. Taylor and Francis Inc.; 2014. p. 1–11.
 11. Deutschman CS, Tracey KJ. Sepsis: Current dogma and new perspectives. Vol. 40, *Immunity*. Cell Press; 2014. p. 463–75.
 12. Martin GS, Mannino DM, Moss M. The effect of age on the development and outcome of adult sepsis. *Crit Care Med*. 2006;34(1):15–21.
 13. Mouton CP, Bazaldua O V, Pierce B, Espino D V. Common infections in older adults. *Am Fam Physician* [Internet]. 2001 Jan 15 [cited 2019 Oct 8];63(2):257–68. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11201692>
 14. Estadísticas Vitales. Boletín técnico (EEVV) [Internet]. 2018 [cited 2019 Oct 8]. Available from: https://www.dane.gov.co/files/investigaciones/poblacion/bt_estadisticasvitales_2018pre-29-junio-2018.pdf
 15. Seymour CW, Liu VX, Iwashyna TJ, Brunkhorst FM, Rea TD, Scherag A, et al. The third international consensus definitions for sepsis and septic shock (sepsis-3). *JAMA - J Am Med Assoc*. 2016; Feb 23;315(8):801–10.
 16. Yarnall AJ, Sayer AA, Clegg A, Rockwood K, Parker S, Hindle J V. New horizons in multimorbidity in older adults. *Age Ageing* [Internet]. 2017 [cited 2020 Dec 16];46(6):882–8. Available from: <http://pmc/articles/PMC5860018/?report=abstract>
 17. Mahoney, F. I. Barthel DW. Functional evaluation: the barthel index. *Md State Med J* [Internet]. 1965 Feb 1 [cited 2020 Jun 22];14:61–5. Available from: <https://europepmc.org/article/med/14258950>
 18. Cid-Ruza J, Damián-Moreno J. Valoración de la discapacidad física: el índice de Barthel. Vol. 71, *Rev Esp Salud Púhca*. 1997.
 19. OMS I Discapacidades. WHO [Internet]. 2016 [cited 2020 Dec 16]; Available from: <http://www.who.int/topics/disabilities/es/>
 20. Inouye SK, Van Dyck CH, Alessi CA, Balkin S, Siegel AP, Horwitz RI. Clarifying confusion: The confusion assessment method: A new method for detection of delirium. *Ann Intern Med* [Internet]. 1990 Dec 1 [cited 2020 Jun 22];113(12):941–8. Available from: <http://europepmc.org/article/med/2240918>
 21. Levey AS, Greene T, Sarnak MJ, Wang X, Beck GJ, Kusek JW, et al. Effect of Dietary Protein Restriction on the Progression of Kidney Disease: Long-Term Follow-Up of the Modification of Diet in Renal Disease (MDRD) Study. *Am J Kidney Dis* [Internet]. 2006 Dec 1 [cited 2020 Jun 22];48(6):879–88. Available from: <http://www.ajkd.org/article/S0272638606013722/fulltext>
 22. KDIGO 2012. Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Off J Int Soc Nephrol* [Internet]. 2013 [cited 2021 Aug 22];3(1). Available from: www.publicationethics.org
 23. Holder AL, Gupta N, Lulaj E, Furgieue M, Hidalgo I, Jones MP, et al. Predictors of early progression to severe sepsis or shock among emergency department patients with nonsevere sepsis. *Int J Emerg Med* [Internet]. 2016 Dec [cited 2019 Oct 10];9(1):10. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26908009>
 24. Knaus WA, Wagner DP, Draper EA, Zimmerman JE, Bergner M, Bastos PG, et al. The APACHE III prognostic system: Risk prediction of hospital mortality for critically III hospitalized adults. *Chest*. 1991;100(6):1619–36.
 25. Tran DD, Groeneveld AB, van der Meulen J, Nauta JJ, Strack van Schijndel RJ, Thijs LG. Age, chronic disease, sepsis, organ system failure, and mortality in a medical intensive care unit. *Crit Care Med* [Internet]. 1990 May [cited 2019 Oct 10];18(5):474–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/2328591>
 26. Pittet D, Thiévent B, Wenzel RP, Li N, Gurman G, Suter PM. Importance of pre-existing co-morbidities for prognosis of septicemia in critically ill patients. *Intensive Care Med* [Internet]. 1993 [cited 2019 Oct 10];19(5):265–72. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8408935>
 27. Novosad SA, Sapiiano MRP, Grigg C, Lake J, Robyn M, Dumyati G, et al. Vital Signs: Epidemiology of Sepsis: Prevalence of Health Care Factors and Opportunities for Prevention. *MMWR Morb Mortal Wkly Rep* [Internet]. 2016 Aug 26 [cited 2020 Sep 9];65(33):864–9. Available from: <http://www.cdc.gov/mmwr/volumes/65/wr/mm6533e1.htm>
 28. Doi K. Role of kidney injury in sepsis [Internet]. Vol. 4, *Journal of Intensive Care*. BioMed Central Ltd.; 2016 [cited 2020 Sep 9]. Available from: <http://pmc/articles/PMC4804517/?report=abstract>
 29. Oltean S, Tatulescu D, Bondor C, Slavcovici A, Cismaru C, Lupşu M, et al. Charlson's weighted index of comorbidities is useful in assessing the risk of death in septic patients. *J Crit Care*. 2012 Aug;27(4):370–5.
 30. Bermejo-Martin J, Martín-Fernandez M, López-Mestanza C, Duque P, Almansa R. Shared Features of Endothelial Dysfunction between Sepsis and Its Preceding Risk Factors (Aging and Chronic Disease). *J Clin Med* [Internet]. 2018 Oct 30 [cited 2020 Sep 9];7(11):400. Available from: <http://pmc/articles/PMC6262336/?report=abstract>
 31. Pisani MA, Kong SYJ, Kasl S V., Murphy TE, Araujo KLB, Van Ness PH. Days of delirium are associated with 1-year mortality in an older intensive care unit population. *Am J Respir Crit Care Med*. 2009 Dec 1;180(11):1092–7.
 32. Salluh JIF, Wang H, Schneider EB, Nagaraja N, Yenokyan G, Damluji A, et al. Outcome of delirium in critically ill patients: Systematic review and meta-analysis. *BMJ*. 2015 Jun 3;350:1–10.
 33. Wu L-W, Chen W-L, Peng T-C, Chiang S-T, Yang H-F, Sun Y-S, et al. All-cause mortality risk in elderly individuals with disabilities: a retrospective observational study. [cited 2020 Apr 10]; Available from: <http://bmjopen.bmj.com/>
 34. Torner N, Izquierdo C, Soldevila N, Toledo D, Chamorro J, Espejo E, et al. Factors associated with 30-day mortality in elderly inpatients with community acquired pneumonia during 2 influenza seasons. *Hum Vaccines Immunother*. 2017;13(2):450–5.
 35. Cillóniz C, Polverino E, Ewig S, Aliberti S, Gabarrús A, Menéndez R, et al. Impact of age and comorbidity on cause and outcome in community-acquired pneumonia. *Chest* [Internet]. 2013 Sep [cited 2019 Oct 10];144(3):999–1007. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23670047>
 36. Quintero-González DC, Ocampo JM, Reyes-Ortiz CA. Factors associated with lethality from pneumonia in an acute care for the elderly unit: a retrospective cohort. *Biomedica* [Internet]. 2020 Dec 2 [cited 2020 Dec 21];40(4):734–48. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/33275351>
 37. Leliggowicz A, Dodek PM, Norena M, Wong H, Kumar A, Kumar A. Association between source of infection and hospital mortality in patients who have septic shock. *Am J Respir Crit Care Med* [Internet]. 2014 May 15 [cited 2020 Apr 10];189(10):1204–13. Available from: <http://www.atsjournals.org/doi/abs/10.1164/rccm.201310-1875OC>
 38. Jaimes, Fabian; Garces, Jenny; Cuervo, Jorge; Ramirez, Federico; Ramirez, Jorge; Quintero, Claudia; Vargas, Andrea; Zapata, Lader; Ochoa, Jorge; Yepes, Mercedes; Leal, Hiuber; Tandioy, Fabio; Estrada JC. NoFactores pronosticos en el síndrome de respuesta inflamatoria sistémica (SRIS): Desarrollo de un índice de severidad / Prognosis factors in SIRS. *Acta méd colomb* [Internet]. 2001 [cited 2019 Oct 8];26(4):149–57. Available from: <http://www.actamedicacolombiana.com/anexo/articulos/04-2001-04.pdf>
 39. Ortíz G, Dueñas C, Rodríguez F, Barrera L, de La Rosa G, Dennis R, et al. Epidemiology of sepsis in Colombian intensive care units. *Biomedica*. 2014;34(1):40–7.
 40. Azoulay E, Adrie C, De Lassence A, Pochard F, Moreau D, Thiery G, et al. Determinants of postintensive care unit mortality: a prospective multicenter study. *Crit Care Med* [Internet]. 2003 Feb [cited 2019 Oct 10];31(2):428–32. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12576947>

