



Acta Agronómica

ISSN: 0120-2812

ISSN: 2323-0118

Universidad Nacional de Colombia

Chan-Cupul, Wilberth; Jiménez-Barrera, Dulce Guadalupe;  
Lezama-Gutiérrez, Roberto; Ruiz-Sánchez, Esaú; Pérez-Zamora,  
Octavio; Buenrostro-Nava, Marco Tulio; Manzo-Sánchez, Gilberto  
Screening of nutritional factors through a Plackett-Burman experimental design  
during the solubilization of tricalcium phosphate by *Penicillium hispanicum*  
Acta Agronómica, vol. 68, no. 2, 2019, April-June, pp. 151-159  
Universidad Nacional de Colombia

DOI: <https://doi.org/10.15446/acag.v68n2.61862>

Available in: <https://www.redalyc.org/articulo.oa?id=169960781010>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UDEM [redalyc.org](https://www.redalyc.org)

Scientific Information System Redalyc  
Network of Scientific Journals from Latin America and the Caribbean, Spain and  
Portugal

Project academic non-profit, developed under the open access initiative

# Screening of nutritional factors through a Plackett-Burman experimental design during the solubilization of tricalcium phosphate by *Penicillium hispanicum*

## Detección de factores nutricionales a través de un diseño experimental de Plackett-Burman durante la solubilización de fosfato tricálcico por *Penicillium hispanicum*

Wilberth Chan Cupul<sup>\*1</sup>, Dulce Guadalupe Jiménez Barrera<sup>1</sup>, Roberto Lezama Gutiérrez<sup>1</sup>, Esaú Ruiz Sánchez<sup>2</sup>, Octavio Pérez Zamora<sup>1</sup>, Marco Tulio Buenrostro Nava<sup>1</sup>, Gilberto Manzo-Sánchez<sup>1</sup>

<sup>1</sup>Universidad de Colima, México. <sup>2</sup>Instituto Tecnológico de Conkal, Yucatán, México.

Rec.: 01.08.2017 Acep.: 12.02.2019.

### Abstract

Phosphorus plays a fundamental role in plant metabolic processes. Lack of phosphorus in agricultural soils is compensated with chemical fertilizers, which increase the cost of production and contaminate the environment. One approach to avoid this situation is the use of P-solubilizing fungi as biofertilizers. In this study, the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$  by five native soil-borne micromycetes (SBM) isolated from the rhizosphere of *Carica papaya* plants was evaluated. The significant nutritional factors involved in the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$  by *Penicillium hispanicum* were determined through a Plackett-Burman experimental design (PBED). Both *Penicillium* SP1 (108.5 mg/L) and *P. hispanicum* (106.5 mg/L) solubilized 100% of  $\text{Ca}_3(\text{PO}_4)_2$  at six days of culture. All studied SBM decreased the pH of the culture medium from 4.15 (control) to <2.45. The PBED revealed that the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$  by *P. hispanicum* was significantly influenced ( $P < 0.05$ ) by low values of  $\text{MnSO}_4$  (0.002 g/L),  $(\text{NH}_4)_2\text{SO}_4$  (0.25 g/L) and pH (4.0) and a high amount of  $\text{Ca}_3(\text{PO}_4)_2$  (0.75 g/L). Total protein content was influenced significantly ( $P < 0.05$ ) by low amounts of glucose (5.0 g/L), yeast extract (0.25 g/L), inoculum ( $1 \times 10^5$  conidia/mL) and pH (4.0) and high amounts of  $\text{MgSO}_4$  (0.15 g/L),  $\text{FeSO}_4$  (0.003 g/L) and  $\text{Ca}_3(\text{PO}_4)_2$  (0.75 g/L). Meanwhile, biomass production was significantly ( $P < 0.05$ ) influenced by a high level of glucose (15.0 g/L) and inoculum amount ( $1 \times 10^7$  conidia/mL). In conclusion, *P. hispanicum* is a good candidate for further field experiments to test  $\text{Ca}_3(\text{PO}_4)_2$  solubilization in poor soils that lack C and N sources.

**Keywords:** Biomass, *Carica papaya*, optimization, rhizosphere, soil borne micromycetes, tricalcium phosphate.

### Resumen

El fósforo es fundamental en los procesos metabólicos de las plantas. La falta de fósforo en los suelos agrícolas se compensa con fertilizantes químicos, incrementando el costo de producción y contaminando el medio ambiente. Una alternativa ante esta situación es el uso de hongos solubilizadores de P como biofertilizantes. En este estudio se evaluó la solubilización del  $\text{Ca}_3(\text{PO}_4)_2$  por cinco micromicetos nativos del suelo (MNS) aislados de la rizosfera de plantas de *Carica papaya*. Además, se determinaron los factores nutricionales significativos implicados en la solubilización de fosfato tricálcico - $\text{Ca}_3(\text{PO}_4)_2$ - por *Penicillium hispanicum* a través de un diseño experimental Plackett-Burman (DEPB). Tanto *Penicillium* SP1 (108.5 mg/L) como *P. hispanicum* (106.5 mg/L) solubilizaron al 100% el  $\text{Ca}_3(\text{PO}_4)_2$ , a los 6 días de cultivo. Todos los MNS evaluados disminuyeron el pH del medio de cultivo de 4.15 a <2.45. El DEPB reveló que la solubilización de  $\text{Ca}_3(\text{PO}_4)_2$  por *P. hispanicum* fue influenciada ( $P < 0.05$ ) por valores bajos de  $\text{MnSO}_4$  (0.002 g/L),  $(\text{NH}_4)_2\text{SO}_4$  (0.25 g/L) y pH (4.0) y un valor alto de  $\text{Ca}_3(\text{PO}_4)_2$  (0.75 g/L). El contenido de proteína fue influenciado ( $P < 0.05$ ) por valores bajos de glucosa (5.0 g/L), extracto de levadura (0.25 g/L), inóculo ( $1 \times 10^5$  conidios/mL) y pH (4.0) y valores altos de  $\text{MgSO}_4$  (0.15 g/L),  $\text{FeSO}_4$  (0.003 g/L) y  $\text{Ca}_3(\text{PO}_4)_2$  (0.75 g/L). Finalmente, la producción de biomasa fue influenciada ( $P < 0.05$ ) por valores altos de glucosa (15.0 g/L) e inóculo ( $1 \times 10^7$  conidios/mL). *Penicillium hispanicum* es un buen candidato para futuros experimentos en campo para la solubilización del  $\text{Ca}_3(\text{PO}_4)_2$  en suelos con limitado C y N.

**Palabras clave:** biomasa, *Carica papaya*, optimización, rizosfera, hongos microscópicos de suelo, fosfato tricálcico.

## Introduction

Phosphorus (P) is one of the most important macronutrients for plant growth and development and is a key nutrient for achieving sustainable agriculture. Phosphorus plays a fundamental role in all major metabolic processes in plants including photosynthesis, macromolecular biosynthesis, energy transfer, signal transduction and respiration. In nature, tricalcium phosphate ( $\text{Ca}_3(\text{PO}_4)_2$ ) is one of the main P reserves; however, cultivated plants are unable to use this chemical form (Sharma et al., 2013).

To release P from  $\text{Ca}_3(\text{PO}_4)_2$ , the activity of soil microorganisms such as bacteria, soil borne micromycetes and mycorrhiza is required. In agricultural soils with low abundance and diversity of P-solubilizing microorganisms, bio-fertilizing microorganisms can be directly applied using a bioaugmentation strategy (Khan et al., 2014). In this sense, soil borne micromycetes have been described as suitable microorganisms for releasing P from inorganic sources such as  $\text{Ca}_3(\text{PO}_4)_2$ ,  $\text{FePO}_4$  and  $\text{AlPO}_4$  (Oliveira-Mendes et al., 2016; Zúñiga-Silva et al., 2016).

Solubilization of inorganic phosphates is gaining greater attention in organic and sustainable agriculture; several soil borne micromycetes have been isolated for this purpose (Hernández-Leal et al., 2011; Hernando-Posada et al., 2012, 2013). These isolates can be inoculated directly into the soil as biofertilizers or can be used in bioreactors to solubilize rock phosphates or other inorganic phosphates in the production of P fertilizers (Khan et al., 2014).

However, to produce P fertilizers from inorganic P sources using fungi, a suitable fungal strain and an optimized culture media are required (Oliveira-Mendes et al., 2016). Some studies have focused on the optimization of inorganic phosphate solubilization using liquid fermentation, considering one independent variable in addition to other variables fixed at a certain level. These single experimental designs are usually laborious and time consuming and screen few experimental factors (Scervino et al., 2011). For instance, Relwani et al. (2008) used Pikovskaya broth to study the influence of C and N source on the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$  by a mutant strain of *Asperigillus tubingensis* (AT1). The study reported that glucose and sucrose promoted  $\text{Ca}_3(\text{PO}_4)_2$  solubilization compared with fructose, lactose and xylose. In addition, potassium nitrate increased  $\text{Ca}_3(\text{PO}_4)_2$  solubilization in comparison with ammonium sulfate, ammonium nitrate, asparagine and tryptophan. In other study, Chun-Qiao et al. (2008) optimized the NBRIP culture medium for rock phosphate solubilization using soil

microfungi isolated from phosphate mines. The authors reported that the optimum culture conditions for rock phosphate solubilization were 32 °C, 160 r.p.m., 2.5 g/L of rock phosphate and 0.5 mg/mL of EDTA. In addition, *Penicillium expansum* resulted in the highest soluble P at a C:N ratio of 40:1. In the studies of both Relwani et al. (2008) and Chun-Qiao et al. (2008) factorial experimental designs were used. Thus, the best culture conditions could not be determined through this experimental design. In comparison, Plackett-Burman experimental design (PBED) represents one means of identifying the main effects and the efficiency of a large number of factors on a specific response (Plackett and Burman, 1946).

Accordingly, we considered this experimental design to be useful in the screening of a large number of factors during the optimization of the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$  by soil-borne micromycetes. The objectives of this study were: (1) to evaluate the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$  by native soil borne micromycetes isolated from the rhizosphere of *Carica papaya* plants, and (2) to determine the significant nutritional factors involved in the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$  by the best P-solubilizing fungi using a Plackett-Burman experimental design.

## Materials and methods

### Fungal sources

Soil-borne micromycetes were isolated from the rhizosphere of *C. papaya* plants, located in an organic orchard in Tecoman, Colima, Mexico (location: 18°50'16.8" N, 103°50'06.4" W). Isolation was carried out by washing the soil particles, which consisted in washing one gram of soil with three liters of sterile distilled water using three sieves (Alcon®, Mexico) of decreasing mesh size, one on top of the other, of 500, 250 and 150 µm, respectively. Washed particles (150 µm) were deposited on sterile filter paper discs for 24 h to remove excess moisture. Ten particles were seeded on potato dextrose agar (PDA) + chloramphenicol (150 ppm, Sigma-Aldrich®, USA). Five Petri dishes were incubated at 25 °C and 75% relative humidity (RH). The fungal colonies were purified by re-isolation in PDA. Five fungal isolates were used to evaluate the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$  in a Pikovskaya culture medium.

### Molecular identification

Only *Penicillium* isolate SP16 was identified at molecular level using the methodology described

by Zúñiga-Silva *et al.* (2016). After genomic DNA extraction, PCR amplification of the internal transcribed spacers (ITS regions: 26S and 16S regions) were carried out using the following primers ITS1 (5'-TCCGTAGGTGAACCTGCGG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3'). Purified PCR products were sequenced by Genomic Biotechnology Center—IPN (Reynosa, Tamaulipas, Mexico) using standard protocols. DNA sequence analyses were performed using the BLAST program, run against the NCBI database (<https://www.ncbi.nlm.nih.gov/>). The Index Fungorum (<http://www.indexfungorum.org/>) was used as a species authority.

### Solubilization of $\text{Ca}_3(\text{PO}_4)_2$ by fungal isolates

Pikovskaya broth was prepared according to Hernández-Leal *et al.* (2011). In Erlenmeyer flasks (250 mL), 120 mL of Pikovskaya broth was added and then autoclaved at 120 °C and 125 lb of pressure during 15 min. Soil-borne micromycetes were inoculated into the flasks using 50 µL of a conidia solution ( $1 \times 10^6$  conidia/mL). Inoculated flasks were incubated for six days at 25 °C, 75% relative humidity (RH) and 12/12 h light/darkness. Five fungal isolates were tested; four independent replicates of each isolate were settled. Every 48 h, aliquots of 10 mL of supernatant were obtained aseptically in a laminar flow cabinet. The supernatant was filtered in nylon membrane disks (0.5 µm), avoiding fungal biomass. Free cell extracts were used to measure soluble P, total protein content and pH.

### Studied variables

Soluble P was measured using a spectrophotometric method during the reduction of phosphomolybdenum complexes to ascorbic acid (Watanabe and Olsen, 1965). Readings were made at 880 nm in a UV-Vis spectrophotometer (Luzeren®, model L6S, St. Louis, MO, USA). The standard curve for soluble phosphorus was:  $\text{P-soluble} = [\text{absorbance} - 0.1000] / 0.009074$ ,  $r^2 = 0.9976$ . As the P standard,  $\text{KH}_2\text{PO}_4$  was used.

Total protein was measured according to the Bradford method. The standard curve to obtain total protein from the free cell extracts was:  $\text{protein content} = [\text{absorbance} - 0.4306] / 0.1188$ ,  $r^2 = 0.9915$ . As the protein standard, egg albumin was used.

To measure the pH of fungal extracts, a potentiometer OHAUS® (model Starter, Parsippany NJ, USA) was used. Finally, for fungal biomass quantification, mycelium from the five fungal screening cultures and the PBED were harvested by filtering the culture medium with

a pre-weighed Whatman No. 1 filter paper and determining the difference in mass after drying over three days at 40 °C.

### Plackett-Burman experimental design

To determine the significant nutritional factors that influence  $\text{Ca}_3(\text{PO}_4)_2$  solubilization and fungal growth of *P. hispanicum* (isolate SP16), a PBED was implemented. The statistical screening was based on the main effects of the experimental factors, although their interaction effects were not considered. Pikovskaya broth was modified to evaluate the influence of different nutritional factors on  $\text{Ca}_3(\text{PO}_4)_2$  solubilization, total protein and biomass production. Eleven factors were assessed: the components of Pikovskaya broth (9 nutrients), inoculum amount (conidia/mL) and pH (Table 1). Each screened factor was set at two levels, coded as high (+1) and low (-1). Four central points (0) were screened by running 16 experiments, as shown in Table 2. The PBED was executed three times, and the average of each experiment was used for the statistical analysis. Factors significant at the 5% level ( $P < 0.05$ ) in the regression analysis were considered to have a significant impact on the response variables. Equation 1 was used to fit the experimental data, which includes the individual effects of each studied factor. The PBED, the response variables were P-solubilization, total protein content and biomass production, which were measured as previously described.

**Table 1.** Original and coded levels of the independent variables used in the Plackett-Burman experimental design.

Code	Factor	Low level (-1)	Level 0	High level (+1)
G	D-Glucose (g/L)	5	10	15
Ye	Yeast extract (g/L)	0.25	0.5	0.75
NH	$(\text{NH}_4)_2\text{SO}_4$ (g/L)	0.25	0.5	0.75
K	KCl (g/L)	0.1	0.2	0.3
Mg	$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (g/L)	0.05	0.1	0.15
Mn	$\text{MnSO}_4 \cdot \text{H}_2\text{O}$ (g/L)	0.002	0.004	0.006
Na	NaCl (g/L)	0.1	0.2	0.3
Fe	$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (g/L)	0.001	0.002	0.003
Ca	$\text{Ca}_3(\text{PO}_4)_2$	0.25	0.5	0.75
I	Inoculum (conidia/mL)	$1 \times 10^5$	$1 \times 10^6$	$1 \times 10^7$
pH	pH	4.0	4.5	5.0

**Table 2.** Plackett-Burman experimental design matrix for evaluating the influence of nutrimental factors for  $\text{Ca}_3(\text{PO}_4)_2$  solubilization, total protein content and biomass.

Exp.	Experimental factors											Responses					
												P-soluble (mg/L)		Total protein (mg/L)		Biomass (mg/flask)	
	G	Ye	NH	K	Mg	Mn	Na	Fe	Ca	I	pH	Observed*	Pre-dicted	Observed*	Pre-dicted	Observed*	Pre-dicted
1	15	0.25	0.75	0.3	0.05	0.006	0.1	0.001	0.25	$1 \times 10^7$	5	4.0	7.3	19.9	12.4	0.39	0.43
2	5	0.25	0.25	0.3	0.15	0.006	0.1	0.003	0.75	$1 \times 10^5$	5	66.1	69.4	182.9	175.4	0.11	0.15
3	15	0.75	0.75	0.1	0.15	0.006	0.1	0.003	0.25	$1 \times 10^5$	4	33.8	37.2	47.0	39.6	0.26	0.30
4	5	0.25	0.75	0.3	0.15	0.002	0.3	0.003	0.25	$1 \times 10^7$	4	36.6	39.9	82.2	74.7	0.28	0.32
5	5	0.75	0.25	0.1	0.05	0.006	0.3	0.003	0.25	$1 \times 10^7$	5	23.7	27.1	59.4	51.9	0.21	0.25
6	5	0.75	0.75	0.3	0.05	0.006	0.3	0.001	0.75	$1 \times 10^5$	4	58.1	61.5	51.8	44.3	0.15	0.19
7	5	0.75	0.75	0.1	0.15	0.002	0.1	0.001	0.75	$1 \times 10^7$	5	70.6	73.9	71.8	64.3	0.25	0.29
8	15	0.25	0.25	0.1	0.15	0.006	0.3	0.001	0.75	$1 \times 10^7$	4	74.5	77.9	52.4	45.0	0.28	0.32
9	5	0.25	0.25	0.1	0.05	0.002	0.1	0.001	0.25	$1 \times 10^5$	4	47.4	50.7	55.9	48.5	0.09	0.13
10	15	0.25	0.75	0.1	0.05	0.002	0.3	0.003	0.75	$1 \times 10^5$	5	66.8	70.1	114.0	106.8	0.22	0.26
11	15	0.75	0.25	0.3	0.05	0.002	0.1	0.003	0.75	$1 \times 10^7$	4	75.8	79.2	58.3	50.8	0.39	0.43
12	15	0.75	0.25	0.3	0.15	0.002	0.3	0.001	0.25	$1 \times 10^5$	5	40.7	44.0	39.9	32.4	0.31	0.35
13	10	0.5	0.5	0.2	0.1	0.004	0.2	0.002	0.5	$1 \times 10^6$	4.5	56.9	53.2	28.6	62.2	0.43	0.29
14	10	0.5	0.5	0.2	0.1	0.004	0.2	0.002	0.5	$1 \times 10^6$	4.5	62.7	53.2	37.7	62.2	0.42	0.29
15	10	0.5	0.5	0.2	0.1	0.004	0.2	0.002	0.5	$1 \times 10^6$	4.5	68.1	53.2	48.6	62.2	0.37	0.29
16	10	0.5	0.5	0.2	0.1	0.004	0.2	0.002	0.5	$1 \times 10^6$	4.5	65.5	53.2	44.4	62.2	0.48	0.29

Prroduction of *Penicillium hispanicum*. \*Observed values were the mean of three replicates. Exp.=Experiments.

$$Y = \beta_0 + \sum \beta_i X_i (i = 1, 2, 3 \dots k), \text{ Eq. 1}$$

where:

Y = Response (soluble P, total protein, biomass)

X = Independent variable

$\beta_0$  = Model intercept

$\beta_i$  = Estimated variable

## Data analysis

An analysis of variance (ANOVA) and a multiple range comparison test (least significant difference,  $P = 0.05$ ) were applied to the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$ , total protein content and biomass production in the screening of the five soil-borne micromycetes. The PBED was designed and analyzed using the StatGraphics Plus® software.

## Results

### Fungal identification

According to sequence analysis, the isolate *Penicillium* SP16 was identified as *P. hispanicum* (Ascomycota: Trichocomaceae) C. Ramírez, A. T. Martínez and Ferrer, with 99% of similarity to the accessions: NR\_138307.1, JX841247.7 and KJ775637.1 reported in the gen bank.

### Solubilization of $\text{Ca}_3(\text{PO}_4)_2$ by soil-borne micromycetes

Soil-borne micromycetes isolated from the *C. papaya* rhizosphere showed statistical differences in the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$ . At 48 h of culture, *P. hispanicum* achieved the highest ( $F = 36.56$ ,  $P = 0.00001$ ) soluble P at 90.4 mg/mL; in contrast, *Penicillium* SP11 (15.5 mg/L) solubilized the lowest amount of  $\text{Ca}_3(\text{PO}_4)_2$  (Table 3). During



the second evaluation period (96 h), *P. hispanicum* maintained the highest  $\text{Ca}_3(\text{PO}_4)_2$  solubilization at 96.6 mg/L compared with that of the other studied fungal isolates ( $F = 37.52$ ,  $P = 0.00001$ ). At the end of experiment, *Penicillium* SP1 (108.8 mg/L) and *P. hispanicum* (106.5 mg/L) solubilized all soluble P contained in  $\text{Ca}_3(\text{PO}_4)_2$ , or 98.6 mg/L, according to the molecular weight of  $\text{Ca}_3(\text{PO}_4)_2$ . In contrast, both *Beauveria* SP9 and *Penicillium* SP18 demonstrated the lowest solubilization of  $\text{Ca}_3(\text{PO}_4)_2$  ( $F = 3828.85$ ,  $P = 0.00001$ ), solubilizing 66.2 and 62.2 mg/L of P, respectively.

**Table 3.** Solubilization of  $\text{Ca}_3(\text{PO}_4)_2$ , total protein content and change in pH in the culture of soil borne micromycetes isolated from *Carica papaya* rhizosphere.

Isolate	P-soluble (mg/mL)		
	48 h	96 h	144 h
<i>Penicillium</i> SP1	0.0 <sup>d</sup>	78.9 ± 3.3 <sup>b</sup>	108.5 ± 8.1 <sup>a</sup>
<i>Beauveria</i> SP9	25.8 ± 0.5 <sup>c</sup>	67.9 ± 2.4 <sup>c</sup>	66.2 ± 5.1 <sup>c</sup>
<i>Penicillium</i> SP11	15.5 ± 0.3 <sup>d</sup>	81.3 ± 4.0 <sup>b</sup>	89.3 ± 6.7 <sup>b</sup>
<i>P. hispanicum</i>	90.4 ± 4.3 <sup>a</sup>	96.6 ± 0.9 <sup>a</sup>	106.5 ± 5.4 <sup>ab</sup>
<i>Penicillium</i> SP18	63.4 ± 6.5 <sup>b</sup>	73.9 ± 1.2 <sup>bc</sup>	62.2 ± 3.3 <sup>c</sup>
Control	ND	ND	ND

Isolate	Total protein (mg/L)		
	48 h	96 h	144 h
<i>Penicillium</i> SP1	17.8 ± 0.53 <sup>bc</sup>	16.8 ± 0.1 <sup>c</sup>	15.9 ± 0.4 <sup>e</sup>
<i>Beauveria</i> SP9	19.0 ± 0.12 <sup>b</sup>	22.8 ± 1.8 <sup>b</sup>	30.7 ± 0.7 <sup>d</sup>
<i>Penicillium</i> SP11	18.5 ± 0.60 <sup>b</sup>	10.8 ± 0.6 <sup>d</sup>	75.2 ± 0.1 <sup>a</sup>
<i>P. hispanicum</i>	11.1 ± 0.50 <sup>d</sup>	19.6 ± 0.6 <sup>c</sup>	11.9 ± 0.3 <sup>f</sup>
<i>Penicillium</i> SP18	22.8 ± 0.16 <sup>a</sup>	28.6 ± 0.5 <sup>a</sup>	47.1 ± 0.2 <sup>b</sup>
Control	—	—	—

Isolate	pH		
	48 h	96 h	144 h
<i>Penicillium</i> SP1	2.54 ± 0.04 <sup>c</sup>	2.67 ± 0.05 <sup>b</sup>	2.45 ± 0.03 <sup>bc</sup>
<i>Beauveria</i> SP9	2.48 ± 0.04 <sup>cd</sup>	2.66 ± 0.05 <sup>b</sup>	2.53 ± 0.04 <sup>b</sup>
<i>Penicillium</i> SP11	2.59 ± 0.08 <sup>c</sup>	2.46 ± 0.02 <sup>b</sup>	2.41 ± 0.02 <sup>bc</sup>
<i>P. hispanicum</i>	2.38 ± 0.02 <sup>d</sup>	2.53 ± 0.02 <sup>b</sup>	2.51 ± 0.02 <sup>b</sup>
<i>Penicillium</i> SP18	2.83 ± 0.05 <sup>b</sup>	2.56 ± 0.03 <sup>b</sup>	2.35 ± 0.01 <sup>c</sup>
Control	4.15 ± 0.04 <sup>a</sup>	4.20 ± 0.23 <sup>a</sup>	4.21 ± 0.04 <sup>a</sup>


Means with the same letter are not significantly different from each other (n=4, mean ± SEM, LDS test,  $\alpha < 0.05$ ).

### Total protein content, biomass production and pH in fungal extracts

At both 48 ( $F = 36.56$ ,  $P = 0.00001$ ) and 96 h ( $F = 37.52$ ,  $P = 0.00001$ ), *Penicillium* SP18 achieved the highest total protein content at 22.8 and 28.6 mg/L, respectively (Table 3). The lowest total protein content was produced by *Penicillium* SP1 at 17.8 and 16.8 mg/L at 48 and 96 h, respectively. At the end of evaluation, *Penicillium* SP11 (75.2 mg/L) achieved the highest total protein content ( $F = 3828.85$ ,  $P = 0.00001$ ) compared with that of the other isolates, while *P. hispanicum* showed

the lowest total protein content (11.9 mg/L). Total protein content is an indirect parameter of the ability of a specific isolate to produce certain fungal enzymes that could be involved in the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$ .

Figure 1 shows the biomass production of the five studied isolates during the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$ . Biomass production ranged from 0.20 to 0.52 mg/flask (120 mL of Pikovskaya culture medium). The isolate with the highest ( $F = 30.06$ ,  $P = 0.00001$ ) biomass production was *Penicillium* SP18 (0.52 mg/flask). In contrast, *Penicillium* SP1 showed the lowest biomass production (0.20 mg/flask).




**Figure 1.** Biomass production in Pikovskaya culture medium by five soil borne Micromycetes in the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$ . Bars with the same letter are not significantly different from each other (n = 4, mean ± SEM, LDS test,  $\alpha < 0.05$ ).

All fungal isolates reduced the pH of the culture medium across time of fermentation (Table 3). The pH of the control was 4.15 at 48 h after incubation; this value increased slightly at 96 and 144 h to 4.20 and 4.21, respectively. However, in the culture of fungal isolates, the pH decreased significantly ( $F = 174.0$ ,  $P = 0.00001$ ) from 4.15 to 2.38 (in *P. hispanicum*) at 48 h of culture. At 96 h of evaluation, the pH of fungal cultures ranged from 2.46 to 2.67, statistically different from the control (pH = 4.20,  $F = 15.06$ ,  $P = 0.00001$ ). Meanwhile, at the end of evaluation (144 h), the pH ranged from 2.35 to 2.53 in the fungal cultures; these values were statistically lower ( $F = 444.59$ ,  $P = 0.00001$ ) than the control (4.21).

### Plackett-Burman experimental design

For *P. hispanicum*,  $\text{Ca}_3(\text{PO}_4)_2$  solubilization, total protein content and biomass production varied widely (Table 2). These variations evidence the importance of culture medium optimization to efficiently solubilize  $\text{Ca}_3(\text{PO}_4)_2$ . In addition, parameters such as total protein production and biomass can reflect the efficiency of fungal growth.

For  $\text{Ca}_3(\text{PO}_4)_2$  solubilization, the highest soluble P was found in the 11<sup>th</sup> trial run (75.8 mg/L of soluble P), whereas the lowest activity was seen in the 1<sup>st</sup> trial run (4.0 mg/L of soluble P). Thus, 19.7% more soluble P was found in the 11<sup>th</sup> trial run compared with the central points (runs: 3, 16, 11 and 9). According to the Pareto chart (Figure 2A) and the ANOVA (Table 4), this level improved in presence of a high amount of  $\text{Ca}_3(\text{PO}_4)_2$  (0.75 g/L) and low values of  $(\text{NH}_4)_2\text{SO}_4$  (0.25 g/L),  $\text{MnSO}_4$  (0.002 g/L) and pH (4.0).



**Figure 2.** Pareto chart of eleven-factor standard effects on: A) solubilization of  $\text{Ca}_3(\text{PO}_4)_2$ , B) total protein content and C) biomass production by *Penicillium hispanicum*

The p-value was used to determine the main nutrimental factors at a confidence level of greater than 95%. According to the ANOVA, the main factors that influenced the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$  by *P. hispanicum* were  $(\text{NH}_4)_2\text{SO}_4$  ( $P = 0.0392$ ),  $\text{MnSO}_4$  ( $P = 0.0186$ ),  $\text{Ca}_3(\text{PO}_4)_2$  ( $P = 0.0009$ ) and pH ( $P = 0.0464$ ) (Table 4). Polynomial Equation 2 represents the model for  $\text{Ca}_3(\text{PO}_4)_2$  solubilization by *P. hispanicum*. The significance of Equation 2 was checked using an F-test, and the value was highly significant [( $P$ -value >  $F$ ) = 0.0029]. The coefficient of determination ( $r^2$ ) of the model [Equation 2] was 0.90194, indicating that 90.19% of the variability in the experiment data could be explained by the estimated model. Also, the standard error was low (4.79).

$$\begin{aligned} \text{Soluble P} = & 53.208 - 0.5652 G + 0.6160 Ye - 4.8586 \\ & NH - 2.9532 K + 3.8678 Mg - 6.4591 Mn + 0.2222 \\ & Na + 0.6160 Fe + 18.8057 Ca - 2.3054 I - 4.5410 pH \end{aligned} \quad \text{Eq. 2}$$

Regarding total protein content, the highest value was found in the 2<sup>nd</sup> trial run at 182.0 mg/L. In the 1<sup>st</sup> trial run, we observed a protein content of 19.9 mg/L. Thus, protein content increased 3.6 fold in the 2<sup>nd</sup> trial run compared with that of the central points (runs: 13<sup>th</sup> to 16<sup>th</sup>). According to the Pareto chart (Figure 2B) and the ANOVA (Table 4), protein content improved significantly in conditions of low glucose (5 g/L,  $P = 0.0107$ ), yeast extract (0.25 g/L,  $P = 0.00396$ ), inoculum amount ( $1 \times 10^5$  conidia/mL,  $P = 0.0164$ ) and pH (4.0,  $P = 0.0189$ ), and high  $\text{MgSO}_4$  (0.15 g/L,  $P = 0.0305$ ),  $\text{FeSO}_4$  (0.003 g/L,  $P = 0.0036$ ) and  $\text{Ca}_3(\text{PO}_4)_2$  (0.75 g/mL,  $P = 0.0049$ ). Polynomial Equation 3 was obtained from the multiple regression analysis and describes the effect of the nutrimental factors on total protein production of *P. hispanicum* during the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$ . Statistical significance of the model was checked using an F-test [( $P$ -value >  $F$ )=0.0538]. The  $r^2$  coefficient of the model [equation 3] was 0.8728, which indicated that 87.28% of the variability in the experimental data could be explained by the estimated model. The standard error of the model was low (8.72), indicating a low variation of the data.

$$\begin{aligned} \text{Protein content} = & 62.1676 - 14.3681 G - 14.9359 YE \\ & - 5.17399 NH + 2.8663 K + 9.75275 Mg - 0.723443 \\ & Mn - 3.01282 Na + 20.9982 Fe + 18.8919 Ca - \\ & 12.2985 I - 11.6758 pH. \end{aligned} \quad \text{Eq. 3}$$

The highest value for biomass production of *P. hispanicum* was found in the central points (13<sup>th</sup> to 16<sup>th</sup> trial run) with a mean of 0.425 mg/flask (120 mL of Pikovskaya culture medium). In contrast, the lowest biomass production was found in the 2<sup>nd</sup> trial run (0.11 mg/flask). The PBED conditions did not lead to an improvement in biomass production. Therefore, the Pikovskaya culture medium was adequate for biomass production of *P. hispanicum* during the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$ . The regression model of PBED, the Pareto chart (Figure 2A) and the ANOVA (Table 4) showed that biomass production was significantly influenced by a high level of glucose (15 g/L,  $P=0.0166$ ) and inoculum amount ( $1 \times 10^7$  conidia/mL,  $P = 0.0242$ ), although biomass production did not improve. Equation 4 describes the biomass production of *P. hispanicum* during  $\text{Ca}_3(\text{PO}_4)_2$  solubilization. The model was significant at 95% [( $P$ -value> $F$ )=0.0285] and had a low determination coefficient ( $r^2=0.4970$ ), explaining only 49.70% of the variability in the data. Also, the standard error was low (0.045).

$$\begin{aligned} \text{Biomass} = & 0.29 + 0.06333G + 0.01666YE + \\ & 0.01333NH + 0.02666K + 0.00333Mg - 0.01166 \\ & Mn - 0.00333Na + 0.0Fe - 0.01166Ca + 0.055I \\ & + 0.00333pH \end{aligned} \quad \text{Eq. 4}$$

**Table 4.** Analysis of variance for studied variables in the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$  by *Penicillium hispanicum*.

Source	F.D.	$\text{Ca}_3(\text{PO}_4)_2$ solubilization		Total protein content		Biomass production	
		F-ratio	P-value	F-ratio	P-value	F-ratio	P-value
Glucose	1	0.17	0.7104	32.51	<b>0.0107</b>	23.67	<b>0.0166</b>
Yeast extract	1	0.20	0.6864	35.13	<b>0.0096</b>	1.64	0.2904
$(\text{NH}_4)_2\text{SO}_4$	1	12.32	<b>0.0392</b>	4.22	0.1323	1.05	0.3811
KCl	1	4.55	0.1226	1.29	0.3380	4.20	0.1329
$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	1	7.81	0.0682	14.98	<b>0.0305</b>	0.07	0.8144
$\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$	1	21.78	<b>0.0186</b>	0.08	0.7927	0.80	0.4362
NaCl	1	0.03	0.8826	1.43	0.3177	0.07	0.8144
$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$	1	0.20	0.6864	69.44	<b>0.0036</b>	0.001	1.0000
$\text{Ca}_3(\text{PO}_4)_2$	1	184.61	<b>0.0009</b>	56.21	<b>0.0049</b>	0.80	0.4362
Inoculum	1	2.77	0.1944	23.82	<b>0.0164</b>	17.85	<b>0.0242</b>
pH	1	10.76	<b>0.0464</b>	21.47	<b>0.0189</b>	0.07	0.8144
Lack of fit	1	23.66	0.0166	34.96	0.0097	47.80	0.0064
Residual	3						
Total	15						

F.D. = Freedom degrees

## Discussion

In the soil the availability and efficiency of nutrients depend to a great extent on the biological activity of soil microorganisms. In this context, biofertilizers are an alternative means of providing plant nutrition (Khan *et al.*, 2014). According to Fuentes-Ramirez and Cabello-Mellado (2005), a biofertilizer is “a product that contains living microorganisms, which exert direct or indirect beneficial effects on plant growth and crop yield through different mechanisms”.

Due to the benefits of biofertilizers, during last two decades, mexican agriculture has developed strategies for the biological fertilization of cultivated plants. Today, biofertilizers formulated from bacteria (i.e., *Rhizobium* spp., *Bacillus* spp., *Pseudomonas* spp., *Azotobacter* spp., and *Azospirillum* spp., among others) and mycorrhizae (i.e., *Glomus intraradices*, *Gigaspora* spp., and *Acaulospora* spp., among others) are commonly found (Armenta-Bojórquez *et al.*, 2010; Malusá and Vassilev, 2014), although P-solubilizing micromycetes are not currently on the market. Therefore, the studies of P-solubilizing micromycetes adapted to specific environmental conditions are of particular interest.

In this sense, we found that *Penicillium* SP1 and *P. hispanicum* are able to solubilize at 100% the  $\text{Ca}_3(\text{PO}_4)_2$  in a Pikovskaya culture media;

both strains achieved better solubilization than a Mexican native strain of *Paecilomyces lilacinus*, which was able to solubilize only 71.28 mg/L of P at 288 h of culture in Pikovskaya culture medium (Hernández-Leal *et al.*, 2011). In another study on native SBM from a Mexican coffee plantation, Hernando-Posada *et al.* (2013) reported that *Paecilomyces marquandii* and *Penicillium janthinelum* released 1.9 and 7.0 mg/L of P from  $\text{Ca}_3(\text{PO}_4)_2$  in the culture medium at 24 days; these values were lower than those achieved by *Penicillium* SP1 and *P. hispanicum*. The ability of a fungal strain to release P from an inorganic source is known to be strain and species dependent and can also be influenced by multiple environmental factors (Zúñiga-Silva *et al.*, 2016).

The PBED was able to show the main nutrimental factors in  $\text{Ca}_3(\text{PO}_4)_2$  solubilization by *P. hispanicum*. Also, a P increase of 19.7% was obtained with low N source ( $(\text{NH}_4)_2\text{SO}_4$ ),  $\text{Mn}^{+2}$  ( $\text{MnSO}_4$ ) and pH. Some authors have reported that the N source mediated the solubilization ability of fungi. Relwani *et al.* (2008) reported that  $\text{KNO}_3$  significantly increased the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$  by a mutant strain of *Aspergillus tubingensis* in comparison with other N sources such as  $(\text{NH}_4)_2\text{SO}_4$ ,  $(\text{NH}_4)\text{NO}_3$ , asparagine and tryptophan. However, the analyzed amount of  $\text{KNO}_3$  was 0.5 g/L, a value higher than the best



solubilization conditions for *P. hispanicum*, which only needed 0.25 g of  $(\text{NH}_4)_2\text{SO}_4$  as an N source.

In a related study, Chun-Qiao *et al.* (2008) reported the different effects of various C:N ratios on rock phosphate solubilization by fungal isolates. The content of soluble P released by *Penicillium expansum* increased with an increasing C:N ratio, up to 104.1 mg/L at a C:N ratio of 40:1 (10 g of glucose and 0.25 g of  $(\text{NH}_4)_2\text{SO}_4$ ). These data are in agreement with what we found in this study using a PBED: a low amount of  $(\text{NH}_4)_2\text{SO}_4$  (0.25 g/L) favored a high P solubilization of  $\text{Ca}_3(\text{PO}_4)_2$ .

In another study, Scervino *et al.* (2011) reported that glucose- and  $(\text{NH}_4)_2\text{SO}_4$ -based media showed the highest values (97.78 mg/L) of  $\text{Ca}_3(\text{PO}_4)_2$  solubilization by *Penicillium purpurogenum* in comparison with using fructose and L-asparagine as C and N sources. According to the molecular weight of added  $\text{Ca}_3(\text{PO}_4)_2$ , *P. purpurogenum* solubilized 100% of P. In addition, P solubilization was optimized by Scervino *et al.* (2011) using a Box-Wilson design that varied the amount of C and N source. According to their results, the optimum amounts of glucose and  $(\text{NH}_4)_2\text{SO}_4$  were 13.66 and 0.34 g/L, respectively.

Recently, Padmavathi (2015) applied a PBED in the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$  by *Aspergillus niger* in which P release was significantly influenced by glucose and  $(\text{NH}_4)_2\text{SO}_4$ . Padmavathi (2015) suggested that the optimum  $(\text{NH}_4)_2\text{SO}_4$  value was 4.0 g/L. Our results with *P. hispanicum* suggest that a low value (0.25 g/L) of  $(\text{NH}_4)_2\text{SO}_4$  increased the P release from  $\text{Ca}_3(\text{PO}_4)_2$ . This difference could be due to the utilized fungal species and also reveals the importance of fungal culture optimization.

Total protein production by *P. hispanicum* improved in presence of low amounts of glucose, yeast extract, inoculum and pH. This suggests that protein production occurs under slight stress conditions because protein production also improved in presence of high amounts of salts such as  $\text{MgSO}_4$ ,  $\text{FeSO}_4$  and  $\text{Ca}_3(\text{PO}_4)_2$ . P-solubilizing fungi are well known to exhibit some enzyme activity, which contributes to P-solubilization; some enzymes include acid and alkaline phosphatase and phytase (Relwani *et al.*, 2008; Gaiind, 2016). Although enzyme activities were not evaluated in this study, total protein production suggests that there was a high amount of protein (according to total enzyme activity) in some of the culture conditions, for example, in the 2<sup>nd</sup> trial experiment. Future studies should evaluate and characterize these important enzymes. Recently, Gaiind (2016) showed that depending on the fungal species, the mechanism for reducing pH and releasing P from inorganic sources may be different. For instance,

in the solubilization of  $\text{Ca}_3(\text{PO}_4)_2$ , *Trichoderma harzianum* increases its citric acid production to reduce the pH of the culture medium, while *Aspergillus niger* increases its phytase activity and decreases its acid phosphate activity.

Mycelial biomass is a significant parameter for evaluating the growth of fungi. According to the PBED, biomass production by *P. hispanicum* did not improve. We found the highest biomass in the central points, which suggest that the Pikovskaya culture medium is adequate for the biomass production of *P. hispanicum*. In the literature, several culture media for rock phosphate and  $\text{Ca}_3(\text{PO}_4)_2$  solubilization have been reported. For instance, Wakelin *et al.* (2004) used a basal medium (0.1 g NaCl, 0.4 g  $\text{NH}_4\text{Cl}$ , 0.78 g  $\text{KNO}_3$ , 0.5 g  $\text{MgSO}_4$ , 0.1 g  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$  and 10 g sucrose L<sup>-1</sup>) for rock phosphate solubilization and reported a high amount of biomass production by *Penicillium bilaiae*, *Penicillium radicum* and *Penicillium sp.1*, over 500, 300 and 250 mg/flask (100 mL of basal medium), respectively. More recently, Zhen *et al.* (2016) reported the biomass production of two P-solubilizing fungi: *Penicillium oxalicum* and *Aspergillus niger*. Under pH stress (1.5 to 6.5) in potato dextrose agar broth (PDAB), both solubilizing fungi showed high biomass production in comparison with *P. hispanicum*, studied in this work. *Penicillium oxalicum* and *A. niger* produced 497.28 and 389.13 mg of biomass/flask (100 mL of PDAB) at pH 4.5 and 2.5, respectively. In the case of *P. oxalicum*, biomass production was not possible at pH 2.5 and 1.5. The decrease in the pH of the culture medium is well associated with organic acid production (Oliveira-Mendes *et al.*, 2016; Zhen *et al.*, 2016); organic acids, in some cases, can reduce the biomass production of P-solubilizing fungi. This could be the reason why *P. hispanicum* had low biomass production despite showing high  $\text{Ca}_3(\text{PO}_4)_2$  solubilization. In addition, the pH values in Pikovskaya culture medium inoculated with *P. hispanicum* were low, between 2.8 to 2.83, during the evaluation period (2 to 6 days).

## Conclusion

*Penicillium hispanicum* was able to solubilize 100% of  $\text{Ca}_3(\text{PO}_4)_2$  at six days of culture in Pikovskaya culture medium. The PBED revealed that P release by *P. hispanicum* was influenced by low values of  $\text{MnSO}_4$ ,  $(\text{NH}_4)_2\text{SO}_4$  and pH and a high amount of  $\text{Ca}_3(\text{PO}_4)_2$ . Thus, *P. hispanicum* is a good candidate to use in field experiments of  $\text{Ca}_3(\text{PO}_4)_2$  solubilization in poor soils with low C and N sources, as glucose (as a C source) did not significantly affect  $\text{Ca}_3(\text{PO}_4)_2$  solubilization and only a low amount of  $(\text{NH}_4)_2\text{SO}_4$  (as an N source) was required.

## Acknowledgments

Thanks to the SEP-PRODEP program (Programa para el Desarrollo Profesional Docente, Secretaría de Educación Pública, Mexico) for the financial support of this study.

## References

- Armenta-Bojórquez, A.D.; García-Gutiérrez, C.; Camacho-Báez, J.R.; Apodaca-Sánchez, M.A.; Gerardo-Montoya, L.; and Nava-Pérez, E. 2010. Role of biofertilizers in the agricultural development in Mexico. *Ra Ximhai*, 6(1), 51-56.
- Chun-Qiao, X.; Ru-An, C.; Xiao-Hui, H.; Wen-Xue, Z.; Guan-Zhou, Q.; and Dian-Zuo, W. 2008. Optimization of rock phosphate solubilization by phosphate solubilizing fungi isolated from phosphate mine. *Ecol Eng*, 33(2), 187-193. <http://dx.doi.org/10.1016/j.ecoleng.2008.04.001>
- Fuentes-Ramírez, L.E.; and Caballero-Mellado, J. 2005. Bacterial biofertilizers. In: Z.A. Siddiqui Ed.) *PGPR: biocontrol and biofertilization*. 1st ed. Springer, The Netherlands, pp 143-172.
- Gaïnd, S. 2016. Phosphate dissolving fungi: mechanism and application in alleviation of salt stress in wheat. *Microbiol Res*, 193, 94-102. <http://dx.doi.org/10.1016/j.micres.2016.09.005>
- Hernández-Leal, T.I.; Carreón, G.; and Heredia-Abarca, G. 2011. *In vitro* phosphate solubilization by a strain of *Paecilomyces lilacinus* Thom. *Agrociencia* 45(8), 881-892.
- Hernando-Posada, R.; Heredia-Abarca, G.; Sieverding, E.; and Sánchez de Prager, M.; 2013. Solubilization of iron and calcium phosphates by soil fungi isolates from coffee plantations. *Arch Agron Soil Sci* 59(2), 185-196. <http://dx.doi.org/10.1080/03650340.2011.610030>
- Hernando-Posada, R.; Sánchez de Prager, M.; Sieverding, E.; Aguilar-Dorantes, K.; and Heredia-Abarca, G.P. 2012. Filamentous and phosphate solubilizing fungi relationships with some edaphic parameters and coffee plantations management. *Rev Biol Trop*, 60(3), 1075-1096. <http://dx.doi.org/10.15517/rbt.v60i3.1759>
- Khan, M.S.; Zaidi, A.; and Ahmad, E. 2014. Mechanism of phosphate solubilization and physiological functions of phosphate-solubilizing microorganism. In: M.S. Khan, A. Zaidi, and J. Musarrat, (Eds.), *Phosphate solubilizing microorganism*, 1st edn. Springer International Publishing, Switzerland, pp 31-62.
- Malusá, E.; and Vassilev, N. 2014. A contribution to set legal framework for biofertilizers. *App Microbiol Biotechnol* 98(15), 6599-6607. <http://dx.doi.org/10.1007/s00253-014-5828-y>
- Oliveira-Mendes, G.; Moreira-De Freitas, A.L.; Liparini-Pereira, O.; Ribeiro-Da Silva, I.; Bojkov-Vassilev, N.; and Dutra-Costa, M. 2016. Mechanism of phosphate solubilizing by fungal isolates when exposed to different P sources. *Annal Microbiol*, 64(1), 239-249. <http://dx.doi.org/10.1007/s13213-013-0656-3>
- Padmavathi, T. 2015. Optimization of phosphate solubilization by *Aspergillus niger* using Plackett-Burman and response surface methodology. *J Soil Sci Plant Nutr*, 15(3), 781-793. <http://dx.doi.org/10.4067/S0718-95162015005000053>
- Plackett, R.L.; and Burman, J.P. 1946. The design of optimum multifactorial experiments. *Biometrika*, 33(4), 305-325.
- Relwani, L.; Krishna, P.; and Reddy, M.S. 2008. Effect of carbon and nitrogen source on phosphate solubilization by a wild-type strain and UV-induced mutants of *Aspergillus tubigensis*. *Curr Microbiol*, 57(1), 401-406. <http://dx.doi.org/10.1007/s00284-008-9212-y>
- Scervino, J.M.; Papinutti, V.L.; Godoy, M.S.; Rodriguez, M.A.; Della, M.; Recchi, M.; Pectenari, M.J.; and Godeas, A.M. 2011. Medium pH, carbon and nitrogen concentration modulate the phosphate solubilization efficiency of *Penicillium purpurogenum* through organic acid production. *J App Microbiol*, 110(1), 1215-1223. <http://dx.doi.org/10.1111/j.1365-2672.2011.04972.x>
- Sharma, S.B.; Sayyed, R.; Trivedi, M.H.; Gobi, T.A. 2013. Phosphate solubilizing microbes: sustainable approach for managing phosphorus deficiency in agricultural soils. *Springerplus* 2, 587-601. <http://dx.doi.org/10.1186/2193-1801-2-587>
- Wakelin, S.A.; Warren, R.A.; Hervey, P.; and Ryder, M.H. 2004. Phosphate solubilization by *Penicillium* spp. closely associated with wheat root. *Biol Fertil Soils*, 40(1), 36-43. <http://dx.doi.org/10.1007/s00374-004-0750-6>
- Watanabe, F.S.; and Olsen, S.R. 1965. Test of an ascorbic acid method for phosphorus in water and NaHCO<sub>3</sub> extracts from soil. *Soil Sci Soc Proc* 26(6), 677-678. <http://dx.doi.org/10.2136/sssaj1965.03615995002900060025x>
- Zhen, L.; Tongshuo, B.; Letian, D.; Fuwei, W.; Jinjin, T.; Shiting, M.; Yunxiao, H.; Shimei, W.; and Shujin, H. 2016. A study of organic acid production in contrast between two phosphate solubilizing fungi: *Penicillium oxalicum* and *Aspergillus niger*. *Sci Rep*, 6(1), 1-8. <http://dx.doi.org/10.1038/srep25313>
- Zúñiga-Silva, J.R.; Chan-Cupul, W.; Kuschik, P.; Loera, O.; Aguilar-López, R.; and Rodríguez-Vázquez, R. 2016. Effect of Cd<sup>2+</sup> on phosphate solubilizing abilities and hydrogen peroxide production of soil-borne Micromycetes isolated from *Phragmites australis*-rhizosphere. *Ecotoxicology*, 25(2), 367-379. <http://dx.doi.org/10.1007/s10646-015-1595-5>