



Acta Agronómica

ISSN: 0120-2812

ISSN: 2323-0118

Universidad Nacional de Colombia

Santos-Pereira, Ivan dos; Campos, Ângela Diniz; Greigh de Brito,
Giovani; Corrêa-Antunes, Luis Eduardo; Martins-Pereira, José Francisco
Cyanogenic glycosides content as graft compatibility indicator in pear/quince combinations
Acta Agronómica, vol. 70, no. 1, 2021, January-March, pp. 73-81
Universidad Nacional de Colombia

DOI: <https://doi.org/10.14482/INDES.30.1.303.661>

Available in: <https://www.redalyc.org/articulo.oa?id=169971425009>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UAEH  redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

Cyanogenic glycosides content as graft compatibility indicator in pear/quince combinations

Contenido de glucósidos cianogénicos como indicador de compatibilidad de injertos en combinaciones de pera/membrillo

Ivan dos Santos Pereira¹, Ângela Diniz Campos², Giovani Greigh de Brito³,
Luis Eduardo Corrêa Antunes⁴, José Francisco Martins Pereira⁵.

¹Empresa Brasileira de Pesquisa Agropecuária EMBRAPA, Brazil. ✉ ivanspereira@gmail.com

²Empresa Brasileira de Pesquisa Agropecuária EMBRAPA, Brazil. ✉ angela.campos@embrapa.br

³Empresa Brasileira de Pesquisa Agropecuária EMBRAPA, Brazil. ✉ giovani.brito@embrapa.br

⁴Empresa Brasileira de Pesquisa Agropecuária EMBRAPA, Brazil. ✉ luis.antunes@embrapa.br

⁵Empresa Brasileira de Pesquisa Agropecuária EMBRAPA, Brazil. ✉ jose.fm.pereira@embrapa.br (Q.E.P.D.)



<https://doi.org/10.15446/acag.v70n1.93516>

2021 | 70-1 p 73-81 | ISSN 0120-2812 | e-ISSN 2323-01180118 | Rec.: 10-02-2021 Acep.: 12-04-2021

Abstract

The objective of this work is to evaluate the concentration of cyanogenic glycosides CGs (amygdalin and prunasin) in the cultivar and rootstock, as an indicator of compatibility in pear grafts. The work consists of two experiments. Experiment 1 evaluated Cascatense/Pyrus Calleryana and Cascatense/EMC combinations. Experiment 2 evaluated the combinations Packham's Triumph/Adams, Santa Maria/Adams, and Rocha/Adams. The experimental design of the two experiments was randomized blocks with four replications. The evaluations were carried out in the spring and summer of the harvest 2014/2015. The results indicated graft compatibility in the Cascatense/P. Calleryana, Packham's Triumph/Adams, and Rocha/Adams combinations; slight incompatibility of Santa Maria/Adams, and severe incompatibility of Cascatense/EMC. The obtained results indicate that differences ≥ 20 mg g⁻¹ of CGs between scion and rootstock were correlated with a drastic reduction in vigor, problems of continuity in graft union, low yield, and early defoliation.

Keywords: amygdalin, Cydonia, grafting, incompatible, prunasin.

Resumen

El objetivo de este trabajo es evaluar la concentración de glucósidos cianogénicos CG (amigdalina y prunasina) en el cultivar y portainjerto, como indicador de compatibilidad en injertos de pera. El trabajo consta de dos experimentos. El experimento 1 evaluó las combinaciones Cascatense/Pyrus Calleryana y Cascatense/EMC. El Experimento 2 evaluó las combinaciones Packham's Triumph/Adams, Santa Maria/Adams y Rocha/Adams. El diseño experimental de los dos experimentos fue de bloques al azar con cuatro repeticiones. Las evaluaciones se realizaron en la primavera y el verano de la cosecha 2014/2015. Los resultados indicaron compatibilidad de injerto en el Cascatense/P. Calleryana; combinaciones de Packham's Triumph/Adams y Rocha/Adams; leve incompatibilidad de Santa Maria/Adams; e incompatibilidad severa de Cascatense / EMC. Los resultados obtenidos indican que diferencias ≥ 20 mg g⁻¹ de CG entre esqueje y portainjerto se correlacionaron con una drástica reducción del vigor, problemas de continuidad en la unión del injerto, bajo rendimiento y defoliación temprana.

Palabras claves: amigdalina, Cydonia, injerto, incompatibilidad, prunasina.

Introduction

The main rootstocks used in the pear seedlings are those of the *Cydonia* genus, especially *Cydonia oblonga* (Mill.). In general, these combinations result in plants of lower vigor, with early entry into production and greater productivity. However, the reduction in vigor that occurs in the pear/quince combinations can be attributed to the graft incompatibility phenomenon (Pina *et al.*, 2012). Graft incompatibility is the abnormal development of the graft due to anatomical, physiological, biochemical, and molecular differences, resulting in damage to the graft union, in addition to delayed vegetative growth, low productivity, and in severe cases, death of the plant (Davies *et al.*, 2017, p. 1024; Dogra *et al.*, 2018).

Graft incompatibility can cause major damage to commercial orchards by negatively affecting plant vegetative growth and thus reducing the potential yield (Machado *et al.*, 2017). But there may be different levels of incompatibility depending on the pear and quince cultivars grafted. Among the main quince rootstocks used in combinations with pear trees are the 'EMC', highly dwarfing, and 'Adams', of greater vigor (Francescato *et al.*, 2010).

The graft incompatibility of pear/quince combinations occurs due to problems in the vascular connection of the graft, especially in the regeneration of the cortex and phloem, occurring elevated percentage of necrotic area in the graft union (Balbi *et al.*, 2019). However, this lack of vascular continuity between pear and quince is attributed, in many cases, to the release of cyanide at the graft interface, a process known as cyanogenesis, which occurs through the hydrolysis of CGs (Gur *et al.*, 1968; Moore, 1986; Nocito *et al.*, 2010).

In pear/quince combinations, the CG prunasin, which occurs in the quince and not in the pear, rises from the rootstock to the cultivar, where it is hydrolyzed by β -glycosidase, causing the release of cyanide at the graft interface (Gur *et al.*, 1968). Thus, the levels of CGs in scion and rootstock can serve as indicators of compatibility (Gur & Blum, 1973; Pereira *et al.*, 2018).

In general, cyanogenic species also have the ability to metabolize cyanide, preventing cell damage. The cyanide detoxification process consists of two main routes. The first route involves the formation of β -cyanoalanine from cyanide and cysteine, a reaction catalyzed by β -cyanoalanine synthase. β -cyanoalanine is later converted to asparagine (Miller & Conn, 1980). The second route converts cyanide to thiocyanate and is catalyzed by rhodanase (Bordo & Bork, 2002). The most common route in plants is β -cyanoalanine (Zagrobelyny *et al.*, 2004). In this context, it can be suggested that the degree of graft compatibility or incompatibility depends both on the levels of CGs and on their ability to metabolize the released cyanide.

The objective of this work was to evaluate the growth and development of the plants, symptoms of graft incompatibility, and the relationship of these parameters with the concentration of CGs amygdalin and prunasin in the cultivar and in the rootstock of pear grafts, establishing a compatibility indicator. The tested hypothesis is that the concentrations of CG in the crown and in the rootstock are directly related to the degree of graft compatibility and to the growth of the plant.

Material and methods

The work was carried out in two private orchards located in the Arroio do Padre, Rio Grande do Sul, Brazil (Coordinates: 31° 26' 35.51" S, 52° 25' 22.10" O and altitude: 251 m). As the orchards had different combinations and ages, they were evaluated as different experiments.

Plant material.

Experiment 1 evaluated combinations between 'Cascatense' (*P. communis* L.) cultivar and *P. Calleryana* (L.) and 'EMC' (*Cydonia oblonga* Mill.) rootstocks. The orchard was implanted in the winter of 2009, with spacing of 1.5 m between plants and 5.0 between lines. Experiment 2 evaluated the *P. communis* (L.) cultivars, 'Packham's Triumph', 'Santa Maria', and 'Rocha', grafted on the 'Adams' quince rootstock (*C. oblonga* Mill.). This orchard was implanted in the winter of 2011, with a spacing of 1.0 m between plants and 5.0 between lines.

Orchard management.

The orchards were conducted on terrace of 0.5 m high and 1.0 m base, with drip irrigation during the summer. The lines and inter-lines were maintained with plant cover (*Avena strigosa*), managed with mowing and herbicide applications.

Fertilization was carried out using urea (45 % N), triple super phosphate (42 % P_2O_5), and potassium chloride (58 % K_2O). With 50 % of N applied in post-harvest, between March 15 and April 15, and the remainder, during the bud swelling period; P and K were applied during the winter dormancy period.

The region has insufficient cold for the evaluated cultivars, with a historical average ranging from 322 to 496 chilling hours, with dormancy breaking with hydrogenated cyanamide and mineral oil being indicated.

Growth, production and compatibility assessments.

Vegetative growth assessments were carried out in October 2014, and consisted of measuring trunk diameter, plant height and vigor. The trunk diameter (cm) was evaluated 5 cm above (scion) and below

(rootstock) of the grafting point, using a digital caliper. The plant height (cm) was determined from the grafting point to the apex of the plant, with a graduated measuring tape. Yield (kg ha⁻¹) was obtained by harvesting fruits of all plants in each plot. The plant vigor was evaluated by classifying the combinations into five classes: 'Very low', 'Low', 'Medium', 'High', and 'Very high', being carried out by three evaluators.

In February, the branches length (cm) of the year was also evaluated by measuring 15 branches per plant. The defoliation was estimated by classifying the combinations into four classes: 0 (without defoliation), 1 (defoliation of medium intensity), 2 (intense defoliation), and 3 (complete defoliation). Three evaluators performed operations.

Characterization of graft compatibility by growth parameters, visual symptoms, and biochemical responses as for symptoms, and combinations were classified (subjectively by three evaluators) according to the intensity of the symptoms in: 'Absent', 'Moderate', and 'Severe'. Taking into account in this classification the presence of swelling in the graft region, diameters of the cultivar and rootstock, the presence of lumps in the region close to the graft union, and density of shoot emission from the rootstock.

Determination of cyanogenic glycosides (CGs).

CGs concentration was determined in the bark tissues from the scion (5 cm above graft union) and rootstock (5 cm below graft union) in the spring (October 2014) and summer (February 2015). The bark samples were immediately frozen in liquid nitrogen, stored at -80 °C and subsequently lyophilized.

The extraction was carried in 500 mg of lyophilized sample, with 8 ml of methanol and 2.000 mg of activated carbon. This mix remained under stirring for 16 h at room temperature. Then, a centrifugation was performed at 5600 g for 20 min and the supernatant was vacuum filtered through a 0.1 µm nylon filter. 20 µL of the filtered solution were subjected to chromatographic analysis. An Inertsil ODS-3 analytical column (150 x 4.6 mm, 4 µm) was used at 25 °C during the analysis. The mobile phase was water-methanol (60:40; v/v) with a flow of 1.3 mL min⁻¹ and UV detection at a wavelength of 254 nm. Were used prunasin and amygdalin standards (Sigma-Aldrich), activated carbon (Sigma-Aldrich), and methanol (Vetec), specifics for HPLC (High Performance Liquid Chromatography). The water used was obtained employing the Milli-Q system purification (Millipore) and filtered through 0.1 mm nylon filter.

Chromatographic separation of the amygdalin and prunasin was successfully performed, with four peaks: Peak 1: water (1.16 min), Peak 2: methanol

(1.38 min), Peak 3: amygdalin (2.44 min), and Peak 4: prunasin (3.44 min). The results were expressed in mg g⁻¹ of the CGs on dry basis.

Statistical analysis.

The experiments were analyzed independently. The experimental design was in randomized blocks with four replications and each experimental unit consisted of five plants. The results were subjected to variance analysis, and variables with significant differences were compared using Tukey test, at 5 % probability of error.

Results and discussion

Experiment 1.

In Experiment 1, the Cascatense/*P. calleryana* showed a larger trunk diameter in the scion and rootstock, as well as a higher plant height and yield (Figures 1a, 1b, 1c). The high vigor of the Cascatense/*P. calleryana*




Figure 1. Growth, yield and biochemical assessments.

*Different lowercase letters differ by Tukey's test at 5 % probability of error.

combination was already an expected result, as *P. calleryana* is considered a vigorous rootstock, that is, it provides intense vegetative growth to the cultivar (Francescato *et al.*, 2010; Melo *et al.*, 2017).

On the other hand, the 'EMC' rootstock is used to reduce the vigor of the cultivar canopy and increase production (Maas, 2008). However, it was found in the present study that in addition to vigor reduction, there was also a decrease in yield, suggesting that graft incompatibility between 'Cascatense' and 'EMC' occurs at a very high level. This fact can be observed both in the vigor and compatibility assessment (Figure 2). Cascatense/EMC combination was classified as 'Very low' vigor and with graft incompatibility symptoms 'Severe' (Figure 2). The incompatibility observed between 'Cascatense' and 'EMC' also explains the high defoliation rate of this combination, classified as 'Intense' (Figure 1d), a phenomenon that also compromises the productive capacity of the next harvest. The incompatibility between 'Cascatense' and 'EMC' has already been noted by Giacobbo *et al.* (2007), who even found the same symptom of marked hypertrophy at the graft union point, in addition to overgrowth and reduced diameter above and below the grafting point. The incompatibility in pear/quince combinations is of the 'localized' type, which manifests itself in the form of structural disorganization in the graft union region (Ermel *et al.*, 1999), mainly due to problems continuity in cortex and phloem (Balbi *et al.*, 2019). In advanced stages, this type of incompatibility can cause the depletion of the root system, which occurs due to poor translocation at the point of union (Errea *et al.*, 1994; Zarrouk *et al.*, 2006).

In the spring, the levels of amygdalin were higher both in the cultivar and in the rootstock from Cascatense/EMC (Figure 1e), while in the summer there were no differences (Figure 1f).

About prunasin, it was observed that in spring the concentrations were higher in the scion and rootstock from Cascatense/*P. calleryana* (Figure 1g). In summer, the prunasin content was higher in the cultivar 'Cascatense' grafted on 'EMC' (Figure 1h).

In general, prunasin concentrations in the cultivar and in the rootstock from two combinations were quite low compared to those of amygdalin, indicating a greater role of amygdalin in the incompatibility process. However, according to Gur *et al.* (1968), the CG responsible for the incompatibility of grafting between pear and quince should be prunasin, which according to the same authors, occurs in quince and not in pear. But, in the present study, in addition to having determined that both CGs are present in the pear and quince tree, the CG that seems to have the greatest relationship with the incompatibility process is amygdalin.

However, it is important to note that amygdalin is a prunasin precursor (Conn, 1980), so depending on the season, the age of the plants, the genotypes studied, among others, it is possible that one or other is determined in greater concentration.

The amygdalin and the β -glycosidase enzyme are compartmentalized in the tissues from cyanogenic plants, and the cyanogenesis process starts only when these compartments rupture, probably due to the grafting process. From that moment, the amygdalin is hydrolyzed by β -glycosidase, generating one molecule of the glucose and another one of prunasin and later, prunasin is hydrolyzed by the same enzyme, producing another molecule of glucose and an α -hydroxylitrile, which finally suffers action of α -hydroxylitrile lyase, resulting in cyanide release (Conn, 1980; Zagrobelny *et al.*, 2004). Thus, perhaps the most appropriate is to consider the cyanogenic potential as the concentration of total CGs (amygdalin + prunasin) in the graft components.




Figure 2. Vigor, graft incompatibility symptoms, and biochemical assessments.

*Vigor classes: 'Very low', 'Low', 'Medium', 'High', and 'Very high'. Grafting incompatibility visual symptoms: 'Absent', 'Light', 'Moderate', and 'Severe'. (ns) does not differ and (**) differs by Tukey's test with 1 % probability of error.

In this sense, in the samples collected in the spring, there were no differences between the levels in the cultivar grafted on the two rootstocks (Figure 1i). However, there was a significant difference in the total content of CGs between rootstocks, with 'EMC' showing 5.6 times more than *Pyrus Calleryana* (Figure 1i). In summer, the total content of CG was higher in the cultivar 'Cascatense' grafted on 'EMC' (Figure 1j). However, there were no differences between the rootstocks of the two combinations (Figure 1j).

The differences between levels of total CGs from scion and rootstock of each combination were observed in the Cascatense/*P. Calleryana*, both in spring and summer, with no significant differences. The fact that probably contributes to its good compatibility. On the other hand, in the combination Cascatense/EMC a significant difference was found in the concentration of total CGs between scion and rootstock (Figure 2). It was found that the rootstock 'EMC' showed a concentration of

total CGs 19.84 mg g⁻¹ higher than 'Cascatense' or 2.4 fold higher. A difference that is possibly associated with the graft incompatibility observed in this combination. Pereira *et al.* (2018) observed in incompatible *Prunus* combinations, a difference of 1.9 fold.

It was also possible to observe that even in the pear/pear combination (*P. communis*/*P. Calleryana*), significant amounts of amygdalin and prunasin were found. Indicating that, differently from what was suggested by Gur *et al.* (1968), the pear has cyanogenic glycosides in its tissues, however, in concentrations lower than that found in quince trees.

Experiment 2.

In Experiment 2, Packham's Triumph/Adams combination showed a larger trunk diameter in the scion and rootstock, in addition to greater plant height and length of branches (Figures 3a, 3b, 3c).




Figure 3. Growth, yield and biochemical assessments.

*Different lowercase letters differ by Tukey's test at 5 % probability of error.

Regarding defoliation, the combination Santa Maria/Adams suffered 'Intense' defoliation, being superior to the others, which had only a defoliation level considered 'Light' (Figure 3d).

The lower vegetative growth of Santa Maria/Adams combination is probably related to a lower affinity between graft components. The effect that is evident from vigor and graft compatibility assessment (Figure 4).

The combination Packham's Triumph/Adams was classified as 'Medium' vigor, while the others presented 'Low' vigor (Figure 4). As for the compatibility assessment, it was found that the combination Packham's Triumph/Adams does not show any incompatibility symptoms, a result that is in line with what was observed by Pinto *et al.* (2009).

Rocha/Adams combination behaved similarly to Santa Maria/Adams for most growth variables but had less pronounced incompatibility symptoms, such as barely noticeable hypertrophy and little bud sprouting from the rootstock. Santa Maria/Adams showed clear symptoms of moderate incompatibility, with hypertrophy at the grafting point, in addition to a smaller trunk diameter on the rootstock and also a high emission of buds from the rootstock (Figure 4).

In the Santa Maria/Adams combination, the signs of grafting incompatibility are clear, but in relation to Rocha/Adams the evaluations carried out do not allow us to state that there is, in fact, grafting incompatibility. However, even if there is incompatibility of grafting in these two combinations, the degree must be light or moderate, with an impact only on growth control, inducing less vigor. Assumption supported by studies carried out in

Vacaria-RS, in which 'Rocha' and 'Santa Maria' grafted on 'Adams' showed adequate growth and yield (Fioravanço *et al.*, 2016; Fioravanço *et al.*, 2017).

The three combinations studied presented low yield, even though significant flowering occurred (data not shown) (Figure 3e). In another region of Brazil, with higher altitude and cold accumulation, 'Rocha' and 'Santa Maria' grafted on 'Adams', showed a yield of 2.35 and 1.52 t ha⁻¹, respectively, also in the third year after planting (Fioravanço *et al.*, 2016; Fioravanço *et al.*, 2017). Yield far superior to that observed in the present experiment. In addition to age, the low yield of the orchard evaluated in this study is likely to have been influenced by climatic factors, such as the lack of cold, as these cultivars need an average of 500-700 chilling hours (7.2 °C) to break bud dormancy properly. However, the accumulation of chilling hours for this region, in that year, was only 270 hours, and in the last 3 years, the average was 362 hours. The insufficiency of cold in pear trees can result in greater floral abortion and, consequently, reduction in production (Pasa *et al.*, 2012).

In the spring, 'Santa Maria' showed a higher amygdalin level than the other cultivars grafted on 'Adams' (Figure 3f). In summer, 'Packham's Triumph' and 'Santa Maria' cultivars showed higher amygdalin levels (Figure 3g). Regardless of time year, there were no differences in the amygdalin concentration of rootstocks from studied combinations (Figures 3f and 3g). For prunasin in the spring, there were no differences between cultivars (Figure 3h). However, the rootstock 'Adams' had a higher concentration of prunasin when grafted under 'Santa Maria' (Figure 3h). In the summer, the combination Rocha/Adams




Figure 4. Vigour, graft incompatibility symptoms and biochemical assessments.

*Vigor classes: 'Very low', 'Low', 'Medium', 'High' and 'Very high'. Grafting incompatibility symptoms: 'Absent', 'Light', 'Moderate', and 'Severe'. (ns) does not differ and (**) differs by Tukey's test at 5 % probability of error.

showed a higher content of prunasin in the cultivar and rootstock, without differing only from the Rocha/Adams in the rootstock (Figure 3i).

In general, there was little difference between amygdalin and prunasin levels, except for the amygdalin in the 'Santa Maria', in spring (Figure 3f). This result was confirmed when evaluating the total CGs, when 'Santa Maria' presented a higher concentration in the spring is 2-fold higher than that determined in 'Packham's Triumph' and 'Rocha' (Figure 3j). However, there were no differences between cultivars or rootstocks in the summer (Figure 3k).

In the analysis of the difference in concentration of total CGs between scion and rootstock of each combination, the main difference was between 'Santa Maria' and 'Adams', in which, 'Santa Maria' had a content of 11.29 mg g^{-1} or 1.9 times greater than 'Adams' (Figure 4). Difference that is probably related to the graft incompatibility symptoms and to the lower performance in the growth parameters of this combination. According to Pereira *et al.* (2018), a difference like this can cause severe graft incompatibility between *Prunus persica* and *Prunus mume*.

The biochemical results associated with the 'Intense' defoliation, the absence of fruit production, the 'Low' vigor and the incompatibility symptoms close to the grafting point, indicate that the incompatibility between 'Santa Maria' and 'Adams' exerts a moderate vigor control in this combination. A similar effect was reported by Pina *et al.* (2012) in *Prunus* combinations.

The Packham's Triumph/Adams combination, on the other hand, proved to be completely compatible, with high level of vigor and similar CG levels between the graft components. However, for Rocha/Adams the occurrence of graft incompatibility is not clear, as some growth assessments indicate less vigor in relation to the Packham's Triumph/Adams, but the symptoms of incompatibility considered in the present study do not allow a confirmation, as well as similar levels of CGs between scion and rootstock.

Cyanogenic glycosides x Grafting incompatibility between pear and quince.

Among studied combinations, in both experiments, combinations with different levels of grafting incompatibility and CG concentrations were identified. Cascatense/P. Calleryana, Packham's Triumph/Adams, and Rocha/Adams, had similar concentrations of CGs between scion and rootstock and showed no graft incompatibility symptoms. On the other hand, Santa Maria/Adams had 'Moderate' differences, resulting in less vigor and moderate incompatibility symptoms. But in the case of the

Cascatense/EMC combination, the difference in the CG content between scion and rootstock was 'High' and the consequences were a 'Severe' incompatibility that compromised the growth and development of the combination.

In view of the results obtained in the present study, in which there is evident relationship between the content of CGs and the growth parameters, which in turn are related to the graft compatibility, a correlation study was carried out between the differences in the total content of CGs from scion and rootstock and the growth variables (Table 1). The results indicate that the difference in CG concentration between scion and rootstock in the spring really influences the growth of plants (trunk diameter, plant height, length of branches, and defoliation). This behavior is probably induced by the different levels of incompatibility of the studied combinations.

Table 1. Pearson correlation coefficient between variables.

Variables	Difference Total CG between scion and rootstock-Spring		Difference Total CG between scion and rootstock-Summer	
	Correlation coefficient	p-value	Correlation coefficient	p-value
Trunk diameter of scion	-0.72	0.1053	-0.40	0.4351
Trunk diameter of rootstock	-0.78	0.0699	-0.11	0.8401
Plant height	-0.77	0.0733	-0.28	0.5911
Length of branches	-0.89	0.0174	-0.35	0.5015
Defoliation	0.84	0.0359	0.18	0.7316
Yield	0.03	0.9569	0.16	0.7657

Conclusions

This study has shown that the graft incompatibility between pear and quince trees occurs in large part, due to the difference in the levels of CGs of both species. In addition, it can be suggested that when the difference between the contents of the scion and the rootstock is small ($\leq 5 \text{ mg g}^{-1}$), there is no effect on grafting compatibility; when the differences increase (> 5 and $< 20 \text{ mg g}^{-1}$), an acceptable level of compatibility for the growth and development of the graft may occur but may result in the reduction of vigor at different levels. Finally, when the difference is high ($\geq 20 \text{ mg g}^{-1}$), in addition to the drastic reduction in vigor, there are problems of continuity in graft union, low yield, and early defoliation.

It is also added that the correct time for the assessment of CG levels is the beginning of spring, since it was at that moment that significant correlations between the levels of CGs of the evaluated genotypes and the growth of plants were identified (Table 1).

From the results presented in this study, a scheme that relates the difference in the concentration of CGs between scion and rootstock with the degree of graft compatibility (Figure 5) is proposed. This scheme will be based on the data obtained in the present study, in which five combinations will be evaluated, in two periods and in two experiments. Thus, it is believed that new studies should be performed, involving a greater number of combinations, in order to make this scheme more robust.




Figure 5. Graft compatibility and difference in total concentration of cyanogenic glycosides, amygdalin, and prunasin.

Acknowledgements

This study was financed by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior-Brasil (CAPES), Empresa Brasileira de Pesquisa Agropecuária (EMBRAPA), and by the project “Desenvolvimento e adaptação de tecnologias para viabilizar produção da pereira no Brasil”.

References

- Balbi, R.V., Pio, R., Farias, D.H., Melo, E.T., Pereira, M.P., & Pereira, F.J. (2019). The cell regeneration and connection of grafting between pear and quince trees are defined by the cortex and phloem. *Scientia Horticulturae*, 257(17), 108662. <https://doi.org/10.1016/j.scienta.2019.108662>
- Bordo, D., & Bork, P. (2002). The rhodanese/Cdc25 phosphatase superfamily. Sequence-structure-function relations. *EMBO Rep.*, 3, 741-746. <https://doi.org/10.1093/embo-reports/kvf150>
- Conn, E.E. (1980). Cyanogenic glycosides. In E.A., Bell, & Charlwood, B.V. (Eds.), *Secondary plant products*, (pp. 461-492). Springer-Verlag Berlin Heidelberg.
- Davies, F.T., Geneve, R.L., Wilson, S.B., Hartmann, H.T., & Kester, D.E. (2017). *Hartmann & Kester's Plant Propagation: Principles and Practices*. 9th ed. Pearson.
- Dogra, K., Kour, K., Kumar, R., Bakshi, P., & Kumar, V. (2018). Graft-incompatibility in horticultural crops. *International Journal of Current Microbiology and Applied Sciences (IJCMAS)*, 7(2), 1805-1820. <https://doi.org/10.20546/ijcmas.2018.702.218>
- Ermel, F.F., Kervella, J., Catesson, A.M., & Poëssel, J.L. (1999). Localized graft incompatibility in pear/quince (*Pyrus communis*/ *Cydonia oblonga*) combinations: multivariate analysis of histological data from 5-month-old grafts. *Tree Physiology*, 19(10), 645-654. <https://pubmed.ncbi.nlm.nih.gov/12651320/>
- Errea, P., Felipe, A.J., & Herrero, M. (1994). Graft establishment between compatible and incompatible *Prunus* spp. *Journal of Experimental Botany*, 45(3), 393-401. <https://doi.org/10.1093/jxb/45.3.393>
- Fioravanco, J.C., Antonioli, L.R., Czermainski, A.B.C., Oliveira, P.R.D., Almeida, G.K., & Ferreira de Araújo, W. (2016). Avaliação agrônômica da Pereira 'Rocha' em Vacaria, RS. Bento Gonçalves, RS: Embrapa Uva e Vinho, Circular Técnica 128. https://www.researchgate.net/publication/305157135_Avaliacao_Agronomica_da_Pereira_'Rocha'_em_Vacaria_RS
- Fioravanco, J.C., & Oliveira, P.R.D. (2017). Avaliação Agrônômica da Pereira 'Santa Maria' em Vacaria, RS. Bento Gonçalves: Embrapa Uva e Vinho, Circular Técnica 137. <https://www.embrapa.br/busca-de-publicacoes/-/publicacao/1075858/avaliacao-agronomica-da-pereira-santa-maria-em-vacaria-rs>
- Francescato, P., Pazzin, D., Gazolla Neto, A., Fachinello, J.C., & Giacobbo, C.L. (2010). Evaluation of graft compatibility between quince rootstocks and pear scions. *Acta Horticulturae*, 872, 253-260. <http://dx.doi.org/10.17660/ActaHortic.2010.872.34>
- Giacobbo, C.L., Fachinello, J.C., & Picolotto, L. (2007). Compatibilidade entre o marmeleiro porta-enxerto cv. EMC e cultivares de pereira. *Scientia Agraria*, 8(1), 33-37. <https://www.redalyc.org/pdf/995/99516333004.pdf>
- Gur, A., & Blum, A. (1973). The role of cyanogenic glycoside in incompatibility between peach scions and almond rootstocks. *Horticultural Research*, 13, 1-10.
- Gur, A., Samish, R.M., & Lifshitz, E. (1968). The role of the cyanogenic glycoside of the quince in the incompatibility between pear cultivars and quince rootstocks. *Horticultural Research*, 8, 113-134.
- Maas, F. (2008). Evaluation of *Pyrus* and quince rootstocks for high density pear orchards. *Acta Horticulturae*, 800, 599-609. https://www.actahort.org/books/800/800_80.htm
- Machado, B.D., Magro, M., Rufato, L., Bogo, A., & Kretschmar, A.A. (2017). Graft compatibility between european pear cultivars and east malling "C" rootstock. *Revista Brasileira de Fruticultura*, 39(3), e-063. <http://dx.doi.org/10.1590/0100-29452017063>
- Melo, E.T., Pio, R., Balbi, R.V., Ferreira, C.A., Mori, F.A. (2017). Anatomic compatibility of pear and quince trees grafted on *Pyrus calleryana* and *Chaenomeles sinensis* rootstocks. *Pesquisa Agropecuária Brasileira*, 52(10), 877-886. <http://dx.doi.org/10.1590/S0100-204X2017001000007>
- Miller, J.M., & Conn, E.E. (1980). Metabolism of hydrogen cyanide by higher plants. *Plant Physiology*, 65(6), 1199-1202. <http://dx.doi.org/10.1104/pp.65.6.1199>
- Moore, R. (1986). Graft incompatibility between pear and quince: the influence of metabolites of *Cydonia oblonga* on suspension cultures of *Pyrus communis*. *American Journal of Botany*, 73(1), 1-4. <https://doi.org/10.1002/j.1537-2197.1986.tb09673.x>
- Nocito, F.F., Espen, L., Fedeli, C., Lancilli, C., Musacchi, S., Serra, S., Sansavini, S., Cocucci, M., & Sacchi, G.A. (2010). Oxidative stress and senescence-like status of pear calli co-cultured on suspensions of incompatible quince microcalli. *Tree Physiology*, 30(4), 450-458. <https://doi.org/10.1093/treephys/tpq006>

- Pasa, M.S., Fachinello, J.C., Schmitz, J.D., Kulkamp de Souza, A.L., & Franceschi, É. (2012). Growth, yield and fruit quality of pear grafted on quince rootstocks and *Pyrus calleryana*. *Revista Brasileira de Fruticultura*, 34(3), 873-880. <https://doi.org/10.1590/S0100-29452012000300029>
- Pereira, I.S., Pina, A., Correa Antunes, L.E., Diniz Campos, Â., & Fachinello, J.C. (2018). Genotypic differences in cyanogenic glycosides levels of compatible *Prunus persica*/P. *persica* and incompatible P. *persica*/P. *mume* combinations. *Bragantia*, 77(1), 1-12. <http://dx.doi.org/10.1590/1678-4499.2016367>
- Pina, A., Errea, P., & Martens, H.J. (2012). Graft union formation and cell-to-cell communication via plasmodesmata in compatible and incompatible stem unions of *Prunus* spp. *Scientia Horticulturae*, 143(16), 144-150. <https://doi.org/10.1016/j.scienta.2012.06.017>
- Pinto Tomaz, Z.F., Couto Rodrigues, A., Veríssimo, V., Marafon, A.C., Herter, F.G., Rufato, A.R. (2009). Compatibilidade de enxertia de cultivares de marmeleiros com pereiras. *Revista Brasileira de Fruticultura*, 31(4), 1211-1217. <http://dx.doi.org/10.1590/S0100-29452009000400041>
- Zagrobelny, M., Bak, S., Vinther Rasmussen, A., Jorgensen, B., Naumann, C.M., & Lindberg Moller, B. (2004). Cyanogenic glucosides and plant-insect interactions. *Phytochemistry*, 65(3), 293-306. <http://dx.doi.org/10.1016/j.phytochem.2003.10.016>
- Zarrouk, O., Pinochet, J., Gogorcena, Y., & Moreno, M.A. (2006). Graft compatibility between peach cultivars and *Prunus* rootstocks. *HortScience*. 41(6), 389-1394. <http://dx.doi.org/10.21273/HORTSCI.41.6.1389>