



Revista Latinoamericana de Hipertensión
ISSN: 1856-4550
latinoamericanadehipertension@gmail.com
Sociedad Latinoamericana de Hipertensión
Venezuela

Seromarkers of synthesis and collagen degradation, electrophysiological heart parameters among patients with syndrome of preexcitation of ventricles

Vasilets, L. M.; Krivaya, A. A.; Vustina, V. V.; Khlynova, O. V.; Nekrutenko, L. A.; Tuev, A. V.

Seromarkers of synthesis and collagen degradation, electrophysiological heart parameters among patients with syndrome of preexcitation of ventricles

Revista Latinoamericana de Hipertensión, vol. 14, no. 5, 2019

Sociedad Latinoamericana de Hipertensión, Venezuela

Available in: <https://www.redalyc.org/articulo.oa?id=170262877009>

Queda prohibida la reproducción total o parcial de todo el material contenido en la revista sin el consentimiento por escrito del editor en jefe.



This work is licensed under Creative Commons Attribution-NoDerivs 4.0 International.

Seromarkers of synthesis and collagen degradation, electrophysiological heart parameters among patients with syndrome of preexcitation of ventricles

Seromarcadores de síntesis y degradación de colágeno, parámetros cardíacos electrofisiológicos entre pacientes con síndrome de preexcitación de ventrículos

L. M. Vasilets

E.A. Wagner Perm State Medical University, Rusia

 <http://orcid.org/0000-0002-5581-3418>

Redalyc: <https://www.redalyc.org/articulo.oa?id=170262877009>

A. A. Krivaya

E.A. Wagner Perm State Medical University, Rusia

 <http://orcid.org/0000-0001-6138-4042>

V. V. Vustina

E.A. Wagner Perm State Medical University, Rusia

 <http://orcid.org/0000-0002-3106-4993>

O. V. Khlynova

E.A. Wagner Perm State Medical University, Rusia

 <http://orcid.org/0000-0002-6447-0277>

L. A. Nekrutenko

E.A. Wagner Perm State Medical University, Rusia

 <http://orcid.org/0000-0002-7941-9634>

A. V. Tuv

E.A. Wagner Perm State Medical University, Rusia

 <http://orcid.org/0000-0001-6473-5661>

ABSTRACT:

: The results of examination of 43 patients with syndrome of preexcitation of ventricles are introduced in the article. The aim of the study was to optimize diagnostics and prediction of the development of arrhythmia in this cohort and to find out the possible role of disorder of fibrous matrix in the development of disorder of cardiac arrhythmias. The main group consisted of 43 patients with syndrome of preexcitation of ventricles (phenomenon WPW, syndrome WPW) and 15 practically healthy people of the same age were in the group of comparison. The average age of the examined patients from the main group was $25,3 \pm 9,15$ years old, among them 29 men (67%) and 14 women (33%). The average age in the comparison group was $27,9 \pm 7,9$ years old. All groups included into the examination were comparable by gender distribution ($p=0,45$, Fisher's test) and by age ($p=0,27$, U-criterion). The highest figures of matrix metal proteinase-9 (MMP-9) were found among patients with syndrome WPW – $96,1 \pm 33,2$ ng/ml and in the group of phenomenon WPW ($54,3 \pm 21,8$ ng/ml; $p=0,0003$). Practically healthy people had minimal values of the MMP-9 – $27,4 \pm 10,9$ ng/ml ($p=0,00014$; $p=0,000002$). Maximal values of tissue inhibitor of matrix metal proteinase (TIMP-1) are introduced in the group of phenomenon WPW and in the group of practically healthy people ($418,5 \pm 69,8$ ng/ml and $461,7 \pm 72,2$ ng/ml; $p=0,27$). Patients with syndrome WPW have lower TIMP-1 – $341,1 \pm 90,1$ ng/ml ($p=0,002$; $p=0,00012$). The level of propeptideprocollagen of the type I (PICP) among patients with syndrome WPW was $179,9 \pm 76,2$ ng/ml and $97,8 \pm 31,7$ ng/ml ($p=0,00014$) in the group of phenomenon WPW. All examined patients according to the level of PICP differed from practically healthy people $69,4 \pm 23,9$ ng/ml ($p<0,05$, U-criterion). The examined patients have changes of fibrous matrix. That includes an increasing the concentration of MMP-9, PICP and reducing TIMP-1. The syndrome WPW is characterized by more expressed disbalance of seromarkers of synthesis and collagen degradation than phenomenon is. It is possible that the examined seromarkers of fibrosis take place in forming disorders of cardiac arrhythmias in actual syndromes.

KEYWORDS: arrhythmia, arrhythmogenesis, markers of fibrosis.

RESUMEN:

En el artículo se presentan los resultados del examen de 43 pacientes con síndrome de preexcitación de ventrículos. El objetivo del estudio fue optimizar el diagnóstico y la predicción del desarrollo de arritmia en esta cohorte y descubrir el posible papel del trastorno de la matriz fibrosa en el desarrollo del trastorno de las arritmias cardíacas. El grupo principal consistió en 43 pacientes con síndrome de preexcitación de ventrículos (fenómeno WPW, síndrome WPW) y 15 personas prácticamente sanas de la misma edad estaban en el grupo de comparación. La edad promedio de los pacientes examinados del grupo principal fue de $25,3 \pm 9,15$ años, entre ellos 29 hombres (67%) y 14 mujeres (33%). La edad promedio en el grupo de comparación fue de $27,9 \pm 7,9$ años. Todos los grupos incluidos en el examen fueron comparables por distribución de género ($p=0,45$, prueba de Fisher) y por edad ($p=0,27$, criterio U). Las cifras más altas de matriz metálica proteinasa-9 (MMP-9) se encontraron entre pacientes con síndrome WPW - $96,1 \pm 33,2$ ng/ml y en el grupo de fenómeno WPW ($54,3 \pm 21,8$ ng/ml; $p=0,0003$). Las personas prácticamente sanas tenían valores mínimos de MMP-9: $27,4 \pm 10,9$ ng/ml ($p=0,00014$; $p=0,000002$). Los valores máximos del inhibidor tisular de la proteinasa de matriz metálica (TIMP-1) se introducen en el grupo de fenómeno WPW y en el grupo de personas prácticamente sanas ($418,5 \pm 69,8$ ng/ml y $461,7 \pm 72,2$ ng/ml; $p=0,27$). Los pacientes con síndrome de WPW tienen menor TIMP-1 - $341,1 \pm 90,1$ ng/ml ($p=0,002$; $p=0,00012$). El nivel de propéptidoprocógeno del tipo I (PICP) entre los pacientes con síndrome WPW fue $179,9 \pm 76,2$ ng/ml y $97,8 \pm 31,7$ ng/ml ($p=0,00014$) en el grupo de fenómenos WPW. Todos los pacientes examinados de acuerdo con el nivel de PICP diferido de personas prácticamente sanas $69,4 \pm 23,9$ ng/ml ($p<0,05$, criterio U). Los pacientes examinados tienen cambios de matriz fibrosa. Eso incluye aumentar la concentración de MMP-9, PICP y reducir TIMP-1. El síndrome WPW se caracteriza por un desequilibrio más expresado de los seromarcadores de síntesis y degradación del colágeno que el fenómeno. Es posible que los seromarcadores examinados de fibrosis tengan lugar en la formación de trastornos de arritmias cardíacas en síndromes reales.

PALABRAS CLAVE: arritmia, arritmogénesis, marcadores de fibrosis.

INTRODUCTION

Most of electro physiologists view the syndrome of Wolff-Parkinson-White (WPW) as a classical substrate for disorders of cardiac arrhythmias. However, an actual syndrome still remains a complicated issue when choosing a tactics for treatment such patients. In most cases its clinical manifestation happens at the age from 10 to 20 years old and very rarely among people of elderly age⁵. In 40-80 per cent the syndrome WPW is clinically developed by various forms of disorders of cardiac arrhythmias (DCA). Approximately half of discovered arrhythmias do not only make worse the quality of life but they are often fatal².

^{1/3} of patients have paroxysms of atrial fibrillation (AF). In most cases it is a reason of cardiovascular accidents. The combination of the AF with high heart rate (HR) and with a short effective refractory period (adiaphoria) (ERP) of additional way of conducting (AWC) creates an ability for development of flutter and fibrillation of ventricles (FV). The conducting the impulses to the ventricles in the ratio of 1:1 up to 340 per minute during episode MA provides transformation of the AF into the FV which is a main mechanism of sudden cardiac death (SCD) of patients with syndrome WPW^{1,6,17,20}.

The possibility to have a sudden death during 10 years with the syndrome of preexcitation of ventricles exceeds a general risk and it is from 0.15 to 0.39 per cent. In some cases the SCD can be the first clinical demonstration of masymptomatic preexcitation of ventricles^{6,8,16}.

Nowadays, the non-invasive and endocardialelectrophysiological study (EPS) of cardiac conduction system has the key point in the diagnosis the syndromes of preexcitation of ventricles and prediction of the DCA.. The results received during the EPS determine the further tactics of treatment of patients^{3,6,18}. However, even the non-invasive EPS has a set of contra-indications and restrictions for usage: advisability of its usage among asymptomatic patients and during childhood is still discussed^{11,12,14,15,19}.

The development of the DCA is connected with progressive structural and functional and electric remodeling of atrial myocardium. The mechanisms of structural remodeling and progression of selective atrial fibrosis are still not studied in detail. An interstitial fibrosis of myocardium is characterized by proffered

accumulation of collagen of the 1st type. The multitude products of metabolism of this protein are terminal propeptide procollagen type I (PICP), matrix metal proteinase-9 (MMP-9), tissue inhibitor of matrix metal proteinase-1 (TIMP-1)^{12,13}. Consequently, the level of PICP, MMP-9 and TIMP-1 can indirectly reflect the degree of synthesis and collagen degradation of type I normally and pathologically.

In that way serumal biomarkers of fibrosis give a certain mark to a structural reconstruction of extracellular matrix that allows to specify mechanisms of arrhythmogenesis, to optimize early diagnosis and to determine tactics of treatment of patients with the DCA^{9,10}.

The above mentioned positions give a basis of the aim and tasks of this examination.

The aim is to examine markers of synthesis and collagen degradation, parameters of electrophysiological condition of myocardium among patients with syndrome and phenomenon WPW and to establish possible correlations between them.

MATERIALS AND METHODS OF EXAMINATION

There were examined 43 patients (the main group) with syndromes of preexcitation of ventricles in the specialized cardiological clinic. Among them there were 29 men and 14 women with the average age of $25,3 \pm 9,15$ years old. Among patients of the main group two subgroups were formed. The patients with phenomenon WPW ($n=16$) were in the first subgroup, the patients with syndrome WPW ($n=27$) were in the second subgroup. The both cohorts were comparable by gender distribution (F-criterion, $p=0,52$) and age (U-criterion, $p=0,42$). The following DCAs were discovered during the examination among patients with syndrome WPW: atrioventricular reentrant tachycardia (AVRT) among 8 people (30%), atrial flutter (AF) among 6 patients (22%), atrial fibrillation and atrial flutter among 13 patients (48%). The comparison group consisted of 15 practically healthy people comparable by sex and age.

The inclusion criteria were defined to form the main group: men and women at the age from 17 to 45 years old, presence among patients the clinical and instrumental data that testified the syndrome or phenomenon WPW. The ECG criterion includes: contraction of the interval $PQ < 120$ msec, presence of the A-wave, confluent character and complex broadening $QRS > 110-120$ msec, change of the segment ST and T-wave discordant to the direction of the complex QRS, presence of the DCA, features of additional conducting ways (ACW) discovered during the EPS.

In order to form homogeneous groups the exclusion criteria were introduced: age younger than 17 and older than 45, illnesses of organs of hepatobiliary zone, diabetes mellitus I and II types (DM), carbohydrate intolerance, hard obstructive pulmonary disease, bronchial asthma, pneumofibrosis, chronic kidney disease, connective tissue diseases, malignant neoplasms, patients with implanted cardioverter defibrillator, electric cardiac pacemaker, hypertensive disease, symptomatic arterial hypertension (AH), patients with chronic cardiac failure (CCF) II FC NYHA, various forms of coronary artery disease (CAD), non-coronarogenic myocardial diseases, acquired affections of valvular apparatus and congenital heart diseases, sick sinus syndrome (SSS) and thyroid function abnormality.

During the examination all patients had general and clinical examination that included analysis of complaints, gathering of medical history and life and physical examination. General and biochemical blood tests and determination of thyroid level were valued. Electrocardiography in 12 leads, long ECG monitoring with defining the cardiac rhythm variability, echocardiography and also electrophysiological study were included in the program of instrumental examination. The quantitative tissue inhibitor of metalloproteinase-1 (TIMP-1) and carboxyterminal propeptide (PIP) - procollagen type I in the blood serum were studied to value the indices of fibrosis factors. The level of TIMP-1 was defined with the help of panels "Human TIMP-1 ELISA" (Bender Med Systems, Austria) by the method of immune-enzyme analysis (IEA) that was realized in several stages. On the first stage antibodies to TIMP-1 adsorbed in the tablet cells are connected with TIMP-1 that is in the samples. Added biotinylated anti-TIMP-1

antibodies are connected with TIMP-1 caught by sorbed antibodies in the wells. After incubation and washing unconjugated biotin conjugate- TIMP-1 is removed from the cells and conjugate streptavidin peroxidase is added. After incubation unconjugated biotin conjugate is removed and substrate solution is added. It forms colouring when interaction with enzyme system. The intensity of colouring is directly proportional to the concentration of TIMP-1. The estimation of PICP was realized by using diagnostic set of "MetraCIPCEIAKit" (QuidelCorporation, the USA). The method is based on the "sandwich" immune-enzyme analysis in micromonoclonal variant with the use of monoclonal anti-PICP antibodies, immobilized in the wells of microplates, rabbit anti-PICP antiserum, conjugate of goat anti-rabbit antibodies with alkaline phosphatase and substratepNPP for quantitative evaluation of PICP in human serum. The antibodies to PICP have 100 per cent of cross-responsiveness with PICP of human blood serum.

Electrophysiological study of the heart conducting system is realized by the method of transesophageal electrostimulation of the auricles with the use of electrostimulator "Astrocard-PolysystemEP/L" (CAS "Meditec", Russia) and multipolar electrode WEFCA-2, Kamenetz-Podolsk. A number of patients had an examination by the method of transvenous electrocardiac stimulation. A universal diagnostics electrocardiac stimulator "Biotok- esophagogastrroduodenoscopy (EGD) 01 ml" (OOO "L.M.E. "Biotok", Russia) and multipolar electrodes RFMarinrMC (Medtronic, the USA) were used. The procedures were carried out according to the standard methods. The generally accepted electrophysiological parameters were registered during the examination.

RESULTS OF EXAMINATION

The highest indices of MMP-9 were defined among patients with syndrome WPW – $96,1 \pm 33,2$ ng/ml, noticeably lower they were in the group with phenomenon WPW ($54,3 \pm 21,8$ ng/ml; $p=0,0003$). Practically healthy people had minimal indices MMP-9 - $27,4 \pm 10,9$ ng/ml ($p=0,00014$; $p=0,000002$). Maximum level TIMP-1 is defined among patients with phenomenon WPW and practically healthy people ($418,5 \pm 69,8$ ng/ml and $461,7 \pm 72,2$ ng/ml; $p=0,27$). Patients with syndrome WPW TIMP-1 had low indices – $341,1 \pm 90,1$ ng/ml ($p=0,002$; $p=0,00012$). The level PICP among patients with syndrome WPW was $179,9 \pm 76,2$ ng/ml and with phenomenon WPW – $97,8 \pm 31,7$ ng/ml ($p=0,00014$). All examined patients according to the level PICP differed from practically healthy coevals by $69,4 \pm 23,9$ ng/ml ($p<0,05$, U-criterion).

All examined patients had electrophysiological study (EPS). Among all patients with syndromes of ventricles' preexcitation the features of anthesystole were discovered. Among patients from the main group the shortening of the interval PQ is observed and besides the patients with syndrome WPW ($106,18 \pm 16,5$ msec) have it more in comparison with the phenomenon WPW ($115,25 \pm 19,54$ msec, $p=0,004$). When having syndromes of preexcitation the average adiaphoria according to the duration is determined, that is from 220 to 300 msec: when having the syndrome WPW it is $240,0 \pm 47,9$ msec and when having the phenomenon WPW it is $241,25 \pm 30,84$ msec. Patients with the syndrome WPW had a shorter adiaphoria ($p=0,049$). Both groups have the inverse correlation with fibrosis bioindicators - MMP-9 and PICP. From the received data we can indicate the influence of markers' disbalance that characterizes homeostasis of extracellular matrix (EN) on electrophysiological properties of the ACW.

The function of the automatism of sinoatrial node was evaluated with the help of the following parameters: time of functional restoration of the sinoatrial node and corrected time of this indicator, values did not exceed the bounds of the set standards. The speeding up of atrioventricular conductivity (AC) was observed: Wenckebach point was $219,63 \pm 30,57$ imp/min, patients with the syndrome WPW had shorter adiaphoria of the atrioventricular node ($278,15 \pm 27,7$ msec). At that the index of adiaphoria of the atrioventricular conductivity correlated with MMP-9 concentration ($R=-0,57$; $p=0,002$). Among patients with phenomenon WPW arrhythmogenic vulnerability of myocardium was low, the DCA was not provoked. Patients with syndrome WPW without MA are provoked by the DCA by type of the AF,

the AVRT with orthodromic and antidromic conductivity on ventricles and their combinations. Thus people with diagnosed additional atrioventricular connection (AAVC) have changes of electrophysiological properties of conductive system of the heart, electric instability of myocardium in the form of high arrhythmogenic vulnerability that can be determined by ascending current through atrioventricular node and by hyperexpression of factors that reflect rearrangement of the EN at the ultrastructural level.

CONCLUSIONS

1. Patients with syndrome of preexcitation of ventricles have changes of fibrous matrix: the increasing of concentration of MMP-9, PICP and reducing of TIMP-1 in the blood serum. The syndrome WPW is characterized by more expressed disbalance of seromarkers of synthesis and collagen degradation than the phenomenon WPW. The effective adaphoria of additional ways of conducting correlated back with MMP-9 and PICP indices.
2. The received results show the possible participation of fibrous disorganization and myocardial inflammation in the formation of the DCA when having syndromes of preexcitation of ventricles.

REFERENCES

1. Ardashev V.N. Treatment of disorders of cardiac arrhythmias. Eds. Ardashev V.N., Ardashev A.V., Steklov V.I., Moscow. ID Medpractice Publ., 2005. 228 p.
2. Bobrov V.A. Diagnostics of syndromes of preexcitation of ventricles. Eds. Bobrov V.A. and others. Ukraine cardiac journal Publ. No. 4. 98-108 Art.
3. Kirkutis A.A. Methodology of usage the transesophageal electrostimulation of the heart. Eds. Kirkutis A.A., Rimsha E.D., Nyavyrauskas Yu.V. Kaunas. 1990. 82 p.
4. Krivaya A.A., Vasilets L.M., Tuev A.V., Khlynova O.V., Parshakova O.A. New methods of prediction disorders of cardiac arrhythmias when having syndromes of preexcitation of ventricles. The family health in the 21st century. 2015. T-1. 201-205 Art.
5. Kruchina T.K. Gender and age peculiarities of clinical flows of WPW among children. Eds. Kruchina T.K., Vasichkina E.S., Egorov D.F., Tatarskii B.A. Russian Vestnik of perinatal medicine and pediatrics Publ., 2012. T-57. No. 1. 3338 Art.
6. Kushakovskii M.S. Cardiac arrhythmia. Eds. Kushakovskii M.S. Saint Petersburg. Foliant Publ., 2012. 642 p.
7. Medvedev M.M. To the question on indices of transesophageal electrophysiological examination Eds. Medvedev M. M. Vestnik of arrhythmology Publ., 2000. No. 20. 55-62 Art.
8. National recommendations on determination the risk and prophylaxis of sudden cardiac death. Eds. Shlyakhto E.V. (Academic of the RAMS), Arutyunova G.P. (Professor) and Belenkova Yu. N. (Academic of the RASM and Fellow of the RAS). Clinical practice Publ., 2012. No. 4. 2-94 Art.
9. Elevated pre-operative serum peptides for collagen I and III synthesis result in post-surgical atrial fibrillation / Michael F. Swartz [et al.] // Journal of the American College of Cardiology. 2012. Vol. 60 (18). 1799-1806 pp.
10. Impact of biomarkers of inflammation and extracellular matrix turnover on the outcome of atrial fibrillation ablation: importance of matrix metalloproteinase-2 as a predictor of atrial fibrillation recurrence / Yasuo Okumura [et al.] // J Cardiovasc Electrophysiol. 2011. 22(9). 987-993 pp.
11. Jaiswal A. Coexistent Brugada syndrome and Wolff-Parkinson-White syndrome: what is the optimal management? / A. Jaiswal, K. Heretis, S. Goldbarg // Indian Pacing and Electrophysiology Journal. 2013. 13 (5). 173-177 pp.
12. Matrix metalloproteinases. Zitka O. [et al.] // Current Medical Chemistry. 2010. No. 17. 3751-3768 pp.
13. Matrix metalloproteinases and their inhibitors in vascular remodeling and vascular disease / J.D. Raffetto [et al.] // Biochem Pharmacol. 2008. 75(2). 346-359 pp.

14. PACES/HRS Expert consensus statement on the management of asymptomatic young patient with a Wolff-Parkinson-White (WPW, ventricular preexcitation) Electrocardiographic pattern / B.C. Cannon [et al.] // Heart Rhythm. 2012. 9. 1006-1024 pp.
15. Heidari S, Imani M, Mostafavi SM. A Validated and Rapid HPLC Method for Quantification of Human Serum Albumin in Interferon beta-1a Biopharmaceutical Formulation. MedBioTech Journal. 2017;1(01):29.
16. Obidoa P, Ugueche U. In-vitro Investigation of Ocular E-Coli Infection via Norfloxacin. Medbiotech Journal. 2017;01(02):96-103.
17. Riahi A. Oil-in-Water Emulsion-Solvent Evaporation Fabrication of Polycaprolactone Containing Piroxicam for Drug Release. Medbiotech Journal. 2018;02(01):14-20.
18. Abdulhameed AA, Al-Hamdi KE, Mathkooor MA. Clinical Evaluation of Efficacy and Safety of Combined Topical Timolol and Oral Propranolol in Children with Infantile Hemangioma. J Clin Exp Invest. 2018;9(1):1-8. <https://doi.org/10.5799/jcei.413049>
19. Takzare, A., Goudarzi, M., Maleki, A., Soltani, A. E., Kaheh, F., Arab, S., & Espahbodi, E. (2018). Paediatric sedation during upper gastrointestinal endoscopy: Comparison of propofol+ ketamine vs sodium thiopental+ fentanyl. Electronic Journal of General Medicine, 15(6).
20. Madhumathi, D., & Sakthi, D. (2019). Evaluation of difference in bacterial contamination of toothbrushes between patients with gingivitis and patients with healthy gingiva-a pilot study. International Journal of Pharmacy Research & Technology, 9(2)38-43.