



Revista Latinoamericana de Hipertensión
ISSN: 1856-4550
latinoamericanadehipertension@gmail.com
Sociedad Latinoamericana de Hipertensión
Venezuela

Evaluation of the effect of adherence to treatment regimen program on quality of life in atrial fibrillation patients hospitalized in Shahid Chamran Hospital in Isfahan in 2017

Yazdi Zadeh, Fatemeh; Moeini, Mahin; Shafiee, Davood

Evaluation of the effect of adherence to treatment regimen program on quality of life in atrial fibrillation patients hospitalized in Shahid Chamran Hospital in Isfahan in 2017

Revista Latinoamericana de Hipertensión, vol. 14, no. 3, 2019

Sociedad Latinoamericana de Hipertensión, Venezuela

Available in: <https://www.redalyc.org/articulo.oa?id=170263176009>

Derechos reservados. Queda prohibida la reproducción total o parcial de todo el material contenido en la revista sin el consentimiento por escrito del editor en jefe. Copias de los artículos: Todo pedido de separatas deberá ser gestionado directamente con el editor en jefe, quien gestionará dicha solicitud ante la editorial encargada de la publicación.



This work is licensed under Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International.

Evaluation of the effect of adherence to treatment regimen program on quality of life in atrial fibrillation patients hospitalized in Shahid Chamran Hospital in Isfahan in 2017

Evaluación del efecto de la adherencia al régimen de tratamiento sobre la calidad de vida en pacientes con fibrilación auricular hospitalizados en el Hospital Shahid Chamran en Isfahan en 2017

Fatemeh Yazdi Zadeh

Department of Adult Health Nursing, School of Nursing and Midwifery, Isfahan University of Medical Science, Isfahan, Iran, Irán

 <http://orcid.org/0000-0003-2640-1040>

Redalyc: <https://www.redalyc.org/articulo.oa?id=170263176009>

Mahin Moeini

Nursing and Midwifery care Research Center, Nursing and Midwifery faculty, Isfahan University of Medical Sciences, Isfahan, Iran, Irán

moeini@nm.mui.ac.ir

 <http://orcid.org/0000-0001-7569-5116>

Davood Shafiee

Heart Failure Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran, Irán

 <http://orcid.org/0000-0003-3884-5183>

ABSTRACT:

Background and objective: Atrial fibrillation is the most common cardiac arrhythmia, which is associated with reduced quality of life due to prolonged treatment and its recurrence. One of the key goals in the care of patients with atrial fibrillation is increasing adherence to treatment regimen and the recommendations provided by the treatment group. Thus, the present study was designed and carried out with the aim of determining the effect of treatment regimen on quality of life in atrial fibrillation patients hospitalized in Shahid Chamran Hospital in Isfahan in 2017.

Background and objective: Atrial fibrillation is the most common cardiac arrhythmia, which is associated with reduced quality of life due to prolonged treatment and its recurrence. One of the key goals in the care of patients with atrial fibrillation is increasing adherence to treatment regimen and the recommendations provided by the treatment group. Thus, the present study was designed and carried out with the aim of determining the effect of treatment regimen on quality of life in atrial fibrillation patients hospitalized in Shahid Chamran Hospital in Isfahan in 2017.

Methodology: This clinical trial study was conducted on patients with atrial fibrillation arrhythmia in the Cardiac Care Unit (CCU) and Post CCU Unit and Internal Heart Surgery Unit of Shahid Chamran Hospital in Isfahan. A total of 50 people were randomly selected as sample of study and assigned to two groups of test and control. The test group received two 45-minute sessions of adherence and educational booklet and they were followed-up for one month through phone call. The control group also received one session of usual care training individually with regard to the illness. Demographic data and quality of life data were collected through Atrial Fibrillation Effects on Quality of Life (AFEQT) before intervention, and one and three month after the intervention. Data were analyzed by descriptive and inferential statistics.

AUTHOR NOTES

moeini@nm.mui.ac.ir

Results: There was no significant difference between two groups in terms of quality of life and demographic information before the intervention. However, significant difference was seen between the two groups in terms of quality of life one month and three months after the intervention.

Conclusion: The results suggest the positive effects of adherence to treatment regimen and follow-up of the patients by experienced nurses on quality of life in these patients one and three months after discharge.

KEYWORDS: Quality of life, adherence to treatment regimen, atrial fibrillation, nurses.

RESUMEN:

Antecedentes y objetivo: la fibrilación auricular es la arritmia cardíaca más común, que se asocia con una reducción de la calidad de vida debido al tratamiento prolongado y su recurrencia. Uno de los objetivos clave en el cuidado de los pacientes con fibrilación auricular es aumentar la adherencia al régimen de tratamiento y las recomendaciones proporcionadas por el grupo de tratamiento. Por lo tanto, el presente estudio se diseñó y llevó a cabo con el objetivo de determinar el efecto del régimen de tratamiento sobre la calidad de vida en pacientes con fibrilación auricular hospitalizados en el Hospital Shahid Chamran en Isfahan en 2017.

Metodología: este estudio clínico se realizó en pacientes con arritmia por fibrilación auricular en la Unidad de Atención Cardíaca (CCU) y en la Unidad de Post CCU y en la Unidad de Cirugía Interna del Corazón del Hospital Shahid Chamran en Isfahan. Un total de 50 personas se seleccionaron al azar como muestra de estudio y se asignaron a dos grupos de prueba y control. El grupo de prueba recibió dos sesiones de adherencia y un folleto educativo de 45 minutos y fueron seguidos durante un mes a través de una llamada telefónica. El grupo de control también recibió una sesión de entrenamiento de atención habitual individualmente con respecto a la enfermedad. Los datos demográficos y los datos de calidad de vida se recopilaron a través de los efectos de la fibrilación auricular en la calidad de vida (AFEQT) antes de la intervención, y uno y tres meses después de la intervención. Los datos fueron analizados mediante estadística descriptiva e inferencial.

Resultados: no hubo diferencias significativas entre los dos grupos en términos de calidad de vida e información demográfica antes de la intervención. Sin embargo, se observó una diferencia significativa entre los dos grupos en términos de calidad de vida un mes y tres meses después de la intervención.

Conclusión: los resultados sugieren los efectos positivos de la adherencia al régimen de tratamiento y el seguimiento de los pacientes por parte de enfermeras experimentadas sobre la calidad de vida en estos pacientes uno y tres meses después del alta.

PALABRAS CLAVE: Calidad de vida, adherencia al régimen de tratamiento, fibrilación auricular, enfermeras.

INTRODUCTION

Nowadays, the cardiovascular diseases are the most important cause of death in the world. Based on American Heart Association in 2010, the general death caused by cardiovascular disease was 235 per 100000 people¹. In Iran, the first and most common cause of death at all ages and in both genders is cardiovascular disease. Shabani et al reported that cardiovascular diseases account for 46% of all deaths in Iran². One of the most common heart disorders is arrhythmia and atrial fibrillation is considered the most common heart arrhythmia. Approximately 2.3 million Americans suffer from this disease³. The prevalence of this disorder is less than 0.1% in people aged below 45 years and 9% in people aged over 80 years. In other words, the prevalence of the disease is directly related to age⁴. Based on the American Society of Cardiovascular Guideline, the risk factors for atrial fibrillation in 2016 included heart failure, valvular disease, hypertension, obesity, and smoking⁵. This disorder control is crucial due to symptoms and complications such as hemodynamic disorder, palpitation, chest pain, shortness of breath, fatigue, dizziness, thromboembolism, increased heart and brain infarctions, and longer hospitalization and recovery period⁶.

Treatment strategies in these patients include the use of anticoagulants, heart rhythm and beat controllers, catheterization, the use of pacemaker, and cardioversion⁷, which alleviate the symptoms and improves the quality of life in these patients⁸. Patients with atrial fibrillation are at increased risk of recurring fibrillation and are always affected by the disease. For this reason, they should undergo long-term treatment, which may reduce the likelihood of recurrence of fibrillation⁹. Adherence to treatment regimen in these patients is very important in preventing progression of the disease and recurrence of the attacks¹⁰. Non-treatment of symptoms puts the individual at risk of recurring fibrillation attack. Several studies report that using

appropriate regimen is crucial in these patients since it prevents recurrence of attacks¹¹. Adherence to treatment regimen in patients means the skills used to reduce stress, observe the proper diet, doing proper physical activities and proper use of prescribed drugs in order to promote appropriate health behaviors in the target population¹². In general, the promotion of health-related behaviors will result in maintaining the performance, strengthening the people autonomy, increasing their quality of life, and reducing health care costs¹³, since the results of studies show that 53% of the reasons for people death are associated with their life style and non-healthy behaviors¹⁴.

Given the long process of treatment, the patients need to be followed-up with the aim of establishing a continuous and dynamic care relationship, which improves the quality of life of patients and reduces the complications of the disease and promotes the health¹⁵.

There are different self-control and training methods in order to help patients to adapt to lifestyle changes and can improve their health and prevent recurrence of complications. By informing the patients, they can help the patients follow the appropriate treatment regimen and reduce the recurrence of the attacks. Non-adherence to treatment regimen leads to harmful consequences for patients, treatment staff and healthcare system. Based on the World Health Organization, almost 50% of patients do not adhere to treatment regimens¹⁶. Evidence suggests that non-adherence to treatment leads to inadequate control of the disease and puts the patients at a greater risk of adverse effects¹⁷. The costs related to direct and indirect consequences of non-adherence to treatment, including hospitalization and disease progression, imposes high burden on the health system¹⁸. The costs imposed on the healthcare system for treatment of atrial fibrillation is about 6.2 billion euros in Europe and 6.65 million? dollars in the United States¹⁹. These costs are spent to treat the atrial fibrillation complications includes stroke, heart failure and hospitalization²⁰.

In a study conducted by Kimmel et al with the aim of evaluating the effect of adherence to the treatment regimen on the control of anticoagulant drugs, results showed that patients who had problems in maintaining adherence to the regimen had significant problems ranging from thrombosis to bleeding in coagulation factors²¹. Another study conducted by Lip et al (2015) showed the importance of emphasizing the use of guidelines to prevent stroke in patients with atrial fibrillation²². In a study conducted by Hessel et al in 2011, the results showed that the intensity of heart rate control had no impact on the quality of life of these patients, but symptoms, gender, age, and severity of the disease affect the quality of life of these patients²³. Another study conducted by Lakkireddy et al in 2013 showed that yoga in atrial fibrillation patients improved the disease symptoms, arrhythmias, heart rate, hypertension, anxiety and depression, and different domains of quality of life²⁴. Moreover, there is a correlation between the adherence to treatment and the quality of their life. In this regard, the World Health Organization defines the quality of life as people perception of the status of life in the form of culture and values governing the community in line with the goals, standards, expectations, and interests of the individual²⁵. Thus, atrial fibrillation is one of the most common heart arrhythmias and one of the important challenges in treatment of this disease is adherence to treatment regimen. Moreover, high costs are imposed on the healthcare system by frequent hospitalization of patients. Few studies have been conducted on atrial fibrillation patients in Iran and foreign countries and contradictory results have been reported by these studies. Given what was stated above, the researcher decided to investigate the effect of a program of adherence to treatment regimen on quality of life of atrial fibrillation patients hospitalized in Shahid Chamran Hospital in Isfahan in 2017.

MATERIALS AND METHODS

The present study was a clinical trial with two-group and three-stage design. It was conducted on atrial fibrillation arrhythmia patients hospitalized in the Cardiac Care Unit (CCU) and Post CCU Unit and

Internal Heart Surgery Unit of Shahid Chamran Hospital in Isfahan in 2018. The sample size was calculated to be 25 people in test group and 25 people in control group based on the 95% confidence level and the test power of 80%. They were selected from patients with atrial fibrillation who met the inclusion criteria of the study. They were randomly assigned to the control and test groups using the odd and even numbers card. The research inclusion criteria included age range of 35-70 years, hospitalized for the first time in a hospital with medical diagnosis and registration of atrial fibrillation in the medical record, lack of limiting motor disorders and non-participation in intervention programs similar to adherence to treatment regimen, the having the reading and writing literacy and the ability to speak Farsi and the willingness to participate in the study.

After obtaining informed written consent from the samples and receiving the ethics code from the Isfahan University of Medical Sciences, the researcher provided the usual care with regard to the disease for control group samples in a 30-minute face to face session. The test group also received two 45-minute sessions of adherence and educational booklet and they were followed-up for one month through phone call. Adherence to treatment regimen intervention included two educational and support sections. In the educational section, the researcher provided the educational content derived from the sources and in accordance with the components of the AFEQT questionnaire, approved by the cardiologist, as a combination of individual training and practice in two consecutive 45-minute sessions on the second and third days of hospitalization during 8-12 am on the patient's bedside. It included information on the type of arrhythmia, the methods of treatment, the drugs used and its complications, the level of activity, the method of coping with mental problems, the importance of performing coagulation tests, the type nutrition. They were provided in the face to face manner and by using the images of educational booklet.

In the supportive section, the researcher provided the content of topics discussed in the sessions, question and answer and the disease control and management power to the control group samples in a weekly phone call between the hours 8 am and 8 pm. The duration of each phone call was 10 minutes based on the patient's need. No intervention was provided for this group during these 8 weeks. Finally, 12 weeks after the beginning of the intervention, the questionnaire was completed by a researcher assistant in the test and control groups. In the case of physical and psychological problems resulting in disability, death of the patient and lack of willingness to continue the study, the person was excluded from the study.

The data collection tool included a two-part questionnaire. The first part consists of demographic characteristics and disease information. The second part consists of a questionnaire on the effects of atrial fibrillation on quality of life (AFEQT). It is a standard questionnaire confirmed by the AFEQT Core team in North America using Cronbach Alpha (0.9). To confirm the validity of this questionnaire, the researcher used the content validity and the opinions of eight faculty members of the Faculty of Nursing and Midwifery and two cardiologists were applied on the questionnaire. Then, Cronbach's alpha coefficient was used to determine the reliability of this questionnaire, so that the questionnaire was provided to ten patients with atrial fibrillation who met the inclusion criteria of the study after applying the necessary modifications in it. It was confirmed with Cronbach's alpha was 0.87. The questionnaire includes 20 questions scored on a 7-point Likert scale in three domains of symptoms and symptoms, activity, treatment concerns. The minimum score was zero and the maximum score was 100. A separate score was calculated for each domain. The higher scores mean the higher quality of life. The questionnaire was completed in three stages by a researcher assistant. Before the intervention, it was completed through questioning in face to face and one and three months after the intervention, it was completed by phone call and questioning. The collected data were analyzed using SPSS21 software. In analyzing the data, independent t-test, Mann-Whitney, Chi-Square, Fisher, covariance and LSD post hoc tests were used

RESULTS

The results showed that in the test group, 44% of the patients were male and 56% were female, and in the control group, 48% of the patients were male and 52% were female. The highest percentage of samples in both groups was self-employed. In both groups, 60% of the samples were married. In the test group, 56% had education lower than diploma, and in the control group, 48% had education lower than diploma. The highest number of samples in both groups was in the age group of 51-55 years and the most of the risk factor in the test group were hypertension (36%), and in the control group, it was smoking (40%). The most common cause of hospitalization was hypertension (28%) in the test group and heart failure (28%) in the control group. The duration of the disease was 1-3 months (32%) in the test group and 3-6 months (40%) in the control group.

Independent t-test showed that the mean of total score of quality of life and its dimensions before intervention was not significant between the two groups ($P>0.05$). However, the covariance analysis showed that the mean of total score of quality of life, and the dimensions of signs and symptoms and physical activity one month after the intervention in the test group was significantly more than that of the control group ($P<0.05$). However, the mean score of quality of life in the dimensions of treatment problems and satisfaction one month after intervention showed no significant difference between two groups ($P>0.05$). Covariance analysis showed that the mean of total score of quality of life and all its dimensions were significantly higher in the test group three months after intervention than that in the control group ($P<0.05$) (Table 2).

TABLE 1 .
Comparing the disease-related demographic information of test and control groups

variable		Test group n	%	Control group n	%	test	χ^2	P
Patient's gender	male female	11 14	44 56	12 13	48 52		08.0	78.0
job	employee Self-employed Retired housewives others	5 10 6 3 1	20 40 24 12 4	4 8 6 7 0	16 32 24 28 0		37.3	50.0
Marital status	single married divorced spouse deceased	2 15 3 5	8 60 12 20	2 15 2 6	8 60 8 24	Chi-square	29.0	96.0
Risk factors	Hypertension High blood fat High blood glucose Smoking others Heart failure	9 6 3 5 2	36 24 12 20 8	6 5 3 10 1	24 20 12 40 4		73.2	60.0
Cause of hospitalization	Hypertension heart attacks ischemic disease others	4 7 5 3 4 2	16 28 20 12 16 8	7 5 4 3 5 1	28 20 16 12 20 4		73.1	89.0
Age range (year)	40-35 45-41 50-46 55- 51 60-56 65-61 Over 65	1 2 3 7 5 3 4	4 8 12 28 20 12 16	2 3 4 5 4 3 4	8 12 16 20 16 12 16	Z	54.0	59.0
Level of education	Under diploma diploma academic	14 9 2	56 36 8	12 10 3	48 40 12	Mann-Whitney	62.0	54.0
Disease duration	1-3 month 3-6 month 6- 12 month 1-2 year 2-5 year 5-10 year 10-15 year Over 15 years	8 3 5 1 2 2 1 3	32 12 20 4 8 8 4 12	4 10 3 2 3 2 0 1	16 40 12 8 12 8 0 4		22.0	83.0

DISCUSSION

The results showed that there was no significant difference between the two groups in terms of demographic information (age, gender, job, level of education, marital status). In addition, no significant difference was found between two groups in terms of information related to disease, such as risk factors, duration of disease and the cause of hospitalization. These results confirm the random allocation of samples in two groups. Moreover, the results of this study showed that the mean score of quality of life and its dimensions before intervention was not significantly correlated between the two groups. However, there is a significant difference between two groups in terms of mean of total score of quality of life one and three months after intervention.

In confirmation agreement with these results, Shojaei et al (2013) showed that the psychological dimension of quality of life could change under severe conditions, including chronic diseases, feeling disability, fear of death, reduced function and many different conditions. Their results also showed that training during discharge and then phone follow-up can increase the level of hope in patients with heart failure and reduce costs to improve the health of patients ¹.

In the present study, adherence to the treatment regimen improved the quality of life of atrial fibrillation patients. One of the important factors playing a major role in controlling the disease is the patient's adherence to the treatment regimen. There is a mutual relationship between adherence to treatment regimen and quality of life of the patients. Receiving the adherence to treatment regimen program in addition to having a significant effect on the general mean of these people's quality of life compared to that in the control group had also a significant effect on the quality of life of the patients in the intervention group. In a study conducted by Roodsari et al. (2013), the results consistent with those of this study were obtained, so that their results showed that there is a significant relationship between the health-related quality of life and adherence to treatment regimen in patients with hypertension ².

In a study conducted by Guo et al in 2017 to evaluate the effect of mobile technology on better management of atrial fibrillation, knowledge, quality of life, drug adherence, and satisfaction with anticoagulants were measured at the beginning of the study, one and three months later by designing a simple program on Android OS and ios in two groups of atrial fibrillation patients. The results showed that a proper clinical and educational process in the management program of these patients would lead to improvement in the continuous adherence to drug therapy, knowledge level and satisfaction with receiving anticoagulants, as well as quality of life in these patients ³².

In this regard, a study carried out by Tekeh Fallah et al in 2016 showed that distance nursing for three months would improve the quality of life of patients with fibroblasts ²⁶. These results were consistent with those of our study. One of the reasons for the consistency might be the duration of follow-up which it was similar in two studies (3 months) ³³. However, the researcher used distance education in this study, but the education was provided by the researcher and in face to face manner in the present study ³⁴. In another study conducted by Lakkireddy et al in 2013, they showed that yoga improved disease symptoms, arrhythmias, heart rate, blood pressure, anxiety, depression, and various domains of quality of life in patients with atrial fibrillation ²⁴. The results of this study also showed that yoga in the life of patients with atrial fibrillation can be a factor affecting their quality of life.

In addition, in another study conducted by Lip et al (2015), the use of guidelines to prevent stroke in patients with atrial fibrillation was emphasized ²². Given the results of this study, one of the reasons for the significant difference between the two test and control groups in the present study might be attributed to the implementation of several educational sessions, follow-up after discharge, and presentation of the educational booklet. Adherence to treatment regimen is a new intervention, which has not been investigated due to its importance in atrial fibrillation. For this reason, few articles were found in this regard. In a study

conducted by Esmkhani et al in 2017, the effect of adherence to the treatment regimen on quality of life in patients with schizophrenia was studied ³⁵. In this clinical trial, 70 patients with schizophrenia were divided into test and control groups. In the test group, six sessions of lecture and discussion were held for 30-60 minutes per week and the control group received the usual cares ³⁶. The results showed that the implementation of adherence to treatment regimen programs improved the quality of life of schizophrenic patients. These results are in line with those of our study ³⁷. However, in the present study, the quality of life of patients was evaluated one and three months later. However, it was examined immediately after the intervention ^{37,39}.

The results of this study also showed that there was no significant difference between the mean scores of the dimensions of quality of life (signs and symptoms, physical activity, treatment problems, and satisfaction with treatment) in patients with atrial fibrillation in the test and control groups before intervention ³⁸. However, there was a significant difference between the mean scores of signs and symptoms in patients with atrial fibrillation in the control and test groups one month after the intervention and three months after the intervention ($p < 0.05$). This suggests that follow-ups after discharge can affect the quality of life of the patient. In this regard, in an 8-month study, Wang et al investigated the effect of drug adherence on depression and quality of life in AIDS patients and showed that quality of life in the test group was significantly higher than that of the control group. In this study, phone intervention increased the quality of life scores in the different dimensions in the intervention group, while in the present study, follow-up treatment with in-person sessions and talking with patients increased the quality of life of patients ^{3,44}. There was no significant difference between test and control groups 1 month after intervention, but there was a significant difference between them 3 months after the intervention ($p < 0.05$). This can indicate that adherence to treatment regimen needs more time to increase the patient's ability. In addition, in this study, there was no significant difference between the test and control groups in terms of mean scores of concerns as one of the dimensions of quality of life of the patients one month after the intervention and three months after the intervention. However, the study conducted by Kirch et al showed that phone follow-up even after 48 hours and providing guidance to the patient's family could improve the quality of patient care. This contradiction at follow up time might be due to the difference in the objectives of the present study, which this study focused more on the quality of life ^{4,43}.

There was no significant difference between the mean scores of satisfaction of the treatment process in patients with atrial fibrillation in the test and control group one month after the intervention, but there was a significant difference between them three months after the intervention ($p < 0.05$). In this regard, Davis et al conducted a research and showed that phone call for six months had an impact on quality of life and the satisfaction of patients with advanced diseases. This might indicate that increasing the duration of phone follow-up can increase the satisfaction of the patients ^{5,42}. In general, the results of this study and other studies confirm the positive effect of adherence to treatment regimen programs in patients with atrial fibrillation. In addition, quality of life is a concept related to the field of nursing and nurses have always tried to improve it ²⁷. By providing health care and participating in nursing research, they tried to improve the quality of life of patients. In this regard, Scott (2004) showed that patients with severe heart failure experience anxiety and depression and loss of control over life caused by the disease ³⁰. In general, a few studies conducted in this regard show that quality of life in patients with heart failure is weaker compared to other chronic diseases ^{28,41}. Another study conducted by Rodriguez et al (2008) found that the training of patients with heart failure was the base of individual management of the heart failure ^{29,40}.

Harrison (2002) showed that providing training by a specialist nurse for the samples and visiting them two weeks after discharge and their follow-up for 12 weeks increased the quality of life of the test groups in the physical and mental dimensions and this increase was significant 6 weeks in the test group, which is in line

with the results of our study. Stewart et al stated that heart failure treatment by a specialist nurse improved the quality of life of the patients ³¹.

CONCLUSION

Thus, it can be concluded that the implementation of an adherence to treatment regimen program based on existing guidelines and follow up of the patients during the hospitalization one and three months after discharge by using experienced nurses is crucial to enhance their knowledge, empower and promote the dimensions of quality of life of the patients. It is important to use this program to improve the quality of life of patients with atrial fibrillation.

AGRADECIMIENTOS

This paper was derived from a master's thesis in Nursing approved by Isfahan University of Medical Sciences with number of 396635. We appreciate the Research Deputy of the Isfahan University of Medical Sciences, the patients and their family members, nursing staff and officials of Shahid Chamran Hospital in Isfahan and other people who helped us in conducting this study.

REFERENCES

1. Go, A.S, Mozaffarian, D, Roger, V.L, Benjamin, E.J, Berry, J.D, Michael J. Blaha, et al, 2014, "Heart disease and stroke statistics", American Heart Association Statistics Committee and Stroke Statistics Subcommittee, 129(3):p.e28-e292, available on line <https://www.ncbi.nlm.nih.gov/>
2. Shabani, R, Gaeini, A, Niko, MR, Nikbakht, H, Sadeghi Far, M, 2010, "Comparison of response to heart rehabilitation program (resistance and endurance exercise simultaneously) in men and women with coronary artery disease". Gilan University of Medical Sciences, issue 74, pp. 48-57, Observed on 2017.02.17, available at gums.ac.ir
3. Krijthe BP, Kunst A, Benjamin EJ, Lip GY, Franco OH, Hofman A et al, 2013, "Projections on the number of individuals with atrial fibrillation in the European Union, from 2000 to 2060, Journal of Eur Heart, 34,p2746–2751 doi:10.1093/eurheartj/ehz280.
4. Ionescu-Ittu, R, Abrahamowicz, M, Jackevicius, C.A, Essebag, V, Eisenberg, M.J, Wynant, W, ETAL, 2012" Comparative Effectiveness of Rhythm Control vs Rate Control Drug Treatment Effect on Mortality in Patients with Atrial Fibrillation", Journal of ARCH INTERN MED/VOL 172(13):997-1004. <http://jamanetwork.com/>
5. Kirchhof, P, Benussi, S, Kotecha, D, Ahlsson, A, Atar, D, Casadei, B, 2016" 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS", Journal of European Heart,1-90, viewed at 14 September 2016 available on line <http://eurheartj.oxfordjournals.org/>
6. Kirchhof, P, Ammentorp, B, Darius, H, Caterina, R.D, Heuzey, J, Schilling, R, 2014" Management of atrial fibrillation in seven European countries after the publication of the 2010 ESC Guidelines on atrial fibrillation: primary results of the prevention of thromboembolic events—European Registry in Atrial Fibrillation (PREFER in AF)", Published by Oxford University Press on behalf of the European Society of Cardiology, 16, P 6–14, doi:10.1093/europace/eut263
7. January, C.T, Wann, L.S, Alpert, J.S, Field, M.E, Calkins, H, Murray, K.T, et al, 2014," 2014 AHA/ACC/HRS Guideline for the Management of Patients With Atrial Fibrillation", Journal of the American College of Cardiology, 129:000–000, available on line <http://circ.ahajournals.org>

8. Lorig, k, Ritter, P.L, Pifer, C, Werner, P, 2014". Effectiveness of the Chronic Disease Self-Management Program for Persons with a Serious Mental Illness", journal of Community mental health, 50(1): p. 96-103, available on line <https://link.springer.com/>
9. Smeulders, E, Haastregt, J, Ambergen T, Stoffers ,H, Janssen-Boyne, J, Uszko-Lencer, N,etal,2010" Heart failure patients with a lower educational level and better cognitive status benefit most from a self-management group programme", Patient education and counseling 81(2):p.214-221, available on line www.elsevier.com
10. McCahona, D, Murraya, E.T, Murrayb, K, Holdera, R.L, Fitzmauricea, D.A, 2011" Does self-management of oral anticoagulation therapy improve quality of life and anxiety ?", Published by Oxford University Press, 28(2):p.134-140, available on line <https://watermark.silverchair.com/>
11. Markou AL, van der Windt A, van Swieten HA, Noyez L, 2008" Changes In Quality Of Life, Physical Activity And Symptomatic Status One Year After Myocardial Revascularization For Stable Angina" European Journal of Cardio-thoracic Surgery;34(5):1009-15. , available on line www.elsevier.com/locate/ejcts
12. World Health Organization 2003, ADHERENCE TO LONG-TERM THERAPIES Rand CS. Measuring adherence with therapy for chronic diseases: implications for the treatment of heterozygous familial hypercholesterolemia. American Journal of Cardiology,1993, 72:68D-74D adherence@who.int
13. Habibi Sola, A, Nikpour, S, Seyed Alshohadaei, M, Haghani, H, 2008, "Relationship between quality of life and physical activities in the elderly people", Iranian Journal of Nursing, Volume 21, Issue 53, pp. 29-39 , observed on 2016.04.11, available at www.sid.ir
14. Samimi, R, Masrur, D, Hosseini, F, Tammadonfar, M, "The Relationship between Lifestyle and General Health of Students", Journal of Nursing, 2006, Volume 19, Issue 48, pp. 83-93.
15. Fairchild, L.S, Varghese, SH.B, Deickman, A, Castelli, D, 2010," Telehealth and Telenursing Are Live: APN Policy and Practice Implications ", The Journal for Nurse Practitioners ,6(2):98-106 available on line <https://ac.els-cdn.com/>
16. World Health Organization 2003. Adherence to long term therapies: evidence for action. http://www.who.int/c hp/knowledge/publications/adherence_report/en/(10 September 2012).
17. Singh, S, 2012," Costs and clinical consequences of suboptimal atrial fibrillation management", Journal of Clinico Econ Outcome Research 4:p79–90.
18. Aslam F, Haque A, Foody JM. 2012"Medication non-adherence: the other drug problem". Hot Topics Cardiol 7:p7–14.
19. Ringborg A, Nieuwlaat R, Lindgren P, Jo"nsson B, Fidan D, Maggioni AP, Lopez-Sendon J, Stepinska J, Cokkinos DV, Crijns HJ,2008," Costs of atrial fibrillation in five European countries: results from the Euro Heart Survey on atrial fibrillation", Published on behalf of the European Society of Cardiology10:p403–411.
20. Camm AJ, Lip GYH, De Caterina R, Savelieva I, Atar D, Hohnloser SH, Hindricks G, Kirchhof P. ESC Committee for Practice Guidelines (CPG), Document Reviewers. 2012.. Focused update of the ESC Guidelines for the management of atrial fibrillation: an update of the 2010 ESC Guidelines for the management of atrial fibrillation developed with the special contribution of the European Heart Rhythm Association",2012,Eur Heart J 33:p2719–2747.
21. Kimmel, S.E, Chen, Zh, Price, M, Parker, C.S, Metlay, J.P, Christie, J.D, etal,2007, "The Influence of Patient Adherence on Anticoagulation Control With Warfarin" journal of Arch Intern Med, vol 167p 229-235 available on line <http://jamanetwork.com/>
22. Lip, G.Y.H, Laroche, C, Popescu, M.L, Rasmussen, L.H, Serdoz, L, Dan, GH,etal,2015" Improved outcomes with European Society of Cardiology guideline-adherent antithrombotic treatment in high-risk patients with atrial fibrillation: a report from the EORP-AF General Pilot Registry" Published on behalf of the European Society of Cardiology, 17, p1777–1786.
23. Hessel F. Groenveld, MD, Harry J. G. M. Crijns, MD, Maarten P. Van den Berg, MD,2011," The Effect of Rate Control on Quality of Life in Patients With Permanent Atrial Fibrillation", Journal of the American College of Cardiology, Vol. 58, doi:10.1016/j.jacc.2011.06.055.

24. Lakkireddy, D, Atkins, D, Pillarisetti, J, Ryschon, K , Bommana, S , Drisko, J, Etal, 2013, "Effect of Yoga on Arrhythmia Burden, Anxiety, Depression, and Quality of Life in Paroxysmal Atrial Fibrillation" Journal of the American College of Cardiology, Vol. 61:1177-1182, available on line <http://dx.doi.org/>
25. WHOQOL Group: The World Health Organization Quality of Life assessment (WHOQOL): position paper from the World Health Organization. Soc Sci Med 1995, 41:1403
26. Yekeh Fallah, L, Najafi, M, Shahrokhi, A, Javadi, HR, Mohammad Pour, A (2016). The Impact of Distance Nursing on Quality of Life in Patients with Atrial Fibrillation Referring to Qazvin Educational Centers. Journal of Qazvin University of Medical Sciences, Volume 20, Issue 1, pp. 56-62.
27. Sharifi Mohammad Hossein, Eftekhari Mohammad Hassan, Ostovan Mohammad Ali, Rezaianazadeh Abbas. Effects of a therapeutic lifestyle change diet and supplementation with Q10 plus L-carnitine on quality of life in patients with myocardial infarction: A randomized clinical trial. J Cardiovasc Thorac Res. 2017;9(1):21–28.
28. Scott Linda, Setter-kline Kay, Britton Agnes, The effects of nursing intervention to enhance mental health and quality of life among individuals with heart failure. Appl Nurs Res. 2004; 17(4):248-256.
29. Rodriguez KL, Appelt J, Switzer GE, Sonel AE, Robert MA. "They diagnosed bad heart": A qualitative exploration of patients' knowledge about and experiences with heart failure. Heart Lung 2008 37:257-65.
30. Harrison MB. Quality of life individuals with heart failure: A randomized trial of the effectiveness of two models of hospital to home transition. Med Care J 2002;40(4):271-82.
31. Stewart M. clinical service frame work for heart failure draft volume 2. 2003. availabe from: <http://www.health.nsw.gov.au>.
32. Yutao Guo, Yundai Chen, Deirdre A. Lane, Lihong Liu, Yutang Wang, Gregory Y.H. Lip, Mobile Health Technology for Atrial Fibrillation Management Integrating Decision Support, Education, and Patient Involvement: mAF App Trial. The American Journal of Medicine (2017) 130,1388–1396
33. Shojaei, A, Nahrir, B, Naderi, N, et al. The effect of patient education and phone follow up by nurse on hope of heart failure patients. Nursing Education Journal. 2013; 2 (1): 16-26
34. Rudsari M, Dokht D, Golchin, D, Yekta, P, Haqani. The relationship between adherence to treatment regimen and the quality of life associated with health in patients with hypertension. Iranian Journal of Nursing. 2013; 26 (85): 44-54.
35. Wang H, Zhou J, Huang L, Li X, Fennie KP, Williams AB. Effects of nurse - delivered home visits combined with telephone calls on medication adherence and quality of life in HIV - infected heroin users in Hunan of China. Journal of clinical nursing. 2010; 19 (3 - 4):380-8.
36. Kirsch SD, Wilson LS, Harkins M, Albin D, Del Beccaro MA. Feasibility of using a pediatric call center as part of a quality improvement effort to prevent hospital readmission. Journal of pediatric nursing. 2015;30(2):333-7.
37. Davis MS, Harrison KL, Rice JF, Logan A, Hess B, Fine PG, et al. A model for effective and efficient hospice care: proactive telephone-based enhancement of life through excellent caring, "telecaring" in advanced illness. Journal of pain and symptom management. 2015;50(3):414-8.
38. Esm Khani, M, Ahangarzadeh Rezaei, S, Khalkhali, HR, The Effect of Adherence to Treatment on the Quality of Life in Patients with Schizophrenia, Journal of Urmia Nursing and Midwifery Faculty, Volume 15, Issue 8, November 2017, pp. 595-604.
39. Kilitci A, Kaya Z, Acar EM, Elmas ÖF. Scrotal Calcinosiis: Analysis of 5 Cases. J Clin Exp Invest. 2018;9(4):150-3.
40. KORKMAZ, Sevda, et al. Sociodemographic and clinical characteristics of patients with violence attempts with psychotic disorders. *European Journal of General Medicine*, 2017, 14(4): 94-98.
41. Mardani, M., Lavasani, S. M., & Omidvari, M. (2014). An investigation into DOW and MOND indices with fuzzy logic based on fire and explosion risk assessment in Iran oil refinery, *UCT Journal of Research in Science, Engineering and Technology*, 2(3): 126-137.
42. Yazdekhesti, A., Erfan, N., & Nazari, N. (2015). Investigating the Relationship between Spiritual Intelligence and Social Adaptation among Girl High School Students in Shahreza City. *UCT Journal of Social Sciences and Humanities Research*, 3(1), 20-23.

43. Ece A, Tünay Z. Successful management of acute bismuth intoxication complicated with acute renal failure, seizures and acute pancreatitis in a child. *J Clin Exp Invest*. 2018;9(3):131-4.
44. Kasimovskaya, N. A., Fattakhutdinova, E. V., Khuziakhmetov, A. N., Kutlin, N. G., Kostina, Y. A., & Kochetkov, I. G. (2018). An attributive component of the condition of people with the diagnosis of " Ischemic heart disease. Effort angina". *Electronic Journal of General Medicine*, 15(6).