



Revista Latinoamericana de Hipertensión  
ISSN: 1856-4550  
latinoamericanadehipertension@gmail.com  
Sociedad Latinoamericana de Hipertensión  
Venezuela

## Investigation of drug -drug interactions status, in University-Based Pharmacies, in Lorestan, IRAN

**Babak, Hadian; Mohsen, Asadbegi; Zokaei, Hamid**

Investigation of drug -drug interactions status, in University-Based Pharmacies, in Lorestan, IRAN

Revista Latinoamericana de Hipertensión, vol. 14, no. 3, 2019

Sociedad Latinoamericana de Hipertensión, Venezuela

**Available in:** <https://www.redalyc.org/articulo.oa?id=170263176020>

Derechos reservados. Queda prohibida la reproducción total o parcial de todo el material contenido en la revista sin el consentimiento por escrito del editor en jefe. Copias de los artículos: Todo pedido de separatas deberá ser gestionado directamente con el editor en jefe, quien gestionará dicha solicitud ante la editorial encargada de la publicación.



This work is licensed under Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International.

## Investigation of drug -drug interactions status, in University-Based Pharmacies, in Lorestan, IRAN

Investigación del estado de las interacciones entre medicamentos y drogas, en farmacias universitarias, en Lorestan, IRAN

*Hadian Babak*

*Assistant Professor of nephrology. Department of internal medicine. Lorestan University of medical Sciences, Khorramabad, Iran, Irán*

 <http://orcid.org/0000-0001-8014-7528>

Redalyc: <https://www.redalyc.org/articulo.oa?id=170263176020>

*Asadbegi Mohsen*

*Pharmacist. Manager of Drug Supervision, Lorestan University of Medical Sciences, Khorramabad, Iran, Irán*

 <http://orcid.org/0000-0001-5674-6109>

*Hamid Zokaei*

*Department of Anesthesiology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran, Irán*

[hamidzokaei@gmail.com](mailto:hamidzokaei@gmail.com)

 <http://orcid.org/0000-0002-0101-0759>

### ABSTRACT:

Drug drug interaction, is an important cause for disturbing of medical management of a disease that usually is preventive, must be continuously investigated for detection of causes. Due to little data about it in recently years in this area, study was done, to investigate of quantity of interactions, and some related factors. In this cross sectional study, all prescriptions data from April to October 2016 were collected. Variable data included: type and name of drugs- the number of drugs in each prescription and characteristics of interactions. We used software of drug interaction fact and lexicomp on desk top DI and textbook 'Drug Interaction Facts'. Statistical analysis done by t test and use of SPSS 19. Finally, 41096 prescriptions were collected. From these data 56% were male. 3043 of subjects had DDI (7.4%). most interaction was in 5<sup>th</sup> degree and majority of interactions were mild. DDI between ammonium chloride and pseudoephedrine -antihypertensive drugs and others-omeprazole and chlorthalidone had the most frequency, respectively. About interaction show, 53% was delay and 43% was rapid. The mean quantity of drugs was less than four type drugs in each prescription. We conclude that, general and even OTC drug had great DDI that must be notice to these drugs in prescriptions. There was no correlation between gender or specialty of physician and frequency of DDI ( $p=0.08$ ). Quantity of drugs in each prescription, correlate with rate of DDI ( $p=.003$ ). Attendance a pharmacist in treatment team of patients could be helpful, for decrease the rate of DDI. In all centers, continuous and more research for detecting DDI is necessary.

**KEYWORDS:** drug interaction, DDI, physician.

### RESUMEN:

La interacción farmacológica, es una causa importante para alterar el manejo médico de una enfermedad que generalmente es preventiva, debe investigarse continuamente para detectar causas. Debido a la poca información al respecto en los últimos años en esta área, se realizó un estudio para investigar la cantidad de interacciones y algunos factores relacionados. En este estudio transversal, se recopilieron todos los datos de las prescripciones de abril a octubre de 2016. Los datos variables incluyeron: tipo y nombre de los medicamentos: la cantidad de medicamentos en cada receta y las características de las interacciones. Usamos software de hechos de interacción de drogas y lexicomp en el escritorio y en el libro de texto "Datos de interacción de drogas". Análisis estadístico realizado mediante la prueba t y uso de SPSS 19. Finalmente, se recogieron 41096 recetas. De estos datos el 56% eran hombres. 3043 de los sujetos tenían DDI (7,4%). la mayor parte de la interacción fue de 5<sup>o</sup> grado y la mayoría de las interacciones

fueron leves. La DDI entre el cloruro de amonio y la pseudoefedrina (medicamentos antihipertensivos y otros), omeprazol y clordiazepóxido tuvieron la mayor frecuencia, respectivamente. Sobre el programa de interacción, el 53% fue retraso y el 43% fue rápido. La cantidad media de medicamentos fue menos de cuatro medicamentos de tipo en cada receta. Llegamos a la conclusión de que, en general, e incluso el medicamento de venta libre tuvo un gran DDI que debe notificarse a estos medicamentos en las recetas. No hubo correlación entre el sexo o la especialidad del médico y la frecuencia de DDI ( $p=0.08$ ). La cantidad de medicamentos en cada receta, se correlaciona con la tasa de DDI ( $p=.003$ ). La asistencia de un farmacéutico en el equipo de tratamiento de pacientes podría ser útil para disminuir la tasa de DDI. En todos los centros, es necesaria una investigación continua y más para detectar DDI.

**PALABRAS CLAVE:** interacción farmacológica, DDI, médico.

## INTRODUCTION:

Medical errors are life threatening events in management of diseases .for increasing safety of patients; physician must be noticed to their causes. DDI (drug drug interactions), is an important cause for disturbing of medical management of a disease.

Patients frequently use more than one medication at a time. Unanticipated, unrecognized, or mismanaged DDIs are important causes of morbidity and mortality associated with prescription drug uses that usually are preventive<sup>1,2</sup>.

Patients, often come to physician with a legacy of drugs, during previous medical experience that if physician not notice to it, risk of DDI may be increased. Some groups of patients, especially old age with multiple long standing disease are predisposed to DDI<sup>3</sup>.

Drug interaction is defined as the pharmacologic or clinical responses to administration of a drug combination, differing from the anticipated known effect of two agents.

Combining drugs may cause pharmacokinetic and/or pharmacodynamics interactions. Pharmacokinetic mechanisms of interaction include alterations of absorption, distribution, biotransformation, or elimination. Pharmacokinetic interactions in general result in an altered concentration of active drug or metabolite in the body, modifying the expected therapeutic response. Interaction must be considered in differential diagnosis of any unusual responses occurring during drug treatment<sup>4,5,6</sup>.

Beside above, DDI may be occurring between prescription drugs but also between food& drug and chemical and drug.it is a dynamic problem , belong to all time and location.it must be continuously investigated for detection of causes and preventive schedule of it. Due to little data about it in recently years in this area, the aim of this study is to investigate of quantity of interactions, and to assay what physicians' prescriptions cause more interactions and to identify common interactions.

## MATERIALS AND METHODS:

In this cross sectional study, all prescriptions data from April to October 2016 were collected. These data were belonging to 2 general hospital pharmacies affiliated to Lorestan University of medical sciences (in capital of province).

All outpatients and inpatients prescriptions were enrolled in study. Variable data included: type and name of drugs - the number of drugs in each prescription - physician specialty and gender, and characteristics of interactions. For detection of DDI, we used software of drug interaction fact and lexicomp on desk top DI. Also, we used reference textbook 'Drug Interaction Facts' published in 2010 for completion of study<sup>7</sup>.

Mean +/- SD and percentage parameters were used.Statistical analysis done by t test and use of SPSS version 19. P value less than 0.05 was considered significant.

## RESULTS

Finally, 41096 prescriptions were collected. From these data 56% were male and others female subjects. Minimum age of patients was 2 and maximum age was 89 years old. Average subject age was 35.5 years old.

Out of all them, 3043 subjects had DDI (7.4%).based on importance, most interaction was in 5<sup>th</sup> degree and respectively, 3<sup>th</sup> and 2<sup>th</sup> degrees were common. Between all DDIs, majority of interactions were mild. About interaction show, 53% was delay and 43% was rapid summarized data was showed in **fig 1**.




FIG 1  
Percentile of DDI characteristics

In this study, most of DDIs belong to combination of pseudo ephedrine and ammonium chloride (in diphenhydramine compound), then respectively: antihypertensive drugs (especially beta blockers and other drugs), omeprazole and chlorthalidone.

In between all prescriptions, NSAIDs – corticosteroids – azithromycin – metoclopramide were the most frequent type of prescribed drugs.

The most quantity of drugs was four type drugs in each prescription. Between one and ten drugs type prescribed in total of papers. Distribution of drugs numbers in each prescription was summarized in **figure 2**.

Between all paper contained DDI, 59% was prescribed by general practitioners and 41% by specialist.




FIG 2.  
Quantity of drug in each prescription

## DISCUSSION:

In term of the DDI severity, interactions can be classified by: Mild interaction denotes that they are not of clinical importance or the effect on the interaction has not been established. Moderate interaction means, that possible changes in therapeutic effect or may cause adverse effects, but can be avoided adjusting the drug doses. Major interaction means, result in potential adverse effects and necessitates individual dose adjustment in the cases.

About time of onset of action: rapid means that the effect will be evident within 24 hr, after administration of drugs and delay reaction means that the effect will not be evident until days or weeks.

About importance of interaction: they divided in five groups. 1. Established: proven to occur in well controlled studies, 2. probable: very likely but not proven clinically, 3. suspected: may occur; need more study, 4. *possible*: could occur but data are limited, 5. Unlikely: doubtful; no good evidence of an altered clinical effect.

At first, in review of literature, we conclude that, there is wide variability between frequencies of DDI in the all centers. In ahmadizadeh study in Iran 77% and in namazi study at least 1 DDI in 43% of all patients reported<sup>8,9</sup>.

In other study (in birjand .Iran), 42% of prescription had DDI. Also, similar study out of Iran showed wide variability in DDI frequency. 18.5% in Greece and 9.8% in Finland and 41% in Nepal<sup>10-13</sup>.

In comparison, we could not detect any rational relation between or result and other study about frequency of DDI. We noticed to methodological effect of studies-different distribution of people age and sex and diversity of type of disease and specialty of physician – time and location of studies for this variation of results. For example some studies, only evaluate patients in critical ward that increase the rate of DDI due to longer admission of patients<sup>14</sup>. Previous study in 2013 showed that drug interactions are somewhat inevitable and like other provinces<sup>15</sup>.

Finally, we suggested that each group of patients must be compare with same groups in other time and other position but no other population.

In our study, the most frequent DDI occurred between ammonium chloride (in diphenhydramine compound) and pseudoephedrine, although these reactions are mild. After it, the most important reaction (related to moderate- severe reactions), was seen between antihypertensive drugs with others. Especially, angiotensin receptor blockers or angiotensin converting enzyme inhibitors and diuretics had more reaction versus other antihypertensive drug. We noticed to multiple prescriptions of these types of drugs, because of high frequency of uncontrolled hypertension in this area. Another frequent DDI was seen between beta blockers (especially, propranolol) and acetaminophen. Notwithstanding, NSAIDs, corticosteroids, azithromycin had the most quantity of prescribed drug in all subjects.

In nabovati study, beta blockers and diuretics and in Farzaneh study, Aspirin had most frequent drug interaction in prescriptions<sup>16,17</sup>, previous study in east of Iran showed that, dexamethasone & ranitidine and then corticosteroids & NSAIDs had majority of DDI<sup>18</sup>. Interaction between beta adrenergic blockers and glimepiride (oral hypoglycemic agents) was the most commonly observed interaction in Jaskumar Nk study<sup>19</sup>.

We conclude that, general and even OTC drug (not specialized drugs), had great DDI and physicians must be notice to these general drugs in prescriptions. in majority of studies, only limited drugs included in DDI that necessities awareness of prescribers<sup>20</sup>.

Similar to other study, there was no significant relation between gender of patients and rate of DDI In our study (Esmail Farzaneh, & Khouri et al study)<sup>17,21</sup>.

Our result showed that, no correlation between gender or specialty of physician and frequency of DDI was documented ( $p=0.08$ ), but in other study, male gender was factor for drug interaction in mousavi study,

specialist had more DDI in their prescription than general practitioners, in Ardabil study, 84.8% of drug interactions have been occurred by male and 15.2% by female physicians and there was significant relation between sex of physician and number of interferences in mousavi study, Medical specialist's prescriptions in comparison with general practitioners had significantly more moderate severity interactions. Similar results reported in podasani study<sup>22,23</sup>.

In our study, DDI severity, classified as: 2.3% in severe category and 17.7% in moderate and 80% in mild category. There is some difference between characteristic of DDI in our study and others. In previous study in Kurdistan 15.6%, 42.6% and 41.8% of interaction were severe, moderate and mild respectively, in gorgan study result was: 35.5%, 63.1% and 1.4% respectively<sup>24,25</sup>.

We suppose that some factor, including: type of disease may be influence on the type of drugs resulting in variation of DDI in different centers. We supposed that, duration time of disease management, variation of patients' age, and other comorbidity, especially: renal or hepatic disease and hemodynamic instability may be influence on the interaction between drugs in different studies<sup>38-40</sup>.

Previous studies showed that 1.3 -2.1 mean drug in each paper is suitable for reducing risk of DDI<sup>26-28</sup>. In this study, mean number of drugs in each prescription was 3.8 that was near to other studies (by Soleymani and mousavi)<sup>29,30</sup>. Previous study by nabovati and et al (in Iran), showed that mean number of drugs in each prescription is high<sup>16</sup>. In our study, quantity of five drugs type in each prescription had more rate of DDI than fewer quantities. Analyses show that quantity of drugs in each prescription, correlate with rate of DDI ( $p=.003$ ). We conclude, the multidrug prescription may be a risk factor for DDI, this result documented in previous studies<sup>31-33</sup>. Patients with 2 or more disorders may be use different type of drugs result in more DDI<sup>34</sup>.

Increased knowledge of practitioners, for Better diagnosis of disease may decrease the quantity of drugs in prescription, although some chronic and end stage patients need more and more drugs<sup>35,36</sup>.

Physician in subspecialty, due to visit of more complicated patients and especially, physician in ICU must be notice to more risk of prescription with DDI.

Finally, for reduction rate of DDI below approaches is recommended:

1. Documentation of all drugs, including over-the- counters.
2. Reduce the number of prescriptions drugs in each prescription.
3. be especially vigilant in high risk situations: elderly, patients in an intensive care unit, or with co-morbid illnesses such as renal or hepatic failure.
4. more knowledge about the pharmacology of the drugs, route of excretion, the type of liver or renal metabolism if it exists, the half- life of the drug, and its bioavailability.

If there is an unexpected deterioration or change in the patient condition, consider drug- drug interactions, which usually present in a subtle manner. We (and others), conclude that use of software for detects of DDI or attendance a pharmacist in treatment team of patients could be helpful, for decrease the rate of DDI<sup>37</sup>.

Drug interactions are tricky and are easy to miss if not quite. More and more research for detecting type and frequency of DDI in much prescription may turn on a light for solving the problem.

## AGRADECIMIENTOS

**Acknowledgment:** The authors would like to thank the Research Vice-Chancellery of Lorestan University of Medical Science for their supports.

## REFERENCES

1. Thomsen LA, Winterstein AG, Søndergaard B, Haugbølle LS, Melander A. Systematic review of the incidence and characteristics of preventable adverse drug events in ambulatory care. *Ann pharmacother*. 2007; 41(9):1411-26.

2. Lazarou J, Pomeranz BH, Corey PN. Incidence of adverse drug reactions in hospitalized patients: A meta#analysis of prospective studies. JAMA 1998; 279:1200#5.
3. Juurlink DN, Mamdani M, Kopp A, Laupacis A, Redelmeier DA. Drug-drug interactions among elderly patients hospitalized for drug toxicity. JAMA. 2003 Apr 2;289(13):1652-8
4. David S. Tatro, Drug interactions facts, Facts and Comparisons Division, J.B. Lippincott Company. 8th ed., 2006; 47-72.
5. Ben D Snyder, Thomas M Polasek, Matthew P Doogue. Drug interactions: Principles and Practice. Aust Prescriber .2012;35:85–8
6. Dennis L Kasper, Anthony S Fauci, Stephen L Hauser, et al. Harrisons principles of internal medicine. McGraw-Hill. 2015. Chapter 5.p:41.
7. Tatro D S. Drug Interaction Facts. The Authority on Drug Interactions .St Louis, Walters Kluwer Health. 2010.
8. Mohammadi M, Esfandnia A, Rezaei S, Ziapour A. Performance evaluation of hospitals under supervision of kermanshah medical sciences using pabonlasoty diagram of a five-year period (2008-2012). Life Science Journal. 2014;11 (1). ):77-81.
10. A.M. Alizadeh, A. Rostamian, Kh. Saeedpour, M. Hemmati , Z. Khorasani, M.A. Mohagheghi, M. Khatami Moghaddam, M. Mousavi. Drug interactions frequency in the bedridden patients in three hospitals of Tehran city .Modern Care, Scientific Quarterly of Birjand Nursing and Midwifery Faculty. 2011;7,(3).pp:22-27.(in Persian).
11. Chatsisvili A, Sapounidis I, Pavlidou G, Zoumpouridou E, Karakousis VA, Spanakis M, Teperikidis L, Niopas I. Potential drug-drug interactions in prescriptions dispensed in community pharmacies in Greece. Pharm World Sci. 2010; 32:187–193. [PubMed]
12. Heikkila T, Lekander T, Raunio H. Use of an online surveillance system for screening drug interactions in prescriptions in community pharmacies. Eur J Clin Pharmacol. 2006; 62:661–665. [PubMed].
13. Sapkota S, Pudasaini N, Singh C, Sagar GC. Drug prescribing pattern and prescription error in elderly: A retrospective study of inpatient record. Asian J Pharm Clin Res 2011;4: 129-32.
14. Mohammad Haji Aghajania, Mohammad Sistanizad, Mohammad Abbasinazari, Mahdieh Abiar Ghamsari, Ladan Ayazkhoo, Olia Safi, Katayoon Kazemi and Mehran Kouchek.Potential Drug-drug Interactions in Post-CCU of a Teaching Hospital . Iranian Journal of Pharmaceutical Research (2013), 12 (1): 243-248.
15. Mohsen Asadbegi , Bahram Delfan , Ahmad Adineh , Raziieh Sohrabnejhad , Reza Sepahvand , Sehar Changizian , Mahmoud Bahmani and Mahmoud Rafeian-Kopae .Drug-drug interactions in prescriptions of Lorestan province, Western of Iran .Journal of Chemical and Pharmaceutical Research, 2015, 7(1):698-700
16. Ehsan Nabovati,Hasan Vakili-Arki,Zhila Taherzadeh, Mohammad Reza Hasibian, Ameen Abu-Hanna,and Saeid Eslami.Drug-drug interactions in inpatient and outpatient settings in Iran: a systematic review of the literature.Daru. 2014; 22(1): 52.
17. Esmaeil Farzaneh , Perham Mohammadi, Fariba Kahn mouei-aghdam, Firouz Amani, Soheila Saeedi, Fatemeh Mohharami, Ghafour Mahmoodi . Study of Drug Interactions and Associated Factors in Prescriptions of General Practitioners in Ardabil City, 2013-2014 .International Journal of Scientific Study .March 2015 .2 ; 12.pp:136-139
18. Mandana Moradi Dirin, Sarah Mousavi, Amir Reza Afshari, Kaveh Tabrizian, Mohammad Hossein Ashrafi Potential drug#drug interactions in prescriptions dispensed in community and hospital pharmacies in East of Iran. Journal of Research in Pharmacy Practice .Jul-Sep 2014;3; 3.pp:104-107
- 19.Patel Jaskumar Nileshekumar, Ansu Anie Sunny , Jaidev Kumar, Mit Kaushikbhai Suthar , Umesh A study on Prevalence of Potential Drug Drug Interactions in Community Pharmacies of Mysore city . International Journal of Research Studies in Biosciences (IJRSB) Volume 4, Issue 5, May 2016, PP 45-49
20. Holm J, Eiermann B, Eliasson E, Mannheimer B. A limited number of prescribed drugs account for the great majority of drug-drug interactions. Eur J Clin Pharmacol. 2014 Nov; 70(11):1375-83.
21. Khouri V, Semnani SH, Roshandel GH. Frequency of drug interactions and associated factors in insured prescriptions in Gorgan City. Med J Tabriz Univ Med Sci 2005;4:29-32

22. Mousavi S, Norouzi M, Ashouri A, Javadi MR, Gholami K, Hadjibabaie M. Study of Potential Drug-Drug Interactions in Prescriptions of University-Based Pharmacies. *J Pharm Care* 2014; 2(2): 60-65.
23. Pudasaini N, Singh C, Sagar GC, Sapkota S. Drug prescribing pattern and prescription errors in elderly: A retrospective study of inpatient record. *Asian J Pharm Clin Res* 2011; 4:83-6..
24. Rashidi K, Senobar Tahae S. Assessment of drug interactions in medical insurance prescriptions in Kurdistan province in 2000. *Scientific Journal of Kurdistan University of Medical Sciences* 2005;10(3):78-84.
25. Abbasi A, Azadfar S, Roshandel GR, Hoseini SM, Golsha R, Khodabakhshi B, Amjadi Kh, Hajimoradloo N. [Prevalence of medicinal drug interactions in medicinal prescriptions in the Golestan province, Iran (2012)]. *J Gorgan Uni Med Sci*. Winter 2017; 18(4):105 -108. [Article in Persian]
26. Mohanty B, Aswini M, Hasamnis A, Patil S, Murty K, Jena S. Prescription pattern in the department of medicine of a tertiary care hospital in South India. *J Clin Diagnos Res* 2010;3:2047-51.
27. Williams D, Bennett K, Feely J. The application of prescribing indicators to a primary care prescription database in Ireland. *Eur J Clin Pharmacol* 2005;61(2):127-33.
28. Zou J, Li L, Zhang C, Yan Y, Gao F, Zhang H. Analysis of outpatient prescription indicators and trends in Chinese Jingzhou Area between September 1 and 10, 2006-2009. *African J Pharm Pharmacol* 2011;5(2): 270-5.
29. Soleymani F, Valadkhani M, Dinarvand R. Challenges and achievements of promoting rational use of drugs in Iran. *Iranian J Publ Health*. 2009; 38(Suppl. 1):166-8.
30. Mousavi S, Norouzi M, Ashouri A, Javadi MR, Gholami K, Hadjibabaie M. Study of Potential Drug-Drug Interactions in Prescriptions of University-Based Pharmacies. *J Pharm Care* 2014; 2(2): 60-65.
31. Qian Y, Ye X, Du W, Ren J, Sun Y, Wang H, et al. A computerized system for detecting signals due to drug#drug interactions in spontaneous reporting systems. *Br J Clin Pharmacol* 2010; 69:67#73.
32. Rafii H, Arab M, Ranjbar H, Sepehri GH, Arab N, Amiri M. The prevalence of potential drug interactions in intensive care unit. *Iran J Crit Care Nurs* 2010; 4:191-6.
33. Astrand E, Astrand B, Antonov K, Petersson G. Potential drug interactions during a three-decade study period: a cross-sectional study of a prescription register. *Eur J Clin Pharmacol*. 2007 Sep;63(9):851-9.
34. Rafeian M. Drug interactions in internal and surgical wards of Kashani Hospital, Shahrekord, 1997. *Tehran Univ Med J* 2001; 59:86-91.
35. Askari M, Eslami S, Louws M, Dongelmans D, Wierenga P, Kuiper R, Abu-Hanna A. Relevance of drug-drug interaction in the ICU - perceptions of intensivists and pharmacists. *Stud Health Technol Inform*. 2012; 180:716–720. [PubMed]
36. Ko Y, Malone DC, Skrepnek GH, Armstrong EP, Murphy JE, Abarca J, Rehfeld RA, Reel SJ, Woosley RL. Prescribers' knowledge of and sources of information for potential drug-drug interactions: a postal survey of US prescribers. *Drug Saf*. 2008; 31:525–536. [PubMed]
37. Ansari J, Drug interaction and pharmacist. *J Young Pharm*. 2010 Jul-Sep; 2(3): 326–331
38. Mirzamasoumzadeh, B., & Mollasadeghi, V. (2013). Effects of osmotic stress on chlorophyll and proline different wheat cultivars, *UCT Journal of Research in Science, Engineering and Technology*, 1(1): 12-13.
39. Yıldırım A, Kapukaya A, Mertsoy Y, Yiğit Ş, Çaçan MA. Treatment of nonseptic bursitis with endoscopic surgery. *J Clin Exp Invest*. 2015;6(3):220-3. <https://doi.org/10.5799/ahinjs.01.2015.03.0522>
40. Al-Sharif, F. M. (2017). Correlation between serum alanine aminotransferase activity and immunologic response and body mass index in obese patients with chronic hepatitis B virus infection. *European Journal of General Medicine*, 14(2),34-37. <https://doi.org/10.29333/ejgm/81879>