



InVet
ISSN: 1514-6634
ISSN: 1668-3498
invet@fvvet.uba.ar
Universidad de Buenos Aires
Argentina

Hantavirus circulation in *Oligoryzomys flavescens* rodents of Buenos Aires City, Argentina

Hercolini, C.; Bruno, A.; Aristegui, E.; De Salvo, M.N.; Vidal, J.; Bellomo, C.; Martinez, V.P.; Brambati, D.F.
Hantavirus circulation in *Oligoryzomys flavescens* rodents of Buenos Aires City, Argentina
InVet, vol. 20, no. 1, 2018
Universidad de Buenos Aires, Argentina

Available in: <https://www.redalyc.org/articulo.oa?id=179158985003>

InVet (Investigación Veterinaria) por Facultad de Ciencias Veterinarias. Universidad de Buenos Aires se distribuye bajo una Licencia Creative Commons Atribución – No Comercial – Sin Obra Derivada 4.0 Internacional.



This work is licensed under Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International.

Hantavirus circulation in *Oligoryzomys flavescens* rodents of Buenos Aires City, Argentina

Hantavirus en roedores de La Ciudad de Buenos Aires circulación de hantavirus en *Oligoryzomys Flavescens* en Ciudad de Buenos Aires, Argentina

C. Herculini
Instituto de Zoonosis Luis Pasteur, Argentina
c.hercolini@gmail.com

Redalyc: <https://www.redalyc.org/articulo.oa?id=179158985003>

A. Bruno
Instituto de Zoonosis Luis Pasteur, Argentina

E. Aristegui
Instituto de Zoonosis Luis Pasteur, Argentina

M.N. De Salvo
Instituto de Zoonosis Luis Pasteur, Argentina

J. Vidal
Instituto de Zoonosis Luis Pasteur, Argentina

C. Bellomo
Instituto Nacional de Enfermedades Infecciosas ANLIS 'Dr. Carlos G. Malbrán', Argentina

V.P. Martinez
Instituto Nacional de Enfermedades Infecciosas ANLIS 'Dr. Carlos G. Malbrán', Argentina

D.F. Brambati
Instituto de Zoonosis Luis Pasteur, Argentina

Received: 28 February 2018
Accepted: 19 December 2018

RESUMEN:

El síndrome pulmonar por hantavirus (SPH) es una enfermedad zoonótica. La principal causa en Argentina es el virus Andes. *Oligoryzomys flavescens* (Sigmodontinae) uno de los reservorios de SPH. El objetivo de esta investigación fue estimar la seroprevalencia de hantavirus en *O. flavescens* en la Ciudad de Buenos Aires. Se colocaron trampas de captura viva en tres parques y una reserva ecológica de la Ciudad para detectar individuos con diagnóstico positivo para virus Andes. Se capturaron un total de 286 roedores, *O. flavescens* fue la especie más capturada (49,65%). Se encontró serología positiva para virus Andes (genotipo Lechiguana) en ejemplares capturados en uno de los sitios en estudio (seroprevalencia = 6,62%) si bien encontramos al hospedador en otros dos parques de la Ciudad. El presente estudio confirma la presencia de roedores infectados con el virus Andes en la Ciudad de Buenos Aires, lo que implica un riesgo de transmisión en un ambiente urbano.

PALABRAS CLAVE: Hantavirus, *Oligoryzomys flavescens*, Ciudad de Buenos Aires.

ABSTRACT:

Hantavirus pulmonary syndrome (HPS) is a zoonotic disease. The main cause in Argentina is Andes virus. *Oligoryzomys flavescens* (Sigmodontinae) was identify as one of the HPS reservoir. The objective of this research was to estimate the seroprevalence of hantavirus in *O. flavescens* in Buenos Aires City. We set rodents live trapping in three parklands and one ecological reserve in Buenos Aires City in order to screen hantavirus Andes infected rodents. A total of 286 rodents were captured, *O. flavescens* was the most frequently captured species (49.65%). Positive serology for Andes virus (Lechiguana genotype) was found in *O. flavescens* in

one site studied (seroprevalence = 6.62%) and we found the host in other two parks within the City. The present study confirms the presence of rodents infected with Andes virus in Buenos Aires City, which implies transmission risk in an urban environment.

KEYWORDS: Hantavirus, *Oligoryzomys flavescens*, Buenos Aires City.

INTRODUCTION

Hantavirus pulmonary syndrome (HPS) is a zoonotic disease caused by viruses belonging to the Hantaviridae family, and mainly harbored by some rodent species. These viruses are transmitted to humans by inhalation of aerosols generated by infected rodents²⁷, although there is also evidence for person to person in the Argentine southern region and Buenos Aires Province^{15, 21}. The main cause of HPS in Argentina and neighboring countries is Andes virus (ANDV). Seven different ANDV genotypes have been characterized from HPS cases in Argentina^{22, 24}.

There are 4 geographically and ecologically distinct HPS endemic areas¹⁶. The Central region, the second in number of cases, comprises Buenos Aires City^{16,14} where one HPS case with no history of travel outside the City was recorded and Lechiguanas genotype was characterized¹¹.

Several species of Sigmodontinae rodents were identified as HPS reservoirs in the central region: *Akodon azarae*, *Necromys obscurus* and *Oligoryzomys flavescens*¹³. Particularly in Buenos Aires City few of these species were reported: *A. azarae*, and *O. flavescens*^{2,4, 17,25}. Recent studies have communicated the presence of *O. flavescens* in Costanera Sur Ecological Reserve^{2,4} and in Presidente Roca Park⁴.

The distribution of *O. flavescens*³¹ showed a wide extension from the north to the center of Argentina⁷, southeast Brazil³² and Uruguay¹². This species is a good colonizer of disturbed ecosystems²⁶, inhabits wild environments such as savannas, marshes and not very dense forests, grass and scrubland, often near water or damp zones⁹, and although it is associated to wild microhabitats it has been captured in peridomestic urban and periurban areas⁵.

Knowing the presence of *O. flavescens* in Buenos Aires City it becomes necessary to study if there is hantavirus circulation between them. Our aim was to estimate the seroprevalence of hantavirus in *Oligoryzomys flavescens* in Buenos Aires City.

Materials and Methods

Rodents were captured in four green open sites of the City during the period 2011-2014: De los Niños Park (NP: 32 hectares, 34°31'43.40"S, 58°27'33.99"O), Ribera Sur Park (RSP: 50 hectares, 34°41'57.73"S, 58°28'04.93"O), Presidente Roca Park (PRP: 154 hectares, 34°40'29.76"S, 58°26'29.79"O) and Costanera Sur Ecological Reserve (CSER: 353 hectares, 34°36'29.18"S, 58°21'03.32"O) (Figure 1). According to the characterization of environmental landscape units by Cavia *et al.* (11), the CSER is a natural reserve and NP and PRP are parklands. We considered RSP as parkland because it shares environmental characteristics with the other two. The CSER is occupied by riparian habitats similar to those developed along the Paraná and de la Plata rivers, like woodlands, riparian thickets, fresh water marshes and flooded grasslands. Parklands are sites of recreation, where areas of spontaneous vegetation and woodlots with planted species are included in a matrix of grass or ornamental lawn³.




FIGURE 1
Studied sites in Buenos Aires City.

Rodents were captured with Sherman live traps in linear transects actively set for four consecutive nights for each sampling (Table I).

TABLE I
Number of traps night per season and year for each studied site

Site	Season - year	Traps night
CSER	spring 2011; summer 2012; autumn 2012; winter 2012; spring 2012; summer 2013; autumn 2013; winter 2013; spring 2013; summer 2014; autumn 2014; winter 2014; spring 2014	11 624
NP	winter 2014	452
RSP	summer 2015	512
PRP	winter 2014	424

The individuals captured were identified according bibliography⁹ and reference material deposited at the Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”. All *O. flavescens* specimens captured were anesthetized with isoflurane and euthanized by cervical dislocation. Body measurements, sex and reproductive status were registered. Blood and lung samples were collected for hantavirus infection analysis, following Mills *et al.*¹⁹. Animals were treated in accordance with the guidelines of the Sociedad Argentina para el Estudio de los Mamíferos⁸.

The relative abundance of each species per study area was estimated with the Trap Success Index (TS)¹⁸. $TS = (\text{number of individuals} / \text{number of traps} * \text{nights}) * 100$. The correlation between the number

of *O. flavescens* captured and number of positive individuals was calculated with Spearman Correlation Coefficient.

In order to screen ANDV antibody-positive individuals, we utilized an enzyme-linked immunosorbent assay based on the test described for human antibodies detection²². Briefly, to determine IgG antibodies against ANDV, rodent blood samples diluted 1:100 were added to 96 wells plate coated with AND recombinant nucleoprotein antigen. To determine ANDV genotype, total RNA extraction was performed on available lung tissues from seropositive rodents (5/9 individuals) using Trizol (Invitrogen) and purified by the RNAid kit (Bio 101). When it was possible (3/5 samples), viral RNA was amplified by RT-PCR and a second round of nested PCR following Ciancaglini *et al.*⁶. Briefly, RT-PCR was performed by One Step RT-PCR kit (QIAGEN) and Taq DNA Polymerase (Invitrogen) following each manufacturer instructions. For the 952 nt S-Segment fragment was used an inner primer to obtain the 530-nt fragments for S-segment (position 22–550, referred to AH1 AND strain, Gene Bank No AF324902) position. For M-segment was amplified the 461nt fragment (position 6 to 467). Amplification products were analyzed on agarose gels and sequenced. Multiple sequence alignment and comparison of nucleotide sequences were conducted using MEGA version 5²⁹

Results

A total of 286 rodents were captured with a total trapping effort of 13012 trap-nights. *O. flavescens* was the most frequently captured species (49.65%) followed by *Mus musculus* (36.71%), *Deltamys kempi* (10.14%) and *Scapteromys aquaticus* (3.50%). Rodents were captured in all sites but NP. In CSER we captured four species of which the trap success of *O. flavescens* and *M. musculus* were the highest. On the contrary, in RSP and PRP we only captured *O. flavescens* (Table II). (Error 2: La referencia: II está ligada a un elemento que ya no existe)

TABLE II
Rodents captured and trap success per studied site

Site	Species	Trap success (%)	N° of individuals captured
CSER	<i>Oligoryzomys flavescens</i>	1.17	136
	<i>Deltamys kempi</i>	0.25	29
	<i>Mus musculus</i>	0.90	105
	<i>Scapteromys aquaticus</i>	0.09	10
NP	without capture	0	0
RSP	<i>Oligoryzomys flavescens</i>	0.39	2
PRP	<i>Oligoryzomys flavescens</i>	0.94	4

Positive serology for ANDV was found in *O. flavescens* captured in CSER where the seroprevalence was 6.62% (95% c.i.: 2.07% - 11.16%). The nine positive rodents corresponded to male individuals, seven of them were sexually active (with testicles in the scrotal position). These positive individuals were caught in different seasons: 3 in winter, 3 in spring, 2 in summer and 1 in autumn (Table III). They were found in a variety of floristic habitats, six of them were captured on the river side and the rest in different habitats of CSER. The Spearman Correlation Coefficient between number of *O. flavescens* captured and number of positive individuals resulted 0.81 ($p < 0.001$, $N = 13$), showing a positive relationship between both variables throughout the seasons. (Error 3: La referencia: III está ligada a un elemento que ya no existe)

TABLE III

Number of captured, positives and seroprevalence of *Oligoryzomys flavescens* in CSER 2011-2014

Period	# <i>O. flavescens</i>	# <i>O. flavescens</i> positive	seroprevalence <i>O. flavescens</i>
spring 2011	7	0	0.00
summer 2012	0	0	0.00
autumn 2012	4	0	0.00
winter 2012	8	1	0.13
spring 2012	3	0	0.00
summer 2013	0	0	0.00
autumn 2013	2	0	0.00
winter 2013	37	2	0.05
spring 2013	33	2	0.06
summer 2014	5	2	0.40
autumn 2014	20	1	0.05
winter 2014	4	0	0.00
spring 2014	13	1	0.08

O. flavescens did not show any seasonal pattern in CSER and was captured in all seasons but summer 2012 and summer 2013. Although, we observed an increase of trap success since winter 2013 with a peak in spring at the same year. On the other hand, after the peak of *O. flavescens*, the trap success of *M. musculus* decreased and we also observed a slight increase of *D. kempfi* trap success and *S. aquaticus* was found for the first time (Figure 2)




FIGURE 2.

Trap success of *Oligoryzomys flavescens* (●), *Mus musculus* (▲), *Deltamys kempfi* (▲) and *Scapteromys aquaticus* (▲) per period in CSER 2011-2014.

ANDV, Lechiguanas genotype, was identified in 3 IgG positive-rodent, showing a 99.8% nucleotide identity between them in the amplified fragment. The 3 individuals were captured in winter 2012, spring 2013 and summer 2014.

Discussion

The present study confirms the presence of rodents infected with Andes virus in Buenos Aires City. This finding is not unexpected because this City is geographically placed in the Central region HPS endemic area.

Positive rodents were captured only in a natural reserve, CSER, which was the most intensely studied site. Since 2014, an exploratory study has been started in parklands in the City. In two of them, RSP and PRP, we found the host. The presence of this hantavirus reservoir species in these two places should be considered as a potential risk per se.

The seroprevalence found in CSER was 6.62%. In similar studies the seroprevalence found in *O. flavescens* was 7.51% in Pre Delta National Park³⁰ and 13.51% in Exaltación de la Cruz²⁸. All the positive individuals were males, according to other authors that found that males are more likely to be infected^{23, 28, 30}. The presence of *O. flavescens* with positive serology for ANDV in all seasons and in different habitats located in CSER, could implies transmission risk for visitants and workers along all year.

Althought it was not part of the objective of this work, we studied the first 15 *D. kempfi* captured for hantavirus infection analysis. All of them were negative for ANDV (data not shown). These results supports, as mention in Vadell *et al.*³⁰ that *D. kempfi* is an uncommon host species for hantavirus and constitutes a spillover host.

We found rodent community changes along the study period in CSER. These changes can be associated with the dredging works and filling in the lagoons with water since 2013². Interestingly, we found the highest peak of *O. flavescens* in 2013/2014, period in which an outbreak of HPS cases was registered in Buenos Aires Province¹¹. On the other hand, Cavia *et al.*⁴ found in CSER a richness of 6 species, and *O. flavescens* and *D. kempfi* were the dominant species; in our study richness was lower (R = 4) and *O. flavescens* and *M. musculus* were the dominant species, which could indicate that rodent community in CSER is not stable.

O. flavescens is an r strategist species, consequently its captured and reproductive activity are strongly conditioned by the density of the rest of the rodent species²⁶. If we take this into account and that we found a positive correlation between number of *O. flavescens* positive and individuals captured, it becomes necessary to continue evaluating the *O. flavescens* dynamics in the assemblages of rodent species. There are several variables in an infection system¹⁰. We found that an increasing population density of *O. flavescens* resulted in an increasing horizontal virus transmission, according to Mills *et al.*²⁰, Hussein *et al.*¹⁰ and Adler *et al.*¹.

Conclusions

Results presented in this work, can be used to support the adoption of preventive measures and optimize the allocation of resources to avoid disease propagation. Hantavirus's epidemiological monitoring in rodents must be continued and new sites should be studied in the City.

Acknowledgements

We thanks the authorities of Costanera Sur Ecological Reserve, Presidente Roca Park, De los Niños Park, Ribera Sur Park, personal of Instituto de Zoonosis Luis Pasteur: Yamila Bechara, Carolina Zotter, Javier Marengo and Javier Donacimiento for field work assistance and Yanina Berra for cooperation in the field work in Costanera Sur Ecological Reserve.

REFERENCES

1. Adler FR, Pearce-Duvel JMC, Dearing MD. How host population dynamics translate into time-lagged prevalence: An investigation of Sin Nombre virus in deer mice. *Bull Math Biol.* 2008; 70:236–252.
2. Arístegui A, Hercolini C, Brambati D, Vidal J, Bruno A, Sassaroli JC, Beaudoin J. Primer registro de *Scapteromys aquaticus* y *Cryptonanus chacoensis* en la Ciudad Autónoma de Buenos Aires, Argentina. *Nótulas Faunísticas: Segunda Serie.* 2014; 169: 1-5.
3. Cavia, R. *Rodents in the city of Buenos Aires: an approach to their control, PhD Thesis, University of Buenos Aires, Buenos Aires.* (in Spanish). 2006.
4. Cavia R, Cueto G R, Suarez OV. Changes in rodent communities according to the landscape structure in an urban ecosystem. *Landscape and Urban Planning.* 2009 90: 11–19.
5. Calderón G, Pini N, Bolpe J, Levis S, Mills J, Segura E, Guthmann N, Cantoni G, Becker J, Fonollat A, Ripoll C, Bortman M, Benedetti R, Sabattini M, Enria D. Hantavirus Reservoir Hosts Associated with Peridomestic Habitats in Argentina. *Emerg Infect Dis.* 1999; 5 (6): 792-797.
6. Ciancaglini M, Bellomo CM, Torres Cabreros CL, Alonso D, Bassi SC, Iglesias AA, Martínez VP. Hantavirus pulmonary syndrome in Tucumán province associated to an unexpected viral genotype. *Medicina (B Aires).* 2017; 77(2):81-84.
7. Cirignoli S, Teta P, Pardiñas UFJ, D'Elía G. Tribu Oryzomyini. In: Barquez RM, Díaz MM, Ojeda RA (eds.) *Mamíferos de Argentina, Sistemática y distribución.* Sociedad Argentina para el Estudio de Mamíferos, Mendoza; 2006. p. 166-175.
8. Giannoni SM, Mera Sierra R, Brengio S y Jiménez Baigorria L. Guía para el uso de animales en investigaciones de campo y en cautiverio. Comisión de Ética de la Sociedad Argentina para el Estudio de los Mamíferos [Internet]. 2003. Available from: <https://www.sarem.org.ar/legislacion/>
9. Gómez Villafaña IE, Miño M, Cavia R, Hodara K, Courtalón P, Suárez O, Busch M. *Roedores: guía de la provincia de Buenos Aires.* L.O.L.A. Buenos Aires. 2005; p 100.
10. Hussein K, Hornfeld B, Evander M, Magnusson M, Olsson G, Ecke F. Dynamics and Drivers of Hantavirus Prevalence in Rodent Populations. *Vector-Borne and Zoonotic Diseases.* 2014; 14(8): 537-551.
11. Iglesias AA, Bellomo CM, Martínez VP. Síndrome pulmonar por hantavirus en Buenos Aires, 2009-2014. *Medicina (B Aires).* 2016; 76: 1-9.
12. Langguth A. Las especies uruguayas del género *Oryzomys* (Rodentia-Cricetidae). *Com. Zool. Mus. Hist. Nat. Mont. Urug.* 1963; 7 (99): 1-22.
13. Levis S, Morzunov SP, Rowe JE, Enria DA, Pini N, Calderon G, Sabattini M, St Jeor SC. Genetic diversity and epidemiology hantavirus in Argentina. *J Infect Dis.* 1998; 177:529-538.
14. Martínez VP, Colavecchia S, García Alay M, Suzuki B, Trinchero A, Busto S, Rabinovich R, Padula P. Hantavirus pulmonary syndrome in Buenos Aires Province. *Medicina (B Aires).* 2001; 61(2):147-56.3.
15. Martinez VP, Bellomo C, San Juan J, Pinna D, Forlenza R, Elder M, Padula PJ. Person-to-person transmission of Andes virus. *Emerg Infect Dis* Dec. 2005; 1(12):1848-53.
16. Martinez VP, Bellomo CM, Cacace ML, Suarez P, Bogni L, Padula PJ. Hantavirus pulmonary syndrome in Argentina, 1995-2008. *Emerg Infect Dis* Dec. 2010; 16(12):1853-60.
17. Massoia E, Fornes A. Rodents collected in the Federal Capital City (Caviidae, Cricetidae, Muridae). *IDIA.* 1967; 47-53.
18. Mills JN, Ellis BA, McKee KT, Maiztegui JI, Childs JE. *Habitat Associations and Relative Densities of Rodent Populations in Cultivated Areas of Central Argentina.* Other Publications in Zoonotics and Wildlife Disease. Lincoln. 1991; p. 82.
19. Mills JN, Yates TL, Childs JE, Parmenter RR, Ksiazek TG. *Guidelines for Working with Rodents Potentially Infected with Hantavirus.* Other Publications in Zoonotics and Wildlife Disease. Lincoln; 1995. p. 84.

20. Mills JN, Ksiazek TG, Peters CJ, Childs JE. Long-term studies of hantavirus reservoir populations in the southwestern United States: A synthesis. *Emerg Infect Dis.* 1999; 5:135.
21. Padula PJ, Edelstein A, Miguel SDL, López NM, Rossi CM, Rabinovich RD. Hantavirus Pulmonary Syndrome Outbreak in Argentina: Molecular Evidence for Person-to-Person Transmission of Andes Virus. *Virology.* 1998; 241: 323–330.
22. Padula PJ, Colavecchia SB, Martínez VP, Gonzalez Della Valle MO, Edelstein A, Miguel SD, Russi J, Riquelme JM, Colucci N, Almirón M, Rabinovich RD. Genetic diversity, distribution, and serological features of hantavirus infection in five countries in South America. *J Clin Microbiol.* 2000; 38(8):3029-35.
23. Padula P, Figueroa R, Navarrete M, Pizarro E, Cadiz R, Bellomo C, Jofre C, Zaror L, Rodriguez E, Murúa R. Infection in Wild Sigmodontine Rodents Transmission Study of Andes Hantavirus. *J Virol.* 2004; 78(21):11972-11979.
24. Padula P, Martinez VP, Bellomo C, Maidana S, San Juan J, Tagliaferri P, Bargardi S, Vazquez C, Colucci N, Estévez J, Almirón M. Pathogenic hantaviruses, northeastern Argentina and eastern Paraguay. *Emerg Infect Dis.* 2007; 13(8):1211-4.
25. Pearson OP. Age structure and reproductive dynamic in a population of meadow mice, *Akodon azarae*. *Physis.* 1967; XXVII (74), 53-58
26. Sánchez López MI. *Factores que limitan la abundancia de los roedores múridos en el delta del Paraná*. PhD Thesis, Universidad de Buenos Aires, Buenos Aires. 1998 p. 150.
27. Schmaljohn CS. Molecular biology of hantaviruses. In Elliott RM (ed). *The Bunyaviridae*. Plenum Press. New York, N.Y. 1996; 63–90.
28. Suárez OV, Cueto GR, Cavia R, Gómez Villafañe IE, Bilenca DN, Edelstein A, Martínez P, Miguel S, Bellomo C, Hodara K, Padula PJ, Busch M. Prevalence of Infection with Hantavirus in Rodent Populations of Central Argentina. *Mem Inst Oswaldo Cruz, Rio de Janeiro.* 2003; 98(6): 727-732.
29. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, and Kumar S. MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Mol Biol Evol.* 2011.
30. Vadell MV, Bellomo C, San Martin A, Padula P, Gomez Villafane I. Hantavirus ecology in rodent populations in three protected areas of Argentina. *Trop Med Int Health.* 2011; 16 (10): 1342–1352.
31. Waterhouse GR. Characters of new species of the genus *Mus*, from the collection of Mr. Darwin. *Proceedings of the Zoological Society of London.* 1837; 5(50-51):15-21.
32. Weksler M, Bonvicino CR. Taxonomy of pigmy rice rats genus *Oligoryzomys* Bangs, 1900 (Rodentia, Sigmodontinae) of the Brazilian cerrado, with the description of two new species. *Arquivos do Museu Nacional, Rio de Janeiro.* 2005; 63:113-130.