



Revista Facultad Nacional de Agronomía Medellín

ISSN: 0304-2847

ISSN: 2248-7026

Facultad de Ciencias Agrarias - Universidad Nacional de Colombia

Sousa, Victor Hugo de Carvalho; Meneses, Anderson Tenório de; Rodrigues, Eduardo Vieira; Gonçalves, Anderson Carlos de Melo; Silva, Toshik Iarley da; Rodrigues, Rummenigge de Macêdo; Santos, Petrônio Donato dos; Souza, Leossávio César de
Seed treatment with silicon on initial growth of soybean (*Glycine max*) cultivars

Revista Facultad Nacional de Agronomía Medellín,
vol. 72, no. 2, 2019, May-August, pp. 8809-8817
Facultad de Ciencias Agrarias - Universidad Nacional de Colombia

DOI: <https://doi.org/10.15446/rfnam.v72n2.73226>

Available in: <https://www.redalyc.org/articulo.oa?id=179960155006>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UAEH
redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Seed treatment with silicon on initial growth of soybean (*Glycine max*) cultivars

Tratamiento de semillas con silicio sobre el crecimiento inicial de cultivares de soja (*Glycine max*)

doi: 10.15446/rfnam.v72n2.73226

Victor Hugo de Carvalho Sousa¹, Anderson Tenório de Meneses¹, Eduardo Vieira Rodrigues¹, Anderson Carlos de Melo Gonçalves², Toshik Iarley da Silva^{3*}, Rummenigge de Macêdo Rodrigues⁴, Petrônio Donato dos Santos⁵ and Leossávio César de Souza¹

ABSTRACT

Keywords:

Beneficial nutrient
Mineral fertilization
Seed treatment
Silicon

Soybean (*Glycine max*) is a crop of high economic power in the world, being used to produce vegetable oil, as well as a source of food for animals and humans. Recent research indicates that nutrient application in the seed stage contributes to an early and productive development of crops. Since silicon (Si), as a nutrient for plants, acts in the cellular structure and the formation and performance of plant architecture, the aim of this study was to evaluate the influence of silicon application via seeds on the initial growth of three soybean cultivars. The experimental design was a randomized block. The treatments were replicated three times and distributed in a 3x5 factorial scheme, composed by three soybean cultivars (C1 - FTR 1186 IPRO, C2 - FTR 1192 IPRO and C3 - FTR 3190 IPRO) and five silicon doses (0, 30, 60, 90 and 120 g per 100 kg of seeds). The data were submitted to ANOVA by F-test and polynomial regression analysis for the silicon doses; the cultivars response was evaluated through the Tukey mean test. The treatment of soybean seeds with silicon did not positively influence the root dry mass of the FTR 1192 IPRO, nor the root length and root diameter of the FTR 1186 IPRO. The application of silicon did not influence the FTR 3190 IPRO.

RESUMEN

Palabras clave:

Elemento benéfico
Fertilización mineral
Tratamiento de semillas
Silicio

La soja (*Glycine max*) es un cultivo de gran poder económico en el mundo, siendo destinada a la producción de aceite vegetal, así como para la alimentación animal y humana. Investigaciones recientes indican que la aplicación de nutrientes en la fase de semilla contribuye al desarrollo inicial y productivo de los cultivos. Dado que el silicio (Si), como nutriente para las plantas, actúa en la estructura celular y en la formación y el desempeño de la arquitectura de la planta, el objetivo de este estudio fue evaluar la influencia de la aplicación de silicio a través de las semillas en el crecimiento inicial de tres cultivares de soja. El diseño experimental fue de bloques aleatorios. Los tratamientos fueron replicados tres veces y se distribuyeron en un esquema factorial 3x5, compuesto por 3 cultivares (FTR 1186 IPRO, FTR 1192 IPRO, FTR 3190 IPRO) y cinco dosis de silicio (0, 30, 60, 90 y 120 g por 100 kg de semillas). Los datos fueron sometidos al análisis de varianza por la prueba F y regresión polinomial para las dosificaciones de silicio, la respuesta de los cultivares fue evaluada por medio de la prueba de medias de Tukey. El tratamiento de las semillas de soja con silicio no influyó positivamente en la masa seca de la raíz del cultivar FTR 1192 IPRO, ni en la longitud y el diámetro de la raíz del cultivar FTR 1186 IPRO. El cultivar FTR 3190 IPRO no fue influenciada por la aplicación de silicio.

¹ Universidade Federal de Paraíba. Rodovia PB-079 – Km 12, CEP 58397-000. Areia, Paraíba, Brazil.

² Universidade Federal de Roraima. Avenida Cap. Ene Garcês, 2413. Boa Vista, Roraima, Brazil.

³ Universidade Federal de Viçosa. Avenida Peter Henry Rolfs, s/n, CEP 36570-900. Viçosa, Minas Gerais, Brazil.

⁴ Universidade Federal de Campina Grande. Rua Manoel Sabiá, 237, CEP 58429-900 Várzea, Sumé, Paraíba, Brazil.

⁵ Universidade Federal do Piauí. Rua Almir Benvindo, CEP 64860-000. Uruçuí, Piauí, Brazil.

* Corresponding author: <iarley.toshik@gmail.com>

The soybean [*Glycine max* (L.) Merrill] is the most important crop cultivated in Brazil. During the period 2015 – 2016, the country's total production was 95.63 million tons, produced on 33.17 million hectares (Embrapa, 2016). Nowadays, the soybean crop is largely cultivated and consumed worldwide, being an important agricultural product destined for animal feed and human food, thus highly demanded in the agrobusiness world (Conab, 2016).

High-quality seeds are necessary to obtain an optimal seedling population. High physiological quality soybean seeds will grow into plants with greater morphological and biochemical performance, thereby increasing the chance of better development and yield (Kolchinski *et al.*, 2005). Brazilian agricultural researchers have developed new soybean cultivars with expressive yield potential, substantiating the crop importance in the international market and export trade.

In the agricultural production system, it is a challenge to uniformly distribute, in the soil, the nutrients that are required in small amounts by the plants. Seed treatments have enabled plants to have higher performance and full genetic potential achievement through the application of pesticides, biological products, plants stimulants, and inoculants, as well as the use of micronutrients, such as silicon (Daronco, 2013). In this sense, seed treatment might be an alternative to maximize agricultural yield (Oliveira *et al.*, 2013), making it necessary the use of techniques directed to improve legume plants mineral nutrition.

The nutritional essentiality of silicon is controversial. Some researchers claim it is essential to plants, while others refer to it as a beneficial nutrient. Some studies concerning silicon have revealed increasing in the physical and structural resistance against biotic and abiotic factors in several species caused by an increase in the cell wall, middle lamella thickness, and rigidity, because of the application of silicon with phenolic compounds and pectins (Datnoff *et al.*, 2007; Currie and Perry, 2007). Moreover, silicon might reduce aluminum toxicity, as well as other micronutrients toxicity, such as manganese, iron, and sodium (Castro and Crusciol, 2013).

The potassium silicate (K_2SiO_3) fertilization brings several benefits to the soybean crop. It makes leaves more upright,

decreasing the plants self-shading. It also enhances the tolerance to lodging and the plants' resistance to several stresses through the accumulation of silicon in their tissues. Also, potassium influences stomatal activity and improve gaseous exchanges in plants while silicon can protect them from pathogens and phytophagous insects, boosting the plants' development and yield (Wise *et al.*, 2007; Zelin *et al.*, 2011).

Machado and Queiroz (2018) reported that the treatment of soybean seeds with silicon increases the vigor of the seeds under laboratory conditions, but the agronomic characteristics in the field were not influenced. Oliveira *et al.* (2015) also pointed out that the application of silicon does not impair the physiological quality of soybean seeds. According to these authors, the agronomic characteristics of the cultivars are not influenced by the doses of silicon, as well as the number of vegetables per plant. Foliar application of potassium silicate in maize, enhanced the plants' photosynthetic efficiency (Sousa *et al.*, 2010). Ferreira (2008) verified positive effects on the photosynthesis, transpiration rate, stomatal conductance and internal CO_2 concentration of cotton plants treated with sodium metasilicate solution. In this sense, the aim of this study was to evaluate the influence of silicon application via seeds on the initial growth of three soybean cultivars (*Glycine max* L.).

MATERIALS AND METHODS

The study was carried out from October to November 2016, at the greenhouse of the Center of Agricultural Sciences, in the Universidade Federal de Paraíba, located in the city of Areia, state of Paraíba, Brazil. According to the climate classification of Köppen-Geiger, the As' climate (hot and humid) is observed in the region, which presents a rainy season concentrated between March and July and an average annual rainfall of 1400 mm (Peel *et al.*, 2007).

The experimental design was a randomized block, and the treatments were replicated three times and distributed in a 3x5 factorial scheme, composed by three soybean cultivars (C1 - FTR 1186 IPRO, C2 - FTR 1192 IPRO and C3 - FTR 3190 IPRO) and five silicon doses (0, 30, 60, 90 and 120 g per 100 kg of seeds). The silicon doses were applied via seeds according to the proposed by Oliveira *et al.* (2014). The silicon doses were prepared with liquid potassium silicate (K_2SiO_3), which presented

the following composition: 130 g L⁻¹ of SiO₂ and 130 g L⁻¹ of K₂O. The potassium silicate was directly applied in 100 g seed in the following proportion: 0, 230, 460, 690, and 920 mL K₂SiO₃ 100 kg seeds⁻¹, equivalent to the silicon doses determined for the experiment. The seeds were immersed in the solution for 12 h.

The soil used was classified as Haplic Planosol (Embrapa, 2014), collected in the 0-30 cm depth, from

the lands of the Center of Agricultural Sciences, in the Universidade Federal de Paraíba. The physicochemical characteristics of the soil are shown in Table 1. The experimental units were 5-liter pots, where the soybeans were cultivated. The soybean seeds were treated with the fungicide thiophanate-methyl (5 g a.i. per 100 kg of seeds) and the insecticide fipronil (50 g a.i. per 100 kg of seeds). The fertilization was done according to the recommended to the soybean crop.

Table 1. Physical and chemical soil analysis report of the Haplic Planosol used in the study.

Depth (cm)	pH H ₂ O	P (mg dm ⁻³)	K ⁺	Ca ²⁺	Mg ²⁺	Na ⁺	Al ³⁺	H+Al	BS (%)	TOC (g kg ⁻¹)
0-10	6.0	0.83	0.02	2.15	0.65	0.02	0.05	3.22	46.9	6.2
10-30	5.8	1.59	0.28	3.35	0.98	0.03	0.05	4.87	48.8	7.0

TOC: Total Organic Carbon; BS: % Base Saturation

In each experimental unit, two seeds were sown at a depth of 1 cm. Following germination, plants were thinned to one plant per pot. The plants were irrigated whenever needed in order to maintain the soil at field capacity. The study was conducted until the V5 phenological stage of soy development, according to the classification of Fehr and Caviness (1977).

The emergence speed index (ESI), plant height (cm), leaf area (cm² plant⁻¹), stem diameter (mm), shoot dry mass (g), root length (cm), root diameter (mm), and root dry mass (g) were measured. Plant height, leaf area, and root length were measured using a millimeter ruler. Stem and root diameter were measured using a pachymeter. Forced air circulation oven and analytical balance (0.001) were used to determine shoot and root dry mass. The emergence speed index was calculated based on Maguire's formula (Maguire, 1962). The leaf area was measured according to the methodology described by Richter *et al.* (2014).

The data were submitted to ANOVA F-test. Tukey test, at 5% of significant level, was used for the soybean cultivars and polynomial regression analysis for the potassium silicate doses. Data analysis was performed using the SAS software (Cody, 2015).

RESULTS AND DISCUSSION

According to the initial development analyses, the emergence speed index presented no significant effect

on the soybean cultivars submitted to different silicon doses. The average emergence speed index was 6.4% (Figure 1). This result might be related to the high-vigor soybean seeds that were used in this study, which led to a faster emergence. Some researchers registered similar results. Pereira *et al.* (2004) found that seed treatment with potassium silicate did not influence the emergence speed index of rice (*Oryza sativa* L.).

In irrigated rice, Vieira *et al.* (2011) observed a decrease in the emergence percentage and the emergence speed index when silicon doses are applied 30 d before sowing. This fact may be due to toxicity generated over extra high calcium silicate doses in the paddy field. These authors stated that highest emergence percentage and emergence speed index were observed in areas that did not receive any calcium silicate dose (0 t ha⁻¹), with an emergence speed index of 3.3 and an emergence percentage of 77%.

Seed treatment with potassium silicate did not influence the soybean plants' height. However, differences were observed between the soybean cultivars (Figure 2). The FTR 1192 IPRO (C2) cultivar presented the highest height, differing only from the FTR 3190 IPRO (C3) cultivar, with mean values of 28.5 and 24.8 cm, respectively. The FTR 1186 IPRO (C1) cultivar average height was 26.7 cm. The FTR 1192 IPRO cultivar has a high potential for rooting and rusticity which may have




Figure 1. Emergence speed index of soybean seedlings submitted to silicon fertilization via seeds.

avored its prominence concerning the FTR 3190 IPRO cultivar.

Oliveira *et al.* (2015) studied the effect of seed treatment with different doses of silicon on the height of two soybean genotypes. The silicon treatment did not influence the height of BMX Turbo RR and NA 5909 RR genotypes

while plants that received no silicon application reached greater heights. Still, Pereira Júnior (2008) registered higher heights in soybean plants treated with silicon, contrasting the results mentioned above. The BRS MG 68 cultivar presented average height of 91, 96 and 94 cm when treated with 400, 450 and 500 kg Si ha⁻¹, respectively.




Figure 2. Plant height of three soybean cultivars submitted to silicon fertilization via seeds. Means followed by the same letter do not differ significantly ($\alpha=0.05$)

Regarding the leaf area, seed treatment with different doses of potassium silicate did not influence the soybean cultivars (Figure 3). According to Takahashi *et al.* (1990), legume plants usually accumulate low levels of silicon in

its cellular structures because they consume silicon in favor of a transpiration flow, which is a slower process when it is compared to water absorption. Besides, silicon is not diffused in the symplastic pathway.




Figure 3. Leaf area of soybean plants submitted to silicon fertilization via seeds.

Comparable results to the ones found in this study were observed by Oliveira *et al.* (2015) that emphasize that the morphological characteristics of the soybean varieties BMX Turbo RR and NA 5909 RR are not influenced with the doses of silicon via seed treatment. Agarie *et al.* (1998) stated that silicon might be associated with leaf retention because of photosynthesis maintenance and chlorophyll distribution under stress conditions, such as high temperatures and low humidity. This remark might have influenced the greater leaf area and dry mass accumulation in the soybean plants submitted to silicon fertilization at the phenological stages mentioned above.

Silicon is deposited in the plants' structure through root or foliar uptake. It stabilizes in the form of hydrated amorphous silica ($\text{SiO}_2 \cdot n\text{H}_2\text{O}$) with polysaccharides (cellulose and hemicellulose) on cell walls. The presence of silicon makes the leaves more upright, which enhances CO_2 uptake, as well as solar radiation exposure. As a result, it might improve soybean photo-assimilates production, directly influencing the crops initial development and yield (Epstein, 1999).

Regarding the stem diameter, potassium silicate treatments did not influence the studied soybean cultivars (Figure 4).

However, several authors registered higher stem diameters in maize plants submitted to silicon fertilization (Neri *et al.*, 2009). These increases are probably related to the monocots plants ability to accumulate more silicon through an active uptake process substantially. The silicon accumulates in the lumen and intercellular spaces of Poaceae plants leaves and culms, such as maize and sugar cane (Epstein, 1999).

When studying the effect of micronutrients fertilization and seed inoculants on BRS 206 soybean cultivar, Souza *et al.* (2009) registered higher stem diameter values compared to the ones found in this study. However, no significant effect was observed between the applied treatments. The higher stem diameter values found by other researchers is related to the different soybean genotypes studied and different phenological stages evaluated.

Regarding the dry shoot mass, significant effect was observed for FTR 1192 IPRO (C2) cultivar, in response to the different doses of potassium silicate applied. The dry shoot mass of C2 cultivar adjusted to the quadratic model, presenting the higher value, approximately 2.1 g, when 120 g of silicon was applied (Figure 5). Because of the greater rooting, when compared with the others, the C2 variety

responded better concerning the increase of dry mass. It may be linked to the fact that with a more developed

root system the plant can absorb more nutrients and, as a result, develop better.




Figure 4. Stem diameter of soybean plants submitted to silicon fertilization via seeds.




Figure 5. Shoot dry mass of soybean plants submitted to silicon fertilization via seeds.

Sousa *et al.* (2010) verified significant variations in the dry culm mass of maize plants fertilized with K_2SiO_3 . Soratto *et al.* (2012) and Figueiredo *et al.* (2007) reported higher shoot dry mass of strawberry and potato plants fertilized with silicon, respectively. Teodoro *et al.* (2015) studied the K_2SiO_3 foliar fertilization effect in the soybean cultivar 5DR615 and verified greater total dry mass production in the crop R6 phenological stage.

Regarding the root parameters, FTR 1186 IPRO (C1) cultivar root length and diameter linearly increased as the K_2SiO_3 doses increased (Figures 6 and 7). The C1 cultivar average values for root length and diameter when treated with 120 g of silicon were 8.7 cm and 4.5 mm, respectively. The FTR 1192 IPRO and FTR 3190 IPRO cultivars root length and diameter presented no significant results in response to the silicon treatment.




Figure 6. Root length of soybean cultivars submitted to silicon fertilization via seeds.

The results mentioned above verified for the C1 cultivar might be related to the SiO_2 accumulation by the roots. Silicon-non-accumulative species are unable to accumulate and translocate high levels of silicon in their shoots; thus they tend to concentrate this element in the root system, where

most of it is deposited in the cell walls (Heine *et al.*, 2005). Soybean plants translocate low levels of silicon to the shoots, up to 30 mg kg^{-1} . This substantiates the fact that in legume plants, such as soybean and bean plant, there is greater silica (SiO_2) accumulation in the roots.




Figure 7. Root diameter of soybean cultivars submitted to silicon fertilization via seeds.

Silicon absorption is mediated by diffusion in soybean and bean plants, through the movement of mono-silicic acid from a high to a low concentration region. This transport process may require energy, due to the silicon's

translocation across the water channel proteins, the aquaporins. In this process, silicon molecules retention by the suberin layer present in the root endodermis can occur, making the roots thicker (Raven, 2001).

Silicon is an important element for the soybeans root system, once enhanced nodulation and nitrogen fixation is observed in silicon fertilized soybean crops (Figueiras, 2007). However, some factors must be observed before applying silicon fertilization, such as the genetic characteristics of the cultivar, its ability to consume, translocate and accumulate the element, as well as the soil and environment conditions.

The silicon doses did not influence the roots' dry mass but the different soybean cultivars presented statistical differences among themselves. The FTR 1192 IPRO (C2) cultivar presented the highest root dry mass compared to the other studied cultivars (Figure 8). These results reveal that same species plants (*G. max*) genotypes, presented different root dry mass accumulation patterns.




Figure 8. Root dry mass of soybean cultivars submitted to silicon fertilization via seeds. Means followed by the same letter do not differ significantly ($\alpha=0.05$).

Paula *et al.* (2007) reported higher roots dry mass in fava d'anta plants (*Demorphanthra mollis* Benth) submitted to potassium silicate fertilization. Still, Ribeiro *et al.* (2011) verified limited root dry mass accumulation in coffee plants submitted to silicon fertilization after 130 of treatment.

CONCLUSIONS

The treatment of soybean seeds with silicon did not positively influence the root dry mass of the FTR 1192 IPRO variety and the root length and root diameter of the FTR 1186 IPRO variety. The application of silicon did not influence the FTR 3190 IPRO cultivar. Regarding the plant height and dry root mass parameters, the FTR 1192 IPRO cultivar presented more precocity compared to the other genotypes.

REFERENCES

Agarie S, Uchida H, Agata W, Kubota F and Kaufman PB. 1998. Effects of silicon on transpiration and leaf conductance in rice plants (*Oryza sativa* L.). *Plant Production Science* 1(2): 89-95. doi: 10.1626/pps.1.89

Castro GSA and Cruscio CAC. 2013. Effects of superficial liming and silicate application on soil fertility and crop yield under rotation. *Geoderma* 195-196: 234-242. doi: 10.1016/j.geoderma.2012.12.006

Cody R. 2015. *An Introduction to SAS University Edition*. SAS Institute. 366 p.

Conab - Companhia Nacional de Abastecimento. 2016. Levantamento de Safra (safra 2016/2017). In <http://www.conab.gov.br>; accessed: November 2017.

Currie HA and Perry CC. 2007. Silica in plants: biological, biochemical and chemical studies. *Annals of Botany* 100(7): 1383-1389. doi: 10.1093/aob/mcm247

Daronco MV. 2013. Óleos essenciais no tratamento de sementes de soja (*Glycine max* L.). Bachelor's dissertation in Agronomy Engineer. Department of Agrarian Studies. Universidade Regional do Noroeste do Estado do Rio Grande do Sul, Ijuí. 50 p.

Datnoff LE, Rodrigues FA and Seebold KW. 2007. Silicon and Plant Nutrition. pp. 233-246. In: Datnoff LE, Elmer WH and Huber DM (eds.). *Mineral Nutrition and Plant Disease*. American Phytopathological Society Press, St. Paul. 278 p.

Embrapa - Empresa Brasileira de Pesquisa Agropecuária. 2014. *Sistema Brasileiro de Classificação de Solos*. Fourth edition. Embrapa, Brasília. 376 p.

Embrapa - Empresa Brasileira de Pesquisa Agropecuária. 2016. Soja em números (safra 2015/2016). Embrapa Soja. 2016. In www.embrapa.br; accessed: October 2016.

- Epstein E. 1999. Silicon. Annual review of plant biology 50(1): 641-664. doi: 10.1146/annurev.arplant.50.1.641
- Fehr WR and Caviness CE. 1977. Stages of soybean development. Special Report. 87. Iowa State University, Ames. 12 p.
- Ferreira SM. 2008. Efeito do silício na cultura do algodoeiro (*Gossypium hirsutum* L.): Aspectos bioquímicos, qualidade de fibra e produtividade. Ph.D's Thesis in Applied Ecology. Escola Superior de Agricultura "Luiz de Queiroz". Universidade de São Paulo, Piracicaba. 68 p.
- Figueiras O. 2007. Silício na agricultura. Pesquisa Fapesp. São Paulo. In: Pesquisa, <http://revistapesquisa.fapesp.br/es/2007/10/01/silicio-en-la-agricultura>; accessed: November 2017.
- Figueiredo FC, Rodrigues CR, Botrel PP and Rodrigues TM. 2007. Benefícios do silício líquido solúvel em olerícolas. Revista Campos & Negócios 3(36): 50-51.
- Heine G, Tikum G and Horst WJ. 2005. Silicon nutrition of tomato and bitter melon with special emphasis on silicon distribution in root fractions. Journal of Plant Nutrition and Soil Science 168(4): 600-606. doi: 10.1002/jpln.200420508
- Kolchinski EM, Schuch LOB and Peske ST. 2005. Vigor de sementes e competição intra-específica em soja. Ciência Rural 35(6): 1248-1256. doi: 10.1590/S0103-84782005000600004
- Machado BR and Queiroz SEE. 2018. Efeito do tratamento de sementes de soja com silício e polímero na qualidade fisiológica das sementes e nas características agrônomicas. Enciclopédia Biosfera 15(27): 1576-1584.
- Maguire JD. 1962. Speed of germination – aid in selection and evaluation for seedling emergence and vigor. Crop Science 2(2): 176-177. doi: 10.2135/cropsci1962.0011183X000200020033x
- Neri DKP, Gomes FB, Moraes JC, Góes GB and Marrocos STP. 2009. Influência do silício na suscetibilidade de *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) ao inseticida lufenuron e no desenvolvimento de plantas de milho. Ciência Rural. 39(6): 1633-1638. doi: 10.1590/S0103-84782009005000111
- Oliveira S, Lemes ES and Tavares LC. 2013. Tratamento de sementes: Ferramenta promissora e eficiente para o agricultor. Seed News 17: 8-11.
- Oliveira S, Lemes ES, Mendonça AO, Ritter R and Meneghello GE. 2014. Efeitos da aplicação de silício via sementes na produtividade e na qualidade de sementes de soja. Enciclopédia Biosfera 10(19): 1-14.
- Oliveira S, Lemes ES, Mendonça AO, Diaz LW, Brunes AP, Leitzke ID and Meneghello GE. 2015. Tratamento de sementes de soja com silício: efeito na qualidade fisiológica e nas características agrônomicas. Revista Cultivando o Saber 8(2): 215-230.
- Paula TOM, Santos AM, Valadares SV, Junior CFC, Fernandes LA, Martins ER and Alves DS. 2007. Influência do silicato no crescimento inicial e produção de flavonoides totais em *Demorphandra mollis* Benth. Revista Brasileira de Biociências 5 Supp. 2: 552-554.
- Peel MC, Finlayson BL and McMahon T.A. 2007. Updated world map of the Köppen-Geiger climate classification. Hydrology and Earth System Sciences 11(5): 1633-1644.
- Pereira HS, Korndörfer GH, Vidal AA and Camargo MS. 2004. Silicon sources for rice. Scientia Agrícola 16(5): 522-528.
- Pereira Júnior P. 2008. Doses de silício na produtividade de soja [*Glycine max* (L.) Merrill] e suas características agrônomicas. Master's Thesis in Plant Science. Universidade Federal de Lavras, Lavras. 37 p.
- Raven JA. 2001. Chapter 3 Silicon transport at the cell and tissue level. pp. 41-55. In: Datnoff LE, Snyder GH and Korndörfer GH (eds.). Studies in plant science. (Vol. 8). Silicon in Agriculture. Elsevier. 403 p. doi: 10.1016/S0928-3420(01)80007-0
- Ribeiro RV, Silva L, Ramos RA, Andrade CA, Zambrosi FCB and Pereira SP. 2011. O alto teor de silício no solo inibe o crescimento radicular de cafeeiros sem afetar as trocas gasosas foliares. Revista Brasileira de Ciência do Solo 35(1): 939-948. doi: 10.1590/S0100-06832011000300028
- Richter GL, Zanon Júnior A, Streck NA, Guedes JVC, Kräulich B, da Rocha TSM, Winck JEM and Cera JC. 2014. Estimating leaf area of modern soybean cultivars by a non-destructive method. Bragantia 73(4): 416-425. doi: 10.1590/1678-4499.0179
- Soratto RP, Fernandes AM, Crusciol CAC and Souza-Schilick GD. 2012. Produtividade, qualidade de tubérculos e incidência de doenças em batata, influenciados pela aplicação foliar de silício. Pesquisa Agropecuária Brasileira 47(7): 1000-1006. doi: 10.1590/S0100-204X2012000700017
- Sousa JV, Rodrigues CR, Luz JMQ, Carvalho PC, Rodrigues TM and Brito CH. 2010. Silicato de potássio via foliar no milho: fotossíntese, crescimento e produtividade. Bioscience Journal 26(4): 502-513.
- Souza LCF, Zanon GD, Pedrosa FF and de Andrade LHL. 2009. Teor de proteína e de óleo nos grãos de soja em função do tratamento de sementes e aplicação de micronutrientes. Ciência e Agrotecnologia 33(6): 1586-1593. doi: 10.1590/S1413-70542009000600018
- Takahashi E, Ma JF and Miyake Y. 1990. The possibility of silicon as an essential element for higher plants. Comments on Agricultural and Food Chemistry 2(2): 99-102.
- Teodoro PE, Ribeiro LP, Oliveira EPD, Corrêa CCG and Torres FE. 2015. Acúmulo de massa seca na soja em resposta a aplicação foliar com silício sob condições de déficit hídrico. Bioscience Journal 31(1): 161-170. doi: 10.14393/BJ-v31n1a2015-22283
- Vieira AR, Oliveira JA, Guimarães RM, Carvalho MLM, Pereira EM and Carvalho BO. 2011. Qualidade de sementes de arroz irrigado produzidos com diferentes doses de silício. Revista Brasileira de Sementes 33(3): 490-500. doi: 10.1590/S0101-31222011000300012
- Wise H, Nikolic M and Romheld V. 2007. Silicon in plant nutrition: Effects on zinc, manganese and boron leaf concentrations and compartmentation. pp. 33-48. In: Sattelmacher B and Horst WJ (eds.). The apoplast of higher plants: Compartment of storage, transport and reactions. Springer, London. 458 p.
- Zelin E, Bussolaro I and Simonetti APMM. 2011. Aplicação de silício no controle de lagartas e produtividade da cultura da soja. Cultivando o Saber 4(1): 171-180.