



Revista Facultad Nacional de Agronomía Medellín

ISSN: 0304-2847

ISSN: 2248-7026

Facultad de Ciencias Agrarias - Universidad Nacional de Colombia

Liriano González, Ramón; Pérez Ramos, Jovana; Pérez Hernández, Yunel;  
Placeres Espinosa, Iraní; Rodríguez Jiménez, Sergio Luis; Peña Alonso, Daniela  
Improvement of the agricultural productivity of lettuce and radish by using efficient microorganisms  
Revista Facultad Nacional de Agronomía Medellín, vol.  
72, no. 3, 2019, September-December, pp. 8937-8943  
Facultad de Ciencias Agrarias - Universidad Nacional de Colombia

DOI: <https://doi.org/10.15446/rfnam.v72n3.76967>

Available in: <https://www.redalyc.org/articulo.oa?id=179961330006>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UAEM  redalyc.org

Scientific Information System Redalyc  
Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

# Improvement of the agricultural productivity of lettuce and radish by using efficient microorganisms

Mejoramiento de la productividad agrícola de la lechuga y el rábano con el uso de microorganismos eficientes

doi: 10.15446/rfnam.v72n3.76967

Ramón Liriano González<sup>1</sup>\*, Jovana Pérez Ramos<sup>1</sup>, Yonel Pérez Hernández<sup>1</sup>, Iraní Placeres Espinosa<sup>1</sup>, Sergio Luis Rodríguez Jiménez<sup>1</sup> and Daniela Peña Alonso<sup>1</sup>

## ABSTRACT

### Keywords:

Biofertilizer  
*Lactuca sativa*  
Organoponic garden  
*Raphanus sativus*

The aim of this work was to evaluate the effect of the application of efficient microorganisms (EM) on the productive performance of lettuce and radish. The experiment was carried out in an organoponic culture in the municipality of Matanzas, Cuba, from December 2017 to January 2018. Five treatments were studied: control (0), EM 8 mL m<sup>-2</sup> at 0 days after the transplant (DAT) (EM 8-0), EM 8 mL m<sup>-2</sup> at 15 DAT (EM 8-15), EM 10 mL m<sup>-2</sup> at 0 DAT (EM 10-0), and EM 10 mL m<sup>-2</sup> at 15 DAT (EM 10-15). A randomized complete block design was set with four replications per treatment. An ANOVA was applied to perform the statistical data analysis, and the Duncan's Multiple Range Test ( $P < 0.05$ ) was used for the comparison of means. The statistic program used was STATISTICA, version 6.0 over Windows. The parameters evaluated in lettuce were the total number of leaves, number of commercial leaves, the diameter of leaf rosette, and yield. The evaluated parameters for radish were fleshy root's diameter and weight, and yield. The results indicated a positive effect on growth-response with the application of the bio-product, which can be considered a promissory alternative for vegetable production in organoponic garden conditions. The application of EM 10 mL m<sup>-2</sup> at 0 and 15 DAT showed the best productive behavior for both crops.

## RESUMEN

### Palabras clave:

Biofertilizante  
*Lactuca sativa*  
Jardín organopónico  
*Raphanus sativus*

El objetivo de este trabajo fue evaluar el efecto de la aplicación de microorganismos eficientes (EM) en el rendimiento productivo de cultivos de lechuga y rábano. El experimento se llevó a cabo en un cultivo organopónico en el municipio de Matanzas, Cuba, desde diciembre de 2017 hasta enero de 2018. Se estudiaron cinco tratamientos: control (0), EM 8 mL m<sup>-2</sup> a los 0 días después del trasplante (DAT) (EM 8-0), EM 8 mL m<sup>-2</sup> a 15 DAT (EM 8-15), EM 10 mL m<sup>-2</sup> a 0 DAT (EM 10-0) y EM 10 mL m<sup>-2</sup> a 15 DAT (EM 10-15). Se estableció un diseño de bloques completos al azar con cuatro repeticiones por tratamiento. Se aplicó un ANOVA para realizar el análisis estadístico de datos, y se utilizó la prueba de rango múltiple de Duncan ( $P < 0.05$ ) para comparar las medias. El programa estadístico utilizado fue STATISTICA, versión 6.0 en Windows. Los parámetros evaluados en lechuga fueron el número total de hojas, el número de hojas comerciales, el diámetro de la roseta y el rendimiento. Los parámetros evaluados para el rábano fueron el diámetro y peso de la raíz carnosa y el rendimiento. Los resultados indicaron un efecto positivo en la respuesta de crecimiento con la aplicación del bio-producto, que puede considerarse una alternativa promisoría para la producción de vegetales en condiciones organopónicas de jardín. La aplicación de EM 10 mL m<sup>-2</sup> a 0 y 15 DAT mostró el mejor comportamiento productivo para ambos cultivos.

<sup>1</sup> Universidad de Matanzas. Autopista a Varadero, km 31/2. Matanzas, Cuba.

\* Corresponding author: <ramon.liriano@umcc.cu>

The use of agrochemicals has grown considerably since the “Green Revolution,” which has caused a reduction in the productivity of soils due to several degrading processes, such as soil compaction, salinization, desertification, and contamination with heavy metal. It provokes severe environmental problems such as eutrophication of water bodies and diseases in humans and animals (Ge *et al.*, 2016; Katiyar *et al.*, 2016). Therefore, the excessive production and application of non-organic synthetic fertilizers cannot be the solution to satisfy the growing need for food to supply the world population.

Several researchers have been focused on friendly, sustainable, and organic agricultural practices; which allow to reducing the costs of production, contributing to the sustainability of agricultural systems and maintaining stable the yield and crop quality (Mesa, 2016). In Cuba, at the end of the 80s, the vegetable production began to develop on a large scale in urban areas, which grew every year to reach 4,200 t in 1994, 480,000 t in 1998 and more than 1,158,452 t in 2015, without considering the productions in gardens and plots, mainly for family consumption (GNAUS, 2015).

This intensive method for vegetable production on an organic substrate has favored obtaining high yields of crops, but at the same time, it requires adequate technological discipline, where the exploitation and management of the substrates result in a vital issue. It is because plants constantly take nutrients from the soil, being removed after harvest; therefore, the productivity of the agricultural systems depend on the different initial nutrient organic resources to guarantee high yields and multiple harvests, at least for two years. At longer periods, nutrients become limited and in consequence, yield and quality decrease.

In this context, organic fertilization should be complemented with biological products based on beneficial microorganisms (the so-called plant probiotics), which stimulate growth and development of plants throughout different mechanisms such as: the synthesis of bioactive compounds like auxin, cytokinin and gibberellin (Khatab *et al.*, 2015; Damam *et al.*, 2016); the solubilization of phosphates (Basu *et al.*, 2017; Ramírez-Gil, 2019), the production of siderophores (Stamenković *et al.*, 2018), the production of lytic enzymes and antibiotics, which play an important role in the organic matter decomposition and bio-control of phytopathogens (Sabaté *et al.*, 2018; Thakur *et al.* 2017).

Those biofertilizers have been pointed out as a viable alternative for achieving an agriculture development with ecological sustainability, based on low costs of production and a minimal impact on the environment (Menendez and Garcia-Fraile, 2017). For this reason, more investigations need to be done in order to study the effect of natural products on crops' growth. The aim of the present work was to evaluate the effect of the application of efficient microorganisms on the productive response of lettuce (*Lactuca sativa* L.) and radish (*Raphanus sativus* L.) at organoponic conditions.

## MATERIAL AND METHODS

### Study area and EM applied

The experiment was carried out in an organoponic system at the municipality of Matanzas, Cuba, from December 2017 to January 2018. Lettuce (*Lactuca sativa* L. variety Riza-15) and radish (*Raphanus sativus* L. variety PS-9) were used in association. In the case of lettuce, the transplanting was used as sowing method, with six rows on the plot at a distance of 15 cm between rows and 15 cm between each plant. Direct sowing method was used for radish, which was planted in one row in the center of the bed with 3 cm between each plant. The average temperature during the experiment (dry season) was 21.7 °C, the relative humidity was 80.9%, and the average rain precipitation was 7.6 mm.

The efficient microorganisms (bio-product) was obtained from the bio-pesticide production laboratory “LABIOFAM” at the province of Matanzas. The microbiological composition analysis of the EM bio-product used was conducted at the Laboratory of Microbiology of the Faculty of Agronomy Sciences of Universidad de Matanzas, Cuba. The isolation of the microbial groups (bacteria, fungi, and yeast) from EM bio-preparation was carried out using 1 mL of the sample following the serial dilutions methodology (Stanier, 1996). Dilutions up to  $10^{-6}$  were prepared in a sterile saline solution (0.9% of NaCl). 0.5 mL of the dilutions was inoculated on Nutrient Agar Medium for bacteria ( $10^{-6}$ ), Saborout Agar Medium for fungi ( $10^{-5}$ ) and Potato Dextrose Agar Medium for yeast ( $10^{-4}$ ). Petri dishes for bacteria were incubated at 37 °C for 24 h, whereas fungi and yeast were grown at 30 °C for 72 h. Table 1 shows the concentration of the main microbial groups found in the initial EM preparation.

**Tabla 1.** Microbiological composition of EM bio-product.

Microbial groups	Concentration (CFU mL <sup>-1</sup> )
Bacteria	13×10 <sup>8</sup>
Fungi	18×10 <sup>5</sup>
Yeast	21×10 <sup>6</sup>

### Experimental design

The experiment was conducted in a randomized complete block design with four replications; each one consists of an organoponic bed of 30.0 m length and 1.20 m width. With five treatments, the experimental units were of 6.0 m<sup>2</sup>. The substrate consisted of 50% soil and 50% organic matter.

The treatments studied were:

0 = Control (without EM application).

EM 8-0 = EM (8 mL m<sup>-2</sup>) at 0 days of transplanting.

EM 8-15 = EM (10 mL m<sup>-2</sup>) at 0 days of transplanting.

EM 10-0 = EM (8 mL m<sup>-2</sup>) at 0 and 15 days of transplanting.

EM 10-15 = EM (10 mL m<sup>-2</sup>) at 0 and 15 days of transplanting.

The application of the EM bio-product was carried out early in the morning (7:00-7:30 am) using manual fumigation equipment (MATABI) of 16 liters of capacity. The application on radish was subordinated to the main crop. Insects, diseases, and weeds were intensively controlled, according to GNAUS (2007). At harvesting (28 days after seed germination in radish and 45 days after transplantation in the case of lettuce), 25 plants were randomly hand-taken in each experimental plot, and it was recorded the following morphometric and yield parameters in each crop:

*Lettuce*: Number of leaves per plant (by direct counting), number of commercial leaves per plant, rejecting those

not suitable for commercialization; leaf rosette diameter using a measuring tape (morphometric parameters) and the yield (kg m<sup>-2</sup>). Plants were weighted on with scale (Sartorius, ALC-110.4).

*Radish*: Fleshy root diameter (using a Vernier caliper), the weight of fleshy root and the yield, being the latter represented as kilogram per linear meter (kg m<sup>-1</sup>) because of its sowing location.

### Statistical analysis

The data were statistically evaluated by the Kolmogorov-Smirnov's and Bartlett's Test to verify normality and variance homogeneity, respectively. An Analysis of Variance (ANOVA) and Duncan's Multiple Ranges Test ( $P < 0.05$ ) (Duncan, 1955) were used for the comparison of means. STATISTICA program version 6.0 was utilized to process the experimental data.

## RESULTS AND DISCUSSION

### Effect the EM in lettuce

The application of EM increased the number of total leaves and the number of commercial leaves of lettuce (Table 2). Regarding the total leaves number, all the treatments with the application of efficient microorganisms displayed higher values compared with the control. Similarly, the number of commercial leaves in 0 (control) was lower than the variants with different dose of EM and times of application.

**Table 2.** Effect of the application of efficient microorganisms on the number of total and commercial leaves of lettuce variety Riza-15.

Treatments	Total leaves per plant	Commercial leaves per plant
0	13.29 b	12.31 b
EM 8-0	17.31 a	16.23 a
EM 8-15	17.59 a	16.59 a
EM 10-0	17.72 a	16.74 a
EM 10-15	18.05 a	17.16 a
± SE x	0.015	0.018

Different letters mean differences among treatments for each parameter ( $P < 0.05$ ) (Duncan, 1955). ± SE x: Standard Error of the Mean.

The higher production of commercial leaves in lettuce after EM inoculation should be positively influenced by growth parameters of the crop, which in turn determined the achieved yield. The beneficial effect of EM on the development of plant morphometric and productivity characters was also reported by several authors in different plant species such as *Spinacia oleracea* L. (Hauka *et al.*, 2016), *Phaseolus vulgaris* L. (Estrada *et al.*, 2017), *Oryza sativa* L. (Ghaffari *et al.*, 2018) and *Rubus glaucus* Benth. cv. Thornless (Robledo-Buriticá *et al.*, 2018).

The EM application increased the diameter of leaf rosette and yield (Table 3). The treatment with EM inoculation of

10 mL m<sup>-2</sup> at 0 and 15 days after transplanting recorded the maximum diameter among the different variants of EM applications, which values ranged between 29.33 up to 36.56 cm. The diameter of leaf rosette in plants control was lower in comparison with all the assayed treatments.

Regarding the yield parameter in the lettuce, the treatments with the higher dose of EM (10 mL m<sup>-2</sup>) and the variant EM 8 mL m<sup>-2</sup> at both 0 and 15 days of transplanting promoted higher yield concerning the control; whereas the variant EM 8 mL m<sup>-2</sup> at 0 days of transplanting (2.29 kg m<sup>-2</sup>) showed no statistical difference with the control (1.85 kg m<sup>-2</sup>).

**Table 3.** Effect of the application of efficient microorganisms on the diameter of leaf rosette and yield of lettuce variety Riza-15.

Treatments	Diameter of leaf rosette (cm)	Yield (kg m <sup>-2</sup> )
0	29.33 c	1.85 b
EM 8-0	33.08 b	2.29 ab
EM 8-15	33.43 b	2.58 a
EM 10-0	34.11 b	2.61 a
EM 10-15	36.56 a	2.67 a
± SE x	0.58	0.31

Different letters mean differences among treatments for each parameter ( $P < 0.05$ ) (Duncan, 1955). ± SE x: Standard Error of the Mean.

The positive response observed on lettuce yield might be related with the increment of the microbiota diversity in the substrate after the application of beneficial microorganisms, which in turn could improve several physiological processes such as photosynthetic activity, growth and productivity of the crops (Pedarza *et al.*, 2010). They pointed out that efficient microorganisms consist of beneficial microorganism mixture (mainly acid lactic-producer and photosynthetic bacteria, yeast, actinomycetes, and fermenting fungi), which can be applied as an inoculant to increase the microbial diversity of soils, followed by an improvement of their quality and health. This EM application allows to enhancing growth, quality, and yield of the crops.

Several researchers have reported good results in plant growth, biochemical parameters, and crop productivity by using efficient microorganisms. Arismendi (2010) observed an enhancement of the weight in lettuce variety Great Lakes 659 after the inoculation with EM.

Similarly, the application of fermented organic matter, along with commercial EM on the lettuce var. Iceberg, improved the plant height, weight, and diameter of the rosette (Pomboza-Tamaquiza *et al.*, 2016). Szczech *et al.* (2016) found a better behavior of germination, fresh mass of transplants, and nitrogen content in lettuce after the application of a combination of beneficial bacterial strains.

#### Effect the EM on radish

The effect of the inoculation of efficient microorganisms on the diameter, weight, and yield of radish fleshy root are shown in Table 4. The application of EM revealed significant root growth-promoting effects, increasing productivity. Regarding the diameter of root and yield, all the treatments with EM showed higher results than the control without statistical difference among them. Regarding the root weight, EM 10-0 (3.89 g) and EM 10-15 (3.95 g) showed the best results in comparison with EM 8-0 and EM 8-15. The control registered the

minimum yield with an average of 0.51 kg m<sup>-1</sup>. The obtained yield with the application of EM was higher than

the reported by the National Group of Urban Agricultural in 2007 ranged between 0.5 to 0.8 kg m<sup>-1</sup>.

**Table 4.** Effect of the application of efficient microorganisms on diameter, fresh weight and yield of fleshy root of radish variety PS-9.

Treatments	Diameter of the root (cm)	Weight of root (g)	Yield (kg m <sup>-1</sup> )
0	2.41 b	2.01 c	0.51 b
EM 8-0	3.24 a	2.38 b	1.16 a
EM 8-15	3.29 a	2.41 b	1.31 a
EM 10-0	3.64 a	3.89 a	1.40 a
EM 10-15	3.68 a	3.95 a	1.49 a
± SE x	0.029	0.017	0.024

Different letters mean differences among treatments for each parameter ( $P < 0.05$ ) (Duncan, 1955). ± SE x: Standard Error of the Mean.

The results observed in the present investigation are in agreement with those reported by Mali *et al.* (2018), who recorded an increment in root weight and yield of radish cv. Japanese white, when applied a combination of organic manure and biofertilizer. Similar trials with *Raphanus sativus* L. reported better results on leaf area, root length, fresh and dry weight, and yield; after supplying vermicompost 12.5 t ha<sup>-1</sup> and a microbial consortium (Pathak *et al.*, 2017). In addition, the positive effect of native microorganisms on vegetable yield have also been reported by Núñez *et al.* (2017) in carrot (*Daucus carota* L.) under organoponic conditions. Those authors observed a good response in yield and its attributes with the inoculation of a biofertilizer, mainly with a dose of 10 mL m<sup>-2</sup> which produced an increment of 0.72 kg m<sup>-2</sup>.

These findings could be explained due to the effect of the efficient microorganisms on plant growth in different ways, for example throughout the production of phytohormones such as auxin and cytokine-like compounds (Berger *et al.*, 2015; Nghia *et al.*, 2017), the solubilization of minerals such as phosphate and nitrogen (Changas-Junior *et al.*, 2015) and indirectly by the production of substances with antibiotic activity which reduce the number of phytopathogen microorganisms (Grosu *et al.*, 2015). Moreover, when EM interact with the organic matter, other beneficial compounds are also released, such as vitamins, organic acids, minerals, and antioxidants (IICA, 2013).

The inoculation of the substrates with efficiency microorganisms may have enhanced the soil microbiological status and the rate of decomposition of the organic matter as well, which in turn could improve the physicochemical and biological properties of soil and the content of humus (Navia-Cuetia *et al.*, 2013; Campo-Martínez *et al.* 2014). These processes release minerals available for plant nutrition, which may stimulate the cellular metabolism, photosynthesis, growth, and development of crops (Kumar *et al.*, 2016; Mani and Anburani, 2018).

## CONCLUSIONS

The application of EM in lettuce-radish association under organoponic conditions improved morphometric parameters and yield. The dose of 10 mL m<sup>-2</sup> at 0 and 15 days of transplanting, showed the best productive performance in both crops. The results obtained indicate the potential of this technology to increase the yield and quality of lettuce and radish.

## REFERENCES

- Arismendi E. 2010. Microorganismos Eficientes, ¿fórmula mágica?. En: [http://www.rapaluruquay.org/organicos/articulos/microorganismos\\_eficientes.html](http://www.rapaluruquay.org/organicos/articulos/microorganismos_eficientes.html) consulta: marzo 2018.
- Basu S, Rabara R and Negi S. 2017. Towards a better greener future - An alternative strategy using biofertilizers. I: Plant growth promoting bacteria. Plant Gene. 12: 43-49. doi: 10.1016/j.plgene.2017.07.004
- Berger B, Wiesner M, Brock AK, Schreiner M and Ruppel S. 2015. *K. radicincitans*, a beneficial bacteria that promotes radish growth under field conditions. Agronomy for Sustainable Development 35(4):1521–1528. doi: 10.1007/s13593-015-0324-z

- Campo-Martínez ADP, Acosta RL, Morales S y Alfonso F. 2014. Evaluación de microorganismos de montaña (MM) en la producción de acelga en la meseta de Popayán. *Bioteconología en el Sector Agropecuario y Agroindustrial* 12(1): 79-87.
- Changas-Junior AF, de Oliveira AG, de Oliveira LA, dos Santos GR, Changas LFB, Lopes da Silva AL and da Luz Costa J. 2015. Production of indole-3-acetic acid by *Bacillus* isolated from different soils. *Bulgarian Journal of Agricultural Science*. 21(2): 282-287.
- Damam M, Kaloori K, Gaddam B and Kausar R. 2016. Plant growth promoting substances (Phytohormones) produced by rhizobacterial strains isolated from the rhizosphere of medicinal plants. *International Journal of Pharmaceutical Sciences Review and Research* 37(1): 130-136.
- Duncan DB. 1955. Multiple range and multiple F test. *Biometrics*. 11(1): 1-42.
- Estrada W, Chávez L, Jerez E, Nápoles MC, Sosa A, Cordoví C y Celeiro F. 2017. Efecto del Azofert® en el rendimiento de variedades de frijol común (*Phaseolus vulgaris* L.) en condiciones de déficit hídrico. *Centro Agrícola* 44(3): 36-42.
- Ghaffari H, Gholizadeh A, Biabani A, Fallah A and Mohammadian M. 2018. Plant growth promoting rhizobacteria (pgpr) application with different nitrogen fertilizer levels in rice (*Oryza sativa* L.). *Pertanika Journal of Tropical Agricultural Science*. 41(2): 715 – 728.
- Ge C, Radnezhad H, Abari MF, Sadeghi M and Kashi G. 2016. Effects of biofertilizers and plant growth promoting bacteria on the growth characteristics of *Asparagus officinalis*. *Applied Ecology and Environmental Research*. 14(3): 547-558. doi: 10.15666/aeer/1403\_547558
- Grosu AI, Siciua OA, Dobre A, Voaides C and Cornea C. 2015. Evaluation of some *Bacillus* spp. strains for the biocontrol of *Fusarium graminearum* and *F. culmorum* in wheat. *Agriculture and Agricultural Science Procedia*. 6: 559-566. doi: 10.1016/j.aaspro.2015.08.085
- GNAUS - Grupo Nacional de Agricultura Urbana. 2007. Manual Técnico para Organopónicos y Huertos Intensivos. Ciudad de La Habana, Cuba. 161 p.
- GNAUS - Grupo Nacional de Agricultura Urbana, Suburbana y Familiar. 2015. Lineamientos de la agricultura urbana, suburbana y familiar para el año 2016. Ministerio de la Agricultura. INIFAT. La Habana, Cuba. p 14.
- Hauka FIA, Moslam TM, Ghanem KhM and El-Shahat MM. 2016. Impact of some plant growth promoting rhizobacteria "pgpr" on organically cultivated spinach plants (*Spinacia oleracea* L.). *Journal of Agricultural Chemistry and Biotechnology* 7(9): 235 – 240.
- IICA - Instituto Interamericano de Cooperación para la Agricultura. 2013. Tecnología de bajo costo: guía de manejo de microorganismos eficientes (ME). Proyecto Red SICTA, Cooperación Suiza en América Central. Managua, Nicaragua. p. 5-6.
- Katiyar D, Hemantaranjan A and Singh B. 2016. Plant growth promoting Rhizobacteria-an efficient tool for agriculture promotion. *Advances in Plants & Agriculture Research* 4(6): 426-434. doi:10.15406/apar.2016.04.00163
- Khatib OH, Nasib MAA, Ghoneimy EA, Abo-Elnasr AA, Hassan HA-A, Hassan MYA and Attitalla IH. 2015. Role of microorganisms in our life's as ecofriendly and replacement for chemical methods. *International Journal of Pharmacy and Life Sciences* 6(2):4221-4229.
- Kumar S, Kumar S, Maji S and Pandey VK. 2016. Effect of inorganic fertilizers and bio-fertilizers on growth, yield and quality of radish (*Raphanus sativus* L.). *International Journal of Plant Sciences (Muzaffarnagar)* 11(1): 71-74. doi: 10.15740/HAS/IJPS/11.1/71-74
- Mali DL, Singh V, Sarolia DK, Teli SK, Chittora A and Dhakar R. 2018. Effect of organic manures and bio-fertilizers on growth and yield of radish (*Raphanus sativus* L.) cv. Japanese white. *International Journal of Chemical Studies* 6(2): 1095-1098.
- Mani AP and Anburani A. 2018. Organic nutrient management technique for enhancing growth and physiological parameters in radish (*Raphanus sativus* L.). *Journal of Phytology* 10(1): 40-42. doi: 10.25081/jp.2018.v10.3461
- Menendez E and Garcia-Fraile P. 2017. Plant probiotic bacteria: solutions to feed the world. *AIMS Microbiology* 3(3): 502-524. doi: 10.3934/microbiol.2017.4.747
- Mesa JR. 2016. Efecto de un biopreparado de producción local a base de microorganismos eficientes sobre diferentes cultivos en la provincia de Cienfuegos. En: *Memorias IV Convención Internacional de Agrodesarrollo*. Cuba.
- Navia-Cuetia CA, Zemanate-Cordoba Y, Morales-Velasco S, Prado FA y Albán-López N. 2013. Evaluación de diferentes formulaciones de compostaje a partir de residuos de cosecha de tomate (*Solanum lycopersicum* L.). *Bioteconología en el Sector Agropecuario y Agroindustrial* 11(spe): 165 – 173.
- Nghia NK, Tien TTM, Oanh NTK and Nuong NHK. 2017. Isolation and characterization of indole acetic acid producing halophilic bacteria from salt affected soil of rice-shrimp farming system in the Mekong Delta, Vietnam. *Agriculture, Forestry and Fisheries* 6(3): 69-77. doi: 10.11648/j.aff.20170603.11
- Núñez DB, Liriano R, Pérez Y, Placeres I y Sianeh Zawolo G. 2017. Respuesta de *Daucus carota*, L. a la aplicación de microorganismos nativos en condiciones de organopónico. *Centro Agrícola* 44(2): 29-35.
- Pathak M, Tripathy P, Dash SK, Sahu GS and Pattanayak SK. 2017. Effect of source of nutrient on growth, yield and quality of Radish (*Raphanus sativus* L.) in radish - coriander cropping sequence. *The Pharma Innovation Journal* 6(12): 496-499.
- Pedraza RO, Teixeira KRS, Fernández A, García de Salamone IB, Azcón R, Vera LDB y Bonilla R. 2010. Microorganismos que mejoran el crecimiento de las plantas y la calidad de los suelos. *Ciencia y Tecnología Agropecuaria* 11(2): 155-164. doi: 10.21930/rcta.vol11\_num2\_art:206
- Pomboza-Tamaquiza P, León-Gordón OA, Villacís-Aldaz LA, Vega J y Aldáz-Jarrín JC. 2016. Influencia del biol en el rendimiento del cultivo de *Lactuca sativa* L. variedad Iceberg. *Journal of the Selva Andina Biosphere* 4(2): 84-92.
- Ramírez-Gil JG. 2019. Dependency, colonization, and growth in *Gmelina arborea* inoculated with five strains of arbuscular mycorrhizal fungi. *Revista Facultad Nacional de Agronomía Medellín* 72(2): 8775-8783. doi: 10.15446/rfnam.v72n2.74691
- Robledo-Buriticá J, Aristizábal-Loaiza JC, Ceballos-Aguirre N and Cabra- Cendales T. 2018. Influence of plant growth-promoting rhizobacteria (PGPR) on blackberry (*Rubus glaucus* Benth. cv. thornless) growth under semi-cover and field conditions. *Acta Agronómica* 67(2): 258-263. doi: 10.15446/acag.v67n2.62572
- Sabaté DC, Pérez C, Petroselli G, Erra-Balsells R and udisio MC. 2018. Biocontrol of *Sclerotinia sclerotiorum* (Lib.) de Bary on common bean by native lipopeptide-producer *Bacillus* strains. *Microbiological Research* 211: 21-30. doi: 10.1016/j.micres.2018.04.003.
- Stanier R. 1996. Microbiología. Segunda edición. Editorial Reverté S.A., Barcelona. 760 p.

Stamenković S, Beškoski V, Karabegović I, Lazić M and Nikolić N. 2018. Microbial fertilizers: A comprehensive review of current findings and future perspectives. *Spanish Journal of Agricultural Research* 16(1): 1-18. doi.org/10.5424/sjar/2018161-12117. doi: 10.5424/sjar/2018161-12117

Szczeczek M, Szafirowska A, Kowalczyk W, Szwejda-Grzybowska J, Włodarek A and Maciorowski R. 2016. The effect of plant growth promoting bacteria on transplants growth and lettuce yield in organic

production. *Journal of Horticultural Research* 24(2): 101-107. doi: 10.1515/johr-2016-0026

Thakur D, Kaur M and Mishra A. 2017. Isolation and screening of plant growth promoting *Bacillus* spp. and *Pseudomonas* spp. and their effect on growth, rhizospheric population and phosphorous concentration of *Aloe vera*. *Journal of Medicinal Plants Studies* 5(1): 187-192. doi: 10.1007/s12088-017-0642-8.

