



Revista Facultad Nacional de Agronomía Medellín

ISSN: 0304-2847

ISSN: 2248-7026

Facultad de Ciencias Agrarias - Universidad Nacional de Colombia

Ogidi, Clement Olusola; George, Oluwatobiloba Hannah;
Aladejana, Oluwatoyin Modupe; Malomo, Olu; Famurewa, Oladiran
Fruit preservation with bioethanol obtained from the fermentation
of brewer's spent grain with *Saccharomyces carlsbergensis*

Revista Facultad Nacional de Agronomía Medellín, vol.
73, no. 3, 2020, September-December, pp. 9321-9331
Facultad de Ciencias Agrarias - Universidad Nacional de Colombia

DOI: <https://doi.org/10.15446/rfnam.v73n3.85316>

Available in: <https://www.redalyc.org/articulo.oa?id=179964927010>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UNAM  redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Fruit preservation with bioethanol obtained from the fermentation of brewer's spent grain with *Saccharomyces carlsbergensis*

Preservación de frutas con bioetanol obtenido a partir de la fermentación de cascarilla de cebada cervecera con *Saccharomyces carlsbergensis*

doi: 10.15446/rfnam.v73n3.85316

Clement Olusola Ogidi^{1*}, Oluwatobiloba Hannah George¹, Oluwatoyin Modupe Aladejana², Olu Malomo¹ and Oladiran Famurewa²

ABSTRACT

Keywords:

Agro-wastes
Bioethanol
Fermentation
Fruit rottenness

Brewer's Spent Grain (BSG) is renewable lignocellulosic biomass generated from the beer brewing process. It serves as a substrate for various biotechnological applications. BSG was used as the main substrate for bioethanol production with *Saccharomyces carlsbergensis* in submerged fermentation. Saccharification and fermentation studies were performed for the production of bioethanol. A sterilized fermenter was loaded with 50 g L⁻¹ of BSG at 29±2 °C and an agitation speed of 180 rpm. pH was adjusted to 6.0 before the addition of 500 mL of yeast culture for 7 days under submerged and optimized conditions. The fermented product was concentrated using a rotary evaporator at 66±1 °C, and ethanol was qualitatively determined by the dichromate method. Bioethanol yield was 22%, with a specific gravity of 0.8 at 28 °C. Fourier-Transform Infrared Spectroscopy (FTIR) confirmed the presence of -CH₃^{3stretch}, -OH^{stretch} and -CH₂^{2stretch} in bioethanol. For the preservative test, *Staphylococcus* spp., *Erwinia* spp., *Lactobacillus* spp., *Bacillus* spp., *Xanthomonas* spp., *Pseudomonas* spp., *Micrococcus* spp. and *Corynebacterium* spp. were the bacteria isolated from fruits examined from different regions of Osun State. The genera of fungi isolated were *Aspergillus*, *Colletotrichum*, *Penicillium*, *Fusarium*, *Alternaria*, *Rhizopus*, *Candida*, *Saccharomyces*, *Geotrichum* and *Pichia*. Bioethanol produced from BSG inhibited the growth of microorganisms with zones of inhibition range from 7.0 mm to 11.5 mm, and thus, selected fruits were preserved. Hence, the fermentation technology of agro-industrial wastes with microorganisms can be adopted to convert waste biomass to useful resources.

RESUMEN

Palabras clave:

Agro-residuos
Bioetanol
Fermentación
Podredumbre de frutos

Cascarilla de cebada cervecera (CCC) es una biomasa lignocelulósica renovable generada a partir del proceso de elaboración de la cerveza, que sirve como sustrato para diversas aplicaciones biotecnológicas. Se usó CCC como sustrato principal para la producción de bioetanol con *Saccharomyces carlsbergensis* en fermentación sumergida. Se realizaron estudios de sacarificación y fermentación para la producción de bioetanol, el fermentador esterilizado se cargó con 50 g L⁻¹ de CCC a 29±2 °C y una velocidad de agitación de 180 rpm. El pH se ajustó a 6,0 antes de la adición de 500 mL de cultivo de levadura durante 7 días en condiciones sumergidas y optimizadas. El producto fermentado se concentró usando un evaporador rotatorio a 66±1 °C y el etanol se determinó cualitativamente por el método de dicromato. El rendimiento de bioetanol fue del 22% con un peso específico de 0,8 a 28 °C. La Espectroscopía Infrarroja por Transformada de Fourier (FTIR) confirmó la presencia de CH₃^{3stretch}, OH y CH₂^{2stretch} en el bioetanol. Para el ensayo de preservación, *Staphylococcus* spp., *Erwinia* spp., *Lactobacillus* spp., *Bacillus* spp., *Xanthomonas* spp., *Pseudomonas* spp., *Micrococcus* spp. y *Corynebacterium* spp. fueron bacterias aisladas de frutas examinadas de diferentes regiones del estado de Osun. Los géneros de hongos aislados fueron *Aspergillus*, *Colletotrichum*, *Penicillium*, *Fusarium*, *Alternaria*, *Rhizopus*, *Candida*, *Saccharomyces*, *Geotrichum* y *Pichia*. El bioetanol producido a partir de CCC inhibió el crecimiento de microorganismos con zonas de inhibición comprendidas entre 7,0 mm y 11,5 mm conservando las frutas seleccionadas. Por lo tanto, se puede adoptar la tecnología de fermentación de desechos agroindustriales con microorganismos para convertir la biomasa residual en recursos útiles.

¹ Biotechnology Unit, Department of Biological Sciences, Kings University, PMB 555, Odeomu, Nigeria.

² Microbiology Unit, Department of Biological Sciences, Kings University, PMB 555, Odeomu, Nigeria.

* Corresponding author: <clementogidi@yahoo.com>

Wheat bran, rice bran, corn cob and wheat straw are examples of agricultural wastes available as carbon sources (Singh *et al.*, 2012). Brewer's Spent Grain (BSG), a waste-product of the mashing process, is one of the initial operations in brewery industries to solubilize malt and grains to ensure adequate extraction of wort (Aliyu and Bala, 2011). BSG is available in larger quantities; approximately 85-90% of the total by-products generated during beer production; however, its central exploitation or disposition has been inadequate (Steiner *et al.*, 2015). Several attempts have been made to use BSG in biotechnological processes, and this is achievable in various researches and industries by adopting solid-state fermentation (SFF) or submerged fermentation (SMF) since BSG contains basic nutrients required for microbial growth (Mussatto, 2014). BSG is a lignocellulosic material with 17% cellulose, 28% non-cellulosic polysaccharides, minerals, vitamins, proteins, amino acids, arabinoxylans, 28% lignin (Ivanova *et al.*, 2017). Therefore, it has a perspective to be recycled and to become useful products. BSG has been utilized as a new and economical medium for the cultivation of microorganisms (Tan *et al.*, 2020).

Agricultural residues are currently utilized for the production of bioethanol to decrease total dependence on forest woody biomass and continuous deforestation. Bioethanol is produced from different agro-wastes using some biotechnological methods such as fermentation with diverse microorganisms (Bušić *et al.*, 2018). The use of carbon sources from renewable biomass is economical for the full exploitation of less expensive sources into the production of beneficial products (Saini *et al.*, 2015). Hence, agro-industrial residues are an attractive alternative to costly raw materials. Bioethanol has been produced by converting sugars directly from BSG or indirectly through starch into alcohol via fermentation followed by distillation (Azhar *et al.*, 2017). Ethanol produced from lignocellulosic biomass raises a global interest because it represents an excellent alternative to petroleum-derived energies and reduces food versus fuel conflict generated by first-generation ethanol (Awoyale and Lokhat, 2019; Prasad *et al.*, 2019).

Ethanol is an excellent preservative agent that protects several surface fruits (external morphology) from

microbial colonization (Dao and Dantigny, 2011). Hitherto, preservation of fresh fruits and vegetables is among the challenges of food products for commercial producers and distributors, particularly in middle-income, low or poor resource countries. Although many fresh fruits are in ideal conditions to hinder microorganisms from colonizing their integument, a lot of challenges in terms of post-harvest, storage, preservation of fruits or vegetables are rising up. It has generated the search of natural preservatives from plants or agro-wastes as an alternative and safer choice since they displayed little or no side effects (Sagar *et al.*, 2018; Saeed *et al.*, 2019). The bioethanol produced from agro-waste using *S. carlsbergensis* in submerged fermentation can be used as a preservative agent, which will not only help to reduce wastes in the environment but will preserve fruits or crops from post-harvest spoilage. Hence, this study aimed to produce bioethanol using BSG as a substrate with *S. carlsbergensis* in submerged fermentation and to assess the preservative potential of bioethanol on some selected fruits.

MATERIALS AND METHODS

Collection of brewer's spent grain and brewer's spent yeast

The brewer's spent grain and brewer's spent yeast were obtained from International Breweries Plc., Ilesha in Osun state, Nigeria. The town is located at longitude 7.6395°N and latitude 4.7588°E.

Collection of fruits from various locations in Osun State

Various types of fruits were collected from different locations in Osun State. This State was grouped into four (4) zones, A: Odeomu/Gbongan axis, B: Ife and its environment, C: Osogbo, and D: Ilesha and its environment. A total of 161 different fruits were collected as pineapple (*Ananas comosus*), orange (*Citrus sinensis*), African star apple (*Chrysophyllum albidum*), tomato (*Solanum lycopersicum*), banana (*Musa acuminata*), lime (*Citrus aurantiifolia*), pawpaw (*Carica papaya*), sour-sop (*Annona muricata*), watermelon (*Citrullus lanatus*), apple (*Malus domestica*), plantain (*Musa paradisiaca*) and almond (*Prunus dulcis*).

Source of *S. carlsbergensis*

S. carlsbergensis was isolated from brewer's spent yeast

using serial dilution method. A loop full from 10^{-6} was cultured on Potato Dextrose Agar (PDA) and incubated at 25 °C for 48 h. The yeast was subcultured into another freshly prepared PDA and incubated at 25 °C for 48 h to get pure isolate of *S. carlsbergensis*. The pure isolate was transferred to yeast broth and incubated for 48 h.

Production of bioethanol from BSG

The method, according to Alam *et al.* (2009), was adopted for the production of bioethanol with a slight modification. The fermenter was sterilized using 3% v/v of hypochlorite. The sterile fermenter was loaded with 50 g L⁻¹ of BSG at 29±2 °C and an agitation speed of 180 rpm. The pH was adjusted to 6.0 before the addition of 500 mL of yeast culture; the fermentation lasted 7 days under submerged and optimized conditions. The aeration and pH were kept stable during fermentation. The fermented product was concentrated using a rotatory evaporator at 66±1 °C, and ethanol was qualitatively determined by the dichromate method.

FTIR spectroscopic of bioethanol from BSG

Structural analysis of the functional group in bioethanol was determined using FT-IR spectroscopy (8400S, Shimadzu Scientific Instruments Inc.). Briefly, bioethanol (1.0 µL) was placed on a fused KBr disc. This was placed on the cell holder, clamped loosely and fixed on the infrared (IR) beam. The running was done at 400 to 4000 per cm wavenumber.

Isolation and identification of microorganisms from fruits

The surface of fruit was sterilized with 1% v/v hypochlorite and rinsed with sterile distilled water to remove normal flora of fruit and other possible microbial contaminants. Fruit samples were observed for 4-5 days to check any sign of spoilage. The rotten part of fruit was aseptically cut and transferred into sterilized peptone water. The sample was shaken vigorously and then allowed to stand for 30 min (Ajayi-Moses *et al.*, 2019). Serial dilution was carried out up to 10^{-4} and 10^{-5} dilution factor. An aliquot of 0.1 mL was aseptically transferred into Petri dish, and molten nutrient agar or PDA was then introduced. The plate solidified at room temperature (29±1 °C), then plates were incubated at 37 °C for 24 h and 2-3 days at 25 °C for bacteria and fungi, respectively. Discrete colonies were counted and recorded as colony-forming

unit per gram (CFU g⁻¹) for bacteria and spore-forming unit per gram (SFU g⁻¹) for fungi. A pure colony was obtained by subcultured, and isolates were tentatively grouped according to their morphological, cultural, and staining characteristics. Biochemical tests such as the catalase test, production of hydrogen sulfide (H₂S), indole, urease, methyl red, oxidase, coagulase, motility, methyl red, Voges-Proskauer, starch hydrolysis and sugars fermentation were carried out using the methods described by Olutiola *et al.* (2000). The results of the biochemical test were compared to Bergey's Manual of Systematic Bacteriology (Krieg *et al.*, 2010). Fungi isolates were identified using cultural and microscopic observations, according to Barnett *et al.* (2000) and Samson *et al.* (2010).

In vitro antimicrobial activity of bioethanol and other preservatives against microorganisms

The antimicrobial activity of the bioethanol against spoilage microorganisms isolated from fruits was performed using agar well diffusion (CLSI, 2014). Suspension of test microorganisms was adjusted by the spectrophotometer to 0.5 McFarland standard. Sterile cotton swabs were dipped in the microbial suspension and spread on the surface of the agar plate. A sterilized cork borer was used for cutting wells in each plate. Bioethanol (50 µL) was introduced and sorbic acid (5.0 mg mL⁻¹), ampiclox (5.0 mg mL⁻¹) and terbinafine (5.0 mg mL⁻¹) were implemented as positive controls against bacteria and fungi, while sterile distilled water was used as the negative control. All the plates were labeled appropriately and incubated at 37 °C during 24 h for bacteria and at 25 °C for 48 h for fungi. The diameter of zones of inhibition around wells was measured in millimeter (mm).

Preservation of fruits using bioethanol from BSG

Most prevalent as well as colonizing microorganisms (1.0×10^5 CFU mL⁻¹ or SFU mL⁻¹) were re-introduced to apparently healthy (absence of diseases, no wound symptoms or lesion) selected fruits (orange, watermelon, pineapple, and tomato). Each fruit was discretely remained in a sterile laminar hood without touching each other. After 24 h, bioethanol (20 mL) was applied on the surface fruits of the group A. The bioethanol was adjusted to 40% v/v since Kalathenos and Russell (2003) revealed that <30% v/v were rarely

biocidal. Ethanol $\geq 70\%$ v/v could cause damage to the fruit integument. Group B was treated with sorbic acid (5% w/v), group C was un-inoculated fruits with microorganism and group D was fruits inoculated with microorganisms but untreated with either bioethanol or sorbic acid. The fruit samples were observed for 7 days at room temperature (29 °C).

Statistical analysis

Data were presented as mean \pm standard deviation from three repetitions. Data obtained in this study were subjected to One-way Analysis of Variance (ANOVA) using Statistical Package for Social Sciences (SPSS) version 20 (USA). For bacteria and fungi count, the mean values were compared by Duncan's new multiple range test (MRT). Differences were considered significant at $P < 0.05$.

RESULTS AND DISCUSSION

Yield and physicochemical parameters of bioethanol from BSG with *S. cerevisiae*

Table 1 shows the physicochemical parameters of bioethanol produced from BSG. The yield obtained was 22%, with a specific gravity of 0.8. These results contrast with the findings of Irfan *et al.* (2014). They reported bioethanol from sugarcane bagasse, rice straw and wheat straw by *S. cerevisiae* with a value of 77 g L⁻¹, 62 g L⁻¹, and 44 g L⁻¹, respectively. Ingale *et al.* (2014) obtained

ethanol of 17.1 g L⁻¹ with a higher yield of 84% from banana pseudostem fermented by *S. cerevisiae* NCIM 3570. The higher proportion of ethanol in their study could be associated with two fungal strains of *Aspergillus* spp., which facilitated the maximal release of sugars to produce more ethanol. The combination of *Aspergillus oryzae* and *S. cerevisiae* NCYC479 produced the highest concentrations of ethanol (37 g L⁻¹) in 10 days from BSG using consolidated bioprocessing (Wilkinson *et al.*, 2017). Moodley and Gueguim Kana (2019) optimized pretreatment techniques to produce 25% more bioethanol from sugarcane leaf waste using *S. cerevisiae* BY4743. *S. cerevisiae* is one of the best yeast widely employed for commercial production of bioethanol. *S. cerevisiae* has been attractive for efficient consolidated bioprocessing and several biotechnological purposes because of its novel amylolytic enzyme combination, relatively high tolerance to osmotic stress and anaerobic conditions. Therefore, it is suitable for large-scale fermentation of agro-wastes into bioethanol (Cripwell *et al.*, 2019). In the study of Wu *et al.* (2020), replacement of ethanol fermentation-associated regulatory gene in *S. cerevisiae* was reported to enhance ethanol production by a 5.30% increase in yield and 12.5% decrease in fermentation time when compared to the original strain. Another method that could increase the yield of ethanol is mixed substrates. Bolade *et al.* (2019) claimed that multi-substrates biomass of agro wastes increases the yield of bioethanol.

Table 1. Physicochemical properties of bioethanol produced from BSG.

Test	Obtained bioethanol	Ethanol 70% v/v (standard)
Yield (%)	22.0 \pm 0.01	-
Specific gravity	0.80 \pm 0.00	0.79 \pm 0.01
Moisture (%)	6.80 \pm 0.00	0.50 \pm 0.00
Flammability	Weak	High

-: yield was not quantified; it was used as control.

BSG is interesting biomass with hydrolyzable fermentable sugars that can be converted to ethanol with different microorganisms through co-fermentation strategies (Rojas-Chamorro *et al.*, 2020). Likewise, cassava peels, potato peels and millet husks with different microbial inoculants such as *S. cerevisiae*, *Rhizopus nigricans*, *Aspergillus niger*, *Spirogyra africana* showed great potential for bioethanol production (Chibuzor *et al.*, 2016).

The large proportion of lignocellulosic materials such as corncob, cornstalk, cornhusk, sugarcane bagasse and sugarcane bark that are creating environmental pollution, can be easily degraded by microorganisms and thus, serve as a substrate for renewable resources (bioethanol).

Figure 1 shows the transmittance and peak representing functional groups in the bioethanol. Table 2 shows

various functional groups found in bioethanol, hydroxyl ($\text{OH}_{\text{stretch}}$), methyl ($\text{CH}_3_{\text{stretch}}$), and alkane ($\text{CH}_2_{\text{stretch}}$). These functional groups in ethanol (alcohol) give it the biocide property, and it is responsible for the antiseptics, disinfection and can be used as a preservative agent (McDonnell and Russell, 1999). Hydroxyl group in

ethanol (alcohol) acts as an antimicrobial agent against microbes except for alcohol-tolerant strains. It causes a partial breakdown of membrane function, inhibiting cell growth or protein synthesis, denaturation of proteins, and membrane damage, which lead to cell perturbations like ion leakage or loss of energy (Horinouchi *et al.*, 2018).




Figure 1. FTIR spectrum of bioethanol produced from BSG

Table 2. The functional group identified in bioethanol produced from BSG.

Sample	Wavenumber (cm^{-1})	Functional group	Name of group
Bioethanol	3444.98	$-\text{OH}_{\text{stretch}}$	Hydroxyl
	2621.36	$-\text{CH}_3_{\text{stretch}}$	Methyl
	1539.45	$-\text{CH}_2_{\text{stretch}}$	Alkane

Total microbial load and occurrence (%) of microorganisms isolated from selected fruits

The colonies counted from each fruit are shown in Table 3. The highest colonies of bacteria count ($1.05 \times 10^5 \text{ CFU g}^{-1}$) were recorded for bananas from Odeomu (A) market ($P < 0.05$). Pineapple from Ilesha had the highest fungal count ($1.96 \times 10^6 \text{ SFU g}^{-1}$). Watermelon from Odeomu (A) had the least bacterial load ($P < 0.05$) of $1.0 \times 10^4 \text{ CFU g}^{-1}$, and African star apple collected from Osogbo (C) had the least fungal count of $1.1 \times 10^5 \text{ SFU g}^{-1}$. Ajayi-Moses *et al.* (2019) reported the highest bacterial count of $5.84 \times 10^5 \text{ CFU g}^{-1}$ for tomatoes and African star apple with the highest fungal count of $3.04 \times 10^5 \text{ SFU g}^{-1}$. Evaluation of microorganisms associated with fruits revealed that

various bacteria and fungi in high densities could spoil fruits. Spoilage microorganisms can also colonize, enter and penetrate plant tissues at fruit development, either through calyx, stem or various specialized water and gas exchange structures of leafy matter. However, successful establishment requires the spoilage microbe to overcome multiple natural protective barriers and other factors (Erkmen and Bozoglu, 2016). Lime and soursop did not show any sign of spoilage, and no microorganisms were isolated (Table 3). Fruits like lime and soursop with higher pKa do not get spoiled but rather get dehydrated, exiting fruit juice, a process attributed to spoilage organisms affecting fruits (Czajkowski *et al.*, 2011).

Table 3. Bacteria and fungi count from fruits obtained from different regions in Osun State.

Fruits	Locations							
	A	B	C	D	A	B	C	D
	Bacteria $\times 10^4$ CFU g ⁻¹				Fungi $\times 10^5$ SFU g ⁻¹			
Pineapple	5.2 \pm 0.0 c	NC	8.5 \pm 0.0 d	3.3 \pm 0.0 b	7.5 \pm 0.0 b	NC	9.0 \pm 1.0 d	19.6 \pm 2.1 c
Orange	8.4 \pm 0.0 d	6.4 \pm 0.3 c	5.3 \pm 0.1 c	-	7.0 \pm 0.1 b	6.6 \pm 0.3 a	7.0 \pm 0.4 c	NC
African star apple	8.4 \pm 0.2 d	NC	5.6 \pm 0.0 c	NC	7.5 \pm 0.2 b	0.0	1.1 \pm 0.0 a	-
Tomato	7.2 \pm 0.4 d	4.7 \pm 0.1 a	5.2 \pm 0.0 c	1.1 \pm 0.0 a	5.5 \pm 0.4 a	7.0 \pm 0.1 a	8.0 \pm 0.3 c	6.5 \pm 0.0 b
Banana	10.5 \pm 1.0 e	0.0	4.2 \pm 0.0 b	1.3 \pm 0.0 a	8.0 \pm 1.0 c	NC	7.5 \pm 0.0 c	2.2 \pm 0.1 a
Lime	-	-	-	-	-	-	-	-
Pawpaw	3.0 \pm 0.0 b	7.7 \pm 0.2 c	1.0 \pm 0.0 a	1.0 \pm 0.0 a	9.0 \pm 0.0 c	6.5 \pm 0.0 a	5.0 \pm 0.0 c	6.2 \pm 0.0 b
Sour sop	-	-	-	-	-	-	-	-
Watermelon	1.0 \pm 0.0 a	5.2 \pm 0.2 b	4.9 \pm 0.0 b	NC	-	7.0 \pm 0.0 b	8.0 \pm 0.0 c	NC
Almond	NC	5.3 \pm 0.0 b	NC	NC	NC	8.0 \pm 0.0 b	NC	NC
Apple	-	-	-	-	-	-	-	-
Plantain	NC	NC	-	1.4 \pm 0.0 a	NC	NC	-	1.7 \pm 0.0 a

Values with different letters along the same column are significantly different ($P < 0.05$).

A: Odeomu/Gbongon axis, B: Ife, C: Osogbo and D: Ilesha.

-: No microbial growth; NC: Fruit samples were not collected.

Table 4 shows the percentage of the occurrence of bacteria. The highest percentage of occurrence (25.6%) was obtained for *Corynebacterium* sp. followed by *Lactobacillus* sp. with 17.9%. The lowest bacteria percentage occurrence (5.1%) was obtained for species of *Micrococcus* and *Staphylococcus*. *Erwinia* spp. were found to be associated with spoilage of pineapple, African star apple, tomato, watermelon and almond. The bacterium *Erwinia carotovora* subsp. *carotovora* is a highly effective spoilage microorganism that produces an increasing amount of pectolytic enzymes to degrade fruit tissues. It causes soft rot on fruits like oranges, tomatoes,

banana, pineapple, and watermelon (Barth et al., 2009; Sharma et al., 2013). Besides, *E. carotovora*, several *Pseudomonas* spp., *Corynebacterium*, *Xanthomonas campestris* and lactic acid bacteria are important spoilage bacteria of fruits (Tournas, 2005; Erkmen and Bozoglu, 2016). Some spoilage microbes are capable of colonizing, creating lesions and damaged healthy plant tissues. The type of microbial spoilage in fruits is based on the pH, nutrient availability, water activity (a_w), temperature, relative humidity, oxidation-reduction potential and content of biological structure of fruits (Erkmen and Bozoglu, 2016).

Table 4. Percentage of occurrence of bacteria isolated from different locations

Bacterial isolates	Pa	O	Asa	T	B	L	P	Wm	Ss	Al	PI	Ap	N	%
<i>Corynebacterium</i> sp.	+	+	+	+	+	-	+	+	-	+	+	-	10	25.6
<i>Lactobacillus</i> sp.	+	+	+	+	+	-	-	-	-	+	+	-	7	17.9
<i>Bacillus</i> sp.	-	-	+	-	+	-	+	-	-	+	+	-	5	12.8
<i>Xanthomonas</i> sp.	+	-	-	+	+	-	-	+	-	+	-	-	5	12.8
<i>Erwinia</i> sp.	+	-	+	+	-	-	-	+	-	+	-	-	5	12.8
<i>Pseudomonas</i> sp.	+	-	-	+	+	-	-	-	-	-	-	-	3	7.7
<i>Micrococcus</i> sp.	+	-	+	-	-	-	-	-	-	-	-	-	2	5.1
<i>Staphylococcus</i> sp.	+	-	-	-	-	-	-	-	-	-	+	-	2	5.1

Pa: Pineapple; O: Orange; Asa: African star apple; T: Tomato; B: Banana; L: Lime; P: Pawpaw; Wm: Watermelon; Ss: Sour sop; Al:

Almond; PI: Plantain; Ap: Apple; N: number of isolates.

+: presence of bacteria; -: absence of bacteria.

Fungi with the highest percentage of occurrence was *Aspergillus niger* (13.9%), followed by *Colletotrichum* sp. and *Penicillium digitatum* with the same value of 11.1%. *Pichia* sp. has the lowest percentage of occurrence of 4.2% (Table 5). The highest occurrence of *A. niger* from examined fruits in Osun State is in concordance with findings of Mailafia *et al.* (2017). Researchers revealed that *Aspergillus* spp. had the highest occurrence in fruits like pineapple, watermelon, oranges, pawpaw, and tomatoes with a frequency of 38%, followed by *Fusarium avenaceum* with occurrence of 31% in pineapple, watermelon, oranges, pawpaw and tomatoes. In comparison, *P. digitatum* and *R. stolonifera* have the least frequency at the same value of 4% for tomato and orange. Other fungal species

identified as agents of spoilage were *Saccharomyces* spp. (10%), *F. solani* (8%), and *A. flavus* (5%). Some of the fungi isolated from fruits were species of *Penicillium*, *Aspergillus*, *Rhizopus*, *Fusarium* and *Mucor iriformis*. Tafinta *et al.* (2013) and Ajayi-Moses *et al.* (2019) isolated similar microorganisms with varying prevalence from banana, pawpaw, orange, tomato, apple, pineapple, watermelon, cucumber, and African star apple. Some of these fungal isolates are known to be pathogenic due to the toxic secondary metabolites produced. *Penicillium expansum* and *Botrytis cinerea* are pathogenic spoilage microorganisms, which cause blue-rot, grey mold in African star apple, cherry, apple, tomato, pears and kiwi fruit called sour sop (Miedes and Lorences, 2004).

Table 5. Percentage of occurrence of fungi isolated from different locations

Fungi isolates	Pa	O	Asa	T	B	L	P	Wm	Ss	Al	Pl	Ap	N	%
<i>Aspergillus niger</i>	+	+	+	+	+	-	+	-	+	+	+	-	10	13.9
<i>Colletotrichum</i> sp.	+	-	+	+	+	-	-	-	+	+	+	-	8	11.1
<i>Penicillium digitatum</i>	-	+	+	+	+	-	+	-	-	-	+	-	8	11.1
<i>Saccharomyces</i> sp.	+	+	+	+	-	-	-	-	-	-	-	-	6	8.3
<i>Penicillium italicum</i>	+	-	-	-	+	-	+	-	-	+	+	-	6	8.3
<i>Candida tropicalis</i>	+	+	+	+	+	-	-	-	-	+	-	+	6	8.3
<i>Geotrichum</i> sp.	+	+	-	+	+	-	+	-	-	-	+	-	6	8.3
<i>Fusarium</i> sp.	+	+	-	+	+	-	-	-	-	-	-	-	5	6.9
<i>Alternaria</i> sp.	+	-	-	+	+	-	-	-	-	-	-	-	4	5.6
<i>Rhizopus</i> sp.	-	-	-	+	+	-	+	-	-	-	+	-	4	5.6
<i>Pichia</i> sp.	-	-	-	-	-	-	+	-	-	-	-	-	3	4.2

Pa: Pineapple; O: Orange; Asa: African star apple; T: Tomato; B: Banana; L: Lime; P: Pawpaw; Wm: Watermelon; Ss: Soursop; Al: Almond; Pl: Plantain; Ap: Apple; N: number of isolates.

+: presence of fungi; -: absence were absent

Likewise, molds of genera *Rhizopus*, *Alternaria* and *Botrytis* produce acidic compounds that cause fruit and vegetable rot with distorted color, texture and or taste (Tournas, 2005). Fernández-Cruz *et al.* (2010), Lewis and Goodrich-Schneider (2012) revealed different species of fungi that produce mycotoxins (aflatoxins, ochratoxin, patulin, fumonisin, alternariol, alternariol methyl ether, and altenuene) in fruits. The problem associated with mycotoxins in fruits includes economic loss, poor organoleptic properties, toxicities (acute to chronic), and a spectrum of effect (mild to severe), including carcinogenicity and death. Spoilage microorganisms exploit fruit components using their extracellular lytic enzymes, pectinases and hemicellulases to degrade fruit

polymers to release intracellular constituents as nutrients for growth (Kalia and Gupta, 2006). Microorganisms from fruits secreted a wide variety of enzymes. Isolation and identification of novel strains of microorganism from fruits can serve as natural origin, a safer and cheaper alternative source of microbial enzymes as promising candidates for biotechnological uses and medical processes (Sharma *et al.*, 2013; Garg *et al.*, 2016).

Antimicrobial and preservative properties of bioethanol from BSG

To minimize wastage recorded on fruits and vegetables and to reduce economic losses associated with microbial spoilage, a reliable and supportive measure

needs to be sourced. Bioethanol from BSG displayed zones of inhibition against bacteria with values ranging from 7.0 to 8.8 mm, sorbic acid was within 6.0 to 13.0 mm, and ampiclox was from 8.0 to 18.5 mm (Table 6). Zones of inhibition by bioethanol against fungi ranged from 7.5 to 11.5 mm, while sorbic acid was 7.5 to 13.0 mm and 10.0 to 17.0 mm for terbinafine (Table 6). Ethanol is used as a disinfectant and acts against microorganisms in two different ways: growth inhibition (bacteriostasis, fungistasis) or lethal action (bactericidal,

fungicidal or virucidal effects) (Maris, 1995). Hence, bioethanol can be used to prevent the growth of spoilage microorganisms on fruits. Ethanol interacts with cell surface followed by penetration into cells and acts on the target site(s) of microorganisms (Maris, 1995). Inactivation of fungal spores, suppression of fungal growth or their germination on fruits by ethanol reflects the inhibitory potential of bioethanol in controlling fruit decaying by fungi and extending the shelf-life of food products (Dao and Dantigny, 2011).

Table 6. Zones of inhibition (mm) displayed by bioethanol and selective preservatives against *microorganisms isolated from fruits.

Bacteria isolates	Bioethanol	Sorbic acid	Ampiclox
<i>Corynebacterium</i> sp.	8.0±0.0	10.0±0.2	17.1±2.2
<i>Lactobacillus</i> sp.	8.0±0.1	12.0±0.5	14.5±1.7
<i>Bacillus</i> sp.	8.8±0.0	11.0±0.0	18.0±1.0
<i>Xanthomonas</i> sp.	-	13.0±0.3	16.7±2.1
<i>Erwinia</i> spp.	-	12.0±0.8	18.5±0.8
<i>Pseudomonas</i> sp.	7.0±0.0	-	17.2±1.1
<i>Micrococcus</i> sp.	-	-	8.0±0.0
<i>Staphylococcus</i> sp.	-	6.0±0.0	8.0±0.0
Fungal isolates	Bioethanol	Sorbic acid	Terbinafine
<i>Aspergillus niger</i>	10.5±1.0	11.0±1.0	16.2±3.0
<i>Colletotrichum</i> sp.	7.5±0.0	13.0±1.7	17.0±2.3
<i>Penicillium italicum</i>	10.5±0.4	7.5±0.1	11.0±1.0
<i>Saccharomyces</i> sp.	11.5±0.8	8.6±0.3	16.0±1.0
<i>Penicillium digitatum</i>	8.0±0.0	10.3±1.2	10.0±0.0





















-: There were no zones of inhibition

Microorganisms able to colonize fruits with spoilage attributes were used as indicator microorganisms.

Table 7 shows different fruits preserved with bioethanol, sorbic acid, un-inoculated fruit with the microorganism, spoiled fruit and open spoiled fruit after 7 days. Dao and Dantigny (2011), found that oranges were protected from fungal infection for 30 days when exposed to 20-100% ethanol. The use of bioethanol as a preservative is important to prevent a higher loss of fruits since it has no side effects on humans. It will proffer the solution to longstanding spoilage of vegetables and fruits, which had been associated with different microorganisms. Ethanol exerts its most effective bactericidal action at ≥40 to 95% v/v against vegetative cells like chemical disinfectants (Kalathenos and Russell, 2003). Findings of Katsinis *et al.* (2008) established effective preservation

with the synergistic effect of conventional chemical preservative (potassium sorbate) and ethanol, which improved the shelf life of bread by suppressing microbial growth (43.5% and 38.5% mold-free shelf life). Bioethanol displayed a higher zone of inhibition (11.5 mm) against *Saccharomyces* sp. than sorbic acid (8.6 mm). Sorbic acids had less or no inhibitory activity against *S. cerevisiae*, *Saccharomycodes ludwigii*, *Zygosaccharomyces bailii*, *Zygosaccharomyces rouxii*, *Schizosaccharomyces pombe* *Brettanomyces* spp., *Pichia membranifaciens*, *Dekkerae* spp, and *Issatchenkia orientalis* at 800 mg kg⁻¹, which is much higher than the permissible limit in foods, but ethanol (>20% v/v) exerts a strong lethal effect on these molds (Loureiro and Malfeito-Ferreira, 2003).

Table 7. Evolution of fruits during 7-day treatment with bioethanol and sorbic acid.

Fruit	Bioethanol	Sorbic acid	Uninoculated	Spoilt fruit	Open spoilt fruit
Orange					
Watermelon					
Pineapple					
Tomato					

The mechanisms of resistance by yeasts to weak organic acids are by inducing the expression of H⁺ ATPases to regulate their cytosolic pH, using their plasma membrane components to modulate the influx of lipophilic weak organic acids (Ullah *et al.*, 2012). ATP binding cassette (ABC) transporter (Pdr12) prevents anion accumulation or degrade sorbic acid to 1,3-pentadiene (Casas *et al.*, 2004). Findings of Linares-Morales *et al.* (2018) revealed that chemical preservatives like organic acids in fruits and vegetables cause disruption of membrane permeability, reduce cell's internal pH, affects metabolic enzymes as well as protein synthesis. However, a microbial product like bioethanol can be encouraged in use as preservative agent since it strongly suppresses microbial activity

on fruits. Microbial by-products are bio-protective or natural preservatives, and thus, exhibit antifungal activities. There is a growing interest in alternatives to preservation other than chemical or synthetic agents with different side effects (Leyva Salas *et al.*, 2017). The microbial growth inhibition or killing action of bioethanol on microorganisms is an indication that; bioethanol produced by *S. cerevisiae* from renewable biomass (lignocellulosic wastes) is needed for commercial purposes.

CONCLUSION

The bioethanol produced from the fermentation of brewer's spent grain with *S. carlsbergensis* inhibited microorganisms associated with post-harvest spoilage

of fruits. The presence of functional groups in bioethanol contributed to its bioactivity, which makes it a potential preservative agent to suppress the colonization of microbial spoilage on fruits, reaching inhibitions of 8.8 and 11.5 mm for bacteria and fungi, respectively. This study proffer solution to the underutilized BSG residue, which can be used for bioethanol and serve as a preservative agent for fruits or vegetables. However, modern facilities to increase the yield of bioethanol from agro wastes need to be considered in subsequent works.

REFERENCES

- Ajayi-Moses OB, Ogidi CO and Akinyele BJ. 2019. Bioactivity of *Citrus* essential oils (CEOs) against microorganisms associated with spoilage of some fruits. *Chemical and Biological in Technologies Agricultures* 6(22): 15. doi: 10.1186/s40538-019-0160-5
- Alam MZ, Kabbashi NA and Hussi SNIS. 2009. Production of bioethanol by direct bioconversion of oil-palm industrial effluent in a stirred-tank bioreactor. *Journal of Industrial Microbiology and Biotechnology* 36(6): 801-808. doi: 10.1007/s10295-009-0554-7
- Aliyu S and Bala M. 2011. Brewer's spent grain: a review of its potentials and applications. *African Journal of Biotechnology* 10(3): 324-331. doi: 10.5897/AJB10.006
- Azhar SHM, Abdulla R, Jambo SA, Marbawi H, Gansau JA, Faik AAM and Rodrigues KF. 2017. Yeasts in sustainable bioethanol production: a review. *Biochemistry and Biophysics Reports* 10: 52-61. doi: 10.1016/j.bbrep.2017.03.003
- Awoyale AA and Lokhat D. 2019. Harnessing the potential of bio-ethanol production from lignocellulosic biomass in Nigeria – a review. *Biofuels, Bioproducts and Biorefining* 13(1): 192-207. doi: 10.1002/bbb.1943
- Barnett JA, Payne RW and Yarrow O. 2000. *Yeast: Characteristics and identification*. Third Edition. Cambridge University Press, London. pp 12-87.
- Barth M, Hankinson TR, Zhuang H and Breidt F. 2009. Microbiological spoilage of fruits and vegetables pp. 135-183. In: Sperber WH and Doyle MP (eds.). *Compendium of the microbiological spoilage of foods and beverages: Food Microbiology and Food Safety*. Springer, New York. doi: 10.1007/978-1-4419-0826-1_6
- Bolade DO, Ana GREE, Lateef SA and Sokan-Adeaga AA. 2019. Exploration of the Bioethanol Yield of Single and Multi-Substrate Biomass from Cassava Processing Wastes. *The Journal of Solid Waste Technology and Management* 45(3): 305-314. doi: 10.5276/JSWTM/2019.305
- Bušić A, Mardetko N, Kundas S, Morzak G, Belskaya H, Ivančić Šantek M, Komes D, Novak S and Šantek B. 2018. Bioethanol production from renewable raw materials and its separation and purification: a review. *Food Technology and Biotechnology* 56(3): 289-311. doi: 10.17113/ftb.56.03.18.5546
- Casas E, de Ancos B, Valderrama MJ, Cano P and Peinado JM. 2004. Pentadiene production from potassium sorbate by osmotolerant yeasts. *International Journal of Food Microbiology* 94(1): 93-96. doi: 10.1016/j.ijfoodmicro.2004.01.001
- Chibuzor O, Uyoh EA and Igile G. 2016. Bioethanol production from cassava peels using different microbial inoculants. *African Journal of Biotechnology* 15(30): 1608-1612. doi: 10.5897/AJB2016.15391
- CLSI - Clinical and Laboratory Standards Institute. 2014. Performance standards for antimicrobial susceptibility testing; twenty-fourth informational supplement. M100 – S24. Wayne, USA. pp. 1-16.
- Cripwell RA, Rose SH, Favaro L and van Zyl W. 2019. Construction of industrial *Saccharomyces cerevisiae* strains for the efficient consolidated bioprocessing of raw starch. *Biotechnology for Biofuels* 12:201. doi: 10.1186/s13068-019-1541-5
- Czajkowski MC, Perombelon JA, Veen JM and Van der W. 2011. Control of blackleg and tuber spoilage of potatoes caused by *Pectobacterium* and *Dickeya* species: a review. *Plant Pathology* 60(6): 999-1013. doi: 10.1111/j.1365-3059.2011.02470.x
- Dao T and Dantigny P. 2011. Control of food spoilage fungi by ethanol. *Food Control* 22(3-4): 360-368. doi: 10.1016/j.foodcont.2010.09.019
- Erkmen O and Bozoglu TF. 2016. Chapter 20: Spoilage of vegetables and fruits. pp 337-363. In: *Food microbiology: principles into practice*. doi: 10.1002/9781119237860.ch20
- Fernández-Cruz ML, Mansilla ML and Tadeo JL. 2010. Mycotoxins in fruits and their processed products: analysis, occurrence and health implications. *Journal of Advanced Research* 1(2): 113-122. doi: 10.1016/j.jare.2010.03.002
- Garg G, Singh A, Kaur A, Singh R, Kaur J and Mahajan R. 2016. Microbial pectinases: an ecofriendly tool of nature for industries. 3 *Biotech* 6: 47. doi: 10.1007/s13205-016-0371-4
- Horinouchi T, Maeda T and Furusawa C. 2018. Understanding and engineering alcohol-tolerant bacteria using OMICS technology. *World Journal of Microbiology and Biotechnology* 34: 157. doi: 10.1007/s11274-018-2542-4
- Ingale S, Joshi SJ and Gupte A. 2014. Production of bioethanol using agricultural waste: banana pseudo stem. *Brazilian Journal of Microbiology* 45(3): 885-892. doi: 10.1590/S1517-83822014000300018
- Irfan M, Nadeem M and Syed Q. 2014. Ethanol production from agricultural wastes using *Saccharomyces cerevisiae*. *Brazilian Journal of Microbiology* 45(2): 457-465. doi: 10.1590/s1517-83822014000200012
- Ivanova K, Denkova R, Kostov G, Petrova T, Bakalov I, Ruscova M and Penov N. 2017. Extrusion of brewers' spent grains and application in the production of functional food. Characteristics of spent grains and optimization of extrusion. *Journal of The Institute of Brewing* 123(4): 544-552. doi: 10.1002/jib.448
- Kalathenos P and Russell NJ. 2003. Ethanol as a food preservative. pp 196-217. In: Russell NJ and Gould GW. (eds.) *Food preservatives*. Springer, Boston. doi: 10.1007/978-0-387-30042-9_10
- Kalia A and Gupta R. 2006. Fruit Microbiology. pp. 1-28. In: Hui YH (ed.). *Handbook of fruit and fruit processing*. First Edition. Blackwell publishing, United Kingdom. doi: 10.1002/9780470277737.ch1
- Katsinis G, Rigas F and Doulia D. 2008. Synergistic effect of chemical preservatives with ethanol on the microbial shelf life of bread by factorial design. *International Journal of Food Science and Technology* 43(2): 208-215 doi: 10.1111/j.1365-2621.2006.01386.x
- Krieg NR, Staley JT, Brown DR, Hedlund BP, Paster BJ, Ward NL, Ludwig W and Whitman WB. 2010. *Bergey's Manual of Systematic Bacteriology*. Second edition. Springer, New York. pp. 40-91. doi: 10.1007/978-0-387-68572-4
- Lewis DC and Goodrich-Schneider R. 2012. Mycotoxins in fruit and fruit products. *Proceedings of the Florida State Horticultural Society*. 125:252-257.

- Leyva Salas M, Mounier J, Valence F, Coton M, Thierry A and Coton E. 2017. Antifungal microbial agents for food biopreservation- A review. *Microorganisms* 5(3): 37. doi: 10.3390/microorganisms5030037
- Linares-Morales JR, Gutiérrez-Méndez N, Rivera-Chavira BE, Pérez-Vega SB and Nevárez-Moorillón GV. 2018. Biocontrol processes in fruits and fresh produce, the use of lactic acid bacteria as a sustainable option. *Frontiers in Sustainable Food System* 2:50. doi: 10.3389/fsufs.2018.00050
- Loureiro V and Malfeito-Ferreira M. 2003. Spoilage yeasts in the wine industry. *International Journal of Food Microbiology* 86(1-2): 23-503. doi: 10.1016/S0168-1605(03)00246-0
- Mailafia S, Okoh GR, Olabode H and Osanupin R. 2017. Isolation and identification of fungi associated with spoilt fruits vended in Gwagwalada market, Abuja, Nigeria. *Veterinary World* 10(4): 393-397. doi: 10.14202/vetworld.2017.393-397
- Maris P. 1995. Modes of action of disinfectants. *Revue scientifique et technique (International Office of Epizootics)* 14 (1):47-55. <https://doi.org/10.1016/j.riep.2017.03.001>
- McDonnell G and Russell AD. 1999. Antiseptics and disinfectants: activity, action, and resistance. *Clinical Microbiology Reviews* 12(1): 147-179. doi: 10.1128/CMR.12.1.147
- Miedes E. and Lorences EP. 2004. Apple (*Malus domestica*) and tomato (*Solanum lycopersicum*) fruits cell-wall hemicelluloses and xyloglucan degradation during *Penicillium expansum* infection. *Journal of Agricultural and Food Chemistry*. 52(26): 7957-7963. doi: 10.1021/jf048890f
- Moodley P and Gueguim Kana EB. 2019. Bioethanol production from sugarcane leaf waste: Effect of various optimized pretreatments and fermentation conditions on process kinetics *Biotechnology Reports* 22: e00329. doi: 10.1016/j.btre.2019.e00329
- Mussatto SI. 2014. Brewer's spent grain: a valuable feedstock for industrial applications. *Journal of the Science of Food and Agriculture* 94(7): 1264-1275. doi: 10.1002/jsfa.6486
- Olutiola PO, Famurewa O and Sonntag HG. 2000. An introduction to General Microbiology, a practical approach. Heidelberg, Nigeria. pp. 1-267.
- Prasad RK, Chatterjee S, Mazumder PB, Gupta SK, Sharma S, Vairale MG, Datta S, Dwivedi SK and Gupta DK. 2019. Bioethanol production from waste lignocelluloses: A review on microbial degradation potential. *Chemosphere* 231: 588-606. doi: 10.1016/j.chemosphere.2019.05.142
- Rojas-Chamorro JA, Romero-García JM, Cara C, Romero I and Castro E. 2020. Improved ethanol production from the slurry of pretreated brewers' spent grain through different co-fermentation strategies. *Bioresource Technology* 296: 122367. doi: 10.1016/j.biortech.2019.122367
- Saeed F, Afzaal M, Tufail T and Ahmad A. 2019. Use of natural antimicrobial agents: a safe preservation approach. In: Var I and Uzunhu S (eds.). *Active antimicrobial food packaging*. IntechOpen, London. doi: 10.5772/intechopen.80869
- Sagar NA, Pareek S, Sharma S, Yahia EM and Lobo MG. 2018. Fruit and vegetable waste: bioactive compounds, their extraction, and possible utilization. *Comprehensive Reviews in Food Science and Food Safety* 17(3): 512-531. doi: 10.1111/1541-4337.12330
- Saini JK, Saini R and Tewari L. 2015. Lignocellulosic agriculture wastes as biomass feedstocks for second-generation bioethanol production: concepts and recent developments. *3 Biotech* 5: 337-353 doi: 10.1007/s13205-014-0246-5
- Samson RA, Houbroken J, Thrane U, Frisvad JC and Andersen B. 2010. Fungi and indoor fungi. CBS-KNAW Fungal Biodiversity Centre, Utrecht. pp.1-390.
- Sharma N, Rathore M and Sharma M. 2013. Microbial pectinase: sources, characterization and applications. *Review in Environmental Science and Biotechnology* 12: 45-60. doi: 10.1007/s11157-012-9276-9
- Singh R, Kapoor V and Kumar V. 2012. Utilization of agro-industrial wastes for the simultaneous production of amylase and xylanase by thermophilic Actinomycetes. *Brazilian Journal of Microbiology* 43(4): 1545-1552.
- Steiner J, Procopio S and Becker T. 2015. Brewer's spent grain: source of value-added polysaccharides for the food industry in reference to the health claims. *European Food Research and Technology* 241(3): 303-315. doi: 10.1007/s00217-015-2461-7
- Tafinta IY, Shehu K, Abdulganiyyu H, Rabe AM and Usman A. 2013. Isolation and identification of fungi associated with the spoilage of sweet orange (*Citrus sinensis*) fruit in Sokoto State. *Nigerian Journal of Basic and Applied Sciences* 21(3): 193-196. doi: 10.4314/njbas.v21i3.4
- Tan YX, Mok WK and Chen WN. 2020. Potential novel nutritional beverage using submerged fermentation with *Bacillus subtilis* WX-17 on brewers' spent grains, *Heliyon* 6(6): e04155. doi: 10.1016/j.heliyon.2020.e04155
- Tournas VH. 2005. Spoilage of vegetable crops by bacteria and fungi and related health hazards. *Critical Reviews in Microbiology* 31(1): 33-44. doi: 10.1080/10408410590886024
- Ullah A, Orij R, Brul S and Smits GJ. 2012. Quantitative analysis of the modes of growth inhibition by weak organic acids in *Saccharomyces cerevisiae*. *Applied Environmental Microbiology* 78: 8377-8387. doi: 10.1128/AEM.02126-12
- Wilkinson S, Smart KA, James S and Cook DJ. 2017. Bioethanol production from brewers spent grains using a fungal consolidated bioprocessing (CBP) approach. *Bioenergy Research* 10(1): 146-157. doi: 10.1007/s12155-016-9782-7
- Wu M, Li H, Wei S, Wu H, Wu X, Bao X, Hou J, Liu W and Shen Y. 2020. Simulating Extracellular Glucose Signals enhances Xylose Metabolism in Recombinant *Saccharomyces cerevisiae*. *Microorganisms* 8(1): 100. doi: 10.3390/microorganisms8010100

