



Agronomía Colombiana

ISSN: 0120-9965

ISSN: 2357-3732

Universidad Nacional de Colombia, Facultad de Agronomía

de Mesquita Alves, Jackson; de Lima, Alex Serafim; Andrade Figueiredo, Francisco Romário; da Silva, Toshik Iarley; Ferreira Cavalcante, Lourival; de Oliveira Mesquita, Francisco; de Mesquita, Evandro Franklin; de Figueiredo Suassuna, Cesenildo
Chlorophyll a fluorescence and development of zucchini plants under nitrogen and silicon fertilization
Agronomía Colombiana, vol. 38, no. 1, 2020, January-April, pp. 45-52
Universidad Nacional de Colombia, Facultad de Agronomía

DOI: <https://doi.org/10.15446/agron.colomb.v38n1.79172>

Available in: <https://www.redalyc.org/articulo.oa?id=180376060006>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UNEM  redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Chlorophyll *a* fluorescence and development of zucchini plants under nitrogen and silicon fertilization

Fluorescencia de la clorofila *a* y desarrollo de plantas de calabacín bajo fertilización con nitrógeno y silicio

Jackson de Mesquita Alves¹, Alex Serafim de Lima², Francisco Romário Andrade Figueiredo^{3*}, Toshik Iarley da Silva⁴, Lourival Ferreira Cavalcante⁵, Francisco de Oliveira Mesquita⁶, Evandro Franklin de Mesquita², and Cesenildo de Figueiredo Suassuna¹

ABSTRACT

Zucchini (*Cucurbita pepo* L.) has a great economic and productive potential in the semi-arid region of Brazil, due to the wide acceptance by consumers and quick financial return. The nitrate (NO₃⁻) assimilation process in C₃ plants such as zucchini is related to photorespiration, and Si accumulated near the stomata reduces the transpiration rate, making the photosynthesis cycle more efficient. The objective of this study was to evaluate the interaction between nitrogen and silicon fertilization on growth, chlorophyll index, and chlorophyll *a* fluorescence of zucchini plants. The treatments were distributed in a split-plot scheme in a randomized block design with three replicates. The plot was arranged by silicon levels (0 and 6 g/plant) and the subplots constituted by five nitrogen levels (30, 60, 90, 120 and 150 kg ha⁻¹). Leaf, stem and total dry masses, chlorophyll *a*, chlorophyll *b*, total chlorophyll, chlorophyll *a/b* ratio and chlorophyll *a* fluorescence were evaluated. The highest dry matter productions in zucchini were obtained in treatments without Si. Si and N application together positively influences the chlorophyll *a/b* ratio of zucchini plants. The interaction between Si and N positively influences the maximum fluorescence, variable fluorescence and quantum yield of photosystem II of zucchini plants.

Key words: *Cucurbita pepo* L., fertilizers, photosynthetic efficiency.

RESUMEN

El calabacín (*Cucurbita pepo* L.) presenta un gran potencial económico y productivo en la región semiárida de Brasil, debido a su amplia aceptación por los consumidores y su rápido rendimiento financiero. El proceso de asimilación de nitrato (NO₃⁻) en plantas C₃ como el calabacín estaría relacionado en parte con la fotorrespiración, y el Si acumulado cerca de los estomas reduce la tasa de transpiración, haciendo que el ciclo de fotosíntesis sea más eficiente. Por lo tanto, el objetivo de este estudio fue evaluar la interacción entre el nitrógeno y la fertilización con silicio en el crecimiento, el índice de clorofila y la fluorescencia de la clorofila *a* en plantas de calabacín. Los tratamientos se distribuyeron en un esquema de parcelas divididas en un diseño de bloques al azar con tres repeticiones. La parcela se formó por niveles de silicio (0 y 6 g/planta) y las subparcelas constituidas por cinco niveles de nitrógeno (30, 60, 90, 120 y 150 kg ha⁻¹), totalizando 30 parcelas experimentales. Se evaluaron masas de hojas, tallos y secas totales, clorofila *a*, clorofila *b*, clorofila total, clorofila *a/b* y fluorescencia de la clorofila *a*. Las mayores producciones de materia seca de calabacín se obtuvieron en tratamientos sin Si. La aplicación de Si y N juntos influye positivamente en la proporción de clorofila *a/b* de las plantas de calabacín. La interacción entre Si y N influye positivamente en la fluorescencia máxima, la fluorescencia variable y el rendimiento cuántico del fotosistema II de plantas de calabacín.

Palabras clave: *Cucurbita pepo* L., fertilizantes, eficiencia fotosintética.

Introduction

Italian zucchini (*Cucurbita pepo* L. - Cucurbitaceae) is a vegetable-like fruit with an erect bushy growing habit,

despite having an herbaceous stem. It is one of the most popular and appreciated vegetables in Brazilian cuisine (Nakada-Freitas, 2014; Azambuja *et al.*, 2015). Zucchini is a summer squash that develops in autumn and spring,

Received for publication: 18 April, 2019. Accepted for publication: 29 February, 2020

Doi: 10.15446/agron.colomb.v38n1.79172

¹ Federal University of Campina Grande, Pombal, Paraíba (Brazil).

² State University of Paraíba, Catole do Rocha, Paraíba (Brazil).

³ Federal Rural University of the Semi-arid, Mossoro, Rio Grande do Norte (Brazil).

⁴ Federal University of Viçosa, Viçosa, Minas Gerais (Brazil).

⁵ Federal University of Paraíba, Areia, Paraíba (Brazil).

⁶ National Institute of the Semi-arid, Campina Grande, Paraíba (Brazil).

* Corresponding author: romarioagroecologia@yahoo.com.br



and also during the winter in places with a warmer climate (Filgueira, 2012).

The effects of nutrition and fertilization management practices on growth, development, and yield of zucchini, as well as other crops, has been one of the most studied agricultural aspects (Domingos *et al.*, 2015). Nitrogen (N), which is responsible for several functions in plants and is directly linked to crop yield, is the main topic for most researchers (Oliveira *et al.*, 2014).

Nitrogen is the second most required nutrient by plants (Viciedo *et al.*, 2017). It is an essential macro-element that participates in the synthesis of amino acids, nucleic acids, secondary metabolites, and components of the chlorophyll molecule. In addition, this nutrient stimulates the photosynthetic capacity of leaves by increasing the content of stromal and thylakoid proteins in chloroplasts (Akram *et al.*, 2011). However, over-fertilization with nitrogen has negative effects, such as disordered plant growth, susceptibility to pathogens and insects, nutrient deficiency due to competition with other nutrients, auto shading, and increased production costs (Porto *et al.*, 2012; Xu *et al.*, 2012).

Rational nitrogen fertilization practices are of great importance, being essential the search for methods that boost the beneficial effects of this nutrient. Therefore, nitrogen application combined with silicon (Si) may provide higher yields (Artigiani *et al.*, 2014), making it a viable alternative. Adequate Si supply may induce growth, resistance to biotic and abiotic stresses, and increase the photosynthetic capacity (Ferraz *et al.*, 2014). In addition, Si application provides a better utilization of some nutrients by plants (Watanabe *et al.*, 2001).

Chlorophyll *a* fluorescence can be a good indicator of the plant's physiological state. The absorbed light energy excites the chlorophyll molecule and the disposing of its available energy occurs through three alternative pathways which compete for the excitation energy (photochemical dissipation, fluorescence or non-photochemical dissipation). Thus, changes in one of these processes, as in photosynthesis for example, shall affect the other ones (Cendrero-Mateo *et al.*, 2016).

Plants spend about 25% of their total energy on NO_3^- assimilation (Rubio-Asensio *et al.*, 2015). In C_3 plants, this process would be partly coupled with photorespiration, which stimulates the transport of chloroplast malate to the cytosol for Nicotinamide Adenine Dinucleotide Hydride (NADH) formation (Bloom, 2015). NADH is used by nitrate

reductase to reduce NO_3^- to NO_2^- , NO_3^- assimilation process and would be related in part to photorespiration. In this regard, Si accumulated near the stomata reduces the rate of transpiration (Oliveira and Castro, 2002), providing a direct relationship between transpiration and chlorophyll fluorescence and Si and N, making the photosynthesis process more efficient.

Chlorophyll fluorescence parameters are widely used in physiological studies since they are obtained by a non-destructive method that allows the qualitative and quantitative analysis of the absorption and use of light energy by the photosynthetic apparatus (Olivos *et al.*, 2017). Therefore, the aim of this study was to evaluate the interaction between nitrogen and silicon fertilization on growth, chlorophyll index and chlorophyll *a* fluorescence of zucchini plants.

Materials and methods

The experiment was carried out under field conditions in the agroecology sector of the Agrarian and Exact Sciences Department (DAE) of the State University of Paraíba (UEPB), Campus IV, in the municipality of Catole do Rocha, Paraíba, Brazil (6°20'38" S, 37°44'48" W, 275 m a.s.l.).

Zucchini seeds cv. Caserta were directly sown in planting pits, spaced at 1 × 1 m. After digging the pits of 30 × 30 × 30 cm dimension, 2 L of tanned cattle manure were added (Filgueira *et al.*, 2012).

The soil of the experimental area was classified as eutrophic Fluvic Neosol, with a sandy loam texture (Embrapa, 2013). Before the installation of the experiment, soil samples were collected at a depth of 0-20 cm for physical and fertility analysis (Tab. 1) and the cattle manure chemical analysis was also performed (Tab. 2), according to the methodology proposed by Embrapa (2013). The soil was prepared by plowing sequenced by harrowing.

The treatments were distributed in a split-plot scheme in a randomized complete block design, with three replicates, and one plant per replicate. The plot was arranged by silicon levels (0 and 6 g/plant) and the subplots were five nitrogen levels (30, 60, 90, 120, and 150 kg ha⁻¹). The nitrogen source used was urea (45% N). The silicon levels were based on an estimate of 60 kg ha⁻¹. The product Bugram Protect®, which contains 100% pure silicon dioxide, pH 6.8 and 0.9% moisture, was used as a silicon source. The N doses were equally divided and applied to cover the soil twice, first at 15 days after planting (DAP) and then at 30 DAP. Foliar fertilization with Si was also divided into two applications,

TABLE 1. Chemical and physical characteristics of the Eutrophic Fluvic Neosol used in the experiment.

Chemical characteristics (Fertility)												
pH	Ca ²⁺	Mg ²⁺	Na ⁺	K ⁺	S	H+Al	CEC	PBS	OC	OM	N	AP
	cmol _c dm ⁻³						cmol _c kg ⁻¹	%			mg dm ⁻³	
6.7	1.49	0.54	0.10	1.72	3.85	0.00	3.85	100%	0.67	1.2	0.07	16.83
Physical characteristics												
Sand	Silt	Clay	EC			GD	PD	TP		FC	PWP	AW
g kg ⁻¹			g dm ⁻³				(%)		g kg ⁻¹			
640.0	206.0	154.0	Sandy loam			1.54	2.68	42.54		146.9	76.60	70.3

CEC: Cation exchange capacity; PBS: Percent base saturation; OC: Organic carbon; OM: Organic matter; AP: Assimilable phosphorus; EC: Exchangeable cations; GD: Global density; PD: Particle density; TP: Total porosity; FC: Field capacity; PWP: Permanent wilting point; AW: Available water.

TABLE 2. Chemical characteristics of the cattle manure used in Catole do Rocha, 2019.

N	P	K ⁺	Ca ²⁺	Mg ²⁺	Na ⁺	Zn ²⁺	Cu ²⁺	Fe ²⁺	Mn ²⁺	SOM	OC	C/N
g kg ⁻¹						mg kg ⁻¹				g kg ⁻¹		
12.76	2.57	16.79	15.55	4.02	5.59	60	22	8550	325	396.0	229.7	18:1

SOM: Soil organic matter; OC: Organic carbon; C/N: carbon-to-nitrogen ratio.

the first at 14 DAP and the second at 28 DAP. Thus, 540 g of silicon dioxide was diluted in 30 L of water in the first application and 540 g was diluted in 40 L of water in the second application. The silicon solutions were uniformly applied on the plants, using a 10 L capacity backpack sprayer, totaling 1.080 kg of silicon dioxide.

Micro-irrigation was performed through a drip irrigation system, spaced at 0.2 m and with a flow rate of 1.7 L per hour, at a service pressure of 147 KPa, using 16 mm drip tapes. The water was supplied through an Amazon well near the experiment site, and it was classified as moderately saline (1.2 dS m⁻¹) (Ayers and Westcot, 1999).

The chlorophyll *a* fluorescence trials were performed at 35 DAP using a modulated fluorometer (Model OS-30p, Hudson, USA). Foliar tweezers were placed for 30 min before the readings to adapt the leaves to the dark. The initial fluorescence (F_0), maximum fluorescence (F_m), variable fluorescence ($F_v = F_m - F_0$) and quantum yield of photosystem II (F_v/F_m) were evaluated. Chlorophyll *a*, *b* and total contents were estimated by the non-destructive method using a portable chlorophyll meter (ClorofiLOG®, model CFL 1030, Porto Alegre, Brazil), and expressed as Falker Chlorophyll Index (FCI). For leaf, stem and total dry mass estimation, each part of the plant was packed in Kraft paper bags and dried in a forced-air circulation oven at 65°C until reaching constant weight. Subsequently, the material was weighed in a precision analytical balance (0.001 g), and the results expressed in g/plant.

The data were submitted to an analysis of variance by the F test at 5% probability. Polynomial regression analysis for

the nitrogen factor and the Tukey test for the silicon factor were performed in case of significance. The statistical program R was used (R Core Team, 2018).

Results and discussion

Leaf dry mass (LDM), stem dry mass (SDM) and total dry mass (TDM) of zucchini were significant at 1% of probability in the interaction between the studied factors, nitrogen (N) and silicon (Si) (Tabs. 3 and 4).

For LDM, an inverse behavior was observed in the treatments with and without Si, respectively. The highest increase in LDM was found in the treatments without Si and at the maximum N dose (150 kg ha⁻¹), which corresponds to 338.6 g/plant. However, the LDM in the presence of Si reached its best value in the order of 265.4 g/plant, at the estimated dose of 87 kg ha⁻¹ (Fig. 1A).

The values for SDM adjusted to a quadratic effect, both in the presence and absence of Si. The highest increase was registered at the N dose of 79.8 kg ha⁻¹ in the treatments without Si, which corresponds to a 21.5% increase when compared to treatments under Si application, with values in the order of 33.3 and 27.4 g/plant, respectively (Fig. 1B).

For TDM, the results were similar to those of LDM. The best performance was obtained at the N dose of 150 kg ha⁻¹ in the treatments without Si, corresponding to 355.5 g/plant, which may be related to the maximum increase observed in LDM (Fig. 1C).

TABLE 3. Summary of the analysis of variance by the mean-square values, from leaf dry mass (LDM), stem dry mass (SDM) and total dry mass (TDM) of zucchini (*Cucurbita pepo* L.) under nitrogen doses and silicon leaf fertilization in Catole do Rocha, 2019.

SV	DF	Mean square		
		LDM	SDM	TDM
Block	1	68.10 ^{ns}	0.85 ^{ns}	67.20 ^{ns}
Si	2	3853.30*	126.12**	5373.70*
Error a	2	180.10	0.10	174.00
N	4	4571.50**	213.17**	4090.60**
Si x N	4	11825.70**	132.69**	10190.10**
Error b	16	239.10	1.15	256.80
CV 1 (%)		5.60	1.19	4.95
CV 2 (%)		6.45	3.99	6.01

SV: Source of variation; DF: Degrees of freedom; CV: Coefficient of variation; ns: Not significant; ** and *: Significant effect by the F test at the level of 1 and 5% probability, respectively.

TABLE 4. Mean values for leaf dry mass (LDM), stem dry mass (SDM) and total dry mass (TDM) of zucchini (*Cucurbita pepo* L.) under nitrogen doses and silicon leaf fertilization in Catole do Rocha, 2019.

Parameter	Silicon (g/plant)	Nitrogen (kg ha ⁻¹)				
		30	60	90	120	150
LDM	0	242 a	250 b	191 b	217 a	355 a
	6	185 b	277 a	257 a	227 a	195 b
SDM	0	26 a	31 a	46 a	25 b	18 b
	6	21 b	26 b	27 b	27 a	23 a
TDM	0	268 a	280 a	236 b	241 a	372 a
	6	206 b	303 a	283 a	254 a	217 b

Means followed by equal letters in the same column do not differ statistically from each other by the Tukey test at 5% probability.

The positive values obtained from nitrogen fertilization and the different morphological parameters may be due to the fact that this nutrient provides increases in the rate of carbohydrates conversion into protein and the size of the cells. This may lead to greater leaf expansion and higher growth rates, which directly influences the production of dry matter. Yasari *et al.* (2009) and Rodrigues *et al.* (2014) also registered positive results for dry mass production with increased N supply in paracress plants (*Acmella oleracea* L.), in which a linear increase was verified, with the highest efficiency registered at 112.5 kg ha⁻¹ of N.

The decrease in dry mass of silicon-applied plants may be associated with the fact that silicon deposited on shoots is condensed by transpiration and subsequently transformed into amorphous silicon located in the cell wall of epidermal and vascular tissues, causing reduced water loss (Alsaedi *et al.*, 2019).




FIGURE 1. (A) Leaf dry mass, (B) stem dry mass, and (C) total dry mass of zucchini (*Cucurbita pepo* L.) under nitrogen doses and silicon leaf fertilization.

According to the results of the ANOVA, it was observed that there was a significant effect of the nitrogen doses on the chlorophyll contents and chlorophyll *a* fluorescence parameters. For the silicon factor, it was observed that there was a significant difference only for the fluorescence variables, except for the variable fluorescence (*F_v*). However, the interaction between the factors significantly affected all variables analyzed (Tab. 5).

Chl *a* and Chl *b* values presented a quadratic effect both in the treatments with and without Si, with the highest index

TABLE 5. Summary of the analysis of variance by the mean-square values, for chlorophyll *a* (Cla), chlorophyll *b* (Cib), total chlorophyll (tCl), chlorophyll *a/b* ratio (Cla/Cib), initial fluorescence (F_0), maximum fluorescence (F_m), variable fluorescence (F_v), and quantum yield of photosystem II (F_v/F_m) of zucchini (*Cucurbita pepo* L.) under nitrogen doses and silicon leaf fertilization in Catole do Rocha, 2019.

SV	DF	Mean square			
		Cla	Cib	tCl	Cla/Cib
Block	1	0.070 ^{ns}	0.050 ^{ns}	1.339 ^{ns}	0.006 ^{ns}
Si	2	0.551 ^{ns}	0.075 ^{ns}	1.254 ^{ns}	0.0002 ^{ns}
Error a	2	0.113	0.007	0.636	0.001
N	4	12.617**	3.654**	9.905**	0.035**
Si x N	4	11.510**	2.167**	41.764**	0.026**
Error b	16	0.168	0.034	0.727	0.004
CV 1 (%)		1.01	0.72	1.80	1.14
CV 2 (%)		1.23	1.58	1.93	2.16
		F_0	F_m	F_v	F_v/F_m
Block	1	11.000 ^{ns}	59.000 ^{ns}	65.100 ^{ns}	0.000001 ^{ns}
Si	2	77724.000**	69216.000**	246.530 ^{ns}	0.019**
Error a	2	8.000	8.000	27.230	0.000034
N	4	3637.000**	3072.000**	921.800**	0.022**
Si x N	4	363.000**	86.000**	784.370**	0.0004**
Error b	16	44.000	18.000	65.580	0.00004
CV 1 (%)		1.31	0.99	6.56	0.80
CV 2 (%)		3.15	1.45	10.17	0.81

SV: Source of variation; DF: Degrees of freedom; CV: Coefficient of variation; ns: Not significant; ** and *: Significant effect by the F test at the level of 1 and 5% probability, respectively.

TABLE 6. Mean values for chlorophyll *a* (Cla), chlorophyll *b* (Cib), total chlorophyll (tCl), chlorophyll *a/b* ratio, initial fluorescence (F_0), maximum fluorescence (F_m), variable fluorescence (F_v), and quantum yield of photosystem II (F_v/F_m) of zucchini (*Cucurbita pepo* L.) under nitrogen doses and silicon leaf fertilization in Catole do Rocha, 2019.

Parameters	Silicon (g/plant)	Nitrogen (kg ha ⁻¹)				
		30	60	90	120	150
Cla	0	32.83 b	29.61 b	35.31 a	34.86 a	34.21 a
	6	34.09 a	32.58 a	34.43 b	30.40 b	33.97 a
Cib	0	11.68 b	10.02 b	12.21 b	12.61 a	11.89 b
	6	12.02 a	10.71 a	12.60 a	10.38 b	12.20 a
tCl	0	45.86 a	40.13 b	42.72 b	46.38 a	46.99 a
	6	42.47 b	45.63 a	47.39 a	39.76 b	44.79 b
Chlorophyll <i>a/b</i> ratio	0	2.81 a	2.96 a	2.89 a	2.76 b	2.88 a
	6	2.84 a	3.04 a	2.73 b	2.93 a	2.78 a
F_0	0	150.67 b	162.67 b	189.00 b	158.00 b	139.00 b
	6	236.33 a	254.33 a	315.33 a	260.00 a	242.33 a
F_m	0	235.67 b	249.33 b	281.00 b	226.67 b	219.00 b
	6	341.67 a	348.00 a	366.67 a	324.33 a	311.33 a
F_v	0	85.00 b	86.67 a	92.00 a	68.67 a	80.00
	6	105.33 a	93.67 a	51.33 b	64.33 a	69.00
F_v/F_m	0	0.67	0.72	0.79	0.73	0.62 a
	6	0.71	0.76	0.84	0.78	0.69 a

Means followed by equal letters in the same column and line do not differ statistically from each other by the Tukey test at 5% probability.

found at 146.1 kg ha⁻¹ of N without Si application, evidencing a 5.6% increase compared to the treatment with Si. This behavior in the Si-containing treatments demonstrates FCI values for Chl *a* and Chl *b* in the order of 35.4 and 33.4, respectively (Fig. 2A).

For Chl *b*, a similar behavior was observed, with the highest increase in the treatments without Si and under the N dose of 137.7 kg ha⁻¹, with an FCI of 12.8. Under Si application, the N dose that provided the highest efficiency was 93.8 kg ha⁻¹ (Fig. 2B). Since N is one of the main components of chlorophyll molecules, an adequate supply of this nutrient maximizes the indices of this photosynthetic pigment (Wang *et al.*, 2014).

The total chlorophyll index increased linearly from the N dose of 51.7 kg ha⁻¹, in the treatments without Si application. Under Si fertilization, the best performance of plants for this variable was found at 103.0 kg ha⁻¹ of N, with a 46.8 FCI (Fig. 2C). However, it can be observed that, for the chlorophyll *a/b* ratio, the Si-containing treatments presented the best results, with the greatest increase at 82.3 kg ha⁻¹ of N (Fig. 2D).

Corroborating this statement, Ávila *et al.* (2010) report a leaf chlorophyll synthesis increase in silicon-fertilized rice plants (*Oryza sativa* L.), especially at high doses of nitrogen. Contrary to this, Rodrigues *et al.* (2016) state that increased potassium silicate doses provide increases in chlorophyll *a* levels in tomato plants (*Solanum lycopersicum* L.) under high and low light conditions. However, the increase in chlorophyll *a/b* ratio may be associated with the greater synthesis of chlorophyll *a*, which is an indication that the plant was not experiencing stress.

It was observed that F₀ and F_m showed a similar behavior, adjusting to a quadratic effect, and the best results were obtained in the treatments under Si fertilization, with values in the order of 288.5 and 355.2 /quantum at the N doses of 91.9 and 72.3 kg ha⁻¹, respectively (Figs. 3A and B). This increase in F₀ indicates that in the aforementioned N dose there was a reduction in the capacity of excitation energy transfer to the photosystem II (SÁ *et al.*, 2015). F_m refers to the state in which the reaction centers of photosystem II reach their maximum capacity (Silva *et al.*, 2014). Silicon accumulated near the stomata reduces leaf transpiration rates (Oliveira and Castro, 2002), a fact that may have benefited




FIGURE 2. (A) Chlorophyll *a*, (B) chlorophyll *b*, (C) total chlorophyll and (D) chlorophyll *a/b* ratio of zucchini (*Cucurbita pepo* L.) under nitrogen doses and silicon leaf fertilization.




FIGURE 3. (A) Initial fluorescence (F_0), (B) maximum fluorescence (F_m), (C) variable fluorescence (F_v), and (D) quantum yield of photosystem II (F_v/F_m) of zucchini (*Cucurbita pepo* L.) under nitrogen doses and silicon leaf fertilization.

nitrate reductase in the photorespiration process, having increased chlorophyll indices in chloroplasts (Bloom, 2015).

The F_v in the treatments with Si showed the best result at the N dose of 30 kg ha⁻¹, with a value of 109.0 quantum electrons (Fig. 3C). For the F_v/F_m variable, the N dose of 89.0 kg ha⁻¹ under Si application provided the highest value (0.81 quantum electrons), a 6.2% increase compared to treatments without Si (76 quantum electrons) (Fig. 3D).

Variable fluorescence refers to active potential energy in photosystem II, which is directly influenced by the maximum fluorescence (Sá *et al.*, 2018). According to these same authors, these increases in fluorescence may be related to the higher content of chlorophyll in the reaction centers as a function of the N doses. Based on the F_v/F_m results, it can be stated that the N combined with Si induced a better energy stability compared to the treatments that did not contain Si (0.81), which is within those indicated as normal (between 0.75 and 0.85) in non-stressed plants (Neves *et al.*, 2019).

Conclusions

The highest dry matter productions in zucchini plants were obtained in treatments without Si. The application

of Si and N together positively influences the chlorophyll *a/b* ratio of zucchini plants. The interaction between Si and N positively influences the maximum fluorescence, variable fluorescence and quantum yield of photosystem II of zucchini plants.

Acknowledgments

The authors thank the Coordination for the Improvement of Higher Education Personnel (CAPES) and the Brazilian National Council for Scientific and Technological Development (CNPq) for financial support of this research.

Literature cited

- Akram, M., M.Y. Ashraf, M. Jamil, R.M. Iqbal, M. Nafees, and M.A. Khan. 2011. Nitrogen application improves gas exchange characteristics and chlorophyll fluorescence in maize hybrids under salinity conditions. *Russ. J. Plant Physiol.* 58(3), 394-401. Doi: 10.1134/S1021443711030022
- Alsaedi, A., H. El-Ramady, T. Alshaal, M. El-Garawany, N. Elhawat, and A. Al-Otaibi. 2019. Silica nanoparticles boost growth and productivity of cucumber under water deficit and salinity stresses by balancing nutrients uptake. *Plant Physiol. Biochem.* 139, 1-10. Doi: 10.1016/j.plaphy.2019.03.008
- Artigiani, A.C.C.A., C.A.C. Crusciol, A.S. Nascente, O. Arf, and R.C.F. Alvarez. 2014. Adubação silicatada no sulco e

- nitrogenada em cobertura no arroz de sequeiro e irrigado por aspersão. *Biosci. J.* 30(1), 240-251.
- Ávila, F.W., D.P. Baliza, V. Faquin, J.L. Araújo, and S.J. Ramos. 2010. Interação entre silício e nitrogênio em arroz cultivado sob solução nutritiva. *Rev. Cienc. Agron.* 41(2), 184-190. Doi: 10.1590/S1806-66902010000200003
- Ayers, R.S. and D.W.A. Westcot. 1999. A qualidade da água na agricultura. 2nd ed. UEPB, Campina Grande, Brazil.
- Azambuja, L.O., C.G.S. Benett, K.S.S. Benett, and E. Costa. 2015. Produtividade da abobrinha 'Caserta' em função do nitrogênio e gel hidrorretentor. *Científica* 43(4), 353-358. Doi: 10.15361/1984-5529.2015v43n4p353-358
- Bloom, A.J. 2015. Photorespiration and nitrate assimilation: a major intersection between plant carbon and nitrogen. *Photosynth. Res.* 123(2), 117-28. Doi: 10.1007/s1120-014-0056-y
- Cendrero-Mateo, M.P., M.S. Moran, S.A. Papuga, K.R. Thorp, L. Alonso, J. Moreno, G. Ponce-Campos, U. Rascher, and G. Wang. 2016. Plant chlorophyll fluorescence: active and passive measurements at canopy and leaf scales with different nitrogen treatments. *J. Exp. Botany* 67(1), 275-286. Doi: 10.1093/jxb/erv456
- Domingos, C.S., L.H.S. Lima, and A.L. Braccini. 2015. Nutrição mineral e ferramentas para o manejo da adubação na cultura da soja. *Sci. Agrar. Paran.* 14(3), 132-140.
- EMBRAPA - Empresa Brasileira de Pesquisa Agropecuária. 2013. Manual de métodos e análise do solo. Centro Nacional de Pesquisa de Solos, Brazil.
- Ferraz, R.N.S., N.E.M. Beltrão, A.S. Melo, I.D. Magalhães, P.D. Fernandes, and M.S. Rocha. 2014. Trocas gasosas e eficiência fotoquímica de cultivares de algodoeiro herbáceo sob aplicação de silício foliar. *Semina: Ciênc. Agrár.* 35(2), 735-748. Doi: 10.5433/1679-0359.2014v35n2p735
- Filgueira, F.A.R. 2012. Novo manual de olericultura: agrotecnologia moderna na produção e comercialização de hortaliças. UFV, Viçosa, Brazil.
- Nakada-Freitas, P.G., M.A. Sandri, A.I.I. Cardoso, and E. Eberhardt Neto. 2014. Controle alternativo de oídio em abobrinha de moita com solução de vinagre. *Nucleus* 11(2), 325-332. Doi: 10.3738/1982.2278.1040
- Neves, J.M.G., L.A. Aquino, P.G. Berger, J.C.L. Neves, G.C. Rocha, and E.A. Barbosa. 2019. Silicon and boron mitigate the effects of water deficit on sunflower. *Rev. Bras. Eng. Agríc. Ambient.* 23(3), 175-182. Doi: 10.1590/1807-1929/agriambi.v23n3p175-182
- Oliosio, G., J. De O. Rodrigues, A.R. Falqueto, F.R. Pires, J.A. Monte, and F.L. Partelli. 2017. Fluorescência transiente da clorofila a e crescimento vegetativo em cafeeiro conilon sob diferentes fontes nitrogenadas. *Coffee Sci.* 12(2), 248-259.
- Oliveira, F.A., J.F. Medeiros, R.C. Alves, P.S.F. Linhares, A.M.A. Medeiros, and M.K.T. Oliveira. 2014. Interação entre a salinidade da água de irrigação e adubação nitrogenada na cultura da berinjela. *Rev. Bras. Eng. Agríc. Ambient.* 18(5), 480-486. Doi: 10.1590/S1415-43662014000500003
- Oliveira, L.A. and N.M. Castro. 2002. Ocorrência de sílica nas folhas de *Curatella Americana* L. e de *Davilla elliptica* St. Hil. *R. Horiz. Ci.* 4(1), 1-16.
- Porto, M.L.A., M. Puiatti, P.C.R. Fontes, P.R. Cecon, J.C. Alves, and J.A. Arruda. 2012. Produtividade e acúmulo de nitrato nos frutos de abobrinha em função da adubação nitrogenada. *Bragantia* 71(2), 190-195. Doi: 10.1590/S0006-87052012005000020
- R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rodrigues, C.R., T.M. Rodrigues, J.M.Q. Luz, V.B.F. Sousa, J.B. Sousa, A.C.P. Nunes, and P.R. Trindade. 2016. Clorofila a e b de tomateiro tratado com silicato de potássio e fungicida. *Global Sci. Technol.* 9(2), 54-64. Doi: 10.14688/1984-3801/gst.v9n2p54-64
- Rodrigues, D.S., M.S. Camargo, E.S. Nomura, V.A. Garcia, J.N. Correa, and T.C.M. Vidal. 2014. Influência da adubação com nitrogênio e fósforo na produção de Jambu, *Acmella oleracea* (L.) R. K. Jansen. *Rev. Bras. Plantas Med.* 16(1), 71-76. Doi: 10.1590/S1516-05722014000100010
- Rubio-Asensio, J.S., S. Rachmilevitch, and A.J. Bloom. 2015. Plant responses to rising CO₂ depend on nitrogen source and night-time CO₂ levels. *Plant Physiol.* 168(1), 156-63.
- Sá, F.V.S., M.E.B. Brito, L.A. Silva, R.C.L. Moreira, P.D. Fernandes, and L.C. Figueiredo. 2015. Fisiologia da percepção do estresse salino em híbridos de tangerineira "Sunki Comum" sob solução hidropônica salinizada. *Comunica Scientiae* 6(4), 463-470. Doi: 10.14295/CS.v6i4.1121
- Sá, F.V.S., H.R. Gheyi, G.S. Lima, E.P. Paiva, R.C.L. Moreira, and L.A. Silva. 2018. Water salinity, nitrogen and phosphorus on photochemical efficiency and growth of west indian cherry. *Rev. Bras. Eng. Agríc. Ambient.* 22(3), 158-163. Doi: 10.1590/1807-1929/agriambi.v22n3p158-163
- Silva, L.A., M.E.B. Brito, F.V.S. Sá, R.C.L.M. Moreira, W.S. Soares Filho, and P.D. Fernandes. 2014. Mecanismos fisiológicos em híbridos de citros sob estresse salino em cultivo hidropônico. *Rev. Bras. Eng. Agríc. Ambient.* 18, S1-S7. Doi: 10.1590/1807-1929/agriambi.v18nsupps1-s7
- Viciedo, D.O., R.M. Prado, R.L. Toledo, L.C.N. Santos, and K.P. Calzada. 2017. Response of radish seedlings (*Raphanus sativus* L.) to different concentrations of ammoniacal nitrogen in absence and presence of silicon. *Agron. Colomb.* 35(2), 198-204. Doi: 10.15446/agron.colomb.v35n2.62772
- Wang, Y., D. Wang, P. Shi, and K. Omasa. 2014. Estimating rice chlorophyll content and leaf nitrogen concentration with a digital still color camera under natural light. *Plant Meth.* 36(10), 1-11. Doi: 10.1186/1746-4811-10-36
- Watanabe, S., T. Fujiwara, T. Yoneyama, and H. Hayashi. 2001. Effects of silicone nutrition on metabolism and translocations of nutrients in rice plants. *Plant Nutrit.* 92, 174-175. Doi: 10.1007/0-306-47624-X_84
- Xu, G., X. Fan, and A.J. Miller. 2012. Plant nitrogen assimilation and use efficiency. 2012. *Ann. Rev. Plant Biol.* 63, 153-182. Doi: 10.1146/annurev-arplant-042811-105532
- Yasari, M.A.E., E. Azadgoleh, S. Mozafari, and M.R. Alashti. 2009. Enhancement of growth and nutrient uptake of rapeseed (*Brassica napus* L.) by applying mineral nutrients and biofertilizers. *Pak. J. Biol. Sci.* 12, 127-133. Doi: 10.3923/pjbs.2009.127.133