



Clínica y Salud

ISSN: 1130-5274

ISSN: 2174-0550

Colegio Oficial de la Psicología de Madrid

Frías, Álvaro; Palma, Carol; Farriols, Núria; Aliaga, Ferrán; Navarro, Sara; Solves, Laia
Parenting Attitudes in People with Obsessive-compulsive
Disorder and Emotional Symptoms in Their Children
Clínica y Salud, vol. 31, no. 2, 2020, May-August, pp. 91-98
Colegio Oficial de la Psicología de Madrid

DOI: <https://doi.org/10.5093/clysa2020a13>

Available in: <http://www.redalyc.org/articulo.oa?id=180663452004>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Parenting Attitudes in People with Obsessive-compulsive Disorder and Emotional Symptoms in Their Children

Frías, Álvaro; Palma, Carol; Farriols, Núria; Aliaga, Ferrán; Navarro, Sara; Solves, Laia
Parenting Attitudes in People with Obsessive-compulsive Disorder and Emotional Symptoms in Their Children
Clínica y Salud, vol. 31, no. 2, 2020
Colegio Oficial de la Psicología de Madrid
Available in: <http://www.redalyc.org/articulo.oa?id=180663452004>
DOI: 10.5093/clysa2020a13

Parenting Attitudes in People with Obsessive-compulsive Disorder and Emotional Symptoms in Their Children

Las actitudes parentales de personas con trastorno obsesivo-compulsivo y los síntomas emocionales de sus hijos

Álvaro Frías a*

Ramon-Llull University, Spain

Carol Palma a

Ramon-Llull University, Spain

Núria Farriols a

Ramon-Llull University, Spain

Ferrán Aliaga a

Ramon-Llull University, Spain

Sara Navarro a

Ramon-Llull University, Spain

Laia Solves b

Consorci Sanitari del Maresme, Spain

Clínica y Salud, vol. 31, no. 2, 2020

Colegio Oficial de la Psicología de Madrid

Received: 15 November 2019

Accepted: 08 April 2020

DOI: 10.5093/clysa2020a13

CC BY-NC-ND

ABSTRACT: This study sought to delineate distinctive parenting attitudes in people with obsessive-compulsive disorder (OCD), specific emotional symptoms in their children, and the association between them. Forty OCD parents and their children were compared with 37 parents with adjustment disorders and their children by using standardized clinical questionnaires. Children of OCD parents exhibited significantly greater (subclinical) emotional symptoms when compared with children of non-OCD parents. After controlling for parents and childrens depression and anxiety symptoms, OCD parents reported significantly poorer parenting attitudes overall relative to non-OCD parents. The presence of sexual/somatic obsessions in OCD parents predicted anxiety symptom severity among their children, but both relationships were mediated by parental involvement. These findings indicate the importance of addressing and treating the distinctive parenting attitudes among people with OCD and its influence on their childrens emotional symptoms.

Keywords: Obsessive compulsive disorder, Adjustment disorders, Emotional symptoms, Parenting attitudes, Involvement.

RESUMEN: Este estudio pretende delimitar las actitudes parentales distintivas en personas con trastorno obsesivo-compulsivo (TOC), los síntomas emocionales específicos en sus hijos y la asociación entre ambos. Cuarenta padres con TOC y sus hijos fueron comparados con 37 padres con trastornos adaptativos y sus hijos mediante el uso de cuestionarios clínicos estandarizados. Los hijos de padres con TOC manifestaron significativamente más síntomas emocionales (nivel subclínico) en comparación con los hijos de padres con trastornos adaptativos. Tras controlar el efecto de los síntomas de depresión y ansiedad de los padres y los niños, los padres con TOC indicaban actitudes parentales significativamente peores que las de los padres con trastornos adaptativos. La presencia de obsesiones sexuales/somáticas en los padres con TOC predecía la severidad de los síntomas de ansiedad en sus hijos, si bien dichas relaciones estaban mediadas por el nivel de implicación parental. Estos hallazgos indican la importancia de abordar y

tratar las actitudes parentales distintivas de las personas con TOC y su influencia en la sintomatología emocional de sus hijos

Palabras clave: Trastorno obsesivo-compulsivo, Trastornos adaptativos, Síntomas emocionales, Actitudes parentales, Implicación.

Introduction

Obsessive-compulsive disorder (OCD) is a mental disorder in which people experience uncontrollable, intrusive thoughts (obsessions), which may be accompanied by distressing and repetitive mental/behavioral acts (compulsions). Obsessions and compulsions negatively affect quality of life among people with OCD (American Psychiatric Association [APA, 1994]). Over the last 15 years, several empirical studies have also shown that OCD has an adverse effect on relatives quality of life and the presence of OCD symptoms has been linked to family accommodation and elevated expressed emotion between family members (Cicek et al., 2013; Gomes et al., 2016; Lebowitz et al., 2012). For instance, the presence of contamination obsessions may lead patients to involve family members in their cleaning rituals (family accommodation) or may result in conflicts with them because their reluctance to participate in or accept these compulsions (high expressed emotion). Also, growing research points out that relatives of people with OCD have high rates of psychopathology (Maliken & Katz, 2013). Regarding parents with OCD, community and family studies have shown that their children are at greater risk of suffering from mental disorders in general when compared with parents without OCD. Current data indicate children have greater prevalence of obsessive-compulsive spectrum disorders, major depression, autism, schizophrenia, and anorexia nervosa (Black et al., 2003; Browne et al., 2015; Meier et al., 2014; Meier, Bulik, et al., 2015; Meier, Peersen, et al., 2015). Also, research focused on the assessment of emotional symptoms has shown that children of parents with OCD are at higher risk of having internalizing, but not externalizing symptoms (Black et al., 2003; Boyd & Tervo-Clemmens, 2013; Challacombe & Salkovskis, 2009). However, these findings have been methodologically limited because they were mostly found when compared with control groups without any other primary psychiatric disorder. Unlike healthy controls, parents with other psychiatric disorders may exhibit emotional symptoms (depression, anxiety), which are also usually observed in parents with OCD beyond their primary diagnosis (Goodwin, 2015). Hence, by including parents with other psychiatric disorders as control groups, the findings obtained could be more specifically linked to parental OCD because other overlapping symptoms may be controlled. Ultimately, the need to address emotional symptoms in children of parents with OCD is warranted because psychological problems are likely to be subtle or subclinical across childhood, possibly manifesting as diagnosable mental disorder only later in their lives (Liu, 2016).




Figure 1
CONSORT Flow Diagram.

As part of this issue, the intergeneration transmission of psychopathology from parents with OCD to children through genetic heritance and/or environmental factors is still a matter of debate. On the one hand, there is preliminary evidence from family studies supporting the presence of genetic load for intergeneration transmission of OCD through altered glutamatergic transmission (Pauls, 2010). On the other hand, the few empirical studies focused on the influence of environmental factors have shown that parenting attitudes among parents with OCD did not influence their childrens risk for OCD or that this association was only evidenced for mothers with OCD (vs. fathers with OCD; Black et al., 2003; Wilcox et al., 2008). In spite of the lack of firm evidence, there is still a need for delineating the link between parenting attitudes in parents with OCD and the risk of psychopathology on their children, including but not limited to childhood OCD. First, because Developmental Psychology in general and attachment theories in particular stress that parenting attitudes represent a key cognitive predisposition underlying child-rearing and childrens emotional development (Leavitt, 1999). Second, because several observational studies have found that specific parenting attitudes (e.g., overprotection) were more prevalent among mothers with OCD when compared with healthy controls, although group differences were partly accounted for by comorbid major depression or general anxiety levels (Challacombe et al., 2016; Challacombe & Salkovskis, 2009; Reich, 2008). Accordingly, there could be only some accurate features of this broad construct linked to parental OCD. This is relevant taking into account that parenting attitudes represents a broad concept covering several aspects such as satisfaction with parenting, involvement, discipline or autonomy (Gerard, 1994). Third, it is also needed because previous studies in other psychiatric samples have reported that parenting attitudes in parents with major depression, borderline personality disorder, or eating disorders may negatively affect childrens emotional development, and vice versa (Bartsch et al., 2015; Tsypes & Gibb, 2015). Overall, by finding specific parenting attitudes in people with OCD and their influence on their childrens emotional symptoms, clinicians could tailor parenting therapies aimed at preventing or ameliorating this environmental risk factor (Challacombe et al., 2017; Reich, 2008).

Within this clinical framework, the main aims of this empirical, exploratory study were: i) to delineate whether children of parents with OCD had significantly greater emotional symptoms when compared with children of parents without OCD; ii) to elucidate whether parents with OCD had significantly poorer parenting attitudes when compared with parents without OCD if so, to verify this after controlling for parents

and childrens differences in severity of depression and anxiety; and iii) within parents with OCD, to determine whether there were any OCD features significantly associated with any emotional symptoms in their children. If so, to elucidate the mediation role of any parenting attitudes on this relationship.

Method

Participants

Detailed data on the selection process is depicted in Figure 1. Forty Caucasian parents with a current primary diagnosis of OCD were consecutively recruited from the Adult Outpatient Mental Health Center of Mataró (Barcelona) from January 2015 to September 2016. A control group composed of 37 Caucasian parents with a current primary diagnosis of adjustment disorder and matched for sociodemographic variables was also collected from this clinical setting. A comparison group of parents with adjustment disorders was chosen because our purpose was to delineate whether parenting attitudes in people with OCD could be disorder-specific. For this reason, and following Challacombe and Salkovskis (2009), we included another clinical group as a control condition. Particularly, a diagnosis of an adjustment disorder represents a broad clinical category of mixed symptoms (e.g., anxiety, depression), which are also usually observed in people with OCD and may be controlled in order to exclude symptom-specific parenting attitudes. The current main diagnoses for both clinical groups were based on the fourth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV; APA, 1994).

Inclusion criteria for both groups were: 1) parents should be primary caregivers and have at least one targeted child living partly or fully with them; 2) children assessed should be aged 7-16 in order to meet age criteria for the questionnaires noted below (see Assessment Instruments). Exclusion criteria were: 1) for both groups, prior diagnoses of neurological disorder, intellectual disability, autism, or schizophrenia according to chart reviews; 2) for both groups, current (hypo)manic episode and/or substance intoxication/withdrawal; 3) for the non-OCD group, a lifetime diagnosis of OCD; 4) for the OCD group, a lifetime diagnosis of adjustment disorder.

Patients from both groups received naturalistic treatment consisting of individual bimonthly cognitive-behavioral therapy implemented by their clinical psychologists. Also, antidepressant prescriptions were as follows when recruited: for the OCD-group, 26 participants took selective serotonin reuptake inhibitors (SSRIs) and 19 received tricyclic antidepressants; for the non-OCD group, 31 took SSRIs, two received tricyclic antidepressants, and six were prescribed selective serotonin noradrenaline reuptake inhibitors.

Patients were initially referred to us by their main clinicians (mostly psychiatrists) once they were initially diagnosed at intake with the

required mental disorders. All patients were from different families. Patients attended one assessment session conducted by the authors of the current research, on average 3.4 (1.7) months after their tenure with the treatment center. The authors are psychiatrists and clinical psychologists widely experienced in OCD. First, assessment interviews were held with each parent to ensure main diagnoses and exclude other current mental disorders. In all cases, diagnoses at intake were confirmed by the researchers using standardized assessment instruments. Then, each parent and his/her child separately completed self-report questionnaires in different rooms. The study was approved by the hospitals institutional review board, and informed consent was obtained from all parents after a full explanation of the nature of the study. Parental consent for children participation in the study was also obtained.

All procedures performed in studies involving human participants and/or animals were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Assessment Instruments

Parents assessment

Structured Clinical Interview for DSM-IV Axis I Disorders. The Spanish translation of the SCID-I/P (research version, patient edition) was used for diagnosing OCD and adjustment disorders. Also, it was employed for ruling out other current mental disorders according to the exclusion criteria (First et al., 1995).

Yale-Brown Obsessive-Compulsive Scale (Y-BOCS). Y-BOCS Spanish version was used to determine the severity and content of any current OC symptoms. Cronbachs alpha coefficients range from .87 to .94. A cut-off score > 15 for the total Y-BOCS is considered clinically significant (Vega et al., 2002).

Parent-Child Relationship Inventory (PCRI). PCRI's Spanish version was used. This is a self-reported questionnaire which was used to measure parental attitudes towards parenting. It comprises 78 items with four response options (totally disagree, disagree, agree, and strongly agree), which are grouped into seven subscales. The seven subscales are support, satisfaction with parenting, involvement, communication, discipline, autonomy, and role orientation, along with a social desirability subscale. Cronbachs alpha coefficients range from .68 to .78 (Roa & Del Barrio, 2001).

Spielberger State-Trait Anxiety Inventory (STAI- S/T). STAI-S/T's Spanish validation was used to assess anxiety severity. Cronbachs alpha coefficients are .87 and .93. A cut-off score of 39-40 has been suggested to detect clinically significant symptoms for the STAI subscales (Spielberger et al., 1982).

Table 1
Sociodemographic Characteristics of Parents with and without OCD and Their Children

Variables	OCD (n = 40)	Non-OCD (n = 37)	Statistics	df	p-value
	n (%)	n (%)			
Age	M = 44.46 (6.68)	M = 42.45 (4.36)	t = -1.46	75	.14
Gender (mothers)	30 (75.0)	28 (75.67)	$\chi^2 = 0.85$	1	.88
Level of education					
Primary	3 (7.5)	2 (5.4)	$\chi^2 = 1.12$	2	.65
Secondary	33 (82.5)	31 (83.8)			
University	4 (10.0)	4 (10.8)			
Marital Status					
Living with the other parent	31 (77.5)	30 (81.08)	$\chi^2 = 1.95$	2	.49
Separated/divorced and living without couple	4 (10.0)	5 (13.51)			
Separated/divorced and living with another couple	5 (12.5)	2 (5.4)			
Occupation					
Employee	18 (45.0)	13 (35.13)	$\chi^2 = 2.99$	2	.16
Unemployed	18 (45.0)	23 (62.16)			
Retired due to disability	4 (10.0)	1 (2.7)			
No. children at home	M = 1.65 (0.56)	M = 1.75 (0.43)	t = 0.88	75	.33
No. children at home aged 7-16	46	40	t = -0.76	75	.37
Age	M = 13.43 (2.94)	M = 12.38 (2.52)	t = -1.77	75	.08
Gender (daughters)	19 (47.5)	17 (45.95)	$\chi^2 = 0.79$	1	.66

Table 2
Clinical Comparison between OCD and non-OCD Parents

	OCD (n = 40)	Non-OCD (n = 37)	Statistics (F)	p-value	Effect size partial χ^2
	M (SD)	M (SD)			
BDI	17.22 (3.93)	20.95 (3.51)	3.38	.05	.04
STAI-S	37.22 (4.04)	30.53 (3.49)	9.90	.002	.10
STAI-T	39.35 (3.14)	31.95 (3.01)	3.83	.04	.05

Note. BDI = Becks Depression Inventory; OCD = obsessive-compulsive disorder;
SD = standard deviation; STAI = Spielberger State-Trait Anxiety Inventory.

Beck Depression Inventory (BDI). The Spanish validation of the 13-item BDI short form was used to assess depression symptom severity. Cronbachs alpha coefficient was .90. Adults with a cut-off score of 13/14 are considered at risk for clinical depression (Conde & Useros, 1975).

Childrens assessment

Childrens Depression Inventory (CDI). CDIs Spanish validation was used to assess depression symptom severity. This self-reported questionnaire provides a total score based on 27 items scored according to one of the following alternatives: 0 = absence of the symptom, 1 = moderate symptom, 2 = severe symptom. Youth with cut- off score 13 are considered at risk for clinical depression. Cronbachs alpha coefficient was .82 (Del Barrio, 1999).

Revised Childrens Manifest Anxiety Scale (R-CMAS). R-CMAS Spanish validation, testing anxiety symptom severity, was used. This self-reported questionnaire consists of 37 items. A total anxiety score is based on 28 items, which are in turn divided into three anxiety subscales: physiological anxiety, restless/hypersensitivity and social concerns/concentration. Also, there are nine additional items that corresponded to a Lie subscale. Score of each subscale was determined by the number of items answered with yes (= 1). Youth with a cut-off score of > 18 for the

total anxiety are considered at risk for anxiety disorders. Cronbachs alpha ranges between .79 and .85 (Reynolds, 1997).

Piers-Harris Childrens Self-Concept Scale (PH-CSCS). PH-CSCS Spanish validation was used to assess self-esteem. This self-reported questionnaire consists of 80 items, each scored 1 (yes) or 0 (no). The item scores are summed to yield a total score and six factor scores: physical appearance, intellectual and school status, freedom from anxiety, behavior, happiness and satisfaction, and popularity. Higher scores for total self-concept and factor scores reflect a more positive self-concept. Cronbachs alpha ranges between .75 and .83 (Cardenal & Fierro, 2003).

Statistical Analysis

Chi-square test and *t*-test were first used to analyze the sociodemographic data. Then the Shapiro-Wilk test was used to confirm the normal distribution of quantitative clinical data for both groups. Also, Levenes test for homogeneity of variance was performed before testing for between-group differences on clinical data.

One-way multivariate analysis of variance (MANOVA) was used to test for significant differences between both groups, either in parents anxiety/depression symptom severity (STAI-S/T, BDI) or in their childrens emotional symptoms (CDI, R-CMAS, PH-CSCS). Then one-way multivariate analyses of covariance (MANCOVA) was conducted to look for between-group differences in parenting attitudes (PCRI), after controlling for general anxiety levels (STAI-S/T), depression symptom severity (BDI), and emotional symptoms in their children (CDI, R-CMAS, PH-CSCS).

Within the OCD group, hierarchical linear regression analyses were then conducted to determine the mediating role of parenting attitudes (PCRI) in the relationship between OCD features (Y-BOCS) and emotional symptoms in their children (CDI, total score from the R-CMAS and the PH-CSCS). Mediation analysis followed the method by Baron and Kenny (1986), which has usually provided the same results compared with other tests (e.g., Sobel test) used to assess indirect effects. The regression analyses at step 1 included each OCD feature (Y-BOCS) as a univariate predictor and each emotional symptom as a dependent variable (CDI, total score from the R-CMAS, and the PH-CSCS). The regression analyses at step 2 included each OCD feature (Y-BOCS) as a univariate predictor and each parenting attitude (PCRI) as a dependent variable. At step 3, bivariate regression analyses included several models resulting from the interaction between each significant OCD feature (Y-BOCS) at both prior steps and each parenting attitude (PCRI). Emotional symptoms (CDI, total score of the R-CMAS and the PH-CSCS) were dependent variables. A particular model would fully support the mediation hypothesis if the OCD feature (Y-BOCS) was no longer significant while the parenting attitude (PCRI) was significant. At step 4, multiple regression analyses including parents psychopathology (BDI, STAI-S) were performed to further ensure the mediation model.

Standardized regression coefficients (β) were used as measures of effect sizes for stepwise linear regression analyses, while partial eta squared (η^2) were employed for MANOVA and MANCOVA. Statistical analyses were performed using the Statistical Package for the Social Sciences (SPSS 18.0). All tests were two-tailed and p -values of less than .05 were considered statistically significant.

Results

Sociodemographic Features of Parents with and without OCD and Their Respective Children

As detailed in Table 1, there were no statistically significant differences in sociodemographic features between parents with OCD ($n_1 = 40$) and non-OCD ($n_2 = 37$) and between their respective children ($p > .05$). Basically, both groups consisted of middle-aged parents, predominantly female, who lived with the other parent. Over half of both parents groups were unemployed at the time of the assessment. The targeted samples of children ($n_1 = 46$ and $n_2 = 40$) were mostly in early adolescence and equally composed by gender.

Clinical Comparison between Parents with OCD and without OCD

Concerning parents psychopathology, the MANOVA reported a statistically significant difference between groups ($F = 14.56$, $df = 13, 63$, $p < .001$, Wilks = .65). As shown in Table 2, univariate analyses showed that depression symptom severity (BDI) was statistically significantly greater in the non-OCD group ($F = 3.38$, $p = .05$), while general anxiety severity (STAI-S/T) was statistically significantly higher in the OCD group ($F = 9.9$, $p = .02$; $F = 3.83$, $p = .04$). For both groups, raw scores on depression severity were clinically significant, while anxiety scores ranged on subclinical intervals.

Table 3
Clinical Comparison between Children of OCD and Non-OCD Parents

Variables	OCD ($n = 46$)	Non-OCD ($n = 40$)	Statistics (F)	p -value	Effect size (partial η^2)
	M (SD)	M (SD)			
CDI	10.22 (1.98)	4.95 (1.93)	65.8	< .001	.43
R-CMAS					
Total anxiety	11.17 (3.37)	6.25 (3.05)	23.42	< .001	.21
Physiological anxiety	4.18 (1.66)	3.63 (2.52)	8.88	.004	.09
Restless/hypersensitivity	6.30 (2.25)	2.1 (2.02)	81.59	< .001	.49
Social concerns/concentration	1.91 (1.60)	0.53 (0.5)	27.56	< .001	.24
Lies	3.48 (1.99)	2.48 (0.5)	9.54	.003	.10
Self-concept (PCRI)					
Overall self-concept	56.91 (5.36)	62.55 (5.10)	11.29	.001	.11
Physical appearance	8 (2.45)	9.95 (1.01)	21.91	< .001	.21
Intellectual and school status	10.3 (2.62)	12.43 (1.51)	20.23	< .001	.19
Freedom from anxiety	7.13 (1.84)	8.28 (1.55)	6.36	.01	.07
Behaviour	14.3 (1.64)	16.48 (.50)	64.29	< .001	.43
Happiness and satisfaction	7.13 (1.77)	7.95 (1.01)	6.66	.01	.07
Popularity	8.39 (0.97)	9.48 (.50)	7.23	.01	.08

Note. BDI = Becks Depression Inventory; OCD = obsessive-compulsive disorder; PCRI, = Parent-Child Relationship Inventory; STAI = Spielberger State-Trait Anxiety Inventory.

Table 4
Comparison between OCD and Non-OCD Parents in Parenting Attitudes, after Controlling for Parents and Childrens Covariates

Variables	OCD (n= 40)	Non-OCD (n= 37)	Statistics (F)	p-value	partial η^2
	M (SD)	M (SD)			
Parenting attitudes (PCRI)					
Support	17.65 (5.03)	29.9 (2.02)	26.50	< .001	.28
Satisfaction	30.57 (5.28)	39.05 (1.01)	63.98	< .001	.48
Involvement	35.13 (2.30)	47.10 (2.02)	161.38	< .001	.70
Communication	25.43 (4.17)	32.48 (1.50)	42.97	< .001	.38
Discipline	25.39 (2.39)	39.53 (1.37)	56.87	< .001	.45
Autonomy	26.22 (2.52)	31.77 (3.55)	10.35	.02	.13
Role orientation	23.30 (3.82)	29.95 (1.01)	260.90	< .001	.79
Social desirability	14.22 (2.27)	12.22 (0.89)	1.36	.24	.02

Note. OCD = obsessive-compulsive disorder; PCRI = Parent-Child Relationship Inventory.

Clinical Comparison between Children from Parents with OCD and without OCD

Regarding childrens emotional symptoms, the MANOVA reported a statistically significant difference between groups ($F = 37.95$, $df = 13$, 72 , $p < .001$, Wilks = .12). As detailed in Table 3, univariate analyses showed that depression symptom severity (CDI) and anxiety levels (R-CMAS) were statistically significantly higher in children of parents with OCD compared to children of parents without OCD ($p < .05$). Also, children of parents with OCD reported statistically significantly lower self-esteem (PH-CSCS; $p < .05$). None of the outcome measures were clinically significant for either group, but scores on depression and anxiety symptom severity were at subclinical levels among children of parents with OCD.

Comparison between Parents with and without OCD in Parenting Attitudes

Concerning parenting attitudes, the MANCOVA reported a statistically significant difference between groups, after controlling for parents and childrens emotional symptoms (BDI, STAI-S/T, CDI, R-CMAS, PH-CSCS; $F = 430.92$, $df = 8$, 66 , $p < .001$, Wilks = .01). As detailed in Table 4, univariate analyses showed that, except for social desirability, the remaining parenting attitudes were statistically significantly poorer in parents with OCD relative to parents without OCD ($p < .05$).

Table 5
Hierarchical Linear Regression Analyses for Those Models Fully Meeting
Mediation Role of Parenting Attitudes in the Relationship between Parents
OCD Features ($n = 40$) and Childrens Emotional Development ($n = 46$)

Model	Step	Outcome variable	Predictor variables	β	t	p	Adjusted R^2
1	Step 1	Anxiety severity (R-CMAS)	Sexual obsessions (Y-BOCS)	.31	2.15	.03	.09*
	Step 2	Parental Involvement (PCRI)	Sexual obsessions (Y-BOCS)	.33	2.34	.02	.11*
	Step 3	Anxiety severity (R-CMAS)	Sexual obsessions (Y-BOCS)	.19	1.37	.17	.19*
			Parental Involvement (PCRI)	.33	2.24	.03	
	Step 4	Anxiety severity (R-CMAS)	Sexual obsessions (Y-BOCS)	.23	1.50	.14	.21*
			Parental Involvement (PCRI)	.44	2.46	.02	
			Depression severity (BDI)	.27	1.10	.27	
			General anxiety (STAI-S)	.25	1.03	.30	
2	Step 1	Anxiety severity (R-CMAS)	Somatic obsessions (Y-BOCS)	.30	2.09	.04	.09*
	Step 2	Parental Involvement (PCRI)	Somatic obsessions (Y-BOCS)	.32	2.26	.02	.16*
	Step 3	Anxiety severity (R-CMAS)	Somatic obsessions (Y-BOCS)	.19	1.34	.19	.19*
			Parental Involvement (PCRI)	.33	2.28	.03	
	Step 4	Anxiety severity (R-CMAS)	Somatic obsessions (Y-BOCS)	.22	1.44	.16	.21*
			Parental Involvement (PCRI)	.43	2.29	.03	
			Depression severity (BDI)	.20	0.80	.43	
			General anxiety (STAI-S)	.26	1.08	.29	

Note. OCD = obsessive-compulsive disorder. Parents measures: BDI = Becks Depression Inventory; PCRI = Parent-Child Relationship Inventory; STAI-S = Spielberger Trait-State Anxiety Inventory; Y-BOCS = Yale-Brown Obsessive-Compulsive Scale. Childrens measures: R-CMAS = Revised Children Manifest Anxiety Scale.

* $p < .05$

Mediation Role of Parenting Attitudes on the Relationship between Parents OCD Features and Children 's Emotional Symptoms

Initial descriptive analyses showed that parents with OCD reported clinically significant OCD symptoms (Y-BOCS total, $M = 22.3$, $SD = 4.91$). Over half of them recognized having either contamination (60%) or symmetry obsessions (60%). Also, parents with OCD usually presented cleaning (65%), arranging (60%), and repetition compulsions (50%).




Figure 2

Full Mediation Model: Effect of Parents Sexual Obsessions on
Childrens Total Anxiety Severity through Parental Involvement, after
Controlling for Parents General Anxiety/Depression Symptoms.

Note. BDI = Becks Depression Inventory; CMAS-R = Revised Childrens Manifest Anxiety Scale;
PCRI = Parent-Child Relationship Inventory; STAI-S = Spielberger State-Trait Anxiety Inventory;
Y-BOCS = Yale-Brown Obsessive-Compulsive Scale. The figure in parenthesis refers to , * $p < .05$.

Hierarchical linear regression analysis was performed to test whether any parenting attitude (PCRI) may mediate the association between parents OCD features (Y-BOCS) and childrens emotional symptoms (CDI, R-CMAS, PH-CSCS). Table 5 shows each step for those models fully meeting the mediation hypothesis. At step 1, the presence of aggressive and somatic obsessions was found to significantly predict both childrens depression ($= .53, p < .001$; $= .42, p = .003$) and total anxiety severity ($= .43, p = .003$; $= .30, p = .04$). The presence of sexual obsessions also significantly predicted childrens total anxiety severity ($= .31, p = .03$).

Neither content nor severity of compulsions were significantly associated with any emotional symptom in children ($p > .05$). Also, childrens self-esteem was not significantly predicted by any OCD feature in parents ($p > .05$). At step 2, the presence of sexual and somatic obsessions was found to significantly predict parental involvement ($= .33, p = .02$; $= .32, p = .02$). Unlike in step 1, aggressive obsessions did not significantly predict any parenting attitude ($p > .05$). At step 3, models including childrens depression severity as a dependent variable did not meet the full mediation hypothesis. However, sexual and somatic obsessions were no longer significant predictors of childrens total anxiety ($= .19, p = .17$; $= .19, p = .19$) when parental involvement was included in both models ($= .32, p = .03$; $= .33, p = .02$). At step 4, both full mediation models were confirmed after controlling for depression and anxiety severity in parents with OCD (Figure 2 and 3).



Figure 3

Full Mediation Model: Effect of Parents Somatic Obsessions on Childrens Total Anxiety Severity through Parental Involvement, after Controlling for Parents General Anxiety/Depression Symptoms.

Note. BDI = Becks Depression Inventory; CMAS-R = Revised Childrens Manifest Anxiety Scale; PCRI = Parent-Child Relationship Inventory; STAI-S = Spielberger State- Trait Anxiety Inventory; Y-BOCS = Yale-Brown Obsessive-Compulsive Scale.

Discussion

This study sought to delineate distinctive parenting attitudes in people with obsessive-compulsive disorder (OCD), specific emotional symptoms in their children, and the association between them. Main findings from the current research showed: 1) children of parents with OCD reported significantly greater anxiety and depression symptoms when compared with children of parents diagnosed with adjustment disorders; 2) after controlling for parents and childrens emotional symptoms, parents with OCD reported significantly poorer overall parenting attitudes relative to parents with adjustment disorders; and 3) within the OCD sample, parental involvement mediated the relationship between sexual/somatic obsessions in parents and anxiety in their children.

Similar to other previous studies, the current research showed that children of parents with OCD reported greater presence of internalizing/emotional symptoms than children of control parents (Black et al., 2003; Boyd & Tervo-Clemmens, 2013; Challacombe & Salkovskis, 2009). This finding may be considered disorder-specific, taking into account that we included a control group composed of parents with a different primary psychiatric disorder who also exhibited depression and anxiety symptoms. Unlike the current study, the research by Challacombe and Salkovskis (2009) did not reveal significant differences in internalizing/emotional

symptoms between children of parents with OCD and children of parents with panic disorder. Regarding the latter study, it is likely that some clinical differences between both psychiatric groups were overlooked because of the small sample sizes ($n < 24$). Apart from this finding, the fact that children of parents with OCD reported subclinical levels for these internalizing/emotional symptoms may also suggest the need to carry out longitudinal assessment in order to identify the subset of children who may eventually develop clinically significant symptoms later on (Liu, 2016).

In accordance with prior research, the current study found that parenting attitudes were poorer in parents with OCD than controls (Challacombe et al., 2016; Challacombe & Salkovskis, 2009; Farrell et al., 2013). However, poor parenting was not limited to specific attitudes. It should be noted that methodological divergences between the current and prior research may account for differences in parenting. Unlike these other studies, our study consisted of not only mothers with OCD, but also fathers with OCD. In addition, our samples of children were older than those used in the other studies. Because of these divergences, further research is warranted aimed at addressing parenting differences by parents gender and childrens age. Also, forthcoming studies should take into account the role of the other parent or caregiver, if present. Regarding this issue, it is likely that some clinical characteristics closely associated with couples (e.g., low marital support) may moderate parenting attitudes among parents with OCD, for example by increasing over-involvement towards children (Taraban et al., 2017). Also, the presence of adequate parenting attitudes (e.g., warmth, optimism, distress tolerance) in the other parent or caregiver may serve as a buffer against poor parenting attitudes in the parent with OCD (Rosa-Alcázar et al., 2019).

Interestingly, when analyzing subtypes of OCD symptoms in parents with OCD, obsessions rather than compulsions were associated with internalizing/emotional symptoms in their children. Specifically, the association between somatic/sexual obsessions in parents with OCD and anxiety symptoms in their children was mediated by parental involvement. To the best of our knowledge, this is the first empirical study which shows the extent to which specific OCD symptoms in parents may predict emotional symptoms in their children. The fact that parenting involvement acted mostly as a mediator variable suggests that the presence of intrusive and unwanted thoughts, commonly associated with their children (e.g., fear of disease contagion or sexual assault), may predispose parents with OCD to higher hyper-responsibility and over-involvement in terms of parenting (Abramowitz et al., 2003). As a result, children may be exposed to a modelling of anxious behavior by parents, which in turn may sensitize children to anxious reactions throughout their lives (Murray et al., 2009). Also, following cognitive models, this pernicious effect within OCD dyads may enhance childrens personal sense of responsibility and an inflated threat perception in their lives (Farrell et al., 2013; Pietrefesa et al., 2010). Ultimately, the aforementioned relationship should be addressed even in mothers with OCD in the

perinatal (postpartum) period, where the presence of intrusive thoughts is highly prevalent (Collardeu et al., 2019).

The current study has several limitations. First, both groups had relatively small sample sizes, though similar to other related studies (Challacombe et al., 2016). In spite of this fact, effect sizes were usually large, which added substantive relevance to these findings. Second, assessment of parenting attitudes relied on a self-report questionnaire instead of using observational methods by independent raters within parent-child dyads (Challacombe & Salkovskis, 2009). To attenuate this limitation, confounding effects from parents depression and anxiety symptomatology were controlled for when analyzing data on parenting. Third, data from categorical diagnoses were not obtained for children cohorts (e.g., OCD) as well as for comorbid disorders in both parents groups (e.g., major depression disorder), except for exclusion criteria. Hence, our results are particularly relevant for dimensional approaches to psychopathology. Fourth, the study sample was a convenience sample of individuals being served by one particular clinic and cannot be seen as definitive nor generalizable to all parenting adults with OCD. Fifth, the choice of the comparison group (parents with an adjustment disorder), whose clinical symptoms were broad and self-limited by definition, limits the extent to which our results on parenting among OCD parents may be assumed to be disorder-specific. Sixth, the cross-sectional nature of the study impedes establishing causal relationships between variables. Hence, longitudinal studies should be prioritized in forthcoming research.

In sum, parents with OCD may exhibit overall poor parenting attitudes, which could be disorder-specific and independent of other co-occurring symptoms. Specifically, some parents' obsessions may predispose children to greater levels (subclinical) of anxiety severity through parental involvement. Clinical implications from these findings are: 1) clinicians should assess parenting attitudes in those patients with OCD that are periodically treated with cognitive-behavior therapy; 2) the presence of intrusive thoughts in parents with OCD, specially somatic/sexual obsessions, should be prioritized as a target intervention, not only for own patients, but also for their children's well-being (Reich, 2008); 3) children of parents with OCD at high-risk for anxiety symptoms should be assessed at least in order to prevent the presence of more relevant psychopathology at mid/long-term; and 4) parenting interventions either in individual or parent-child format should be developed by implementing clinical trials in patients with OCD.

References

- Abramowitz, J. S., Schwartz, S. A., Moore, K. M., & Luenzmann, K. R. (2003). Obsessive-compulsive symptoms in pregnancy and the puerperium: A review of the literature. *Journal of Anxiety Disorders*, 17, 461-478. [https://doi.org/10.1016/S0887-6185\(02\)00206-2](https://doi.org/10.1016/S0887-6185(02)00206-2)
- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). American Psychiatric Association.

- Baron, R. M., & Kenny, D. A. (1986). The moderator-mediator variable distinction in social psychological research conceptual, strategic, and statistical considerations. *Journal of Personality & Social Psychology*, 51, 1173-1182. <https://doi.org/10.1037/0022-3514.51.6.1173>
- Bartsch, D. R., Roberts, R. M., Davies, M., & Proeve, M. (2015). The impact of parental diagnosis of borderline personality disorder on offspring: Learning from clinical practice. *Personality and Mental Health*, 9, 33-43. <https://doi.org/10.1002/pmh.1274>
- Black, D. W., Gaffney, G. R., Schlosser, S., & Gabel, J. (2003). Children of parents with obsessive-compulsive disorder a 2-year follow-up study. *Acta Psychiatrica Scandinavica*, 107, 305-313. <https://doi.org/10.1034/j.1600-0447.2003.02182.x>
- Boyd, R. C., & Tervo-Clemmens, B. (2013). Exploring maternal and child effects of comorbid anxiety disorders among African American mothers with depression. *Journal of Depression & Anxiety*, 2(129). <https://doi.org/10.4172/2167-1044.1000129>
- Browne, H. A., Hansen, S. N., Buxbaum, J. D., Gair, S. L., Nissen, J. B., Nikolajsen, K. H., Schendel, D. E., Reichenberg, A., Parner, E. T., & Grice, D. E. (2015). Familial clustering of tic disorders and obsessive-compulsive disorder. *JAMA Psychiatry*, 72, 359-366. <https://doi.org/10.1001/jamapsychiatry.2014.2656>
- Cardenal, V., & Fierro, A. (2003). Componentes y correlatos del autoconcepto en la escala de Piers-Harris. *Estudios de Psicología*, 24, 101-111. <https://doi.org/10.1174/021093903321329094>
- Challacombe, F., & Salkovskis, P. (2009). A preliminary investigation of the impact of maternal obsessive-compulsive disorder and panic disorder on parenting and children. *Journal of Anxiety Disorders*, 23, 848-857. <https://doi.org/10.1016/j.janxdis.2009.04.002>
- Challacombe, F. L., Salkovskis, P. M., Woolgar, M., Wilkinson, E. L., Read, J., & Acheson, R. (2016). Parenting and mother-infant interactions in the context of maternal postpartum obsessive-compulsive disorder: Effects of obsessional symptoms and mood. *Infant Behavior and Development*, 44, 11-20. <https://doi.org/10.1016/j.infbeh.2016.04.003>
- Challacombe, F. L., Salkovskis, P. M., Woolgar, M., Wilkinson, E. L., Read, J., & Acheson, R. (2017). A pilot randomized controlled trial of time-intensive cognitive-behaviour therapy for postpartum obsessive-compulsive disorder: Effects on maternal symptoms, mother-infant interactions and attachment. *Psychological Medicine*, 47(8), 1478-1488. <https://doi.org/10.1017/S0033291716003573>
- Cicek, E., Cicek, I.E., Kayhan, F., Uguz, F., & Kaya, N. (2013). Quality of life, family burden and associated factors in relatives with obsessive-compulsive disorder. *General Hospital Psychiatry*, 35, 253-258. <https://doi.org/10.1016/j.genhosppsych.2013.01.004>
- Collardeu, F., Corbyn, B., Abramowitz, J., Janssers, P.A., Woody, S., & Fairbrother, N. (2019). Maternal unwanted and intrusive thoughts of infant-related harm, obsessive-compulsive disorder and depression in the perinatal period: Study protocol. *BMC Psychiatry*, 19, 94. <https://doi.org/10.1186/s12888-019-2067-x>

- Conde, V., & Useros, E. (1975). Spanish adaptation of the Becks depression behavioural assessment scale. *Revista de Psiquiatría y Psicología Médica de Europa y América Latina*, 12, 217-236.
- Del Barrio, V. (1999). El Childrens Depression Inventory [CDI; Kovacs, 1992]. Su aplicación en población española. *Clinica y Salud*, 10, 393-416.
- Farrell, L. J., Hourigan, D., & Waters, A. M. (2013). Do mothers enhance responsibility in children with obsessive compulsive disorder? A preliminary study of mother-child interactions during a problem solving discussion. *Journal of Obsessive-Compulsive and Related Disorders*, 2, 78-84. <https://doi.org/10.1016/j.jocrd.2012.12.001>
- First, M. B., Spitzer, R. L., Gibbon, M., & Williams, J. B. W. (1995). *Structured Clinical Interview for DSM-IV (SCID-I)*. Research version. Biometrics Research Department, New York Psychiatric Institute. <https://doi.org/10.1037/t07827-000>
- Goodwin, G. M. (2015). The overlap between anxiety, depression, and obsessive-compulsive disorder. *Dialogues in Clinical Neuroscience*, 17, 249-260.
- Gomes, J. B., Cordioli, A. V., Bortoncello, C. F., Braga, D. T., Gonçalves, F., & Heldt, E. (2016). Impact of cognitive-behavioral group therapy for obsessive-compulsive disorder on family accommodation: A randomized clinical trial. *Psychiatry Research*, 246, 70-76. <https://doi.org/10.1016/j.psychres.2016.09.019>
- Leavitt, L. A. (1999). Research to practice: emotional development and maternal/infant attachment. *Journal of Pediatric Health Care*, 13, S4-7. [https://doi.org/10.1016/S0891-5245\(99\)90048-2](https://doi.org/10.1016/S0891-5245(99)90048-2)
- Lebowitz, E. R., Panza, K. E., Su, J., & Bloch, M. H. (2012). Family accommodation in obsessive-compulsive disorder. *Expert Review of Neurotherapeutics*, 12, 229- 238. <https://doi.org/10.1586/ern.11.200>
- Liu, R. T. (2016). Taxometric evidence of a dimensional latent structure for depression in an epidemiological sample of children and adolescents. *Psychological Medicine*, 46, 1265-1275. <https://doi.org/10.1017/S0033291715002792>
- Maliken, A. C., & Katz, L. F. (2013). Exploring the impact of parental psychopathology and emotion regulation on evidence-based parenting interventions: A transdiagnostic approach to improving treatment effectiveness. *Clinical Child and Family Psychology Review*, 16, 173-186. <https://doi.org/10.1007/s10567-013-0132-4>
- Meier, S. M., Bulik, C. M., Thornton, L. M., Mattheisen, M., & Mortensen, P. B. (2015). Diagnosed anxiety disorders and the risk of subsequent anorexia nervosa: A Danish population register study. *European Eating Disorders Review*, 23, 524-530. <https://doi.org/10.1002/erv.2402>
- Meier, S. M., Petersen, L., Pedersen, M. G., Arendt, M. C., Nielsen, P. R., Mattheisen, M., Mors, O., & Mortensen, P. B. (2014). Obsessive-compulsive disorder as a risk factor for schizophrenia: A nationwide study. *JAMA Psychiatry*, 71, 1215-1221. <https://doi.org/10.1001/jamapsychiatry.2014.1011>
- Meier, S. M., Petersen, L., Schendel, D. E., Mattheisen, M., Mortensen, P. B., & Mors, O. (2015). Obsessive-compulsive disorder and autism spectrum disorders: Longitudinal and offspring risk. *PLoS One*, 10, e0141703. <https://doi.org/10.1371/journal.pone.0141703>

- Murray, L., Creswell, C., & Cooper, P. J. (2009). The development of anxiety disorders in childhood: an integrative review. *Psychological Medicine*, 39, 1413-1423. <https://doi.org/10.1017/S0033291709005157>
- Pauls, D. L. (2010). The genetics of obsessive-compulsive disorder: A review. *Dialogues in Clinical Neurosciences*, 12, 149-163.
- Pietrefesa, A. S., Schofield, C. A., Sochting, I., & Coles, M. E. (2010). Obsessive beliefs in youth with OCD and their mothers. *Journal of Cognitive Psychotherapy*, 24, 187-197. <https://doi.org/10.1891/0889-8391.24.3.187>
- Reich, G. (2008). Family dynamics and family therapy of obsessive-compulsive disorder. *Praxis Der Kinderpsychologie Und Kinderpsychiatrie*, 57, 486-498. <https://doi.org/10.13109/prkk.2008.57.6.486>
- Reynolds, R. (1997). *Escala de Ansiedad Manifiesta en Niños, R-CMAS*. Manual Moderno.
- Roa, L., & Del Barrio, V. (2001). Adaptation of the Parent Child Relationship Inventory (PCRI-M) for Spanish population. *Revista Latinoamericana de Psicología*, 33, 329-341.
- Rosa-Alcázar, Á., Parada-Navas, J. L., Olivares-Olivares, P. J., Bernal Ruiz, C., & Rosa-Alcázar, A. I. (2019). Estilos educativos parentales y emociones como predictores de respuestas obsesivo-compulsivas en población adolescente. *Terapia psicológica*, 37, 241-254. <https://doi.org/10.4067/S0718-48082019000300241>
- Spielberger, C. D., Gorsuch, R. R., & Lushene, R. E. (1982). *STAI: Cuestionario de Ansiedad Estado/Rasgo*. Técnicos Especialistas Asociados (TEA).
- Taraban, L., Shaw, D. S., Leve, L. D., Wilson, M. N., Dishion, T. J., Natsuaki, M. N., Neiderhiser, J. M., & Reiss, D. (2017). Maternal depression and parenting in early childhood: Contextual influence of marital quality and social support in two samples. *Developmental Psychology*, 53, 436-449. <https://doi.org/10.1037/dev0000261>
- Tsypes, A., & Gibb, B. E. (2015). Peer victimization mediates the impact of maternal depression on risk for suicidal ideation in girls but not boys: A prospective study. *Journal of Abnormal Child Psychology*, 43, 1439-1445. <https://doi.org/10.1007/s10802-015-0025-8>
- Vega, J. M., Sal, Y., Rosas, H. J., Mazzotti Suárez, G., Vidal, H., Guimas, B., Adrianzén, C., & Vivar, R. (2002). Validation of a version in Spanish of the Yale-Brown Obsessive-Compulsive Scale. *Actas Españolas de Psiquiatria*, 30, 30-35.
- Wilcox, H. C., Grados, M., Samuels, J., Riddle, M. A., Bienvenu, O. J. 3rd, Pinto, A., Cullen, B., Wang, Y., Shugart, Y. Y., Liang, K. Y., & Nestadt, G. (2008). The association between parental bonding and obsessive compulsive disorder in offspring at high familial risk. *Journal of Affective Disorders*, 111(1), 31-39. <https://doi.org/10.1016/j.jad.2008.01.025>

Notes

Para citar este artículo: Frías A., Palma C., Farriols N., Aliaga F., Navarro S. y Selves L (2020). Parenting attitudes in people with obsessive-compulsive disorder and emotional symptoms in their children. *Clínica y Salud*, 31(2), 91-98. <https://doi.org/10.5093/clysa2020a13>

Author notes

Correspondencia: afrias@csgm.cat (A. Frías).

Conflict of interest declaration

Conflict of Interest

The authors of this article declare no conflict of interest.