



Acta Scientiarum. Biological Sciences

ISSN: 1679-9283

ISSN: 1807-863X

actabiol@uem.br

Universidade Estadual de Maringá

Brasil

Duarte, Ana Carolina Oliveira; Oliveira, Cynthia de; Ramos, Sílvia Junio; Castro, Evaristo Mauro de; Siqueira, José Oswaldo; Guilherme, Luiz Roberto Guimarães  
Lanthanum content and effects on growth, gas exchanges, and chlorophyll index in maize plants  
Acta Scientiarum. Biological Sciences, vol. 40, 2018, January-December 2019  
Universidade Estadual de Maringá  
Brasil

DOI: <https://doi.org/10.4025/actascibiols.v40i1.38469>

Available in: <https://www.redalyc.org/articulo.oa?id=187158163042>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UDEM  redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative



## Lanthanum content and effects on growth, gas exchanges, and chlorophyll index in maize plants

Ana Carolina Oliveira Duarte<sup>1\*</sup>, Cynthia de Oliveira<sup>2</sup>, Sílvia Junio Ramos<sup>3</sup>, Evaristo Mauro de Castro<sup>1</sup>, José Oswaldo Siqueira<sup>3</sup> and Luiz Roberto Guimarães Guilherme<sup>2</sup>

<sup>1</sup>Departamento de Biologia, Universidade Federal de Lavras, Cx. Postal 3037, 37200-000, Lavras, Minas Gerais, Brasil. <sup>2</sup>Departamento de Ciência do Solo, Universidade Federal de Lavras, Minas Gerais, Brasil. <sup>3</sup>Instituto Tecnológico Vale-Desenvolvimento Sustentável, Nazaré, Belém, Pará, Brasil. \*Author for correspondence. E-mail: [acoliveiraduarte@gmail.com](mailto:acoliveiraduarte@gmail.com)

**ABSTRACT.** Rare earth elements, such as lanthanum (La), have been applied to agriculture via fertilizers, aiming to increase the productivity and crop quality, such results observed mainly in China. However, the knowledge about the effect of La on maize growth, as well as for other species, despite the growing interest it is still limited. The aim of this study was to evaluate the effects of La on maize growth, La content, photosynthetic rate and chlorophyll content in maize plants in response to La treatment (0, 25, 50, 100, 150, 300 and 600  $\mu$ M) in nutrient solution for three weeks. The plants were placed in germinated pots using a split-root technique. One of the pots in the germinated set was filled with a complete nutrient solution without La, while another was filled with a nutrient solution without phosphorus but containing different concentrations of La. It was verified that roots of maize plants can accumulate approximately sixty percent more La than shoots. Moreover, low La concentrations stimulated an increase in chlorophyll index, resulting in a slight increase in shoot biomass. At higher levels, La didn't reduce growth but caused a decrease in both photosynthetic rate and chlorophyll index.

**Keywords:** crops; photosynthesis; rare earth elements; *Zea mays*.

## Teor de lantânio e efeitos no crescimento, trocas gasosas e teor de clorofila em plantas de milho

**RESUMO.** Elementos terra rara, como lantânio (La), são aplicados à agricultura via fertilizantes, com o objetivo de aumentar a produtividade e a qualidade das culturas agrícolas, fato observado principalmente na China. Entretanto, o conhecimento sobre o efeito de La no crescimento do milho, bem como para outras culturas agrícolas, ainda é limitado apesar do interesse crescente. O presente estudo objetivou avaliar o efeito da aplicação de La sobre o crescimento, teor de La, fotossíntese e teor de clorofila em plantas de milho, expostas a concentrações crescentes de La (0, 25, 50, 100, 150, 300 e 600  $\mu$ M) em solução nutritiva por três semanas. As plantas foram colocadas em vasos geminados usando uma técnica de raiz divisória. Um dos potes no conjunto geminado foi preenchido com solução nutritiva completa que não continha La, enquanto outra foi preenchida com solução nutritiva sem fósforo, mas contendo diferentes concentrações de La. Foi verificado que as raízes das plantas de milho podem acumular sessenta por cento mais La quando comparado à parte aérea. Além disso, as baixas concentrações de La estimularam o aumento do teor de clorofila, resultando em um aumento na biomassa da parte aérea. Em concentrações mais elevadas, a aplicação de La não reduziu o crescimento das plantas de milho, mas causou diminuição na taxa fotossintética e no índice de clorofila.

**Palavras-chave:** culturas agrícolas; elementos terras raras; fotossíntese; *Zea mays*.

### Introduction

Rare earth elements (REE) are a group of seventeen chemical elements in the periodic table that exhibit similar chemical and physical properties (International Union of Pure and Applied Chemistry [IUPAC], 2005). Current uses of REE include automotive and petroleum refining catalysts, rechargeable batteries for hybrid and electric vehicles, and numerous medical and electronic

devices (Alonso et al., 2012, Chakhmouradian & Wall, 2012; Massari & Ruberti, 2013). In addition, these elements have been applied to agriculture, aiming to increase the productivity and quality of agricultural crops, a practice primarily observed in China (Xiong, 2000), where commercial fertilizers are enriched with REE (Pang, Li, & Peng, 2013). Studies also show that REE are present in phosphate fertilizers (Otero, Vitória, Soler, & Canals, 2005;

Turra, Fernandes, & Bacch, 2011) due to their presence in phosphatic rocks, which are used in fertilizer production (Al-Thyabat & Zhang, 2015; Ramos et al., 2016).

Finding ways to increase crop productivity is a constant concern among all of those involved in production chains. It is highly probable that the Brazilian agriculture industry is indirectly applying REE in the long term via phosphate fertilizers. In general, the increase in production due to the use of REE is not conclusive, making it necessary to elucidate their mechanisms in plants (Skovran & Martinez-Gomez, 2015). In this way, it is evident that the functions and physiological and biochemical pathways of these elements in plants are still uncertain. Therefore, it is important to determine the effects (beneficial or harmful) and the safe concentration limits of REE, primarily with respect to species of agricultural importance.

It is probable that the influences of lanthanum (La) on cells are numerous and complex, including ionic charge and homeostasis or binding to aromatic rings and phospholipids (Liu & Hasenstein, 2005). Hypotheses that support the beneficial effects of REE on plant metabolism include stimulation of the antioxidant system and increase in nutrient absorption (D'Aquino, Pinto, Nardic, Morganad, & Tommasi, 2009). Recently, it was verified that La influences the mitotic index of soybean plants, increases photosynthetic rate and chlorophyll index, and develops cell cycle abnormalities that benefit biomass production (Oliveira et al., 2015). La was also introduced as a calcium antagonist, since the ionic radii of La and Ca are similar, and La competes for sites with which Ca would normally bind (Hu, Ding, Chen, Wang, & Dai, 2002; Xie et al., 2002; Wang et al., 2016). In addition, studies show that La and cerium (Ce) can bind to the porphyrin of chlorophyll, forming La-chlorophyll or Ce-chlorophyll (Hong, Wei, & Zhao, 2005). Many of the contradictions found in these studies are due to the stage of vegetative growth and the different concentrations applied. Therefore, it is important to determine the real effects of La, as well as the concentration limits between these contrasting situations.

Brazil stands out as one of the world's largest maize producers, with the crop being widely cultivated and possessing high environmental adaptability. The maize crop also receives a large amount of phosphate fertilizer, which may indirectly carry REE, such as La. This can introduce considerable levels of REEs into soils and crops. Consequently, elucidating the effects of REEs on

plant growth and development is of fundamental importance. The aim of this study was to verify the effect of La on maize growth, La content, photosynthetic rate, and chlorophyll index of maize plants in response to La treatment.

## Material and methods

The experiment was conducted under greenhouse conditions, where upon maize seeds (2B604Hx hybrid) were sowed in vermiculite. After 14 days, the seedlings were transplanted for acclimatization to pots containing Hoagland nutrient solution (Hoagland & Arnon, 1950) at 40% of their ionic strength and later transferred to germinated pots using a split-root technique. One of the pots in the germinated set was filled with a complete nutrient solution without La, while another was filled with a nutrient solution without phosphorus but containing different concentrations of La. In the pots containing La, phosphorus was not added in order to avoid precipitation of these two elements, and to keep La constantly available for absorption (Ding, Liang, Zhang, Yan, & Zhang 2005). The maize plants were exposed to the following La concentrations: 0, 25, 50, 100, 150, 300 and 600  $\mu\text{M}$ , with 4 replicates of each being employed. After three weeks of La exposure, the photosynthetic rate ( $A$ ), transpiration ( $E$ ) and stomatal conductance ( $g_s$ ) were determined using an infrared gas exchange analyser – IRGA (LI-6400XT Infra Red Gas Analyzer, Li-Cor, Nebraska, USA). The evaluations were conducted between 8:00 and 10:00 for all four replicates of each La concentration. In addition, an indirect measurement of chlorophyll content was performed using the portable chlorophyll, SPAD-502 (Konica Minolta, Tokyo, Japan).

The roots and shoots (of the pots containing La) were harvested separately and washed in distilled water and dried in an oven with forced air circulation at 60°C until reaching constant weight. Each part was subsequently ground in a knife mill (Willey type), and 500 mg was weighed for the analytical measurements of La. Samples were digested using the alkaline fusion technique, and the La quantification was performed through ICP-MS, essentially as described previously (Oliveira et al., 2015). A certified reference material (AquaticPlant-BCR670, Institute for Reference Materials and Measurements, IRMM, Geel, Belgium) was included for quality control. Blank and certified reference samples were along with every batch of fusion. The mean La concentration was  $481 \pm 0.23 \text{ mg kg}^{-1}$  in the standard reference material and  $487 \text{ mg kg}^{-1}$  the value found in the certified reference

material (AquaticPlant-BCR670). Thus, the mean recovery in the certified samples (98.8%) has reliable analytical data accuracy for La analysis.


All results were analysed using analysis of variance (ANOVA), and submitted to regression analysis at a 0.05 significance level of probability using probability using Sisvar 5.3 (Build77) statistics software (Ferreira, 2011), and the graphs were made in the SigmaPlot Programme (version 12.5, Systat Software, Chicago, IL, USA).

## Results and discussion

Lanthanum caused a significant increase in shoot biomass yield at lower rates, and decreased shoot biomass was not noted at higher levels of La (Figure 1A). The La concentrations of 25 and 50  $\mu\text{M}$  led to an increase in both maize biomass and chlorophyll index, although increases in photosynthetic rate, transpiration and stomatal conductance were not verified. Emmanuel, Anandkumar, Natesan, and Maruthamuthu (2010) observed an increase in biomass production and length in maize plants submitted to concentrations of La. Xie et al. (2002) also observed that low concentrations of La promoted higher growth and production in rice plants, which is observed because La can regulate the physiological activities of the plant, such as the production of superoxide dismutase (SOD) and peroxidase (POD) enzymes. The activities of such enzymes as SOD, peroxidase, ascorbate, carotenoids and flavonoids could have been stimulated by La, helping to control reactive oxygen species, thereby reducing the oxidative stress affecting the membranes of several organelles and, in this way, promoting plant growth (Wang et al., 2011).

Lanthanum content in maize roots and shoots increased significantly when plants were exposed to increasing concentration of La (Figure 1B). There was no presence of La in the roots that developed in the germinated pot without addition of this element. In general, La content was higher in roots than shoots; only 3% of La was translocated into the shoots (Figure 1C). These results are consistent with others observed in prior studies, which report that the REE tend to remain in the roots, and less than 5% is translocated into the shoots (Hu et al., 2002). Similar results have been reported with low REE concentration in the shoot (Oliveira et al., 2015). The lower La content in the shoots may be associated with the presence of apoplastic barriers in the roots. Diatloff, Asher, and Smith (1995) reported that the Caspary bands in the root endoderm of the plant provide a barrier for the diffusion of La from the cortex apoplast to the vascular tissue, preventing

the transport of the absorbed La from the roots to the shoots of the plants.




**Figure 1.** Variation in dry matter of shoot (A), in lanthanum index in shoots and roots (B) and translocation of La (C) in maize plants (*Zea mays*) submitted to La concentrations in the culture solution.

There was a decrease in photosynthetic rate, beginning at 50  $\mu\text{M}$  of La (Figure 2A). There was no significant difference in transpiration (Figure 2B), while for stomatal conductance, there was a reduction at intermediate concentration applied (Figure 2C). The relative chlorophyll index showed a significant increase at 25 and 50  $\mu\text{M}$  of La and a reduction at higher La concentrations together with chlorosis in the leaves. According to Xiong (2000), supplying REE to plants could increase the intensity of photosynthesis and the net photosynthetic rate by 11-31%, depending on whether a mixture of REEs or individual elements is added. Zhou et al. (2011) evaluated gas exchange in *Salvia miltiorrhiza* with increasing La concentrations and observed that La increased the photosynthetic rate, stomatal conductance, transpiratory rate and biomass. For soybean plants, Oliveira et al. (2015) also reported an increase in photosynthetic rate and chlorophyll

content in the presence of 5 and 10  $\mu\text{M}$  of La. It is believed that this phenomenon occurs because REE may influence gas exchange, especially stomatal conductance, raising the photosynthetic rate due to a greater assimilation of  $\text{CO}_2$  for the photosynthetic process (Wang, Wang, Zhou, & Huang, 2014). In *Juglans nigra*, a reduction of photosynthetic activity was observed with increasing concentrations of La (Nicodemus, Salifu, & Jacobs, 2009), which was also observed in this study. As mentioned before, there was a slight increase of the biomass at the lowest concentration (25  $\mu\text{M}$ ) which correlated with the corresponding photosynthetic rate and chlorophyll index. However, when the concentration exceeds 50  $\mu\text{M}$ , the inverse response was observed, with the reduction of the biomass, probably due to the reductions in the gas exchanges and consequently, the reduction of photoassimilates.

The effects of REE on photosynthesis are also related to the development of chloroplasts, chlorophyll content and enzymatic activity, since

La may participate in the degradation of chlorophyll, an essential pigment to the photosynthetic process. Several authors assume that chlorophyll content and photosynthetic rate can be regulated by REE by increasing the light absorption efficiency, regulating the excitation energy distribution of photosystems I and II, and promoting Hill reaction activities (Liu, Wang, & Chen 2012). An increase in these variables could also be attributed to several photosynthetic processes, including the stimulation of photochemical activities, gas exchanges and  $\text{CO}_2$  fixation (Zhang, Li, Guo, & Tezuka, 2009). According to Hu, Richter, Sparovek, and Schnug (2004), the effects of REE application on photosynthetic rate, chlorophyll content, and photosynthetic enzyme activity are the primary reasons for increased wheat yield. However, since the results observed in the present study indicated a reduction of photosynthetic rate and a small increase in chlorophyll index, La concentration was most likely not efficient in maximizing photosynthetic processes.



**Figure 2.** Variation in photosynthetic rate (A), transpiration (B), stomatal conductance (C) and SPAD index of chlorophyll (D) in maize plants (*Zea mays*) submitted to La concentrations in the culture solution.

The results of Maksimovi, Kastori, Putnik-Deli, and Borišev (2014) show that total leaf area, dry biomass, photosynthetic pigment concentration, and proline decreased, indicating that maize plants were under stress, especially when yttrium (Y) was applied. The decrease in chlorophyll index could be attributed to the disturbance in the biosynthesis of the pigment, the degradation due to lipid peroxidation (Wang, Shi, Liu, Wang, & Qiao, 2012), or the damages caused by the entry of REE into the chloroplast (Fu et al., 2014). In the present study, the other factor that may have contributed to the reduction of photosynthesis was the reduction of the chlorophyll index, an essential pigment to this metabolic process. Kopsell (2011) and Wang et al. (2011) also observed that La and Ce reduced the chlorophyll content in *Hydrilla verticillata*. However, different results were shown by Hong, Wei, and Zhao (2002), where a La concentration of 500  $\mu\text{g mL}^{-1}$  increased the growth, chlorophyll index, photosynthetic rate, and uptake of nitrogen and phosphorus in spinach plants. Thus, when the photosynthetic rate, transpiration, and chlorophyll are impaired, they could be associated with plant growth.

## Conclusion

The application of La up to 25  $\mu\text{M}$  in the nutritive solution shows a slight increase in maize growth, accompanied by the stimulated photosynthetic rate, as well as the chlorophyll index. At higher levels, La did not reduce the growth but caused a decrease in both photosynthetic rate and chlorophyll index in maize plants.

These results provide important information in terms of La effects on maize growth. Further studies are needed to consolidate such information and it is suggested that the researches focus at concentrations up to 25  $\mu\text{M}$ .

## Acknowledgements

The authors are grateful to the *Instituto Tecnológico Vale* (ITV) and the *Universidade Federal de Lavras* (UFLA) (cooperation agreement ITV/UFLA 020 Vale Terras Raras) for providing financial support.

## References

- Alonso, E., Sherman, A. M., Wallington, T. J., Everson, M. P., Field, F. R., Roth, R., & Kirchain, R. E. (2012). Evaluating rare earth element availability: A case with revolutionary demand from clean technologies. *Environment Science Technology*, 46(6), 3406-3341. doi: 10.1021/es203518d.
- Al-Thyabat, S., & Zhang, P. (2015). REE extraction from phosphoric acid, phosphoric acid sludge, and phosphogypsum. *Mineral Processing and Extractive Metallurgy Review Journal*, 124(3), 1-8. doi: 10.1179/1743285515Y.0000000002
- Chakhmouradian, A. R., & Wall, F. (2012). Rare earth elements: Minerals, mines, magnets (and more). *Elements*, 8(5), 333-340. doi: 10.2113/gselements.8.5.333
- D'aquino, L., Pinto, M. C., Nardic, L., Morganad, M., & Tommasi, F. (2009). Effect of some light rare earth elements on seed germination, seedling growth and antioxidant metabolism in *Triticum durum*. *Chemosphere*, 75(7), 900-905. doi: 10.1016/j.chemosphere.2009.01.026
- Diatloff, E., Asher, C. J., & Smith, F. (1995). W. Foliar application of rare earth elements to maize and mungbean. *Australian Journal of Experimental Agriculture*, 39(2), 189-194. doi: 10.1071/EA98149
- Ding, S. M., Liang, T., Zhang, C., Yan, J., & Zhang, Z. (2005). Accumulation and fractionation of rare earth elements (REEs) in wheat: controlled by phosphate precipitation, cell wall absorption and solution complexation. *Journal of Experimental Botany*, 56(420), 2765-75. doi: 10.1093/jxb/eri270
- Emmanuel, E. S. C., Anandkumar, B., Natesan, M., & Maruthamuthu, S. (2010). Efficacy of rare earth elements on the physiological and biochemical characteristics of *Zea mays* L. *Australian Journal of Crop Science*, 4(4), 289-294.
- Ferreira, D. F. (2011). Sisvar: a computer statistical analysis system. *Ciência e Agrotecnologia*, 35(6), 1039-1042. doi: 10.1590/S1413-70542011000600001
- Fu, Y. Y., Li, F. F., Xu, T., Cai, S. J., Chu, W. Y., Qiu, H., Xu, Q. S. (2014). Bioaccumulation, subcellular, and molecular localization and damage to physiology and ultrastructure in *Nymphoides peltata* (Gmel.) O. Kuntze exposed to yttrium. *Environmental Science and Pollution Research*, 21(4), 2935-2942. doi: 10.1007/s11356-013-2246-0
- Hoagland, D. R., & Arnon, D. I. (1950). The water-culture method for growing plants without soil. *California Agricultural Experiment Station Circular*, 347, (1)1-39.
- Hong, F., Wei, Z., & Zhao, G. (2005). Mechanism of Lanthanum effect on chlorophyll of spinach. *Science China*, 45(2), 166-176. Doi: 10.1360/02yc9019.
- Hu, X., Ding, Z., Chen, Y., Wang, X., & Dai, L. (2002). Bioaccumulation of lanthanum and cerium and their effects on the growth of wheat (*Triticum aestivum* L.) seedlings. *Chemosphere*, 48(6), 621-629. doi: 10.1016/S0045-6535(02)00109-1
- Hu, Z., Richter H., Sparovek, G., & Schnug, E. (2004). Physiological and biochemical effects of rare earth elements on plants and their agricultural significance: A Review. *Journal of Plant Nutrition*, 27(1), 183-220. doi: 10.1081/PLN-120027555
- International Union of Pure and Applied Chemistry [IUPAC]. (2005). *Recommendations*. Norfolk, UK: IUPAC.

- Kopsell, D. E. (2011). Leaf tissue pigments and chlorophyll fluorescence parameters vary among sweet corn genotypes of differential herbicide sensitivity. *Pesticide Biochemistry and Physiology*, 99(2), 194-199. doi: 10.1016/j.pestbp.2010.12.004
- Liu, D., Wang, X., & Chen, Z. (2012). Effects of rare earth elements and REE-binding proteins on physiological responses in plants. *Protein Peptide Letter*, 19(2), 198-202. doi: 10.1081/PLN-120027555
- Liu, M., & Hasenstein, K. (2005). H. La<sup>3+</sup> uptake and its effect on the cytoskeleton in root protoplasts of *Zea mays* L. *Planta*, 220(5), 658-666. doi: 10.1007/s00425-004-1379-2
- Maksimovi, I., Kastori, R., Putnik-Deli, M., & Borišev, M. (2014). Effect of yttrium on photosynthesis and water relations in young maize plants. *Journal of Rare Earths*, 32(4), 372-378. doi: 10.1016/S1002-0721(14)60080-6
- Massari, S., & Ruberti, M. (2013). Rare earth elements as critical raw materials: Focus on international markets and future strategies. *Resources Policy*, 38(1), 36-43. doi: 10.1016/j.resourpol.2012.07.001
- Nicodemus, M. A., Salifu, K. F., & Jacobs, D. F. (2009). Influence of lanthanum level and interactions with nitrogen source on early development of *Juglans nigra*. *Journal of Rare Earth*, 27(2), 270-279. doi: 10.1016/j.resourpol.2012.07.001
- Oliveira, C., Ramos, S. J., Siqueira, J. O., Faquin, V., Castro, E. M., Amaral, ... Guilherme, L. R. G. (2015). Bioaccumulation and effects of lanthanum on growth and mitotic index in soybean plants. *Ecotoxicology and Environmental Safety*, 122(1), 136-144. doi: 10.1016/j.ecoenv.2015.07.020
- Otero, N., Vitória, L., Soler, A., & Canals, A. (2005). Fertiliser characterisation: Major, trace and rare earth elements. *Applied Geochemistry*, 20(8), 1473-1488. doi: 10.1016/j.apgeochem.2005.04.002
- Pang, X., Li, D., & Peng, A. (2013). Application of rare-earth elements in the agriculture of China and its environmental behavior in soil. *Environmental Science and Pollution Research*, 9(2), 143-148. doi: 10.1007/BF02987718
- Ramos, S. J., Dinali, G. S., de Carvalho, T. S., Chaves, L. C., Siqueira, J. O., & Guilherme, L. R. G. (2016). Rare earth elements in raw materials and products of the phosphate fertilizer industry in South America: Content, signature, and crystalline phases. *Journal of Geochemical Exploration*, 168(1), 177-186. doi: 10.1016/j.gexplo.2016.06.009
- Skovran, E., & Martinez-Gomez, N. C. (2015). Just add lanthanides. *Science*, 348(623), 862-863. doi: 10.1126/science.aaa9091
- Turra, C., E. E., Fernandes, A. N., & Bacch, M. A. (2011). Evaluation on rare earth elements of Brazilian agricultural supplies. *Journal of Environmental Chemistry and Ecotoxicology*, 3(4), 86-92. doi: 10.1590/1806-9657rbcs20150413
- Wang, C., Shi, C., Liu, L., Wang, C., & Qiao, W. (2012). Lanthanum Element Induced Imbalance of Mineral Nutrients, HSP 70 Production and DNA-Protein Crosslink, Leading to Hormetic Response of Cell Cycle Progression in Root Tips of *Vicia faba* L. seedlings. *Dose-response*, 10(1), 96. doi: 10.2203/dose-response.11-041.
- Wang, L., Wang, W., Zhou, Q., & Huang, X. (2014). Combined effects of lanthanum (III) chloride and acid rain on photosynthetic parameters in rice. *Chemosphere*, 112(2), 355-361. doi: 10.1016/j.chemosphere.2014.04.069
- Wang, C., Lu, X., Tian, Y., Cheng, T., Hu, L., Chen, F., ... & Wang, X. (2011). Lanthanum resulted in unbalance of nutrient elements and disturbance of cell proliferation cycles in *V. faba* L. seedlings. *Biological Trace Element Research*, 143(2), 1174-1181. doi: 10.1007/s12011-010-8939
- Wang, L., Cheng, M., Chu, Y., Li, X., Chen, D. D. Y., Huang, X., & Zhou, Q. (2016). Responses of plant calmodulin to endocytosis induced by rare earth elements. *Chemosphere*, 154(2), 408-415. doi: 10.1016/j.chemosphere.2016.03.106.
- Xie, Z. B., Zhu, J. G., Chu, H. Y., Zhang, Y. L., Zeng, Q., & Ma H. L. (2002). Effect of Lanthanum on Rice Production, Nutrient Uptake, and Distribution. *Journal of Plant Nutrition*, 25(10), 2315-2331. doi: 10.1081/PLN-120014078
- Xiong, B. K. (2000). Rare earth element research and applications in Chinese agriculture and forest (in Chinese). *Beijing: Metallurgical Industry Press*, 67(1), 1-151.
- Zhang, R. H., Li, J., Guo, S. R., & Tezuka, T. (2009). Effects of exogenous putrescine on gas-exchange characteristics and chlorophyll fluorescence of NaCl-stressed cucumber seedlings. *Photosynthesis Research*, 100(3), 155-162. doi: 10.1007/s11120-009-9441-3
- Zhou, J., Guo, L., Zhang, J., Zhou, S., Yang, G., Zhao, M., & Huang, L. (2011). Effects of LaCl<sub>3</sub> on photosynthesis and the accumulation of tanshinones and salvianolic acids in *Salvia miltiorrhiza* seedlings. *Journal of Rare Earths*, 29(5), 49-49. doi: 10.1016/S1002-0721(10)60486-3

Received on August 1, 2017.

Accepted on May 18, 2018.

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.