



Cultivos Tropicales
ISSN: 1819-4087
Ediciones INCA

Cruz-Suárez, Arasay Santa; Nápoles-García, María Caridad; Morales-Guevara, Donaldo
El déficit hídrico en los cultivos y la acción de los microorganismos
Cultivos Tropicales, vol. 43, no. 3, 2022, pp. 1-8
Ediciones INCA

Available in: <https://www.redalyc.org/articulo.oa?id=193275342013>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UDEM [redalyc.org](https://www.redalyc.org)

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative



Water deficit in crops and the action of microorganisms

El déficit hídrico en los cultivos y la acción de los microorganismos

^{ID}Arasay Santa Cruz-Suárez*, ^{ID}María Caridad Nápoles-García, ^{ID}Donaldo Morales-Guevara

Instituto Nacional de Ciencias Agrícolas (INCA), carretera San José-Tapaste, km 3½, Gaveta Postal 1, San José de las Lajas, Mayabeque, Cuba. CP 32 700

ABSTRACT: One third of the planet's surface is considered arid or semi-arid, while most of the remaining surface is subject to temporary periods of water deficit. Drought is considered to be one of the world's major natural disasters, the most frequent and persistent, the one with the greatest negative effect on agricultural production and also the one that causes real adverse impacts on the environment. Water is the main limiting factor for plant growth on earth, acting as a first degree selective force for the evolution and distribution of plant species. Soil microorganisms are known to contribute a wide range of essential services to the sustainability of all ecosystems. They act as the main drivers of nutrient cycling; they regulate soil organic matter dynamics, carbon sequestration and greenhouse gas emissions; they modify soil physical structure and water regime. Inoculation of some microorganisms improves the efficiency of nutrient uptake, promotes plant growth and yield, thus attenuating the adverse effects of stress. The present work was aimed to deepen the knowledge of water deficit effects on crops and the role of some microorganisms in the mitigation of this stress on plants.

Key words: water, stress, inoculation, sustainability.

RESUMEN: Una tercera parte de la superficie del planeta se considera como árida o semiárida, mientras que la mayoría de la superficie restante está sujeta a períodos temporales de déficit hídrico. La sequía es considerada como uno de los mayores desastres naturales del mundo, el más frecuente y persistente, el de mayor efecto negativo para la producción agrícola y también como la causante de impactos adversos reales en el medio ambiente. El agua constituye el principal factor limitante del crecimiento de las plantas en la tierra, actuando como una fuerza selectiva de primer grado para la evolución y distribución de las especies vegetales. Se conoce, que los microorganismos del suelo contribuyen con un amplio rango de servicios esenciales a la sostenibilidad de todos los ecosistemas. Ellos actúan como los principales agentes impulsores del ciclo de nutrientes; regulan la dinámica de la materia orgánica del suelo, el secuestro de carbono y la emisión de gases de efecto invernadero; modifican la estructura física del suelo y el régimen de agua. La inoculación de algunos microorganismos mejora la eficiencia de la toma de nutrientes, promueve el crecimiento y rendimiento de las plantas, atenuando de esta forma los efectos adversos del estrés. El presente trabajo estuvo encaminado a profundizar en el conocimiento de los efectos del déficit hídrico en los cultivos y el papel de algunos microorganismos en la mitigación de este estrés sobre las plantas.

Palabras clave: agua, estrés, inoculación, sostenibilidad.

INTRODUCTION

Water is the essential component of life; in plants, it constitutes approximately 85 to 90 % of their mass and is present in all the vital processes that take place within them (1). Agriculture consumes, annually, 70 % of the total water used for all uses (2) and during crop establishment, its

success will depend largely on the access to water available (3). It is a widely known fact that each stage of crop development is individually sensitive to environmental factors and, in particular, to the lack of soil moisture that can cause water stress (4). For these reasons, mankind is facing a new challenge, that of producing more food with less water.

*Author for correspondence: arasay@inca.edu.cu

Received: 15/12/2020

Accepted: 27/09/2021

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial (BY-NC 4.0). <https://creativecommons.org/licenses/by-nc/4.0/>



The functions of microorganisms in the soil and the possibility of plants to associate with many of them are dissimilar, which provide them with the capacity to adapt to the environmental conditions where they develop; they act directly and indirectly on the growth of plants and their capacity to tolerate adverse conditions of biotic and abiotic stress, their functions differ, from biofertilizers, phytostimulators, biocontrollers, bioprotectors, to the combination of several of them (5). Hence, their use is of great importance in the search for new alternatives to mitigate the consequences caused by water scarcity.

The present work aims to deepen the knowledge of water deficit effects on crops and the role of some microorganisms in the mitigation of this stress on plants.

Water deficit

Water deficit is the abiotic stress with the greatest impact on plant growth (6). From an ecophysiological point of view, it occurs when there is not enough water in plant root zone to meet their needs at a given time and place, i.e., when the water transpired exceeds the water absorbed (7). From a physiological point of view, the word stress refers to some kind of suffering, i.e., an alteration of normal functioning with consequences on plant growth or development (8).

The lack of water availability that causes stress may be due to an edaphic water deficit, to an excessive atmospheric evaporative demand, or to the interaction of both factors (9). This type of hydroedaphic or agricultural drought stress, as it is known, is considered a transitory episode for plants (10).

There are two criteria, based on authors who affirm, taking into account thermodynamic theoretical considerations, that water stress is caused by a decrease in soil water potential (Ψ_s); and those who affirm, based on empirical evidence, that it is caused by a decrease in edaphic moisture content, referred to in terms of the fraction of transpirable edaphic water (FTSW) (11). It is reasonable to think that both positions may have some reason, since insufficient water availability could be due to a decrease in Ψ_s as the soil dries out, but also, to a decrease in the soil's capacity to supply water, the latter aspect that could depend on its FTSW. Not only does water deficit occur when there is little water in the environment (12), it can also be the result of low rainfall, excessive salinity, and extreme temperatures, hot or cold (13).

Effects of water deficit on plants

Water deficit has important consequences for plant physiology and morphology. It causes, in the short and medium term, a decrease in water potential, relative water content and stomatal conductivity (14). This effect results in negative impacts on plant growth and production, causing anatomical changes due to modifications in cell size, senescence, and even death in many species (15). The magnitude of the damage caused depends mainly on its duration, intensity and the time in which it occurs within the crop cycle (16).

When at the beginning of a productive crop cycle of the first stage of seed germination is compromised, with moisture deficiencies present in the soil during sowing, imbibition is affected and, consequently, the activation of metabolic processes, such as rehydration, repair mechanisms (membranes, proteins and DNA), cell elongation and the appearance of the radicle (11). In a research carried out in wheat (*Triticum aestivum* L.), simulating drought with polyethylene glycol, it was determined that as drought increases, seedling germination decreases by 50 % (17).

Some investigations in several crops make reference to the fact that the physiological process of photosynthesis is also affected by the different levels of drought (18). Other manifestations are a decrease in protein synthesis and photorespiration, photosynthate translocation and growth regulators are affected (19) and gas exchange is reduced, limiting biomass production (20). These events result in a reduction of plant organ growth (21) which, if maintained for prolonged periods, leads to a modification of the specific leaf area (22), resulting in less leaf area exposed to solar radiation (23). An example of this is a study carried out on cotton plants (*Gossypium hirsutum* L.), indicating that in environments of hydric deficit there is a strong decrease in leaf expansion (24). In rice (*Oryza sativa* L.), less leaf growth was also found in plants subjected to drought in the vegetative phase, due to a reduction in the expansion rate of the aerial part and the development of the leaf area, caused by a decrease in the turgor pressure of the leaves (25).

Regarding plant reproduction, a phase negatively influenced by water shortage is flowering. It seems to be the most sensitive in most crops where the final product is grain or fruit (4). As water stress becomes more severe, the reduction in photosynthetic activity causes a decrease in the rate of cell division and, consequently, the initiation of new leaf primordia is halted (26). The balance between productions of assimilates and the demand for the development of reproductive organs is severely affected by the reduction of the photosynthetically active leaf area (27). In this regard, in plants of sweet orange [*Citrus sinensis* (L.) Osbeck] cv. 'Valencia late', an increase in floral abortion was observed in plants that received less water, and moderate stress increased flower and fruit abscission (28).

In relation to the production stage of a crop, photosynthesis plays an important role (29), since grain yield is potentially influenced by the duration of the rate of carbohydrate accumulation (30). Water deficit can affect the use of carbohydrates, basically by changing the efficiency with which photoassimilates are converted into the development of new plant parts (31). The yields to be obtained in a crop will depend on production, the latter being another factor vulnerable to drought. It has been proven in grape plants (*Vitis vinifera* L.) that the lowest yields are obtained when the plants do not receive irrigation, in the stage between reproductive sprouting and greening (22). Similarly, in pomegranate (*Punica granatum* L.) plants grown under severe stress, yield and fruit quality were found to be affected (32).

Plant response to water deficit

Throughout evolution, plants have developed several strategies facing water deficit that allow them to alleviate, tolerate and adapt to such circumstances (33). Stress conditions produce two types of response in organisms: those that tend to avoid or prevent stress (avoidance mechanisms) and those that allow them to resist it (tolerance mechanisms) (34). Another type of mechanism, not considered by many authors as a true resistance to water deficit, is the elusive or drought escape mechanism (35). Within this strategy, some annual plants complete their life cycle before water stress (in spring), and therefore produce seeds that can withstand the unfavorable period for a prolonged period of time (36). However, this escape strategy may include species that enter partial vegetative dormancy during drought (35).

In avoidance mechanisms, plants minimize the penetration of stress into their tissues. Species following this strategy maximize water uptake, for example, through deep root systems, or minimize water losses through rapid and sensitive stomatal closure to slight decreases in tissue water content or water potential, the formation of small leaves, low transpiration rates, and changes in xylem hydraulic properties (37). Therefore, within this strategy there are two avoidance mechanisms: one, by water wastage, which allows maintaining the tissues hydrated in full drought, as long as the access to soil water and its internal distribution through the xylem is not limiting, and the other, by water saving. Both maintain the plants in the turgor state, and when water deficit conditions are accentuated, the wasters cannot maintain the high transpiration rates, so they become water savers or the individual dies (38).

Regarding tolerance mechanisms, they are very specific to the deformation that occurs. Different modes of resistance are triggered, and even plastic deformations are tolerated. These species have mechanisms that minimize or eliminate the deformation they may suffer as a consequence of stress, reaching a thermodynamic equilibrium with the stress without suffering damage. An important aspect of this strategy is the stress repair mechanisms, which the plant has to activate when the stress has ceased to act (12). The mechanisms that confer tolerance to drought are based on structural stabilization by preferential hydration, whereas the mechanisms that confer tolerance to desiccation are based on the replacement of water by molecules that form hydrogen bonds (39).

At the cellular and molecular level, one of the main responses of plants to adapt to the environment in the face of this limitation is the modification of specific genes. This action depends on the nature, duration, and severity of the stress (37). Genes induced by water deficit are categorized into three groups: 1) genes encoding proteins with known structural or enzymatic functions; 2) proteins with as yet unknown functions; and 3) regulatory proteins (40). But the function of these genes is not only to protect plant cells from dehydration, but they also act in the regulation of other

genes that translate into signals in response to water stress (41). In situations of water deficiency, certain enzymes and proteins are affected and lose their biological activity. Therefore, the maintenance of proteins in their functional form and the prevention of protein aggregation is particularly important for the life of the cell under stress conditions (39).

It has been detected that when some plants develop under water deficiency, they accumulate higher amounts of ABA, proline and some soluble sugars. Following this process, chemical signals are sent from the roots to the leaves by increasing the concentration in the cytoplasm of ABA, pH, ethylene precursor and malate, in immediate response to water stress, which causes stomatal closure (42). Studies have also demonstrated the importance of this acid in root growth in drought environments by regulating the negative effect of ethylene on root development (43).

Other biochemical responses generated, due to the decrease of the osmotic potential through accumulations of inorganic (K^+ , Ca^{2+} , Mg^{2+}) and organic (proline, aspartic acid, proteins and sugars) solutes (44), K^+ and Ca^{2+} outflows in the leaf mesophyll, regulate the response to drought (45). The stomatal response allows an increase in transpiration efficiency (46), which is why, when there is a significant decrease in leaf water potential, there is an increase in resistance to water loss (47), coupled with a decrease in net CO_2 fixation (48).

On the other hand, it has been observed that complete recovery of photosynthesis is slow as the plant approaches the point of permanent wilting, varying with plant species, soil type, and method of irrigation water application (49).

Many of these adaptations that plants have developed are related to a greater capacity to absorb water, or to a more efficient use of this resource (50). This is the case of leaf senescence, which constitutes the major regulator in the endogenous activity of the associated processes and contributes to reduce water loss and induces plant dormancy for survival (36). It has been demonstrated that, in the presence of soil water deficit, leaf expansion accelerates senescence (51). The increase of this process in the vegetative phase becomes a potential limitation for crop productivity (52). However, when water tension in the stomata increases, CO_2 assimilation is reduced and the plant prioritizes the synthesis of primary metabolites and, therefore, fruit growth (53).

Role of some microorganisms against water deficit in plants

Most plants adapted to diverse ecological niches are associated with soil microorganisms, which play a key role in the protection against environmental stress to which they may be subjected (54).

Among the PGPR (Plant Growth Promoting Rhizobacteria) genera most commonly used in crops of economic importance in the face of water deficit are: *Bacillus*, *Rhizobium*, *Azospirillum*, *Azotobacter*, *Pseudomonas* (55), *Herbaspirillum* (56), *Bradyrhizobium*

(57), *Enterobacter* (58), *Achromobacter*, *Serratia* (59), *Klebsiella* and *Beijerinckia* (60). Among the positive effects associated with these microorganisms are biological nitrogen fixation, production of plant growth regulating substances such as auxins, cytokinins and gibberellins, increased root growth and pathogen control (61). These characteristics, added to their tolerance to osmotic stress, the endophytic capacity that in some cases provide morphophysiological changes, such as accumulation of osmolytes, stomatal regulation and decrease in membrane potential, contribute to the mitigation of water stress in host plants (62).

Several mechanisms by which rhizobacteria induce tolerance to water deficit stress have been proposed, including the production of phytohormones such as abscisic acid, gibberellic acid, cytokinins, and indolacetic acid; the synthesis of ACC (1-aminocyclopropane-1-carboxylate) deaminase, which reduces the level of ethylene in roots; the induction of induced systemic tolerance; and the presence of exopolysaccharides (Figura 1) (63).

Some of these bacteria form resistance structures to favor their survival in circumstances of drought stress (60). The production of ACC deaminase can be incorporated into the ethylene production pathway, capturing the amino acid ACC (ethylene precursor) and decreasing the levels of this hormone. This action promotes greater plant tolerance to water stress (64).

Eighty percent of dinitrogen (N_2) fixing bacteria produce IAA (3-indolacetic acid) (65). This growth regulator leads to an increase in total phenols, calcium content and the activity of the polyphenol oxidase enzyme that protects the crop against pathogens. It also improves plant growth by eliminating reactive oxygen species, which are formed in

the plant as a result of water stress (66). It has been observed that bacterial production of IAA has high sensitivity in plant roots inoculated with this hormone. For example, root length and height in guinea grass (*Megathyrsus maximus*) were favored by inoculation with *Bacillus* sp. strains in two trials conducted in the presence of water stress (67).

Plants inoculated with this type of rhizobacteria that promote growth under abiotic stress due to drought, improve the water status of the leaves, since they tend to have higher values of water use efficiency, that is, the ratio between dry mass gain and water consumed (68). In this sense, it is likely that rhizobia produce antioxidant enzymes (catalase), exopolysaccharides and other substances to be able to survive in extreme environments, especially in drought (69).

Other soil microorganisms that establish symbiotic relationships and increase nutrient assimilation and tolerance to various types of biotic and abiotic stresses in plants are arbuscular mycorrhizal fungi (AMF) (70). These fungi promote resistance to water deficiencies in the host plant, which is a consequence of different mechanisms, ranging from a physical response to a biochemical response (71). The mitigation of the negative effect of water stress by mycorrhization is the result of modifications of the water balance (transpiration and efficient use of water) and nutritional balance, specifically for P (phosphorus), N (nitrogen) and K (potassium) (72). The development of extraradical mycelium allows the roots to have greater access to soil water and thus increase their hydration (73).

The mycorrhizal association alters water relations, independently of the plant stage, which favors plant establishment, vigor, productivity and survival in an




Figure 1. Water deficit tolerance mechanisms induced by plant growth-promoting rhizobacteria

environment with limited water conditions (71). It has been demonstrated that mycorrhizal plants subjected to water deficit conditions recover faster and resist drought conditions longer (74). A study carried out on corn shows that when the crop is subjected to drought and is mycorrhized by *Glomus intraradices* and *Glomus mosseae*, it recovers faster and presents higher values of leaf water potential and CO₂ assimilation rate (75).

CONCLUSIONS

In recent years, considerable progress has been made in the knowledge of the responses induced in plants by water stress. Likewise, in the beneficial effect that microorganisms exert on crops, by promoting the growth of the aerial and root part in times of drought stress. The understanding of the different biochemical and physiological mechanisms involved in stress tolerance allows us to have a better vision of the development obtained by many plants and their interaction with microorganisms throughout their evolutionary process, due to climate change and other environmental events occurring worldwide.

BIBLIOGRAPHY

- Turner NC, Begg JE. Plant-water relations and adaptation to stress. *Plant and soil*. 1981;58(1):97-131.
- II G d'experts intergouvernemental sur l'évolution du climat WG, Change IP on C, I IP on CCWG, Staff IP on CC. *Climate Change 2007 - Impacts, Adaptation and Vulnerability: Working Group II Contribution to the Fourth Assessment Report of the IPCC* [Internet]. Cambridge University Press; 2007. 765 p. Available from: https://books.google.com/cu/books?hl=es&lr=&id=TNo-SeGpn7wC&oi=fnd&pg=PA81&dq=WMO-UNEP-+Climate+Change+2007:+Impacts,+Adaptation+and+vulnerability.+&ots=vR6AnfYvRA&sig=9EJ8YM6oqC-RFDXzOJxDZtUw84E&redir_esc=y#v=onepage&q=WMO-UNEP-%20Climate%20Change%202007%3A%20Impacts%2C%20Adaptation%20and%20vulnerability.&f=false
- González LM, Estrada A, Zaldivar N, Argente L. Tolerancia a la sequía en diferentes variedades de trigo sobre la base de algunas variables del régimen hídrico y la concentración de pigmentos en estadio de plántula. *Revista Ciencias Técnicas Agropecuarias*. 2007;16(1):45-9.
- de la Casa A, Ovando G. Integración del Índice de Vegetación de la Diferencia Normalizada (NDVI) y del Ciclo Fenológico de Maíz para Estimar el Rendimiento a Escala Departamental en Córdoba, Argentina. *Agricultura Técnica*. 2007;67(4):362-71. doi: [10.4067/S0365-28072007000400004](https://doi.org/10.4067/S0365-28072007000400004)
- Kumar A, Verma JP. The role of microbes to improve crop productivity and soil health. In: *Ecological wisdom inspired restoration engineering* [Internet]. Springer; 2019. p. 249-65. Available from: https://link.springer.com/chapter/10.1007/978-981-13-0149-0_14
- Gao J-P, Chao D-Y, Lin H-X. Understanding Abiotic Stress Tolerance Mechanisms: Recent Studies on Stress Response in Rice. *Journal of Integrative Plant Biology*. 2007;49(6):742-50. doi: <https://doi.org/10.1111/j.1744-7909.2007.00495.x>
- Sans JF, Gil HM. Respuesta de las plantas al estrés hídrico. In: *La ecofisiología vegetal : una ciencia de síntesis*, 2003, ISBN 84-9732-267-3, págs. 253-286 [Internet]. Thomson-Paraninfo; 2003 [cited 28/02/2022]. p. 253-86. Available from: <https://dialnet.unirioja.es/servlet/articulo?codigo=830199>
- Salisbury FBR, Cleon W. *Fisiología de las plantas I Células: agua, soluciones y superficies/por Frank B. Salisbury y Cleon W. Ross*.
- Campbell GS, Norman JM. *An Introduction to Environmental Biophysics* [Internet]. Springer Science & Business Media; 1998. 224-225 p. Available from: https://books.google.es/books?hl=es&lr=&id=v6UpE6lThCwC&oi=fnd&pg=PR5&dq=Norman+JM.+Environmental+biophysics&ots=JXeeUlbW&sig=RuU6q1hj_lb2uT-yd7PBjB_L3RY#v=onepage&q=Norman%20JM.%20Environmental%20biophysics&f=false
- Maliva R, Missimer T. *Arid Lands Water Evaluation and Management* [Internet]. Germany: Springer Science & Business Media; 2012. 1075 p. Available from: <https://books.google.es/books?hl=es&lr=&id=Tx-gqLgdz0YC&oi=fnd&pg=PR5&dq=Aridity+and+drought.+En:+Arid+lands+w%C3%A1ter+evaluation+and+management,+environmental+science+and+engineering.+Springer-Verlag+Berlin+Heidelberg,+Germany.+2012&ots=Z-Clq5xSPz&sig=WFuSMgwIxOSx4ijYAtcDG1FuCwv#v=onepage&q&f=false>
- Dubreucq B, Berger N, Vincent E, Boisson M, Pelletier G, Caboche M, et al. The Arabidopsis AtEPR1 extensin-like gene is specifically expressed in endosperm during seed germination. *The Plant Journal*. 2000;23(5):643-52. doi:<https://doi.org/10.1046/j.1365-313x.2000.00829.x>
- Levitt J. *Responses of plants to environmental stresses. Physiological ecology*. Academic Press, New York; 1980.
- Nilsen ET, Orcutt DM. *Physiology of plants under stress. Abiotic factors*. John Wiley and sons. Inc. 1996;333.
- Medrano H, Bota J, Cifre J, Flexas J, Ribas-Carbó M, Gulías J. Eficiencia en el uso del agua por las plantas. *Investigaciones geográficas (Esp)*. 2007;(43):63-84.
- Morales CG, Pino MT, del Pozo A. Phenological and physiological responses to drought stress and subsequent rehydration cycles in two raspberry cultivars. *Scientia Horticulturae*. 2013;162:234-41. doi: [10.1016/j.scienta.2013.07.025](https://doi.org/10.1016/j.scienta.2013.07.025)
- Ballester C, Castel J, Intrigliolo DS, Castel JR. Response of Clementina de Nules citrus trees to summer deficit irrigation. Yield components and fruit composition. *Agricultural Water Management*. 2011;98(6):1027-32. doi:[10.1016/j.agwat.2011.01.011](https://doi.org/10.1016/j.agwat.2011.01.011)
- González LM, Argente L, Zaldivar N, Ramírez R. Efecto de la sequía simulada con PEG-6000 sobre la germinación y el crecimiento de las plántulas de dos variedades de trigo. *Cultivos tropicales*. 2005;26(4):49-52.
- Erice G, Louahia S, Irigoyen JJ, Sánchez-Díaz M, Alami IT, Avice J-C. Water use efficiency, transpiration and net CO₂ exchange of four alfalfa genotypes submitted to

- progressive drought and subsequent recovery. Environmental and Experimental Botany. 2011;72(2): 123-30. doi:[10.1016/j.envexpbot.2011.02.013](https://doi.org/10.1016/j.envexpbot.2011.02.013)
19. Coombs J, Hall DO, Long SP, Scurlock JMO. Técnicas en fotosíntesis y bioproductividad. eds. UNEP, CP. Chapingo, México; 1988 p. 42-51.
20. Erice G, Irigoyen JJ, Sánchez-Díaz M, Avise J-C, Ourry A. Effect of drought, elevated CO₂ and temperature on accumulation of N and vegetative storage proteins (VSP) in taproot of nodulated alfalfa before and after cutting. Plant Science. 2007;172(5):903-12. doi:[10.1016/j.plantsci.2006.12.013](https://doi.org/10.1016/j.plantsci.2006.12.013)
21. Quiñones A, Martínez-Alcántara B, Legaz F. Influence of irrigation system and fertilization management on seasonal distribution of N in the soil profile and on N-uptake by citrus trees. Agriculture, Ecosystems & Environment. 2007;122(3):399-409. doi:[10.1016/j.agee.2007.02.004](https://doi.org/10.1016/j.agee.2007.02.004)
22. Ferreyra E. R, Sellés V. G, Ruiz S. R, Sellés M. I. Efecto del estrés hídrico aplicado en distintos periodos de desarrollo de la vid cv. chardonnay en la producción y calidad del vino. Agricultura Técnica. 2003;63(3):277-86. doi:[10.4067/S0365-28072003000300007](https://doi.org/10.4067/S0365-28072003000300007)
23. Álvarez S, Navarro A, Bañón S, Sánchez-Blanco MJ. Regulated deficit irrigation in potted Dianthus plants: Effects of severe and moderate water stress on growth and physiological responses. Scientia Horticulturae. 2009;122(4):579-85. doi:[10.1016/j.scienta.2009.06.030](https://doi.org/10.1016/j.scienta.2009.06.030)
24. Ball RA, Oosterhuis DM, Mauromoustakos A. Growth Dynamics of the Cotton Plant during Water-Deficit Stress. Agronomy Journal. 1994;86(5):788-95. doi:<https://doi.org/10.2134/agronj1994.00021962008600050008x>
25. Siopongco JD, Yamauchi A, Salekdeh H, Bennett J, Wade LJ. Growth and Water Use Response of Doubled-Haploid Rice Line to Drought and Rewatering during the Vegetative Stage. Plant Production Science. 2006;9(2):141-51. doi:[10.1626/pp.9.141](https://doi.org/10.1626/pp.9.141)
26. Kumar A, Singh DP, Singh P. Influence of water stress on photosynthesis, transpiration, water-use efficiency and yield of Brassica juncea L. Field Crops Research. 1994;37(2):95-101. doi:[10.1016/0378-4290\(94\)90037-X](https://doi.org/10.1016/0378-4290(94)90037-X)
27. Gerik TJ, Faver KL, Thaxton PM, El-Zik KM. Late Season Water Stress in Cotton: I. Plant Growth, Water Use, and Yield. Crop Science. 1996;36(4):crops1996.0011183X003600040017x. doi: <https://doi.org/10.2135/cropsci1996.0011183X003600040017x>
28. Garzón Correa DL, Vélez-Sánchez JE, Orduz Rodríguez JO. Efecto del déficit hídrico en el crecimiento y desarrollo de frutos de naranja Valencia (Citrus sinensis Osbeck) en el piedemonte del Meta, Colombia. Acta Agronómica. 2013;62(2):136-47.
29. Wulschleger SD, Oosterhuis DM. Photosynthetic Carbon Production and Use by Developing Cotton Leaves and Bolls. Crop Science. 1990;30(6):1259-61. doi:<https://doi.org/10.2135/cropsci1990.0011183X003000060021x>
30. Crafts-Brandner SJ, Poneleit CG. Selection for Seed Growth Characteristics: Effect on Leaf Senescence in Maize. Crop Science. 1992;32(1):127-31. doi:<https://doi.org/10.2135/cropsci1992.0011183X003200010028x>
31. Jordan WR. Whole Plant Response to Water Deficits: An Overview. In: Limitations to Efficient Water Use in Crop Production [Internet]. John Wiley & Sons, Ltd; 1983 [cited 18/04/2022]. p. 289-317. doi:[10.2134/1983.limitationsto efficientwateruse.c18](https://doi.org/10.2134/1983.limitationsto efficientwateruse.c18)
32. Mellisho CD, Egea I, Galindo A, Rodríguez P, Rodríguez J, Conejero W, et al. Pomegranate *Punica granatum* L.) fruit response to different deficit irrigation conditions. Agricultural Water Management. 2012;114:30-6. doi:[10.1016/j.agwat.2012.06.010](https://doi.org/10.1016/j.agwat.2012.06.010)
33. Whitmore AP, Whalley WR. Physical effects of soil drying on roots and crop growth. Journal of Experimental Botany. 2009;60(10):2845-57. doi:[10.1093/jxb/erp200](https://doi.org/10.1093/jxb/erp200)
34. Larcher W. Physiological Plant Ecology: Ecophysiology and Stress Physiology of Functional Groups. Springer Science & Business Media; 2003. 540 p.
35. Valladares F, Vilagrosa A, Peñuelas J, Ogaya R, Camarero JJ, Corcuera L, et al. Estrés hídrico: ecofisiología y escalas de la sequía. In Madrid: Ministerio de Medio Ambiente; 2004. Available from: www.irmase.csic.es/users/jefer/pdf/Valladares%20F.doc
36. Lelièvre F, Seddaiu G, Ledda L, Porqueddu C, Volaire F. Water use efficiency and drought survival in Mediterranean perennial forage grasses. Field Crops Research. 2011;121(3):333-42. doi:[10.1016/j.fcr.2010.12.023](https://doi.org/10.1016/j.fcr.2010.12.023)
37. Moreno F. LP. Respuesta de las plantas al estrés por déficit hídrico. Una revisión. Agronomía Colombiana. 2009;27(2):179-91.
38. Kozłowski TT, Kramer PJ, Pallardy SG. The Physiological Ecology of Woody Plants [Internet]. Academic Press; 1991. 678 p. Available from: <https://books.google.es/books?hl=es&lr=&id=iSTOcsNbVxMC&oi=fnd&pg=PP1&dq=The+physiological+ecology+of+woody+plants.+Academic+Press.&ots=zP86EJN18B&sig=11deEoDix0YktiZj8dtNpjpVGKg#v=onepage&q=The%20physiological%20ecology%20of%20woody%20plants.%20Academic%20Press.&f=false>
39. Hoekstra FA, Golovina EA, Buitink J. Mechanisms of plant desiccation tolerance. Trends in Plant Science. 2001;6(9):431-8. doi:[10.1016/S1360-1385\(01\)02052-0](https://doi.org/10.1016/S1360-1385(01)02052-0)
40. Bhatnagar-Mathur P, Vadez V, Sharma KK. Transgenic approaches for abiotic stress tolerance in plants: retrospect and prospects. Plant Cell Reports. 2008;27(3):411-24. doi:[10.1007/s00299-007-0474-9](https://doi.org/10.1007/s00299-007-0474-9)
41. Reddy AR, Chaitanya KV, Vivekanandan M. Drought-induced responses of photosynthesis and antioxidant metabolism in higher plants. Journal of Plant Physiology. 2004;161(11):1189-202. doi:[10.1016/j.jplph.2004.01.013](https://doi.org/10.1016/j.jplph.2004.01.013)
42. Schachtman DP, Goodger JQD. Chemical root to shoot signaling under drought. Trends in Plant Science. 2008;13(6):281-7. doi:[10.1016/j.tplants.2008.04.003](https://doi.org/10.1016/j.tplants.2008.04.003)
43. Sharp RE, LeNoble ME. ABA, ethylene and the control of shoot and root growth under water stress. Journal of

- Experimental Botany. 2002;53(366):33-7. doi:[10.1093/jexbot/53.366.33](https://doi.org/10.1093/jexbot/53.366.33)
44. Yu Z, Chen W, Zhang Q, Yang H, Tang J, Weiner J, et al. Salt tolerance and stress level affect plant biomass-density relationships and neighbor effects. *Acta Oecologica*. 2014;58:1-4. doi:[10.1016/j.actao.2014.04.001](https://doi.org/10.1016/j.actao.2014.04.001)
 45. Mak M, Babla M, Xu S-C, O'Carrigan A, Liu X-H, Gong Y-M, et al. Leaf mesophyll K⁺, H⁺ and Ca²⁺ fluxes are involved in drought-induced decrease in photosynthesis and stomatal closure in soybean. *Environmental and Experimental Botany*. 2014;98:1-12. doi:[10.1016/j.envexpbot.2013.10.003](https://doi.org/10.1016/j.envexpbot.2013.10.003)
 46. Sánchez-Díaz M, Aguirreolea J. Movimientos estomáticos y transpiración. Azcon-Bieto, J. y M. Talon. Fundamentos de fisiología vegetal. 1ª ed. Ed. McGraw-Hill/ Interamericana de España, Madrid. 2000;31-43.
 47. Taiz L, Zeiger E, Møller IM, Murphy A. Plant physiology and development. [Internet]. Sinauer; 2015 [cited 25/04/2022]. 733-734 p. Available from: <https://www.cabdirect.org/cabdirect/abstract/20173165866>
 48. Wu Y, Huang M, Warrington DN. Growth and transpiration of maize and winter wheat in response to water deficits in pots and plots. *Environmental and Experimental Botany*. 2011;71(1):65-71. doi:[10.1016/j.envexpbot.2010.10.015](https://doi.org/10.1016/j.envexpbot.2010.10.015)
 49. Mota F da. Meteorología agrícola. Nobel São Paulo; 1983.
 50. Langridge P, Paltridge N, Fincher G. Functional genomics of abiotic stress tolerance in cereals. *Briefings in Functional Genomics*. 2006;4(4):343-54. doi:[10.1093/bfpgp/eli005](https://doi.org/10.1093/bfpgp/eli005)
 51. Sivakumar MVK, Shaw RH. Relative Evaluation of Water Stress Indicators for Soybeans1. *Agronomy Journal*. 1978;70(4):619-23. doi:<https://doi.org/10.2134/agronj1978.00021962007000040022x>
 52. Carlesso R. Influence of soil water deficits on maize growth and leaf area adjustments. Michigan State University; 1993.
 53. Horner JD. Nonlinear effects of water deficits on foliar tannin concentration. *Biochemical Systematics and Ecology*. 1990;18(4):211-3. doi: [10.1016/0305-1978\(90\)90062-K](https://doi.org/10.1016/0305-1978(90)90062-K)
 54. Lugtenberg BJ, Malfanova N, Kamilova F, Berg G. Plant growth promotion by microbes. *Molecular microbial ecology of the rhizosphere*. 2013;2:561-73.
 55. Vimal SR, Singh JS, Arora NK, Singh S. Soil-Plant-Microbe Interactions in Stressed Agriculture Management: A Review. *Pedosphere*. 2017;27(2):177-92. doi:[10.1016/S1002-0160\(17\)60309-6](https://doi.org/10.1016/S1002-0160(17)60309-6)
 56. Santana SRA, Voltolini TV, Antunes G dos R, da Silva VM, Simões WL, Morgante CV, et al. Inoculation of plant growth-promoting bacteria attenuates the negative effects of drought on sorghum. *Archives of Microbiology*. 2020;202(5):1015-24. doi:[10.1007/s00203-020-01810-5](https://doi.org/10.1007/s00203-020-01810-5)
 57. Puente ML, Zawoznik M, de Sabando ML, Perez G, Gualpa JL, Carletti SM, et al. Improvement of soybean grain nutritional quality under foliar inoculation with *Azospirillum brasilense* strain Az39. *Symbiosis*. 2019;77(1):41-7. doi:[10.1007/s13199-018-0568-x](https://doi.org/10.1007/s13199-018-0568-x)
 58. Jochum MD, McWilliams KL, Borrego EJ, Kolomiets MV, Niu G, Pierson EA, et al. Bioprospecting Plant Growth-Promoting Rhizobacteria That Mitigate Drought Stress in Grasses. *Frontiers in Microbiology*. 2019;0:1-9. doi:[10.3389/fmicb.2019.02106](https://doi.org/10.3389/fmicb.2019.02106)
 59. Barra PJ, Inostroza NG, Mora ML, Crowley DE, Jorquera MA. Bacterial consortia inoculation mitigates the water shortage and salt stress in an avocado *Persea americana* Mill.) nursery. *Applied Soil Ecology*. 2017;111:39-47. doi:[10.1016/j.apsoil.2016.11.012](https://doi.org/10.1016/j.apsoil.2016.11.012)
 60. Loredó-Ostí C, López-Reyes L, Espinosa-Victoria D. Bacterias promotoras del crecimiento vegetal asociadas con gramíneas: Una revisión. *Terra Latinoamericana*. 2004;22(2):225-39.
 61. Peñín López A. Rizobacterias promotoras del crecimiento vegetal [Internet]. de la Laguna; 2017. Available from: <https://riull.ull.es/xmlui/bitstream/handle/915/5766/Rizobacterias+promotoras+del+crecimiento+vegetal.pdf?sequence=1>
 62. Murcia Linares A, Bustos SC. Efecto de la inoculación de bacterias promotoras del crecimiento vegetal en maracuyá y badea cultivadas en condiciones de estrés hídrico. *Biología* [Internet]. 2017; Available from: <https://ciencia.lasalle.edu.co/biologia/14>
 63. Vurukonda SSKP, Vardharajula S, Shrivastava M, SkZ A. Enhancement of drought stress tolerance in crops by plant growth promoting rhizobacteria. *Microbiological Research*. 2016;184:13-24. doi:[10.1016/j.micres.2015.12.003](https://doi.org/10.1016/j.micres.2015.12.003)
 64. Van de Poel B, Van Der Straeten D. 1-aminocyclopropane-1-carboxylic acid (ACC) in plants: more than just the precursor of ethylene! *Frontiers in Plant Science*. 2014;5:1-11. doi:[10.3389/fpls.2014.00640](https://doi.org/10.3389/fpls.2014.00640)
 65. Ahmad F, Ahmad I, Khan MS. Screening of free-living rhizospheric bacteria for their multiple plant growth promoting activities. *Microbiological Research*. 2008;163(2):173-81. doi:[10.1016/j.micres.2006.04.001](https://doi.org/10.1016/j.micres.2006.04.001)
 66. Yang J, Kloepper JW, Ryu C-M. Rhizosphere bacteria help plants tolerate abiotic stress. *Trends in Plant Science*. 2009;14(1):1-4. doi:[10.1016/j.tplants.2008.10.004](https://doi.org/10.1016/j.tplants.2008.10.004)
 67. Mangmang J, Deaker R, Rogers G. Germination Characteristics of Cucumber Influenced by Plant Growth-promoting Rhizobacteria: *International Journal of Vegetable Science*. 22(1):66-75.
 68. Naveed M, Hussain MB, Zahir ZA, Mitter B, Sessitsch A. Drought stress amelioration in wheat through inoculation with *Burkholderia phytofirmans* strain PsJN. *Plant Growth Regulation*. 2014;73(2):121-31. doi: [10.1007/s10725-013-9874-8](https://doi.org/10.1007/s10725-013-9874-8)
 69. Vanderlinde EM, Harrison JJ, Muszyński A, Carlson RW, Turner RJ, Yost CK. Identification of a novel ABC transporter required for desiccation tolerance, and biofilm formation in *Rhizobium leguminosarum* bv. *viciae* 3841. *FEMS Microbiology Ecology*. 2010;71(3):327-40. doi:[10.1111/j.1574-6941.2009.00824.x](https://doi.org/10.1111/j.1574-6941.2009.00824.x)
 70. Smith SE, Read DJ. *Mycorrhizal Symbiosis*. Academic Press; 2010. 815 p.
 71. Augé RM. Water relations, drought and vesicular-arbuscular mycorrhizal symbiosis. *Mycorrhiza*. 2001;11(1):3-42. doi:[10.1007/s005720100097](https://doi.org/10.1007/s005720100097)

72. Ruiz-Lozano JM. Arbuscular mycorrhizal symbiosis and alleviation of osmotic stress. New perspectives for molecular studies. *Mycorrhiza*. 2003;13(6):309-17. doi:[10.1007/s00572-003-0237-6](https://doi.org/10.1007/s00572-003-0237-6)
73. Augé RM. Arbuscular mycorrhizae and soil/plant water relations. *Canadian Journal of Soil Science*. 2004;84:373-81. doi:[10.4141/S04-002](https://doi.org/10.4141/S04-002)
74. Azcón R, Barea JoséM. Mycorrhizal dependency of a representative plant species in mediterranean shrublands *Lavandula spica* L.) as a key factor to its use for revegetation strategies in desertification-threatened areas. *Applied Soil Ecology*. 1997;7(1):83-92. doi:[10.1016/S0929-1393\(97\)00013-9](https://doi.org/10.1016/S0929-1393(97)00013-9)
75. Amerian MR, Stewart WS, Griffiths H. Effect of two species of arbuscular mycorrhizal fungi on growth, assimilation and leaf water relations in maize *Zea mays*). *Aspects of Applied Biology*. 2001;63:71-6.