



Germination of rice seeds (*Oryza sativa* L.) at different temperatures

Germinación de semillas de arroz (*Oryza sativa* L.) a diferentes temperaturas

^{ID}Lázaro A. Maqueira López*, ^{ID}Rogelio Morejón Rivera, ^{ID}Osmany Roján Herrera,
^{ID}Yaisys Blanco Valdés, ^{ID}Ana Isabel Izquierdo Collazo

Unidad Científico Tecnológica de Base "Los Palacios". Instituto Nacional de Ciencias Agrícolas (INCA), carretera San José-Tapaste, km 3½, Gaveta Postal 1, San José de las Lajas, Mayabeque, Cuba. CP 32 700

ABSTRACT: The research was developed in the Base Scientific Science Unit, Los Palacios, Pinar del Río, belonging to the National Institute of Agricultural Sciences. The objective was to evaluate the influence of temperatures on the germination of four rice cultivars. Seeds of *Oryza sativa* L. (rice) were analyzed from the cultivars, INCA LP-5, Reforma, INCA LP-2 y Jucarito 104 (J-104). The germination temperatures considered in each trial were 10, 15, 20, 25, 30, 35 and 40 °C. The experimental design used in all cases was completely randomized, each trial was evaluated individually and was repeated twice, we worked with the means of these repetitions. Germinated seeds were counted from their establishment until stabilization, with the data obtained was determined: Percentage of germination, seeds germinated per day and the Maguire Index. The results were subjected to an analysis of variance (ANOVA) with a confidence level of 95 %. The results of this work indicate that there is a sensitivity in the germination process of different rice cultivars to temperature effects. Each cultivar shows a greater or lesser germination resistance depending on the temperature. Therefore, temperature effect in the process is closely related to the genetic material with which it is working. In addition, with a germination speed increase in the increase in temperature, an optimal value is reached and then decreases, regardless of the cultivar.

Key words: cereal, vigor, reserve.

RESUMEN: La investigación se desarrolló en la Unidad Científico Tecnológica de Base, Los Palacios, Pinar del Río, perteneciente al Instituto Nacional de Ciencias Agrícolas. El objetivo fue evaluar la influencia de las temperaturas en la germinación de cuatro cultivares de arroz. Se analizaron semillas de *Oryza sativa* L. (arroz) de los cultivares INCA LP-5, Reforma, INCA LP-2 y Jucarito 104 (J-104). Las temperaturas de germinación consideradas en cada ensayo fueron 10, 15, 20, 25, 30, 35 y 40°C. El diseño experimental utilizado en todos los casos fue completamente aleatorizado, cada ensayo fue evaluado individualmente y repetido dos veces, se trabajó con las medias de dichas repeticiones. Se contaron las semillas germinadas desde su establecimiento hasta la estabilización, con los datos obtenidos se determinó: porcentaje de germinación, semillas germinadas por día, Índice Maguire (IM) y número medio de días germinados (NMDG). Los resultados fueron sometidos a un análisis de varianza (ANOVA) con un nivel de confianza del 95%. Los resultados de este trabajo indican que existe sensibilidad en el proceso de germinación de diferentes cultivares de arroz ante los efectos de la temperatura. Cada cultivar muestra una mayor o menor resistencia de germinación en función de la temperatura. Por lo tanto, el efecto de la temperatura en el proceso está estrechamente relacionado con el material genético con que se esté trabajando. Además, con el incremento en la velocidad de germinación a partir del aumento de la temperatura, se alcanza un valor óptimo para después disminuir, independientemente del cultivar.

Key words: cereal, vigor, reserva.

*Author for correspondence. lazaro@inca.edu.cu

Received: 20/01/2021

Accepted: 25/08/2021

Conflict of interest: The authors declare that they have no conflict of interest.

Authors' contribution: **Conceptualization-** Lázaro A Maqueira López, Yaisys Blanco Valdés. **Research-** Lázaro A Maqueira López, Osmany Roján Herrera, Rogelio Morejón Rivera. **Methodology-** Lázaro A Maqueira López, Osmany Roján Herrera, Rogelio Morejón Rivera, Yaisys Blanco Valdés. **Supervision-** Lázaro A Maqueira López. **Writing of initial draft-** Lázaro A Maqueira López, Yaisys Blanco Valdés, Osmany Roján Herrera, Rogelio Morejón Rivera, Ana Isabel Izquierdo Collazo. **Writing of the final draft-** Lázaro A Maqueira López, Yaisys Blanco Valdés.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial (BY-NC 4.0).

<https://creativecommons.org/licenses/by-nc/4.0/>



INTRODUCTION

Rice, wheat and maize are among the most important grains for human consumption worldwide.

Rice (*Oryza sativa*) is consumed by more than half of the population and provides 27 % of the food calories consumed by the population. Approximately 900 million of the world's poor depend on rice as consumers and/or producers (1).

In Cuba, rice occupies only 7.6 % of the 2,733.5 million hectares (ha) of land in the country of cultivated land (2). The agricultural yield in the country for the period 2013 to 2016 was 3.07 t ha⁻¹ although yield potentials of between 6.4 to 7.7.5 t ha⁻¹ have been between 6.4 to 7.4 t ha⁻¹ for cold planting and 4.4 to 5.3 t ha⁻¹ for cold planting 5.3 t ha⁻¹ for spring plantings. In the last five years, the country's production behavior in the country is similar to that of the aforementioned period. In order to meet the demands of the Cuban population, it has been necessary to import more than 50 % of the rice consumed in the country (3). All of the above is the reason why strategies are being followed to increase yields and reduce imports. The implementation of actions is strengthened where, genetic improvement of cultivars; seed production and the achievement of a correct phytotechnical with emphasis on achieving germination that guarantees an adequate plant density are among the most important (4).

Rice, wheat and and maize are among the most important cereals for human consumption worldwide. worldwide. Rice (*Oryza sativa*) is consumed by more than half of the world's population and provides 27 % of the calories provide 27 % of the calories of the food consumed by the population. Approximately 900 million of the world's poor depend on rice as consumers and/or producers (1) consumers and/or producers (1).

However, the response of rice cultivars to different environmental to different environmental conditions and to the great variability of climate is now a vital element in the is now a vitally important element in achieving high productivity in the face of climate change effects.

productivity in the face of the effects of climate change (5). In general, crop production is affected by climate variability and especially by the effect of temperatures. It is therefore important to know the most sensitive developmental stages and phenological stages, as well as the detrimental effect of temperatures on the processes (6). Of the phases and stages of cultivation, the germination process is considered one of the most affected by temperature due to its influence on the activity of enzymes regulating the speed of biochemical reactions occurring in the seed after rehydration (7).

At present, there is little information on temperature consequences on the germination of rice cultivars obtained in the country and a precise understanding of germination dynamics at different temperatures is required to ensure a number of seedlings in production practice. In in addition, this experiment is important to promote programs for the improvement of cultivars according to climate adaptation in the agricultural sector. Therefore, this work was carried out with the objective of evaluating the influence of temperatures on the germination of four rice cultivars.

MATERIALS AND METHODS

Experiments were carried out at the Base Scientific Science Unit, Los Palacios, Pinar del Río, belonging to the National Institute of Agricultural Sciences. Seeds of *Oryza sativa* L (rice) of four cultivars were analyzed: INCA LP-5, Reforma, INCA LP-2 and Jucarito 104 (J-104) (Table 1) (8, 9). These were supplied by the INCA seed group.

Once the crop was harvested, and with the purpose of lowering its humidity to 12 %, seeds were dried in an oven with forced air at 30 °C. Seed moisture was determined with a Japanese Kett J5 moisture meter. They were then stored in tightly covered plastic containers.

In the experiment, the germination temperatures considered in each trial were 10, 15, 20, 25, 30, 35 and 40 °C. In each trial, the experimental unit corresponded to glass petri dishes 140 mm in diameter and 20 mm high, with 2 layers of filter paper moistened with distilled water on the bottom of the dishes and 40 seeds inside. Four

Table 1. Main biological characteristics of the rice (*Oryza sativa* L.) cultivars studied in the experiments

Characteristics	Cultivars			
	INCA LP-5	Reforma	INCA LP-2	Jucarito 104
Length of cycle (days)				
Low rainy season	128	122	134	147
Rainy" season	110	105	114	119
Classification according to cycle	Short	Short	Medium	Medium
Plant height (cm)	90	90-110	88	85
Panicle length (cm)	23	26	24	23
Grains per panicle	90	150	90	112
Mass of thousand grains (g)	29.5	26.6	29	28
Yield (t ha ⁻¹)				
Low rainy season	8.2	7.5	7.8	8.6
Rainy" season	5.7	5.7	6.7	5.9

replicates per treatment were established. Once seeds were placed on the plates, they were placed in a germination chamber regulated at the respective temperature and in dark conditions. The experimental design used in all cases was completely randomized; each trial (temperatures 10, 15, 20, 25, 30, 35 and 40 °C) was evaluated individually and was repeated twice, working with the means of these replicates.

The germinated seeds were counted from establishment to stabilization, and the data obtained were used to determine:

- The germination percentage (GP): germinated seed was considered as that with a radicle length greater than or equal to 2 mm (10).

$$GP = \frac{\text{Number of germinated seeds}}{\text{Number of sowed seeds}} * 100 \quad (1)$$

- Germinated seeds per day (SGD): Germinated seeds were counted daily and the criterion used was the appearance of the radicle greater than or equal to 2 mm.
- Maguire index (IM): It represents the germination rate calculated through a weighted time (in days) of accumulated germination. Where G is the percentage of seedlings that germinate during the time interval t (11).

$$IM = \sum_{t=1}^n (G/t) \quad (2)$$

- Mean number of days germinated (NMDG) where: Ni is the number of seeds germinated within consecutive time intervals Ti is the time in days elapsed between the start of the test and the end of the interval, n is the total number of seeds germinated (12).

$$NMDG = \sum_{i=1}^n \frac{N_i * T_i}{n} \quad (3)$$

The assumptions of normality and homogeneity of variance (Bartlett's test and Akolmogorov-Smirnov, respectively) were checked for the data of each variable evaluated. The results (PG, MI) were subjected to an analysis of variance (ANOVA) with a confidence level of 95 %. Comparison of means was performed using Tukey's Multiple Range Test ($P > 0.05$) (13). For this purpose, the STATGRAPHICS Plus 5.0 Centurion program on Windows, version XV (14) was used.

RESULTS AND DISCUSSION

The final germination percentages are shown in Table 2. All the cultivars showed significant differences in the temperature range from 10 to 40 °C. The highest values were observed in the range of 10 to 40 °C. The highest values were observed between 20 and 30 °C, while the lowest values were observed at 10 °C. INCA LP-2 was the cultivar that best responded to effects of the different temperatures studied, with germination percentages ranging from 53 to 97 %. The cultivar INCA LP-5 was the most affected by the effects of temperatures of 10 and 15 °C, since only 13 and 30 % germinated, respectively.

The influence of temperature on the germination percentage of seeds of crops of agricultural interest has been reported by several authors, who emphasize that the percentage of seed emergence guarantees 50 % of the success of production and point out that germination only occurs properly within a certain temperature range (15, 16). In studies carried out in peanut cultivation, it was demonstrated in all the genotypes studied that the germination response increased with the rise in temperature above 14 °C. For the range between 16 and 32 °C, germination percentages were obtained that meet the standard for commercialization of common seed for peanut (> 80 %); which was in all genotypes equal to or higher than 90 % (17). However, it should be noted that in the results of the present study with rice cultivars, there is a differential and particular response of the cultivars to germination temperature since there is variation in final percentages.

With the Maguire index (Figure 1), where the accumulated germination was weighted over time, it was possible to differentiate the role of temperature with respect to the germination process of the different cultivars. Figure 1 shows that there are statistically significant differences between the different temperatures for each cultivar.

In general, an increase in the values of the index is observed with increasing temperature, and then decreases, so that a well-defined maximum of 30 °C was observed for all the cultivars under study. However, the cultivars at 10 and 15 °C had a low index (between 0.53 and 6.4) and showed in this sense a considerable affectation in germination speed as a response to the presence of low temperatures. Although the values in INCA LP-2 are higher

Table 2. Rice seed germination percentages at different temperatures

Temperature (°C)	Cultivars			
	INCA LP-5	Reforma	INCA LP-2	J-104
10	13.13 e	28.13 de	53.13 d	53.13 c
15	30.63 d	40.00 cd	71.25 c	40.00 b
20	98.13 a	90.00 a	81.88 b	91.88 a
25	100.00 a	98.75 a	96.88 a	99.38 a
30	84.40 b	63.13 b	96.88 a	93.13 a
35	72.50 c	49.38 bc	93.75 a	48.13 b
40	39.38 d	14.38 e	68.75 b	20.63 c
SE x	0.06***	0.05***	0.03***	0.06***

Different letters in the same column indicate significant differences (Tukey's test, $p \leq 0.05$)




Figure 1. Maguire index of rice seeds of four cultivars (INCA LP-5, Reforma, INCA LP-2 and Jucarito 104) at different temperatures (10, 15, 20, 25, 30, 35 and 40 °C)

than those ones of the rest. It may be related to the genetic characteristics of this cultivar. This cultivar could be taken into account for further studies in which tolerance of cultivars to relatively low temperatures is sought and for the selection of progenitors in the rice-breeding program seeking tolerance to abiotic stress. In this regard, some authors state that even when it is impossible to eliminate all stress factors to which crops are subjected, it is of vital importance the knowledge of the way in which these cause their effects in the physiological processes of plants. The identification of crops and/or cultivars that survive and tolerate damages that can act more severely to others of the same species, since it is possible to modify them genetically or minimize the effects that they are capable of producing (18).

Another aspect to highlight is that in the cultivar J-104 the values of this index decrease considerably at temperatures of 35 and 40 °C (values between 3 and 7 approximately), aspect that can also be related to the genetic characteristics of this cultivar and demonstrates its susceptibility to high temperatures during the germination process. In the climatic conditions of Cuba, it is of great interest the problem that exists regarding the decrease of germination in the different existing cultivars at temperatures lower than 15 °C and above 30°C. This is because the sowing of the crop is done in two seasons, "cold" and "spring", so that the existing climatic conditions during the germination stage could increase or decrease the days for a given cultivar, especially for its greater or lesser susceptibility to temperatures during the germination process. There are also studies where seed germination indexes are used for the early selection of genotypes tolerant to certain stresses

(16, 19). Several authors highlight the importance of this index (IM), since it weights and combines the parameters germination capacity, speed and uniformity to evaluate the quality of germination; the higher the result, the higher the quality of the seed and the response of the cultivars to the conditions in the germination process (20).

In aspects pointed out by some authors, it is highlighted that the poor germination of seeds at high temperatures is related to the synthesis of proteins in the embryo, which influences its germination speed. In this regard, it is reported that several processes occur in the seed that depend solely on the reserves, the gibberellic acid of the embryo acts on the aleurone layer where the enzyme amylase is activated, which initiates the degradation of reserve substances contained in the endosperm and cotyledon. From the digestion of reserve tissues, various compounds are released to be reused in multiple synthesis processes. Complex molecules such as celluloses, hemicelluloses, starches, amylopectins, lipids, lignins, proteins, nucleic acids, vitamins and hormones are degraded to simple molecules by specific enzymes (21). Other authors emphasize the role of the pericarp in the thermo inhibition of seeds, pointing out that this is partly due to the presence of inhibitors in the tissue or also due to the fact that in the presence of high temperatures the oxygen requirements are relatively high and do not allow the pericarp to maintain these demands (22).

The results of the present work coincide with studies carried out by other authors, who report that in most cases the germination speed increases with increasing temperature although very high temperatures also tend to decrease it. With studies using various temperatures within

a certain range (10-40 °C), the authors state that it is possible to find the optimum germination temperature. They emphasize in their results that temperatures above 32 °C can decrease the germination speed index of seeds. However, regardless of the genotype, increasing the temperature up to 38 °C significantly decreases the germination speed (22, 23). Other authors (24) also exposed the negative effect of high temperatures on germination speed. They pointed out that once the optimum temperature level is reached, where germination speed is higher, a decrease occurs as temperatures approach their maximum limit where irreversible damage to seeds occurs.

The average number of days to germination in the rice cultivars studied varied considerably according to the germination temperature (Table 3). In general, a different behavior to that observed for germination speed (Maguire index) was observed, since as temperature increased, the mean number of days to germination decreased for all the cultivars under study. The cultivar INCA LP-2 showed a shorter time (approximately 7 days) for the germination process at low temperatures (10 °C) compared to the rest of the cultivars studied, which required 10 days. However, at high temperatures (40 °C) the result was different.

The NMDG has been used by several authors to highlight differences in the seed germination process in certain studies. When comparing the germination of sorghum seeds of different caliber, it was observed that seeds of larger caliber presented the lowest values of this variable (10). On the contrary, in a study carried out in the cultivation of corn where three cultivars were used and their seeds were submitted before the germination test to different immersion times in water at 100 °C (0, 2, 4, 6 and 8 seconds). As a result, there was no significant response between treatments in terms of NMDG, since the values ranged between four and five days (25). However, the results of this study clearly show the resistance to germination of each cultivar at the different temperatures to which seeds were subjected, due to the significant differences between treatments. This aspect is highlighted by other authors (26) who have determined the NMDG for this purpose.

CONCLUSIONS

The results of this work indicate that there is sensitivity in the germination process of different rice cultivars to the effects of temperature. Each cultivar shows greater or lesser germination resistance as a function of temperature. Therefore, the action of temperature on the process is closely related to the genetic material being worked with. In addition, with an increase in germination speed as the temperature increases, an optimum value is reached and then decreases, regardless of the cultivar.

BIBLIOGRAPHY

1. Saito, K.; Asai, H.; Zhao, D.; Laborte, A.G. y Grenier, C. "Progress in varietal improvement for increasing upland rice productivity in the tropics", *Plant Production Science*, vol. 21, no. 3, 3 de julio de 2018, pp. 145-158, ISSN 1343-943X, DOI [10.1080/1343943X.2018.1459751](https://doi.org/10.1080/1343943X.2018.1459751), [Consultado: 18 de julio de 2023], Disponible en: <https://doi.org/10.1080/1343943X.2018.1459751>.
2. Inst. Oficina Nacional de Estadísticas e Información *Anuario Estadístico de Cuba. Año 2016* [en línea], Oficina Nacional de Estadísticas e Información, La Habana, Cuba, 2017, http://www.onei.gob.cu/sites/default/files/00_anuario_estadistico_2016.pdf, [Consultado: 18 de julio de 2023], Disponible en: <http://www.onei.gob.cu/node/13608>.
3. Julián Herrera Puebla; Guillermo Hervis Granda; Felicita González Robaina; y Carmen Duarte Díaz "Estudio sobre el balance hídrico del arroz en Cuba | Ingeniería Agrícola", vol. 9, no. 3, 2019, pp. 48-59, [Consultado: 18 de julio de 2023], Disponible en: <https://ojs.edicionescervantes.com/index.php/IAgric/article/view/1143>.
4. Pérez, Á.L.P.; López, L.A.M.; Noval, W.T. de la.; Pérez, A.S.; Mesa, S.A.P. y Lescano, G.G. "Comportamiento de las variables clima en áreas de la Empresa Agroindustrial de Granos Los Palacios y su influencia en el rendimiento de cultivares de arroz", *Avances*, vol. 17, no. 2, 17 de agosto de 2015, pp. 184-193, ISSN 1562-3297, [Consultado: 18 de julio de 2023], Disponible en: <http://www.ciget.pinar.cu/ojs/index.php/publicaciones/article/view/11>.

Table 3. Mean number of days to germination (NMDG) of rice seeds at different temperatures

Temperature (°C)	Cultivars			
	INCA LP-5	Reforma	INCA LP-2	J-104
10	10.00 a	10.00 a	6.70 a	10.00 a
15	5.90 b	6.00 b	4.65 b	5.58 b
20	3.54 d	3.87 e	3.91 c	4.28 c
25	4.24 c	4.62 c	3.46 d	4.01 d
30	2.09 f	2.20 f	1.68 g	1.91 g
35	2.53 e	1.94 g	1.95 f	2.57 e
40	2.44 e	4.00 d	2.79 e	2.33 f
SE x	0.50***	0.48***	0.31***	0.50***

Different letters in the same column indicate significant differences (Tukey's test, $p \leq 0.05$)

5. FAO. 2018. *El cambio climático, la agricultura y la seguridad alimentaria*. [en línea], junio de 2018, Disponible en: https://www.google.com/cu/url?sa=t&rct=j&q=&esrc=s&source=web&cd=3&cad=rja&uact=8&ved=0ahUKEwjVzcxxeXaAhVD0IMKH6ODcAQFggYMAI&url=http%3A%2F%2Fwww.fao.org%2F3%2Fa-i5349s.pdf&usg=AOvVaw3_qKHYNbE2MVCnP-jBivC.
6. Chaves-Barrantes, N.F. y Gutiérrez-Soto, M.V. "Respuestas al estrés por calor en los cultivos. II. Tolerancia y tratamiento agronómico", *Agronomía Mesoamericana*, vol. 28, no. 1, 2017, pp. 255-271, [Consultado: 18 de julio de 2023], Disponible en: <https://www.redalyc.org/journal/437/43748637021/html>.
7. Obregón P. *La germinación* [en línea], *Monografías.com. Agricultura y ganadería*, 2007, 4 de enero de 2018, Disponible en: <http://www.monografias.com/trabajos70/germinacion/germinacion.shtml>.
8. Puldon Padron Violeta; Faure. B; Cantillo P. I; y Toledo. D *Catálogo de Variedades del Instituto de Investigaciones de Granos Arroz, Maíz, Frijol, Soya.* | ISBN 978-959-285-060-6 - Libro [en línea], Ediciones IIGranos, Artemisa, 2018, (ser. TECNOLOGÍA (CIENCIAS APLICADAS)), p. 93, ISBN 978-959-285-060-6, [Consultado: 18 de julio de 2023], Disponible en: <https://isbn.cloud/9789592850606/catalogo-de-variedades-del-instituto-de-investigaciones-de-granos-arroz-maiz-frijol-soya/>.
9. Valdés, E.; Díaz Valdés, E.C.; Blanco Reinoso, G.; González Cepero, M.C.; Pérez León, N. de J. y Díaz Solís, S.H. *Cultivares Cubanos de Arroz* [en línea], INCA, 2018, (ser. TECNOLOGÍA (CIENCIAS APLICADAS)), p. 25, ISBN 978-959-7258-00-1, [Consultado: 18 de julio de 2023], Disponible en: <https://isbn.cloud/9789597258001/cultivares-cubanos-de-arroz/>.
10. Ruiz-Sánchez, M.; Muñoz-Hernández, Y.; Guzmán, D.; Velázquez-Rodríguez, R.; Díaz-López, G.S.; Martínez, A.Y. y Almeida, F.M. "Efecto del calibre semilla (masa) en la germinación del sorgo", *Cultivos Tropicales*, vol. 39, no. 4, diciembre de 2018, pp. 51-59, ISSN 0258-5936, [Consultado: 18 de julio de 2023], Disponible en: http://scielo.sld.cu/scielo.php?script=sci_abstract&pid=S0258-59362018000400007&lng=es&nrm=iso&tlng=es.
11. Ferraz, M.; Franco, C.; Batista, G. s. y Pivetta, K. "Salinity on the germination of seed and index of germination speed of three ornamental species", *Ornamental Horticulture*, vol. 22, 1 de septiembre de 2016, pp. 196-201, DOI 10.14295/oh.v22i2.919.
12. Duarte, E.; González, B. y Rocha Peña, M.A. "Análisis físico y germinación de semillas de «Austrochthamalia teyucuaensis» H. A. Keller, endémica del bosque atlántico y en peligro de extinción", *Agrociencia*, *Agrociencia*, vol. 53, no. 3, 2019, pp. 403-416, ISSN 1405-3195, [Consultado: 18 de julio de 2023], Disponible en: <https://dialnet.unirioja.es/servlet/articulo?codigo=6933814>.
13. Duncan, D.B. "Multiple Range and Multiple F Tests", *Biometrics*, vol. 11, no. 1, 1955, pp. 1-42, ISSN 0006-341X, DOI 10.2307/3001478, [Consultado: 18 de julio de 2023], Disponible en: <https://www.jstor.org/stable/3001478>.
14. *Statistical Graphics Crop. STATGRAPHICS® Plus* [en línea], 2000, Disponible en: <http://www.statgraphics.com/statgraphics/statgraphics.nsf/pd/pdpricing>.
15. Kayongo SY y Andersson P (PDF) FARMER PARTICIPATORY EVALUATION OF BEAN (*Phaseolus vulgaris* L) VARIETIES FOR SEED PRODUCTION IN TESO- KARAMOJA SUB-REGION, UGANDA", *International Journal of Research In Agriculture and Food Sciences*, vol. 2, no. 3, 2014, pp. 58-60, [Consultado: 18 de julio de 2023], Disponible en: https://www.researchgate.net/publication/282076329_FARMER_PARTICIPATORY_EVALUATION_OF_BEAN_Phaseolus_vulgaris_L_VARIETIES_FOR_SEED_PRODUCTION_IN_TESO_KARAMOJA_SUB-REGION_UGANDA.
16. Aflaki, F.; Sedghi, M.; Pazuki, A. y Pessarakli, M. "INVESTIGATION OF SEED GERMINATION INDICES FOR EARLY SELECTION OF SALINITY TOLERANT GENOTYPES: A CASE STUDY IN WHEAT", *Emirates Journal of Food and Agriculture*, 15 de febrero de 2017, pp. 222-226, ISSN 2079-0538, DOI 10.9755/ejfa.2016-12-1940, [Consultado: 18 de julio de 2023], Disponible en: <https://www.ejfa.me/index.php/journal/article/view/595>.
17. Caroca R; Zapata N; y Vargas M "EFECTO DE LA TEMPERATURA SOBRE LA GERMINACIÓN DE CUATRO GENOTIPOS DE MANÍ (*Arachis hypogaea* L.)", *Chilean journal of agricultural & animal sciences*, vol. 32, no. 2, 2016, pp. 94-101, [Consultado: 18 de julio de 2023], Disponible en: https://www.scielo.cl/scielo.php?script=sci_arttext&pid=S0719-38902016000200002.
18. Torres, W. "Aspecto de la biología de las plantas ante condiciones de estrés abiótico." [en línea], ed. Cornide, M.T., *Genética, genómica y fitomejoramiento.*, vol. 1, edit. Felix Varela, La Habana (Cuba), 2018, pp. 211-255.
19. Vitória, R.Z.D.; Oliveira, F.D.T.G.D.; Posse, S.C.P.; Arantes, S.D.; Schmildt, O.; Viana, A.; Malikuski, R.G. y Barros, B.L.A. "Qualidade fisiológica de sementes de aroeira em função da maturação dos frutos sob diferentes temperaturas de germinação.", *Nucleus*, vol. 15, no. 2, 30 de octubre de 2018, pp. 575-582, ISSN 1678-6602, 1982-2278, DOI 10.3738/1982.2278.2870, [Consultado: 18 de julio de 2023], Disponible en: <http://www.nucleus.feituverava.com.br/index.php/nucleus/article/view/2870/2729>.
20. Romero-Rangel, S.; Rubio-Licona, L.E.; Chávez-Serrano, L.; Rojas-Zenteno, E.C. y García-Pineda, M. "COMPORTAMIENTO GERMINATIVO Y CRECIMIENTO TEMPRANO DE PINUS DEVONIANA Y PINUS PSEUDOSTROBUS (PINACEAE)", *BIOCYT Biología Ciencia y Tecnología*, vol. 10, no. 37-39, 11 de septiembre de 2017, ISSN 2007-2082, DOI 10.22201/fesi.20072082.2017.10.61719, [Consultado: 18 de julio de 2023], Disponible en: <https://www.revistas.unam.mx/index.php/biocytt/article/view/61719>.
21. Morales-Santos, M.E.; Peña-Valdivia, C.B.; García-Esteva, A.; Aguilar-Benítez, G.; Kohashi-Shibata, J.

- Morales-Santos, M.E.; Peña-Valdivia, C.B.; García-Esteva, A.; Aguilar-Benítez, G. y Kohashi-Shibata, J. "Características físicas y de germinación en semillas y plántulas de frijol (*Phaseolus vulgaris* L.) silvestre, domesticado y su progenie", *Agrociencia*, vol. 51, no. 1, febrero de 2017, pp. 43-62, ISBN 1405-3195, [Consultado: 18 de julio de 2023], Disponible en: <http://www.scielo.org.mx/scielo.php?script=sci_abstract&pid=S1405-31952017000100043&lng=es&nrm=iso&tlng=es>.
22. Drew, R.L.K. y Brocklehurst, P.A. "Investigations on the Control of Lettuce Seed Germination at High Temperatures", *Journal of Experimental Botany*, vol. 35, no. 156, 1984, pp. 986-993, ISSN 0022-0957, [Consultado: 18 de julio de 2023], Disponible en: <<https://www.jstor.org/stable/23691115>>.
 23. Grey, T.L.; Beasley, J.P.; Webster, T.M. y Chen, C.Y. "Peanut Seed Vigor Evaluation Using a Thermal Gradient", *International Journal of Agronomy*, vol. 2011, 4 de agosto de 2011, p. e202341, ISSN 1687-8159, DOI [10.1155/2011/202341](https://doi.org/10.1155/2011/202341), [Consultado: 18 de julio de 2023], Disponible en: <<https://www.hindawi.com/journals/ija/2011/202341/>>.
 24. Finch-Savage, W.E. "The use of population-based threshold models to describe and predict the effects of seedbed environment on germination and seedling emergence of crops", *Haworth Press, New York, USA.*, 2004, (ser. applications to agriculture), pp. 51-96, Disponible en: <[https://www.scirp.org/\(S\(351jmbntvnsjt1aadkposzje\)\)/reference/ReferencesPapers.aspx?ReferenceID=1626981](https://www.scirp.org/(S(351jmbntvnsjt1aadkposzje))/reference/ReferencesPapers.aspx?ReferenceID=1626981)>.
 25. González-Rodríguez, F.; Borges-Gómez, L.; Pinzón-López, L.; Magaña-Magaña, M.; Sanginés-García, R. y Urrestarazu-Gavilán, M. "Inmersión de semillas de maíz en agua caliente en la producción de germinados para forraje", *Agronomía Mesoamericana*, vol. 25, no. 1, junio de 2014, pp. 54-62, ISSN 1659-1321, [Consultado: 18 de julio de 2023], Disponible en: <http://www.scielo.sa.cr/scielo.php?script=sci_abstract&pid=S1659-13212014000100006&lng=en&nrm=iso&tlng=es>.
 26. Asier Jáñez Ortiz de Landaluze *Desarrollo de los protocolos de germinación y viabilidad de semillas de dos especies de interés para la conservación en la capv: una rutácea, *Haplophyllum linifolium* y una liliácea, *Lilium pyrenaicum gouan** [en línea] [Tesis Maestría], Universidad del País Vasco, Leioa, 2015, p. 51, [Consultado: 18 de julio de 2023], Disponible en: <<https://www.vitoria-gasteiz.org/docs/wb021/contenidosEstaticos/adjuntos/es/31/30/83130.pdf>>.



Available in:

<https://www.redalyc.org/articulo.oa?id=193279251001>

How to cite

Complete issue

More information about this article

Journal's webpage in redalyc.org

Scientific Information System Redalyc
Diamond Open Access scientific journal network
Non-commercial open infrastructure owned by academia

Lázaro A. Maqueira López, Rogelio Morejón Rivera,
Osmany Roján Herrera, Yaisys Blanco Valdés,
Ana Isabel Izquierdo Collazo

**Germinación de semillas de arroz (*Oryza sativa* L.) a
diferentes temperaturas**

**Germination of rice seeds (*Oryza sativa* L.) at different
temperatures**

Cultivos Tropicales
vol. 44, no. 4, e01, 2023
Ediciones INCA,
ISSN-E: 1819-4087