



Chemical characterization and evaluation of the biological activity of Spirulina extracts

Caracterización química y evaluación de la actividad biológica de extractos de Spirulina

^{ID}Anaysa Gutierrez Almeida*, ^{ID}Miriam de la C. Núñez Vázquez, ^{ID}Yanelis Reyes Guerrero, ^{ID}Geydi Pérez Domínguez, ^{ID}Lisbel Martínez González

Instituto Nacional de Ciencias Agrícolas (INCA), carretera San José-Tapaste, km 3½, Gaveta Postal 1, San José de las Lajas, Mayabeque, Cuba. CP 32 700

ABSTRACT: The increase in the yield and productivity of crops, without affecting the environment, is one of the challenges that farmers experience today. The use of Spirulina (*Arthrospira platensis*) extracts as a plant biostimulant is one of the most viable options to use for these purposes. The objective of this work was to obtain, characterize and evaluate the biological activity of some alcoholic extracts of Spirulina. For this, extracts were prepared from Spirulina powder using ethanol as solvent in two concentrations (70 and 90 %), two mass:solvent ratios (1:20 and 1:10) and two maceration times (10 and 21 days) and the content of proteins, phenols and flavonoids of each was determined by spectrophotometric determinations. Biological activity was evaluated using a rice seed germination assay from seed imbibition for 24 h and the germination process was carried out in Petri dishes with distilled water for seven days. The results obtained from the chemical characterization will allow to adapt the extraction parameters depending on the components that are to be favored and the physiological effect that is to be achieved, since the concentration of proteins, phenols and flavonoids in the extracts significantly influenced the percentage end of germination of rice seeds.

Key words: *Arthrospira platensis*, composition, germination, rice.

RESUMEN : El incremento en el rendimiento y la productividad de los cultivos, sin afectar el medio ambiente, es uno de los desafíos que experimentan los agricultores en la época actual. El empleo de extractos de Spirulina (*Arthrospira platensis*) como bioestimulante vegetal, es una de las opciones más viables a utilizar con estos fines. El objetivo del presente trabajo fue la obtención, caracterización y evaluación de la actividad biológica de algunos extractos alcohólicos de Spirulina. Para ello, se prepararon los extractos a partir de Spirulina en polvo utilizando etanol como solvente en dos concentraciones (70 y 90 %), dos relaciones masa-solvente (1:20 y 1:10) y dos tiempos de maceración (10 y 21 días) y se determinó el contenido de proteínas, fenoles y flavonoides de cada uno mediante determinaciones espectrofotométricas. La actividad biológica fue evaluada utilizando un ensayo de germinación de semillas de arroz a partir de la imbibición de semillas durante 24 h y el proceso de germinación se llevó a cabo en placas Petri con agua destilada durante siete días. Los resultados obtenidos de la caracterización química permitirán adecuar los parámetros de extracción en función de los componentes que se quieran favorecer y el efecto fisiológico que se quiera lograr, puesto que, la concentración de proteínas, fenoles y flavonoides en los extractos influyó significativamente en el porcentaje final de germinación de las semillas de arroz.

Palabras clave: *Arthrospira platensis*, composición, germinación, arroz.

*Autor para correspondencia: anaysa@inca.edu.cu

Received: 08/07/2023

Accepted: 12/05/2024

Conflict of interest: Authors declare that they have no conflict of interest.

Authors' contribution: **Conceptualization-** Miriam de la C. Núñez Vázquez, Yanelis Reyes Guerrero, Anaysa Gutierrez Almeida.

Research- Anaysa Gutierrez Almeida, Miriam de la C. Núñez Vázquez, Yanelis Reyes Guerrero, Geydi Pérez Domínguez, Lisbel Martínez González. **Supervision-** Miriam de la C. Núñez Vázquez, Yanelis Reyes Guerrero. **Methodology** - Anaysa Gutierrez Almeida, Miriam de la C. Núñez Vázquez, Yanelis Reyes Guerrero, Geydi Pérez Domínguez, Lisbel Martínez González. **Writing of the initial draft** - Anaysa Gutierrez Almeida. **Revision and editing of initial draft** - Yanelis Reyes Guerrero, Miriam de la C. Núñez Vázquez, Anaysa Gutierrez Almeida. **Final writing and editing-** Miriam de la C. Núñez Vázquez, Yanelis Reyes Guerrero, Anaysa Gutierrez Almeida. **Review and approval of final draft and data curation-** Miriam de la C. Núñez Vázquez, Yanelis Reyes Guerrero, Anaysa Gutierrez Almeida.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial (BY-NC 4.0). <https://creativecommons.org/licenses/by-nc/4.0/>



INTRODUCTION

In recent years, natural products based on algae and cyanobacteria are being used as agrochemical substitutes and have acquired great importance due to the benefits they have on crops and the reduced impact they have on the environment. It has been proven that their application increases certain metabolic and physiological expressions in plants. These products, such as algae and cyanobacteria extracts, are generally obtained by the use of a solvent and an adequate extraction process, and are mainly marketed as biofertilizers due to their high content of macro and micronutrients or as biostimulants because they contain, among other compounds, plant growth promoting hormones (1-4).

Spirulina (*Arthrospira platensis*), one of the most widely used for these purposes, is a microscopic blue-green cyanobacterium, where the blue color comes from the phycocyanin present and the green from the chlorophyll, and it was considered a microalgae until very recently. It derives its name from the spiral nature of its filaments and has become an object of scientific study due to its bioavailability of nutrients, since 85-95 % are assimilable (5).

It has approximately 60-70 % of its dry mass in proteins with high bioavailability, it also contains chlorophylls, as well as phenolic compounds and flavonoids that can act as natural antioxidants (6-8). It is the terrestrial and aquatic organism with the highest protein content and the best aminogram and digestibility (7); therefore, it is widely used as a source of amino acids for humans, animals and plants. It also contains essential polyunsaturated fatty acids and vitamins, as well as xanthines, phycobiliproteins (9,10), carbohydrates, nitrogen, phosphorus, potassium, calcium, iron, manganese, zinc (10). It also has a high content of vitamins B12, B1, B2, B6 and E, biotin, pantothenic acid, folic acid, inositol and niacin (10), halogenated compounds, polyketides, agar agar, alginic acid and carrageenan (11), great richness in α - and β -carotenes (7,12), phycocyanin, considerable amounts of α -linolenic acid (polyunsaturated fatty acid with different beneficial effects) (13), a high concentration of phytohormones, trace elements, antioxidants and polysaccharides, therefore, it is an excellent biological supplement (14).

The effects that the Spirulina application has caused in different plant species have been reported by several authors. Thus, in *Amaranthus gangeticus*, it has been found that imbibition of seeds and foliar application of Spirulina extracts increased protein (14) and iron levels in plants (15). Similarly, it was reported that imbibition of *Phaseolus aureus* and *Solanum lycopersicum* L. seeds in extracts of this cyanobacterium increased Zn levels in plants (16). In the species *Solanum melongena* L., the application of a commercial fertilizer based on Spirulina increased plant yield without affecting foliar levels of N, P, K and Na or quality indicators (17). In addition, it was found that Spirulina platensis extract has positive effects on wheat and barley seed germination, as well as on root and stem lengths (18).

In Cuba, Spirulina has been cultivated for more than three decades; however, its use for agricultural purposes has been

very limited and there is no information available about the use of extracts for these purposes. For this reason, the objective of the present work was to obtain, characterize and evaluate the biological activity of some alcoholic extracts of Spirulina.

MATERIALS AND METHODS

The experiment was carried out at the Department of Plant Physiology and Biochemistry of the National Institute of Agricultural Sciences (INCA). Extracts were prepared from Spirulina powder from the Génix Company of LABIOFAM S.A., using ethanol (absolute ethanol M= 46.07) as solvent.

Two ethanol concentrations (70 and 90 %), two mass-solvent ratios (1:20 and 1:10) and two maceration times (10 and 21 days) were studied for the elaboration of the extracts. The extracts obtained were as follows:

1. Mass-solvent ratio 1:20 using EtOH 90 % for 21 days.
2. Mass-solvent ratio 1:20 using EtOH 90 % for 10 days.
3. Mass-solvent ratio 1:20 using EtOH 70 % for 21 days.
4. Mass-solvent ratio 1:20 using EtOH 70 % for 10 days.
5. Mass-solvent ratio 1:10 using EtOH 90 % for 21 days.
6. Mass-solvent ratio 1:10 using EtOH 90 % for 10 days.
7. Mass-solvent ratio 1:10 using EtOH 70 % for 21 days.
8. Mass-solvent ratio 1:10 using EtOH 70 % for 10 days.

Chemical characterization of the extracts

The chemical characterization of the extracts was carried out by biochemical analysis methods with spectrophotometric determinations of proteins, flavonoids and phenols from the representation of the standard curve corresponding to each technique with the measurement of four repetitions of absorbances of each standard. Three calibration curves were obtained, where absorbance values of each standard were plotted as a function of concentration (19).

The quantification of proteins of each extract was performed using the Micro-Lowry method (20), the phenol content according to the Folin-Ciocalteu method (21) and flavonoids were determined using a spectrophotometric method (22).

Evaluation of the biological activity of the extracts

To determine whether the composition of proteins, phenols and flavonoids influenced the biological activity of the extracts, biological evaluation was performed on the extracts with the lowest (A) and highest (B) content of these compounds. For this purpose, an experiment was carried out in which rice seeds cv. INCA LP-7, for 24 h, with different concentrations (5; 1; 0.5; 0.05; 0.005 mg L⁻¹) of A and B extracts.

For germination, seeds were placed in Petri dishes (20 seeds per dish and four dishes per treatment) containing distilled water. The treatments were as follows:

1. Imbibition with distilled water (Control).

2. Imbibition with 5 mg L⁻¹ of extract A.
3. Imbibition with 1 mg L⁻¹ of the extract A.
4. Imbibition with 0.5 mg L⁻¹ of extract A.
5. Imbibition with 0.05 mg L⁻¹ of extract A.
6. Imbibition with 0.005 mg L⁻¹ of the extract A.
7. Imbibition with 5 mg L⁻¹ of extract B.
8. Imbibition with 1 mg L⁻¹ of extract B.
9. Imbibition with 0.5 mg L⁻¹ of extract B.
10. Imbibition with 0.05 mg L⁻¹ extract B.
11. Imbibition with 0.005 mg L⁻¹ of extract B.

The plates were placed in the germination chamber at 28 °C for seven days, evaluating the number of germinated seeds per plate at 24, 48, 72 and 144 hours, determining the final germination percentage and germination speed, and after ten days, the dry mass of the radicles (25 radicles per treatment, five samples of five radicles each).

For data processing, means, standard deviations and confidence intervals at $\alpha=0.05$ were calculated using Excel software.

RESULTS AND DISCUSSION

Chemical characterization of the extracts

Table 1 shows the content of proteins, phenols and flavonoids present in the extracts analyzed.

As shown in Table 1, the highest protein concentrations were obtained when 70 % EtOH was used (extracts 3, 4, 7, 8), regardless of the mass-solvent ratio used, so that a lower ethanol concentration favors the presence of proteins in the extracts. Note that with a mass-solvent ratio of 1:20 and a 10-day maceration (extract 4), which is the most economical extract, a good protein content was obtained, suggesting that this extract could be used for crop biofortification.

With the phenol content, something similar occurred, while the highest concentrations of flavonoids were obtained in the extracts obtained with 90 % EtOH (extracts 1, 2, 5, 6), independently of the mass-solvent ratio and maceration time.

It follows that extraction with EtOH 70 % favored the content of soluble proteins and total phenols in the extracts,

which is logical since the higher amount of water favors polarity. According some authors (23), the extraction of antioxidant compounds from Spirulina biomass was more effective with a 70 % ethanol solution. Likewise, antioxidant activity was favored in Spirulina extracts obtained with ethanol as solvent over other solvents such as H₂O, methanol and petroleum ether (24). On the other hand, the extraction of total phenolic compounds was favored in extracts prepared with 50 % ethanol over those prepared with 50 % dimethyl sulfoxide (25).

Maceration with 90 % EtOH solution increased the content of total flavonoids, which could be useful to favor crops subjected to environmental stress, since the role of these compounds in increasing antioxidant activity is known (26). Other authors have obtained high concentrations of flavonoids, alkaloids and saponins from Spirulina biomass with absolute ethanol extraction (27).

It is interesting to note that the concentrations of the compounds evaluated did not differ significantly when comparing the two maceration times used. The time factor becomes more significant when using methods such as ultrasonication or microwave-assisted extraction, not so much maceration (28).

The presence of proteins, phenols, flavonoids, etc. in Spirulina extracts favors seed germination, the formation of healthy and balanced plants, higher crop yields, longer post-harvest life, greater leaf vigor, less incidence of diseases and greater resistance to stress due to climatic factors or drought (29-31). It should be taken into account that in the extracts, as mentioned above, there are many other compounds that were not determined in this work and that can influence their biological activity.

Nevertheless, with this limited chemical characterization, it was decided to select extracts 1 and 3 as the extracts with the lowest and highest protein and phenol content, respectively, to evaluate their biological activity as a stimulator of rice seed germination.

Biological activity of the extracts

The speed and final percentage of seed germination for each of the treatments are shown in Table 2. As can be seen, there was an influence on the final germination percentage with two concentrations (5 and 0.5 mg L⁻¹) of

Table 1. Concentration of total soluble proteins, phenols and flavonoids of the different alcoholic extracts of Spirulina

Extracts ^o	Proteins (µg µL ⁻¹)	Phenols (µg µL ⁻¹)	Flavonoids (µg µL ⁻¹)
1	5.5663 ± 1.1764	0.6513 ± 0.0420	3.1232 ± 0.1822*
2	6.7619 ± 0.8062	0.6255 ± 0.0339	2.9988 ± 0.1405*
3	26.4102 ± 1.7561*	1.3309 ± 0.0184*	1.8640 ± 0.1030
4	13.0988 ± 2.2376*	0.7109 ± 0.0181	0.6049 ± 0.0849
5	10.2957 ± 1.9360	0.8627 ± 0.0389	4.4198 ± 0.1523*
6	7.7716 ± 0.5229	0.7421 ± 0.0426	3.3150 ± 0.0842*
7	17.8415 ± 0.3408*	1.1201 ± 0.0181*	1.8506 ± 0.0719
8	21.0298 ± 2.4684*	1.0707 ± 0.0244*	1.4563 ± 0.0473

1. Mass ratio: solvent 1:20 using EtOH 90 % for 21 days. 2. Mass ratio: solvent 1:20 using EtOH 90 % for 10 days. 3. Mass ratio: solvent 1:20 using EtOH 70 % for 21 days. 4. Mass ratio: solvent 1:20 using EtOH 70 % for 10 days. 5. Mass ratio: solvent 1:10 using EtOH 90% for 21 days. 6. Mass ratio: solvent 1:10 using EtOH 90 % for 10 days. 7. Mass ratio: solvent 1:10 using EtOH 70 % for 21 days. 8. Mass ratio: solvent 1:10 using EtOH 70 % for 10 days. *Means differing significantly from the control treatment according to confidence interval at $\alpha=0.05$

extract 3, although there were no significant differences in the germination speed, so that the higher concentration of soluble proteins and total phenols presented by this extract significantly increased the final germination percentage of rice seeds cv. INCA LP-7.

The results of radicle dry mass showed that there were no significant differences between treatments; however, the immersion of the seeds in 5 mg L⁻¹ and 0.05 mg L⁻¹ of extract 3 increased the dry mass of the radicles by 8.2 and 10.3 %, respectively. Note that the treatment with 5 mg L⁻¹ of this extract was one of those that significantly increased the final germination percentage; confirming the influence that a higher concentration of soluble proteins and total phenols exerted on the germination and initial growth of rice seedlings.

Microalgae and their extracts are natural stimulants that accelerate seed germination and increase seedling vigor when used at relatively low doses (32). Several studies describe their beneficial effects on germination percentage, rate and mean germination time, as well as plumule and radicle length. These results are attributed to the activation of key enzymatic pathways for germination physiology. For example, α -amylase is an enzyme synthesized in the aleurone layer and its gene expression is regulated by

gibberellins. This enzyme is responsible for the mobilization of reserve substances, such as starch, from the endosperm to support embryo growth and differentiation (33). To demonstrate this hypothesis, the significant effects that an extract of *Spirulina platensis* exerted on peanut (*Arachis hypogaea* L.) crop on indicators such as seed germination percentage, radicle length, and protein and carbohydrate contents were recently reported (32). In another study, enhanced vigor, quality and germination of black bean (*Vigna mungo* L.) seeds were reported to be correlated with increased gibberellic acid and α -amylase enzyme activity, when soaked in 1.5 % *Spirulina platensis* extract for 3 hours (34). On the other hand, it was found that the use of a methanolic extract (0.25 %) favored almost all growth indicators, nutrient content, yield components and phytohormone level of *Lupinus luteus* (35). In another investigation, it was found that when tomato seeds were treated with 0.2 g 100 mL of *Spirulina* extract, a germination rate of 86.7 % was obtained (36), and concentrations of 25 and 50 % of this extract had a positive effect on seed germination and the development of lettuce seedlings (37).

The results obtained in this research, although still preliminary, showed the importance of ethanol concentration

Table 2. Effect of different concentrations of two alcoholic extracts of *Spirulina* on seed germination of rice cv. INCA LP-7 (Means $\pm$ confidence intervals)

Concentrations mg L ⁻¹	Alcoholic extracts of <i>Spirulina</i>	Final germination	Germination rate (seeds germinated day ⁻¹)
0	-	72.50 $\pm$ 2.83	11.95 $\pm$ 0.39
5	Extract 1	72.50 $\pm$ 6.33	10.95 $\pm$ 1.59
1		75.00 $\pm$ 8.95	10.62 $\pm$ 0.92
0.5		82.50 $\pm$ 10.20	12.91 $\pm$ 1.29
0.05		78.33 $\pm$ 5.66	12.05 $\pm$ 0.83
0.005		75.00 $\pm$ 4.00	11.08 $\pm$ 0.78
5	Extract 3	82.50 $\pm$ 6.33*	12.40 $\pm$ 1.42
1		71.25 $\pm$ 4.69	9.29 $\pm$ 0.80
0.5		83.33 $\pm$ 71.48*	11.44 $\pm$ 0.92
0.05		68.75 $\pm$ 14.07	10.04 $\pm$ 2.52
0.005		67.50 $\pm$ 8.49	9.54 $\pm$ 0.82

Extract 1: Mass ratio: solvent 1:20 using EtOH 90 % for 21 days. Extract 3: Mass-solvent ratio 1:20 using EtOH 70 % for 21 days *Means differing significantly from control treatment according to confidence interval at $\alpha=0.05$

Table 3. Effect of different concentrations of two alcoholic extracts of *Spirulina* on the dry mass of radicles of rice seedlings cv. INCA LP-7 (Means $\pm$ confidence intervals)

Concentrations mg L ⁻¹	<i>Spirulina</i> alcoholic extracts	Dry mass of radicles (mg plant ⁻¹)
0	-	4.37 $\pm$ 0.54
5	Extract 1	4.00 $\pm$ 0.23
1		4.50 $\pm$ 0.69
0.5		4.02 $\pm$ 0.57
0.05		4.15 $\pm$ 0.34
0.005		4.44 $\pm$ 0.69
5	Extract 3	4.73 $\pm$ 0.35
1		3.92 $\pm$ 0.60
0.5		3.81 $\pm$ 0.30
0.05		4.82 $\pm$ 0.49
0.005		4.04 $\pm$ 0.27

Extract 1: Mass ratio: solvent 1:20 using EtOH 90 % for 21 days. Extract 3: Mass-solvent ratio 1:20 using EtOH 70 % for 21 days

and maceration time in the composition and activity as agricultural biostimulant of Spirulina extracts; therefore, it is necessary to continue deepening in this sense, with a view to obtaining biologically active extracts that can be satisfactorily used in agriculture.

CONCLUSIONS

- The chemical characterization of the alcoholic extracts of Spirulina allowed the selection of two extracts for the evaluation of their biological activity.
- A higher concentration of soluble proteins and phenols in the extracts favored germination and dry mass of rice seedling radicles.

RECOMMENDATION

- To continue to characterize the extracts and evaluate the biological activity of the extracts in other crops.

BIBLIOGRAPHY

1. Dmytryk A, Samoraj M, Moustakas K, Witek-Krowiak A, Chojnacka K. Bioactive fatty acids and compounds from *Spirulina (Arthrospira) platensis*: Potential as biostimulants for plant growth. *Sustain Chem Pharm* [Internet]. 2022 Dec 1 [cited 2024 Nov 25];30:100899. Available from: <https://www.sciencedirect.com/science/article/pii/S2352554122003035>
2. Kumar A, Ramamoorthy D, Verma DK, Kumar A, Kumar N, Kanak KR, et al. Antioxidant and phytonutrient activities of *Spirulina platensis*. *Energy Nexus* [Internet]. 2022 Jun 16 [cited 2024 Nov 25];6:100070. Available from: <https://www.sciencedirect.com/science/article/pii/S277242712200033X>
3. Das P, Khan S, Chaudhary AK, AbdulQuadir M, Thafer MI, Al-Jabri H. Potential Applications of Algae-Based Bio-fertilizer. In: *Biofertilizers for Sustainable Agriculture and Environment*. Springer; 2019. p. 41-65.
4. Plaza BM, Gómez Serrano C, Ación Fernández FG, Jiménez Becker S. Effect of microalgae hydrolysate foliar application (*Arthrospira platensis* and *Scenedesmus* sp.) on *Petunia x hybrida* growth. *Journal of Applied Phycology*. 2018 Feb 19 [cited 2024 Nov 25]; Available from: <https://repositorio.ual.es/handle/10835/8381>
5. Belaustegui I. *Espirulina: propiedades y beneficios* - Jesús Sierra [Internet]. 2023 [cited 2024 Nov 25]. Available from: <https://jesussierra.com/espirulina/>
6. Kaur M, Bhatia S, Bayram I, Decker EA, Phutela UG. Oxidative stability of emulsions made with self-extracted oil from euryhaline microalgae *Spirulina* and *Scenedesmus*. *Algal Res* [Internet]. 2023 Sep 1 [cited 2024 Nov 25];75:103280. Available from: <https://www.sciencedirect.com/science/article/pii/S2211926423003132>
7. Manrich A, de Oliveira JE, Martins MA, Capparelli ML. Physicochemical and Thermal Characterization of the *Spirulina platensis*. *Journal of Agricultural Science and Technology* [Internet]. 2020;298-307. Available from: <http://www.davidpublisher.com/Public/uploads/Contribute/601b72a5d8536.pdf>
8. Munawaroh H, Fathur R, Gumilar G, Aisyah S, Yuliani G, Mudzakir A, et al. Characterization and physicochemical properties of chlorophyll extract from *Spirulina* sp. *J Phys Conf Ser* [Internet]. 2019 Nov 1;1280:022013. Available from: https://www.researchgate.net/publication/337443355_Characterization_and_physicochemical_properties_of_chlorophyll_extract_from_Spirulina_sp
9. Papadaki S, Kyriakopoulou K, Tzovenis I, Krokida M. Environmental impact of phycocyanin recovery from *Spirulina platensis* cyanobacterium. *Innov Food Sci Emerg Technol* [Internet]. 2017 Dec 1 [cited 2024 Nov 25];44:217-23. Available from: <https://www.sciencedirect.com/science/article/pii/S1466856416305264>
10. Campanella L, Crescentini G, Avino P. Chemical composition and nutritional evaluation of some natural and commercial food products based on *Spirulina*. *Analisis* [Internet]. 1999 Jul 1;27:533-40. Available from: https://www.researchgate.net/publication/245275925_Chemical_composition_and_nutritional_evaluation_of_some_natural_and_commercial_food_products_based_on_Spirulina
11. Ronga D, Biazzi E, Parati K, Carminati D, Carminati E, Tava A. Microalgal Biostimulants and Biofertilisers in Crop Productions. *Agronomy* [Internet]. 2019 Apr [cited 2024 May 14];9(4):192. Available from: <https://www.mdpi.com/2073-4395/9/4/192>
12. Wuang SC, Khin MC, Chua PQD, Luo YD. Use of *Spirulina* biomass produced from treatment of aquaculture wastewater as agricultural fertilizers. *Algal Res* [Internet]. 2016 Apr 1 [cited 2024 Nov 25];15:59-64. Available from: <https://www.sciencedirect.com/science/article/pii/S2211926416300480>
13. Sathasivam R, Radhakrishnan R, Hashem A, Abd_Allah EF. Microalgae metabolites: A rich source for food and medicine. *Saudi J Biol Sci* [Internet]. 2019 May 1 [cited 2024 Nov 25];26(4):709-22. Available from: <https://www.sciencedirect.com/science/article/pii/S1319562X17302784>
14. L A, P K, G SB. Evaluation of *Spirulina platensis* as microbial inoculants to enhanced protein levels in *Amaranthus gangeticus*. *Afr J Agric Res* [Internet]. 2016 Apr 14 [cited 2024 Nov 25];11(15):1353-60. Available from: <http://academicjournals.org/journal/AJAR/article-abstract/1221D1E58055>
15. Anitha L, Kalpana P, Bramari SG. BIOFORTIFICATION OF AMARANTHUS GANGETICUS USING SPIRULINA PLATENSIS AS MICROBIAL INOCULANT TO ENHANCE IRON LEVELS. *Int J Res Appl Nat Soc Sci* [Internet]. 2014 [cited 2024 Nov 25];2:103-10. Available from: https://www.academia.edu/7025447/BIOFORTIFICATION_OF_A_MARANTHUS_GANGETICUS_USING_SPIRULINA_PLATENSIS_AS_MICROBIAL_INOCULANT_TO_ENHANCE_IRON_LEVELS
16. Anitha L, Bramari GS, Kalpana P. Effect of supplementation of *Spirulina platensis* to enhance the zinc status in plants of

- Amaranthus gangeticus, Phaseolus aureus and tomato. Advances in Bioscience and Biotechnology. 2016;7(6):289-99. Available from: https://www.researchgate.net/publication/304608048_Effect_of_Supplementation_of_Spirulina_platensis_to_Enhance_the_Zinc_Status_in_Plants_of_Amaranthus_gangeticus_Phaseolus_aureus_and_Tomato
17. Dias G, Rocha R, Araújo J, Lima J, Guedes W. Growth, yield, and postharvest quality in eggplant produced under different foliar fertilizer (*Spirulina platensis*) treatments. Semina Ciênc Agrár [Internet]. 2016 Dec 14;37(6):3893. Available from: https://www.researchgate.net/publication/311851452_Growth_yield_and_postharvest_quality_in_eggplant_produced_under_different_foliar_fertilizer_Spirulina_platensis_treatments
18. Akgül F. Effect of *Spirulina platensis* (Gomont) Geitler Extract on Seed Germination of Wheat and Barley. Alinteri J Agric Sci [Internet]. 2019 Dec 31 [cited 2024 Nov 26];34(2):148-53. Available from: <https://dergipark.org.tr/en/pub/alinterizbd/issue/51098/639000>
19. Silva A de S e, de Magalhães WT, Moreira LM, Rocha MVP, Bastos AKP. Microwave-assisted extraction of polysaccharides from *Arthrospira (Spirulina) platensis* using the concept of green chemistry. Algal Res [Internet]. 2018 Nov 1 [cited 2024 Nov 26];35:178-84. Available from: <https://www.sciencedirect.com/science/article/pii/S2211926418301668>
20. Waterborg JH. The Lowry Method for Protein Quantitation. In: Walker JM, editor. The Protein Protocols Handbook [Internet]. Totowa, NJ: Humana Press; 2009 [cited 2024 Nov 26]. p. 7-10. Available from: https://doi.org/10.1007/978-1-59745-198-7_2
21. Singleton VL, Rossi JA. Colorimetry of Total Phenolics with Phosphomolybdic-Phosphotungstic Acid Reagents. Am J Enol Vític [Internet]. 1965 Jan 1 [cited 2024 Nov 26];16(3):144-58. Available from: <https://www.ajevonline.org/content/16/3/144>
22. Quettier-Deleu C, Gressier B, Vasseur J, Dine T, Brunet C, Luyckx M, et al. Phenolic compounds and antioxidant activities of buckwheat (*Fagopyrum esculentum* Moench) hulls and flour. J Ethnopharmacol. 2000 Sep;72(1-2):35-42.
23. Cepoi L, Ludmila R, Vera M, Angela C, Chiriac T, Sadovnic D. Antioxidative activity of ethanol extracts from *Spirulina platensis* and *Nostoc linckia* measured by various methods. Analele Univ Din Oradea Fasc Biol [Internet]. 2009 Nov 1;TOM XVI. Available from: https://www.researchgate.net/publication/40422973_Antioxidative_activity_of_ethanol extracts_from_Spirulina_platensis_and_Nostoc_linckia_measured_by_various_methods
24. Varsale A, Singh V, Mali S, Parihar P, Mane R. Phytochemical analysis, antioxidant and antifungal activity of different solvent extracts of *Spirulina platensis* collected from Rankala Lake, Kolhapur, Maharashtra. Phytochem Anal [Internet]. 2019;10(1):36-42. Available from: <https://storage.unitedwebnetwork.com/files/521/ba6ab112b0e28c164f459c61b622daf1.pdf>
25. Martí-Quijal FJ, Ramon-Mascarell F, Pallarés N, Ferrer E, Berrada H, Phimolsiripol Y, et al. Extraction of Antioxidant Compounds and Pigments from *Spirulina (Arthrospira platensis)* Assisted by Pulsed Electric Fields and the Binary Mixture of Organic Solvents and Water. Appl Sci [Internet]. 2021 Jan [cited 2024 Nov 26];11(16):7629. Available from: <https://www.mdpi.com/2076-3417/11/16/7629>
26. Agustina S, Aidha NN, Oktarina E, Kurniati NF. Evaluation of Antioxidant and Wound Healing Activities of *Spirulina* sp. Extract. Egypt J Chem [Internet]. 2021 Aug 1 [cited 2024 Nov 26];64(8):4601-10. Available from: https://ejchem.journals.ekb.eg/article_162338.html
27. Yuniati R, Zainuri M, Kusumaningrum H. Qualitative Tests of Secondary Metabolite Compounds in Ethanol Extract of *Spirulina platensis* from Karimun Jawa Sea, Indonesia. Biosaintifika J Biol Biol Educ [Internet]. 2020 Dec 25 [cited 2024 Nov 26];12(3):343-9. Available from: <https://journal.unnes.ac.id/nju/biosaintifika/article/view/23153>
28. Kong S, Wiratni W, Budiman A. Protein Extraction from *Spirulina platensis* by Using Ultrasound Assisted Extraction: Effect of Solvent Types and Extraction Time. Key Eng Mater [Internet]. 2021 Jan 11;872:33-7. Available from: https://www.researchgate.net/publication/348402116_Protein_Extraction_from_Spirulina_platensis_by_Using_Ultrasound_Assisted_Extraction_Effect_of_Solvent_Types_and_Extraction_Time
29. Rahim A, Çakir C, Ozturk M, Şahin B, Soulaïmani A, Sibaeueih M, et al. Chemical characterization and nutritional value of *Spirulina platensis* cultivated in natural conditions of Chichaoua region (Morocco). South Afr J Bot [Internet]. 2021 Sep 1 [cited 2024 Sep 23];141:235-42. Available from: <https://www.sciencedirect.com/science/article/pii/S0254629921001770>
30. Silva SC, Almeida T, Colucci G, Santamaria-Echart A, Manrique YA, Dias MM, et al. *Spirulina (Arthrospira platensis)* protein-rich extract as a natural emulsifier for oil-in-water emulsions: Optimization through a sequential experimental design strategy. Colloids Surf Physicochem Eng Asp [Internet]. 2022 Sep 5 [cited 2024 Nov 26];648:129264. Available from: <https://www.sciencedirect.com/science/article/pii/S0927775722010196>
31. López-Padrón I, Martínez-González L, Pérez-Domínguez G, Reyes-Guerrero Y, Núñez-Vázquez M, Cabrera-Rodríguez JA. Las algas y sus usos en la agricultura. Una visión actualizada. Cultiv Trop [Internet]. 2020 Aug 5 [cited 2024 May 14];41(2):e10-e10. Available from: <https://ediciones.inca.edu.cu/index.php/ediciones/article/view/1554>
32. Sivalingam KM. Isolation, identification and evaluation of *Spirulina platensis* for its effect on seed germination of groundnut (*Arachis hypogaea* L.), Wolaita Sodo, Southern Ethiopia. J. Algal Biomass Utln. 2020;11(2):34-42. Available from: <https://storage.unitedwebnetwork.com/files/521/4ca5cd59fe556a59877dcb110401ee19.pdf>
33. Oliveira GE, Pinho RGV, Andrade T de, Pinho ÉV de RV, Santos CD dos, Veiga AD. Physiological quality and amylase enzyme expression in maize seeds. Ciênc E Agrotecnologia [Internet]. 2013 Feb [cited 2024 Sep 23];37:40-8. Available from: <https://www.scielo.br/j/cagro/a/JJpGvzwnJkQ9Zx5JLQRN9xR/?lang=en>

34. Field Crop Research Institute, VAAS, Vietnam, Thinh NQ. Influences of seed priming with *Spirulina platensis* extract on seed quality properties in black gram (*Vigna mungo* L.). *Vietnam J Sci Technol Eng* [Internet]. 2021 Mar 31 [cited 2024 Nov 26];63(1):36-41. Available from: <https://vietnamscience.vjst.vn/index.php/VJSTE/article/view/388/268>
35. Shedeed ZA, Gheda S, Elsanadily S, Alharbi K, Osman MEH. *Spirulina platensis* Biofertilization for Enhancing Growth, Photosynthetic Capacity and Yield of *Lupinus luteus*. *Agriculture* [Internet]. 2022 Jun [cited 2024 Nov 26];12(6):781. Available from: <https://www.mdpi.com/2077-0472/12/6/781>
36. Iuliana Neag E, Stupar Z, Roman C. EFFECT OF SPIRULINA SPP. EXTRACTS ON TOMATO AND ONION SEED GERMINATION |. *AGRICULTURA* [Internet]. 2022 [cited 2024 Nov 26];124(3-4). Available from: <https://journals.usamvcluj.ro/index.php/agricultura/article/view/14437>
37. Akgül F, Akgül R. The effect of *Spirulina platensis* (Gomont) Geitler extracts on seed germination of *Lactuca*. *Alinteri Ziraat Bilim Derg* [Internet]. 2019 Dec 31 [cited 2024 Nov 26];34(2):148-53. Available from: <https://ageconsearch.umn.edu/record/296790?v=pdf>



Available in:

<https://www.redalyc.org/articulo.oa?id=193281883005>

How to cite

Complete issue

More information about this article

Journal's webpage in redalyc.org

Scientific Information System Redalyc
Diamond Open Access scientific journal network
Non-commercial open infrastructure owned by academia

Anaysa Gutierrez Almeida, Miriam de la C. Núñez Vázquez,
Yanelis Reyes Guerrero, Geydi Pérez Domínguez,
Lisbel Martínez González

**Caracterización química y evaluación de la actividad
biológica de extractos de Spirulina**
**Chemical characterization and evaluation of the
biological activity of Spirulina extracts**

Cultivos Tropicales
vol. 46, no. 1, e05, 2025
Ediciones INCA,
ISSN-E: 1819-4087