



Revista Colombiana de Obstetricia y Ginecología
ISSN: 0034-7434
Federación Colombiana de Obstetricia y
Ginecología; Revista Colombiana de Obstetricia y
Ginecología

Incidence of postpartum hemorrhage based on the use of uterotonics. Maternal outcomes in an intermediate complexity hospital in Bogotá, Colombia, 2016¹

López-García, Luis Felipe; Ruiz-Fernández, Diana Paola; Zambrano-Cerón, Carlos Germán; Rubio-Romero, Jorge Andrés

Incidence of postpartum hemorrhage based on the use of uterotonics. Maternal outcomes in an intermediate complexity hospital in Bogotá, Colombia, 2016¹

Revista Colombiana de Obstetricia y Ginecología, vol. 68, no. 3, 2017

Federación Colombiana de Obstetricia y Ginecología; Revista Colombiana de Obstetricia y Ginecología

Available in: <http://www.redalyc.org/articulo.oa?id=195255662007>

DOI: 10.18597/rcog.2916

Reporte de caso

Incidence of postpartum hemorrhage based on the use of uterotonics. Maternal outcomes in an intermediate complexity hospital in Bogotá, Colombia, 2016¹

Luis Felipe López-García ² luflopezga@unal.edu.co

Universidad Nacional de Colombia, Colombia

Diana Paola Ruiz-Fernández ³

Universidad Nacional de Colombia, Colombia

Carlos Germán Zambrano-Cerón ³

Universidad Nacional de Colombia, Colombia

Jorge Andrés Rubio-Romero ⁴

Universidad Nacional de Colombia, Colombia

ABSTRACT

Objective: To describe the incidence of obstetric postpartum haemorrhage (PPH), defined by the use of uterotonic drugs, as well as the interventions performed, and maternal outcomes in the first 24 hours.

Materials and methods: A series of cases of pregnant women who presented PPH after vaginal delivery or cesarean section between February 1 and October 31, 2016, in a public intermediate complexity institution in Bogotá. Consecutive sampling was used. Sociodemographic, clinical and risk factors for PPH were measured. Variables measured were the cause of PPH, the degree of shock, estimated postpartum bleeding, frequency of activation of the "obstetric red code," interventions performed, admission to an intensive care unit, blood transfusions, and maternal mortality. A descriptive analysis was performed.

Results: Out of 1633 births (1080 vaginal deliveries and 553 cesarean deliveries), 35 cases (2.1%) (26 and 9, respectively) were identified and the "obstetric red code" was activated in 11 cases (0.67%). There was no maternal mortality. The main cause of PPH was uterine hypotonia in 29/35 (82%), and 82.8% of the cases resolved with medical treatment. Uterine tamponade was required in 4/35 (11.4%) women to control bleeding, and surgical management was required in 2/35 (5.6%) women: haemostatic sutures in 1/35 (2.8%) and hysterectomy in 1/35 (2.8%).

Conclusions: The use of more reproducible criteria for the identification of severe PPH and timely initiation of treatment could be safer and more effective in terms of maternal outcomes.

Key words: Postpartum haemorrhage++ delivery++ uterine balloon tamponade++ treatment outcome.

INTRODUCTION

Primary postpartum haemorrhage (PPH) is traditionally defined as an estimated blood loss greater than 500 cc in the first 24 hours in cases of vaginal delivery, and of more than 1000 cc following cesarean section ¹. It occurs in 2-4% of all vaginal deliveries and 6-7% of cesarean deliveries. It is severe in 1% of cases, with a blood loss greater than 1,000 cc ². It is

Revista Colombiana de Obstetricia y Ginecología, vol. 68, no. 3, 2017

Federación Colombiana de Obstetricia y Ginecología; Revista Colombiana de Obstetricia y Ginecología

Received: 10 February 2017
Accepted: 25 September 2017

DOI: 10.18597/rcog.2916

CC BY-NC-ND

the most frequent cause of maternal mortality (19%) and morbidity in the world, accounting for 8% and 19.7% of maternal deaths in developed and developing countries, respectively ³. In Colombia, PPH accounts for 23% of all maternal deaths ⁴. Prevention and treatment of primary postpartum haemorrhage are critical in diminishing maternal mortality ⁵. Causes of obstetric bleeding after childbirth have been divided into four main groups as follows: tone, trauma, tissue, and thrombin, the most frequent being uterine atony at 70% ⁶⁻⁹.

Severe obstetric bleeding after childbirth is a life-threatening emergency, classified as a near-miss (extreme maternal morbidity) and its occurrence is mandatory to be reported to the epidemiological surveillance system ¹⁰. Given that the diagnosis of severe PPH based on the visual estimation of blood loss is not reliable and tends to underestimate the magnitude of the bleeding, several methods have been proposed based on more accurate clinical findings for estimating blood loss ¹¹. All these methods are aimed at triggering a series of sequential, coordinated and timely actions by an interdisciplinary team in order to achieve the core objective of diminishing maternal morbidity and mortality from this cause. In Colombia, these actions are known as the "obstetric red code" ¹¹. In this approach, the initial step to trigger actions is to make the diagnosis of the presence of cardiovascular shock. This management algorithm incorporates the recommendations from the World Health Organisation (WHO) guidelines ⁵ and of the Royal College of Gynaecology and Obstetrics ¹². Effective implementation of these actions requires the activation of the rapid response team and priority allocation of hospital resources until the condition is resolved.

Delay in the identification of severe PPH is the main cause of maternal morbidity and mortality secondary to haemorrhage ^{11,13,14}. The "obstetric red code" approach to diagnosis and treatment, though effective for bleeding control and improvement of patient survival once it is activated, is not very sensitive when it comes to identifying and reporting all cases of primary obstetric haemorrhage, considering that the presence of PPH could be judged as "not sufficiently" severe to warrant reporting. This means that although management by the treating staff is triggered, there is no formal activation of the obstetric red code in the institution. This "not sufficiently severe" PPH may be associated with significant maternal morbidity and risk of dying, and also with under-recording and under-reporting of haemorrhage cases to the epidemiological surveillance systems in Colombia.

The objective of this study is to explore the incidence of primary obstetric haemorrhage from the perspective of a more reproducible definition based on the use of one or more uterotonic medications in additional doses than those used routinely for active childbirth management, regardless of the activation of the obstetric red code; and to describe the interventions implemented for bleeding control, as well as the maternal outcomes with treatment initiated within the first 24 hours

and maintained during 7 days in the postpartum period, and to evaluate the magnitude of under-recording.

MATERIALS AND METHODS

Case series of women with primary obstetric haemorrhage as part of a cohort of pregnant women who presented for vaginal or cesarean delivery between February 1 and October 31, 2016, at a public intermediate complexity hospital located in Bogotá that serves patients covered by the Colombian Social Security System (SSSC) and is a training centre for undergraduate and graduate medical students. Women with antepartum diagnosis of uterine rupture, placenta accreta, placenta praevia, coagulation disorders or disseminated intravascular coagulation were excluded, considering their indication for definite treatments other than the administration of uterotonics in the presence of PPH.

Cases of obstetric haemorrhage were defined as women requiring the use of additional medications (oxytocin, methylergonovine or misoprostol) apart from the dose of 10 units of oxytocin for the active management of childbirth during vaginal or cesarean delivery, as judged by the physician in charge of the obstetric service in accordance with institutional protocols and guidelines.

Sampling was consecutive. Based on the incidence of postpartum haemorrhage reported in the literature of 2-4% in vaginal delivery and 6-7% in cesarean delivery ^{13,15} and the reported frequency of 2,200 births per year at the institution, a sample size of 90 cases of obstetric haemorrhage was estimated, with a distribution of 50 cases after vaginal delivery and 40 cases after cesarean section during the study period, according to the historical proportion of cesarean sections performed at the institution ¹⁶.

Procedure. Patients were identified by means of an active search in the institutional pharmacy of any released doses of the available uterotonics (oxytocin, methylergonovine or misoprostol) additional to the 10 units of oxytocin used for active childbirth management, and in the records of mandatory reporting of obstetric red code team activations. Clinical records were reviewed up to patient discharge and the 7-day follow-up visit. Data were collected using specific forms and were kept in a secure area under lock and key, with access restricted to the study personnel. They were then entered into an Excel database and analysed using EPIINFO 7.0, after quality verification.

The variables measured included maternal age, medical history, pregnancy-associated diseases, parity, a history of postpartum bleeding, weeks of gestational age at the time of delivery, presence of prenatal visits, haemoglobin concentration (g/dL) at the time of the last prenatal visit, uterine height, labour induction, delivery route, episiotomy, perineal tears, and neonatal weight and length. Outcome variables were the presence of obstetric bleeding defined by the use of medications (oxytocin, methylergonovine or misoprostol) additional to the dose of 10 units of oxytocin used for active management of childbirth during

vaginal or cesarean delivery and during the next 24 hours after childbirth; cause of bleeding (hypotony, trauma, presence of tissue or postpartum coagulation abnormality; degree of shock estimated in accordance with the criteria defined in the "red code" protocol¹¹; estimated postpartum bleeding; and activation of the obstetric red code team. Interventions defined for obstetric bleeding control were: medical management (use of the uterotonics described), advanced medical management (uterotonics plus uterine tamponade), and surgical management (laparotomy for haemostatic uterine sutures or hysterectomy). Maternal outcomes included in the assessment were mortality, hysterectomy, admission to the intensive care unit, and need for blood transfusion.

Statistical analysis. Quantitative variables are expressed in the form of central trends, means or medians, and scatter measures according to the results of normalcy tests, and qualitative variables are expressed in absolute numbers and proportions.

Ethical considerations. This research project was submitted to the Ethics Committee of Universidad Nacional de Colombia Medical School for evaluation and was approved as set forth in Minutes 143 of September 10, 2015, and by the Ethics Committee of Hospital de Engativa ESE, on December 18, 2015.

RESULTS

During the study period there were in total 1,633 births: 1,080 vaginal deliveries and 553 cesarean sections, for a proportion of 33.8% cesarean deliveries. A total of 35 cases of obstetric haemorrhage were identified as defined by the additional use of uterotonics, for an overall incidence of 2.1% during the study period. By delivery route, the incidence of postpartum obstetric haemorrhage was 2.4% (26 cases) following vaginal delivery, and 1.6% (9 cases) following cesarean section. According to reporting based on obstetric red code activation, the incidence of postpartum obstetric haemorrhage was 0.83% (9/1080) after vaginal delivery and 0.36% (2/553) after cesarean section. There were no maternal deaths. Table 1 shows the demographic characteristics of the cases and the causes of PPH. In 24 out of 35 patients (68.6%), uteronic medications were administered on top of active childbirth management, and there was no activation of the institutional response protocol or the obstetric red code.

Six patients had a medical history and, of them, three were obese, one presented with repeated urinary tract infections, one was drug-dependent and another one had hypertriglyceridemia. There were no patients with anaemia before childbirth. Pregnancy-related diseases found in 11 patients included preeclampsia, hyperthyroidism, gestational diabetes and gestational hypertension. Table 1 shows the causes of postpartum hemorrhage. There were three cases of retained placenta as the cause of obstetric haemorrhage following vaginal delivery (11.5%), and no case of tissue retention following cesarean delivery. There were three trauma cases, one of them following cesarean section as a result of a cervical tear

after a failed instrumentation attempt. There were no cases attributable to coagulation disorders.

Table 1
Demographic characteristics of obstetric postpartum haemorrhage cases at Hospital de Engativá, Bogotá, Colombia, 2016

Variable	n = 35 (100 %)
Age*	24 (20-30)
Women with a medical history	6 (17)
Parity	
Nulliparous	19 (54.3)
Multiparous	16 (45.7)
Gestational age (weeks)*	38 (35-39)
Prenatal visits	
No	4 (11.5)
Yes	31 (88.5)
Haemoglobin concentration on the last prenatal visit (g/dL) *	13.3 (11.3-16.8)
Pregnancy-associated disease	11 (31.4)
Uterine height greater than 35 cm	2 (11.4)
History of postpartum bleeding	0 (0)
Labour induction	16 (45.7)
Route of childbirth	
Vaginal	26 (65.4)
Cesarean section	9 (34.6)
Episiotomy (n = 26)	5 (14.2 %)
Estimated postpartum bleeding (mL) (n = 32)*	800 (500-1,000)
Cause of postpartum haemorrhage	
Hypotony	29 (82)
Trauma	3 (8.5)
Tissue	3 (8.5)
Coagulation	0 (0)
Weight of the neonate (g)*	3,024 (2,630-3,490)
Length of the neonate (cm)*	51.2 (49-53)

In 29 of 35 patients with PPH, bleeding was controlled with medical management (Figure 1) based on the use of oxytocin at a dose of 20-60 IU (median 40 IU). In 19 of these 29 women (65.5%), methylergonovine had to be added at a median dose of 0.2 mg, and 12 women (41.3%) also required rectal administration of a median dose of 1,000 mg of misoprostol. In 11 cases in which obstetric red code activation was required, medications were administered concurrently.




Figure 1
Flow-chart of patients and maternal outcomes according to the route of delivery and activation of the obstetric red code in women with postpartum haemorrhage at Hospital de Engativá, Bogotá, Colombia, 2016
PPH: Obstetric haemorrhage. ICU: Intensive Care Unit. PRBCs: Packed Red Blood Cells

Of the remaining 6 patients with obstetric haemorrhage who required additional interventions, 4 patients after vaginal delivery responded to hydrostatic tamponade, and 2 who presented bleeding following caesarean section required surgical management: one haemostatic suture (B-Lynch) and one subtotal hysterectomy.

Haemoglobin levels were under 11 in 14 (63%) out of 22 patients in whom they were measured 6 hours after the bleeding episode. Of the women in whom the rapid response protocol was not activated, 63% developed postpartum anaemia; 50% required blood product transfusions, and 16% had to be admitted to the ICU. Advanced medical management of obstetric haemorrhage with hydrostatic tamponade^{1,5,11} was used with 100% effectiveness in 4 women following vaginal delivery.

Responses to the treatments instituted according to the degree of shock at the time of case identification are shown in Table 2. The stratified analysis shows that a smaller proportion of women with the highest degree of shock responded to medical management and had a higher frequency of transfusions.

Table 2

Clinical response according to the degree of shock in patients with postpartum obstetric haemorrhage at Hospital de Engativá, Bogotá, Colombia, 2016

Degree of shock n = 35 (100%)	Response to medical management n (%)	Need for advanced management* n (%)	Need for surgical management n (%)	Need for transfusion n (%)
COMPENSATED Heart rate 60-90 bpm Systolic blood pressure >90 mm/Hg n = 17 (48.5)	14 (82.3)	2 (11.7)	1 † (5.8)	2 (11.7)
MILD Heart rate 91-100 bpm, Systolic blood pressure 80-90 mm/Hg n = 5 (14.2)	5 (100)	0 (0)	0 (0)	1 (20)
MODERATE Heart rate 101-120 bpm, Systolic blood pressure 70-79 mm/Hg n = 9 (25.7)	7 (77.7)	2 (22.3)	0 (0)	4 (44.4)
SEVERE Heart rate >120 bpm, Systolic blood pressure <70 mm/Hg n = 4 (11.4)	3 (75)	0 (0)	1 ‡ (25)	2 (50)

DISCUSSION

The incidence of obstetric haemorrhage in this study was 2.1% lower than the one reported in the world literature (4-6%), as well as the one reported previously for the same institution ^{1,3,8,13}. The incidence determined on the basis of the use of additional medications other than the ones used for active management of childbirth, after excluding cases in which preventive use of postpartum uterotonics was documented, was 3.5 higher than the reported incidence, according to the case definition used by the Colombian Epidemiological Surveillance System (SIVIGILA). In women who required the additional use of uterotonics, there was maternal morbidity and transfusion requirement in 50% of the cases, and invasive management in 16%. This observation points to the benefit of using more repeatable methods for diagnosing PPH, such as the use of uterotonic medications, and of assessing different cut-off points based on progressive needs.

Cesarean section has been reported as a risk factor, and a higher incidence of PPH is described in association with cesarean delivery ¹². However, in this study, the incidence of post-cesarean PPH was lower compared to the incidence following vaginal delivery (1.6% vs. 2.4%).

These differences may be attributed to the exclusion of cases with PPH causes identified antepartum, including placenta praevia and placenta accreta, and to the inclusion of a population with low-risk characteristics for PPH.

The low frequency of red code activation reflects the challenge for the staff responsible for childbirth when faced with the need of making a clinical diagnosis of severe obstetric haemorrhage^{13,14}. A review of the literature including 36 studies assessed visual estimation of blood loss¹⁴ and observed that clinical estimation was inaccurate when compared to the use of collection bags. However, the use of collection bags did not prevent delays in decision-making or obstetric bleeding progression. The advantages of false positive results with the use of this method for estimating blood loss would be early management in patients with no severe but abnormal bleeding¹³, and avoiding the need for institutional protocol activation, notification and overuse of resources in response to this obstetric emergency, based on the clinical judgement of the treating staff. In contrast, a false negative result leads to late initiation of management because of the underestimation of the blood loss, resulting in serious consequences and increased maternal morbidity.

Our observations regarding the cause of PPH are consistent with a study conducted in low income countries, in which 82% of the cases of obstetric haemorrhage reported were attributable to uterine atony¹⁵.

A treatment protocol for obstetric haemorrhage has been consolidated as a sequence of interventions of increasing complexity and degree of invasion^{5,11,12}. Reported success rates of initial medical management with uterotonics and massage is 60-70%⁶. In this study, medical management following obstetric red code activation was successful in 6/11 cases (54.5%), and in 29/35 (82%) in those cases in which the obstetric red code was not activated. This difference could be due to differences in bleeding volume and severity that lead treating staff to the selective activation of the institutional protocol: in women with larger estimated bleeding, predefined coordinated actions are implemented together with initiation, although late, of advanced medical or surgical management, while in women with abnormal bleeding requiring the use of additional uterotonics besides the ones used in active childbirth management, and no red code activation, the response to medical treatment is close to 80%. Along the same lines, the stratified analysis of the women's response to the treatment based on the degree of shock showed that, the higher the degree of shock, the lower the proportion of cases that responded to medical treatment, and the higher the frequency of transfusions.

The need for transfusion and ICU admission of patients with PPH in whom the obstetric red code was not activated points to the importance of assessing the impact of prompt identification and activation of the rapid response protocol triggered by the use of a specific uterotonic drug, or by the use of one or more of those medications, because the little validity of clinical estimation of bleeding and the time that it takes to recognise the signs of shock, even in cases of haemorrhage considered as

non-severe¹³ have repercussions in terms of response to treatment and maternal outcomes.

The effectiveness of hydrostatic tamponade found in our case series is consistent with the reports of several observational studies showing that hydrostatic tamponade resolves 74% to 90% of all obstetric haemorrhages that do not respond to medical management during vaginal delivery¹⁷⁻²⁰, and it is recommended as the first line of advanced management when the cause of bleeding is atony¹². In women with obstetric haemorrhage following cesarean section, the second line treatment was surgical in accordance with the WHO recommendations⁵ and because of the availability of the surgical resource at the institution. This finding is also related to the institutional protocol¹¹ which recommends that women in severe shock according to the guidelines must be taken to surgery as soon as possible and given blood products for prompt resuscitation.

The strength of this study is the active identification of the cases through the use of reliable sources such as release of medications from the pharmacy in order to control for measurement, reporting and poor classification biases. This allowed the identification of women with obstetric bleeding who would not have been detected using the activation of the rapid response team or the need for blood product transfusions as criteria. The observational nature of the study, the fact that it is based on records, and the low observed incidence of the event, are limitations of this work.

CONCLUSIONS

The findings of this study point to the need of assessing the effectiveness and safety of more reproducible diagnostic methods in order to determine the right timing for initiating treatment per protocol in women with severe postpartum haemorrhage, regardless of the route of delivery, and to encourage activation of the rapid response team, given the evidence of morbidity and adverse maternal outcomes in those women in whom the obstetric red code was not activated.

REFERENCES

1. American College of Obstetricians and Gynecologists. ACOG Practice Bulletin: Clinical Management Guidelines for Obstetrician-Gynecologists Number 76, October 2006: postpartum hemorrhage. *Obstet Gynecol.* 2006;108:1039-47.
2. Say L, Chou D, Gemmill A, Tunçalp O, Moller A-B, Daniels J, et al. Global causes of maternal death: A WHO systematic analysis. *Lancet Glob Health.* 2014;2:e323-333. [https://doi.org/10.1016/S2214-109X\(14\)70227-X](https://doi.org/10.1016/S2214-109X(14)70227-X)
3. Carroli G, Cuesta C, Abalos E, Gulmezoglu AM. Epidemiology of postpartum haemorrhage: A systematic review. *Best Pract Res Clin Obstet Gynaecol.* 2008;22:999-1012. <https://doi.org/10.1016/j.bpob-gyn.2008.08.004>

4. Francois KE, Foley MR. Antepartum and postpartum hemorrhage. En: Gabbe SG, Niebyl JR, Simpson JL, Landon MB, Galan HL, Jauniaux ERM, et al, Editors. Obstetrics. Normal and Problem Pregnancies. Seventh edition; 2017. p. 395-424.
5. WHO Recommendations for the Prevention and Treatment of Postpartum Haemorrhage. Geneva: World Health Organization; 2012. (WHO Guidelines Approved by the Guidelines Review Committee) [visitado 2017 Sep 15]. Disponible en: <http://www.ncbi.nlm.nih.gov/books/NBK131942/>
6. OB Hemorrhage Toolkit V 2.0 California Maternal Quality Care Collaborative [visitado 2017 Sep 15]. Disponible en: <https://www.cmqc.org/resources-tool-kits/toolkits/ob-hemorrhage-toolkit>
7. Tsu VD. Postpartum haemorrhage in Zimbabwe: a risk factor analysis. Br J Obstet Gynaecol. 1993;100:32733. <https://doi.org/10.1111/j.1471-0528.1993.tb12974.x>
8. Al-Zirqi I, Vangen S, Forsen L, Stray-Pedersen B. Prevalence and risk factors of severe obstetric haemorrhage. BJOG Int J Obstet Gynaecol. 2008;115:1265-72. <https://doi.org/10.1111/j.1471-0528.2008.01859.x>
9. Ayadi AME, Nathan HL, Seed PT, Butrick EA, Hezelgrave NL, Shennan AH, et al. Vital sign prediction of adverse maternal outcomes in women with hypovolemic shock: The role of Shock Index. PLOS ONE. 2016;11:e0148729. <https://doi.org/10.1371/journal.pone.0148729>
10. PRO Morbilidad materna extrema.pdf [visitado 2017 Sep 15]. Disponible en: <http://Vigilancia/sivigila/Protocolos%20SIVIGILA/PRO%20Morbilidad%20Materna%20Extrema.pdf>
11. Vélez-Álvarez GA, Agudelo-Jaramillo B, Gómez-Dávila JG, Zuleta-Tobón JJ. Código rojo: guía para el manejo de la hemorragia obstétrica. Rev Colomb Obstet Ginecol. 2016;60:34-48.
12. Postpartum haemorrhage, prevention and management (Green-top Guideline No. 52). Royal College of Obstetricians and Gynaecologists [visitado 2017 Sep 15]. Disponible en: <https://www.rcog.org.uk/en/guidelines-research-services/guidelines/gtg52/>
13. Rubio-Romero JA, Guevara-Cruz OA, Gaitán-Duarte H. Validity of visual estimation of blood loss as diagnostic tool in severe post partum hemorrhage in an university hospital. Bogotá. 2007. Rev Fac Med. 2010;58:173-84.
14. Rubio-Romero JA, Gaitán-Duarte HG, Rodríguez-Malagón N. Concordancia entre la estimación visual y la medición del volumen recolectado en una bolsa del sangrado intraparto en mujeres con parto normal en Bogotá, Colombia, 2006. Rev Colomb Obstet Ginecol. 2016;59:92-102.
15. Ngwenya S. Postpartum hemorrhage: Incidence, risk factors, and outcomes in a low-resource setting. Int J Womens Health. 2016;8:647-50. <https://doi.org/10.2147/IJWH.S119232>
16. Jiménez-Hernández DL, Guevara-Rodríguez AP, Zuleta-Tobón JJ, Rubio-Romero JA. Tasa de cesáreas por grupos de Robson en una institución de mediana complejidad de la ciudad de Bogotá, 2012-2014. Rev Colomb Obstet Ginecol. 2016;67:101-11. <https://doi.org/10.18597/rcog.381>
17. Kandeel M, Sanad Z, Ellakwa H, El Halaby A, Rezk M, Saif I. Management of postpartum hemorrhage with intrauterine balloon tamponade using a

- condom catheter in an Egyptian setting. *Int J Gynaecol Obstet off Organ Int Fed Gynaecol Obstet*. 2016;135:272-5. <https://doi.org/10.1016/j.ijgo.2016.06.018>
18. Goldrath MH. Uterine tamponade for the control of acute uterine bleeding. *Am J Obstet Gynecol*. 1983;147:869-72. [https://doi.org/10.1016/0002-9378\(83\)90237-5](https://doi.org/10.1016/0002-9378(83)90237-5)
19. Keriakos R, Mukhopadhyay A. The use of the Rusch balloon for management of severe postpartum haemorrhage. *J Obstet Gynaecol*. 2006;26:335-8. <https://doi.org/10.1080/01443610600595077>
20. Condous GS, Arulkumaran S, Symonds I, Chapman R, Sinha A, Razvi K. The “tamponade test” in the management of massive postpartum hemorrhage. *Obstet Gynecol*. 2003;101:767-72

Notes

- 1 Paper submitted for the title as specialist in Obstetrics and Gynaecology, Universidad Nacional de Colombia, Bogotá (Colombia).

Conflict of interest declaration

Conflict of interest: none declared.