



Revista Colombiana de Obstetricia y Ginecología

ISSN: 0034-7434

ISSN: 2463-0225

Federación Colombiana de Obstetricia y Ginecología;
Revista Colombiana de Obstetricia y Ginecología

Eslava-Schmalbach, Javier; Eslava-González, Laura

Iodine deficiency in indigenous pregnant women in Colombia: an equity and public health concern

Revista Colombiana de Obstetricia y Ginecología, vol. 72, no. 1, 2021, January-March, pp. 7-11

Federación Colombiana de Obstetricia y Ginecología; Revista Colombiana de Obstetricia y Ginecología

DOI: <https://doi.org/10.18597/rcog.3675>

Available in: <https://www.redalyc.org/articulo.oa?id=195266777001>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UAEV
redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative



Iodine deficiency in indigenous pregnant women in Colombia: an equity and public health concern

Javier Eslava-Schmalbach, M. D., M. Sc., Ph. D.¹; Laura Eslava-González, B. Sc.²

Iodine deficiency, goiter, hypothyroidism and congenital hypothyroidism have existed as public health problems for many years, and attempted global and local solutions have consisted of interventions at a community and individual level. In Colombia, the frequency of endemic goiter has been historically high. In school children in Caldas, Rueda-Williamson reported a prevalence of 83.1% of endemic goiter in 1945 and of 33.9% in 1952 (1). Later, the same author reported that the implementation of salt iodization in the department of Caldas – one of the first places to introduce fortification on a experimental basis with the aim of preventing endemic goiter - a program created in 1963 by the National Institute of Nutrition, reduced the frequency of goiter in this population of school children by up to 1.8% in 1965 (2). However, by 1984, 21 years after the formal implementation of iodization in Colombia, the frequency of endemic goiter was reported at 52% in Chameza and 13% in Yopal, two municipalities in the department of Casanare, with high average TSH values in both populations (3). Later, the neonatal screening program for congenital hypothyroidism was implemented in 2000 (4). In 2015, the National Nutritional Situation Survey (ENSIN) showed an 16.6% iodine deficiency in children of indigenous ethnicity 1 to 4

years of age, more than twice the deficiency found for children of the same age with no ethnic claim, for which the reported prevalence of iodine deficiency was 7.5% (Table 1) (5). These figures suggest that this nutritional deficiency continues to be a public health problem in many regions of the country.

This issue of the *Colombian Journal of Obstetrics and Gynecology* (*Revista Colombiana de Obstetricia y Ginecología*) features the study conducted by Herrera-Murgueitio *et al.* in 2019 with the aim of identifying the prevalence of iodine deficiency in indigenous pregnant women in five departments of Colombia: Amazonas, Cauca, Córdoba, La Guajira and Meta. In a random sample of a universe of 5,925 pregnant women, the authors found a prevalence of iodine deficiency of 33.2%. In this same group of pregnant women, chronic malnutrition was 27.9% (min-max: 16.6-36.3), the median poverty line was 34.8% (min-max: 25.4-53.0), median extreme poverty was 11.1% (min-max: 9.1-26.7) and median illiteracy was 21.6% (min-max: 17.2-40.9). The prevalence of goiter was 34.4%.

It has been shown clearly that iodine deficiency has an impact on the health of pregnant women and, in particular, on that of their children, considering that it is a cause of abnormal neurological development of the fetus that may result in cognitive, psychomotor, and perceptual impairment, as well as infantile hearing loss (6-8). Studies in laboratory animals (Wistar rats) have shown evidence of molecular changes in the hippocampus of progeny born to mothers subjected to

¹ Health Equity Group Leader, School of Medicine, Universidad Nacional de Colombia; Hospital Universitario Nacional de Colombia, Bogotá (Colombia).

² Behavioral Neurosciences, Simon Fraser University, Burnaby (Canada).

iodine deficiency during pregnancy. The cubs showed a reduction in factors that play a key role in synapsis reconstruction and regulation, cell differentiation, and memory consolidation and maintenance in the hippocampus. These changes resulted in poor performance in spatial memory tests such as the Morris water maze, showing permanent memory and learning changes in the cubs (6). The hippocampus has been described as especially important for consolidation of long-term memory and learning (7).

The permanent neurological effects that have been observed in children born to mothers who had iodine deficiency during pregnancy depend on the severity of the deficiency. It has been suggested that more severe deficiencies may result in more significant changes, including cretinism, a disease characterized by congenital hypothyroidism, mental retardation, deafness, speech difficulties and psychomotor deficits (8,9). It has also been suggested that more severe deficiencies during pregnancy lead to lower scores in intelligence quotient tests in children. Moderate or mild deficiencies may lead to perception and cognition deficits, an even the development of attention deficit and hyperactivity disorder (8). In contrast, the treatment of these deficiencies during the first trimester of pregnancy may improve neurological development and prevent permanent neurological consequences (8-10).

As highlighted by Herrera-Murgueitio *et al.*, this is the first approach to the exploration of iodine deficiency and the presence of goiter in indigenous pregnant women as a population in which significant inequities associated with maternal mortality have been described in Colombia. Two prior studies had shown that maternal mortality gaps were found in the departments with the highest poverty levels (11) or with the most profound basic unmet needs (12). The indigenous pregnant women described in the study live in some of those departments.

Likewise, the ENSIN-2015 survey reveals profound differences in social determinants of health as they relate to indigenous ethnicity, and reflects that

iodine deficiency is a proxy for a structural problem that alienates these populations from potential developments that occur in urban areas or in big cities. This alienation, according to the level of iodine deficiency, places the pregnant women of the study in a similar situation as that in 1952, when experimental research in Colombia (2) managed to lower endemic goiter to similar levels as those found among indigenous pregnant women in the study by Herrera-Murgueitio *et al.* for 2019. Other authors show a very close relationship between maternal iodine deficiency and the deficiency in their children, given that they find themselves in the same environment, with the same structural deficiencies and food insecurity (13).

The ENSIN-2015 survey shone the light on the problems of housing, access to water systems, low education in mothers of children under 5 years, informal work, high food insecurity and global malnutrition among the indigenous population, when compared to the group with no ethnic claim (Table 1).

A previous study on social determinants of health associated with early death due to malnutrition in children under 5 years (14) proposes an explanatory model that includes the following: 1) socioeconomic and political conditions; 2) the environment with its environmental conditions, rurality, armed conflict, food availability, safe water and access to utilities; 3) parents and caregivers, which includes ethnicity, education, work and housing conditions; 4) individual-related aspects. Many of these variables are shown in Table 1 and reflect the level of vulnerability of the indigenous population in Colombia, not only for congenital hypothyroidism but also for premature malnutrition-related mortality.

On the other hand, an exploration conducted by the Institute for Health Metrics and Evaluation (IHME) shows Disability Adjusted Life Years (DALY) for Colombia in relation to iodine deficiency-related disorders. In this regard, although Colombia showed a drop in DALYs between 2005 and 2010, the values were again similar to those reported in 1990 (15). In other words, a 30-year regression.

Table 1.
Deficits in different social determinants of health assessed in the ENSIN-2015

Assessed determinant	Indigenous (%) (95% CI)	No ethnic claim (%) (95 % CI)
Quantitative housing deficit	*10.9 (8.1-14.6)	6.5 (5.8-7.2)
Qualitative housing deficit	51.2 (43.7-58.6)	18.7 (17.4-20)
With public water supply systems	42.2 (34.5-50.3)	79.8 (78.3-81.2)
Mothers of children under 5 years of age, with completed secondary education and incomplete higher education	25 (18.8-32.5)	54.7 (52.5-56.8)
Employment informality	*81 (74.1-86.5)	43.6 (41.7-45.4)
Household in the lowest quartile of the wealth index	67.8 (61.7-73.3)	22.5 (20.9-24.2)
Household with food insecurity	76.9 (74.5-79.1)	52.3 (54.2-55.9)
Self-consumption	50.2 (43.9-56.5)	14.9 (13.9-16.0)
Coping strategies in the past 7 days due to food insecurity	71.3 (66.2-75.9)	49.9 (48.7-51.1)
Households with children under 18 years of age who sent a child to neighbors or relatives due to the lack of food	15 (12.0- 18.6)	7.7 (7.2-8.2)
Minimum acceptable food intake in children under 2 years of age	24.7 (18.6-32.1)	37.4 (34.6-40.3)
Iodine deficiency in children from 1 to 4	16.6 (12.3-21.9)	7.5 (6.4-8.7)
Global malnutrition in children from 0 to 4 years of age	7.2 (5.1-10.2)	3.0 (2.4-3.7)
Pregnant women out of the total, from 10 to 19 years of age	24.8 (16.1-36.2)	19.2 (16.2-22.5)

Source: Authors, from data taken from ENSIN-2015 (5).

* Variation coefficient 15 to 20 %, acceptable accuracy, according to ENSIN-2015.

Apart from salt iodization and congenital hypothyroidism screening, Colombia launched a strategy for the prevention and control of micronutrient deficiencies (16), iodine included. However, according to the study by Herrera-Murgueitio et al., the strategy has not yet reached the regions inhabited by the indigenous population, or at least has not been sustainable, as was the case with the declaration for the elimination of iodine deficiency disorders in Colombia in 1998 (17).

The results found by Herrera-Murgueitio et al. could be extrapolated not only to the 5,925 indigenous pregnant women who are part of the study universe (at least 2,000 households in a situation of food and nutritional insecurity) but also to other marginalized populations in the inner cities or the

rural areas which experience similar problems of food insecurity during gestation and in their daily lives, as they are subjected to the same impact of social determinants. Congenital hypothyroidism is a preventable, avoidable and by all means intolerable condition because of its potential consequences (18).

The salt iodization policy needs to be revisited and the impact on the people that benefit from it or who do not have access to it needs to be closely re-examined. Monitoring of urine iodine, iodine intake, or thyroglobulin levels (19), or other forms of evaluation (20) must be part of the regular prenatal care of pregnant women, developing children and the general population. Be as it may, outcomes will also be a reflection of the underlying social determinants of health.

REFERENCIAS

1. Rueda-Williamson R, Pardo Téllez F, Piedrahíta Hoyos F, Ariza Macías J, Uribe Naranjo L. La efectividad de la yodación de la sal en la prevención del bocio endémico en Colombia: resultados en escolares de Caldas. *Rev Soc Colomb Endocrinol*. 1966;4(2):271-6.
2. Rueda-Williamson R, Pardo Téllez F. La prevención del bocio endémico en Colombia. *Bol Sanit Panam*. 1966:495-503.
3. Carrillo J, Mockus I, Casadiego C, Salcedo L, Tarazona B, Galarza L, et al. Evaluación de desórdenes por deficiencia de Yodo (DDI) en dos poblaciones de Casanare, Colombia, 1984-1986. *Temas de Pediatría*. Vevey: Nestlé, Servicio de Información Científica; 1986.
4. Instituto Nacional de Salud. Manual de tamizaje neonatal por el laboratorio. Bogotá: Instituto Nacional de Salud; 2014. 80 p.
5. Gobierno de Colombia. Encuesta Nacional de Situación Nutricional ENSIN 2015. Instituto Colombiano de Bienestar Familiar, Instituto Nacional de Salud, Universidad Nacional de Colombia; 2015.
6. Liu Y, Zhang L, Li J, Shan Z, Teng W. Maternal marginal iodine deficiency affects the expression of relative proteins during brain development in rat offspring. *J Endocrinol*. 2013;217(1):21-9. <https://doi.org/10.1530/JOE-12-0410>
7. Gerges NZ, Alkadhi KA. Hypothyroidism impairs late LTP in CA1 region but not in dentate gyrus of the intact rat hippocampus: MAPK involvement. *Hippocampus*. 2004;14: 40-45. <https://doi.org/10.1002/hipo.10165>
8. Caron P. Neurocognitive outcomes of children secondary to mild iodine deficiency in pregnant women. *Ann Endocrinol*. 2015;76(3):248-52. <https://doi.org/10.1016/j.ando.2015.01.001>
9. Zimmermann MB. The role of iodine in human growth and development. *Semin Cell Dev Biol*. 2011;22(6):645-52. <https://doi.org/10.1016/j.semcdb.2011.07.009>
10. Lavado-Autric R, Ausó E, García-Velasco JV, Arufe MC, Escobar del Rey F, Berbel P, et al. Early maternal hypothyroxinemia alters histogenesis and cerebral cortex cytoarchitecture of the progeny. *J Clin Invest*. 2003;111(7):1073-82. <https://doi.org/10.1172/JCI16262>
11. Cárdenas-Cárdenas LM, Cotes-Cantillo K, Chaparro-Narváez PE, Fernández-Niño JA, Paternina-Cañedo A, Castañeda-Orjuela C, et al. Maternal mortality in Colombia in 2011: A two level ecological study. *PLoS One*. 2015;10(3):1-14. <https://doi.org/10.1371/journal.pone.011894>
12. Sandoval-Vargas Y, Eslava-Schmalbach J. Inequidades en mortalidad materna por departamentos en Colombia para los años 2000-2001, 2005-2006 y 2008-2009. *Rev Salud Pública*. 2013;15(4):577-89.
13. Wong EM, Sullivan KM, Perrine CG, Rogers LM, Peña-Rosas JP. Comparison of median urinary iodine concentration as an indicator of iodine status among pregnant women, school-age children, and nonpregnant women. *Food Nutr Bull*. 2011;32(3):206-12. <https://doi.org/10.1177/156482651103200304>
14. Agudelo Ibáñez DR. Mortalidad por desnutrición infantil en menores de 5 años y sus determinantes a nivel municipal en Colombia 1998-2016. Bogotá: Universidad Nacional de Colombia; 2019.
15. Institute for Health Metrics and Evaluation. GBD Compare | IHME Viz Hub. Global Burden of Disease. 2017. Disponible en: <https://vizhub.healthdata.org/gbd-compare/>
16. Ministerio de Salud y Protección Social. Estrategia Nacional para la prevención y control de las deficiencias de micronutrientes en Colombia 2014-2021. Bogotá: Ministerio de Salud y Protección Social; 2015. 72 p.
17. Ministerio de Salud y Protección Social. Definición del problema para el análisis del impacto normativo relacionado con sal para consumo humano. Bogotá: Ministerio de Salud y Protección Social; 2019.
18. Rahmani K, Yarahmadi S, Etemad K, Koosha A, Mehrabi Y, Aghang N, et al. Congenital hypothyroidism: Optimal initial dosage and time of initiation of treatment: A systematic review. *Int J Endocrinol Metab*. 2016;14(3). <https://doi.org/10.5812/ijem.36080>
19. Katko M, Gazso AA, Hircsu I, Bhattoa HP, Molnar Z, Kovacs B, et al. Thyroglobulin level at week 16 of pregnancy is superior to urinary iodine concentration

in revealing preconceptional and first trimester iodine supply. *Matern Child Nutr.* 2018;14(1). <https://doi.org/10.1111/mcn.12470>

20. Zimmermann MB, Andersson M. Assessment of iodine nutrition in populations: Past, present, and future. *Nutr Rev.* 2012;70(10):553-70. <https://doi.org/10.1111/j.1753-4887.2012.00528.x>