



Revista Ciência Agronômica

ISSN: 0045-6888

ISSN: 1806-6690

Universidade Federal do Ceará

Medeiros, Wilker Nunes; Valente, Domingos Sárvio Magalhães; Queiroz, Daniel
Marçal de; Pinto, Francisco de Assis de Carvalho; Assis, Igor Rodrigues de
Apparent soil electrical conductivity in two different soil types¹
Revista Ciência Agronômica, vol. 49, no. 1, January-March, 2018, pp. 43-52
Universidade Federal do Ceará

DOI: 10.5935/1806-6690.20180005

Available in: <http://www.redalyc.org/articulo.oa?id=195358115005>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

UAEM 

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Apparent soil electrical conductivity in two different soil types¹

Condutividade elétrica aparente do solo em dois tipos de solo

Wilker Nunes Medeiros^{2*}, Domingos Sárvio Magalhães Valente², Daniel Marçal de Queiroz², Francisco de Assis de Carvalho Pinto² and Igor Rodrigues de Assis²

ABSTRACT - Mapping the apparent soil electrical conductivity (EC_a) has become important for the characterization of the soil variability in precision agriculture systems. Could the EC_a be used to locate the soil sampling points for mapping the chemical and physical soil attributes? The objective of this work was to examine the relations between EC_a and soil attributes in two fields presenting different soil textures. In each field, 50 sampling points were chosen using a path that presented a high variability of EC_a obtained from a preliminary EC_a map. At each sampling point, the EC_a was measured in soil depths of 0-20, 0-40 and 0-60 cm. In addition, at each point, soil samples were collected for the determination of physical and chemical attributes in the laboratory. The EC_a data obtained for different soil depths was very similar. A large number of significant correlations between EC_a and the soil attributes were found. In the sandy clay loam texture field there was no correlation between EC_a and organic matter or between EC_a and soil clay and sand content. However, a significant positive correlation was shown for the remaining phosphorus. In the sandy loam texture field the EC_a had a significant positive correlation with clay content and a significant negative correlation with sand content. The results suggest that the mapping of apparent soil electrical conductivity does not replace traditional soil sampling, however, it can be used as information to delimit regions in a field that have similar soil attributes.

Key words: Precision agriculture. Management zones. Soil sampling.

RESUMO - O mapeamento da condutividade elétrica aparente do solo (CE_a) tornou-se importante para caracterizar a variabilidade do solo em sistemas de agricultura de precisão. A CE_a pode ser usada para definir os pontos de amostragem para mapeamento de atributos químicos e físicos do solo? O objetivo deste trabalho foi avaliar as relações entre CE_a e atributos do solo em duas áreas com texturas de solo diferentes. Em cada área, 50 pontos de amostragem foram escolhidos considerando uma linha de caminamento que apresentou alta variabilidade de CE_a , obtida através de um mapeamento preliminar do atributo. Em cada ponto de amostragem a CE_a foi mensurada nas profundidades de 0-20, 0-40 e 0-60 cm do solo, recolhendo-se também amostras de solo para determinação dos atributos físicos e químicos em laboratório. Os dados de CE_a mensurados em diferentes profundidades do solo foram muito similares. Observou-se um grande número de correlações significativas entre CE_a e os atributos do solo. Na área de solo arenoso, não houve correlação entre CE_a e matéria orgânica ou entre CE_a e teor de argila e de areia. Entretanto, uma correlação positiva significativa foi observada para o fósforo remanescente. Na área de solo argiloso, a CE_a teve correlação positiva significativa com o teor de argila e correlação negativa significativa com o teor de areia. Os resultados sugerem que o mapeamento da CE_a não substitui a amostragem tradicional do solo, porém, pode ser utilizada como informação para delimitar regiões que apresentam atributos de solo semelhantes em uma área.

Palavras-chave: Agricultura de precisão. Zonas de manejo. Amostragem do solo.

DOI: 10.5935/1806-6690.20180005

*Autor para correspondência

Recebido para publicação em 26/02/2015; aprovado em 26/04/2017

¹Parte da Dissertação do primeiro autor apresentada no Programa de Pós-Graduação em Engenharia Agrícola/DEA/UFV

²Departamento de Engenharia Agrícola, Universidade Federal de Viçosa/UFV, Avenida Peter Henry Rolfs, s/n - Campus Universitário, Viçosa-MG, Brasil, 36.570-900, wilkermedeiros@yahoo.com.br, valente@ufv.br, queiroz@ufv.br, facpinto@ufv.br, igor.assis@ufv.br

INTRODUCTION

Globally, Brazil is an important grain producer. This position was achieved with the aid of research and development programs including plant breeding, mechanization, and incorporating new methods and techniques of cultivation. It is important to highlight the intensification of no-till planting systems, and more recently, the use of precision agriculture systems.

The use of precision agriculture has become an important tool for producing food to meet the demands of the global population, to improve economic return and to preserve the environment. It is described as a set of tools applied to agriculture that enables site specific management of agriculture areas. The application of precision agriculture is generally conducted by considering the spatial and temporal variability of crop fields to increase production.

Mapping of the apparent soil electrical conductivity (EC_a) has been adopted in precision agriculture systems to characterize the variability of chemical and physical soil attributes. Research has shown that apparent soil electrical conductivity is highly correlated with different chemical and physical soil properties and also with crop yields (ALCÂNTARA; REIS; QUEIROZ, 2012; FAROOQUE *et al.*, 2012; GHOLIZADEH *et al.*, 2012; MOLIN; FAULIN, 2013; MORARI; CASTRIGNANÒ; PAGLIARIN, 2009; SUDDUTH *et al.*, 2005; TERRÓN *et al.*, 2011; VALENTE *et al.*, 2012). Because of these correlations, areas with similar soil characteristics are identified more quickly and at a lower cost and treated in a site-specific way.

However, apparent soil electrical conductivity is influenced by a large number of soil attributes, which complicates the interpretation of the generated maps of this soil attribute. For instance, it may be influenced by static or dynamic soil characteristics such as salinity, texture, mineralogy, moisture content, density, temperature and organic matter content (EKWUE; BARTHOLOMEW, 2010; FRIEDMAN, 2005; JOHNSON *et al.*, 2003; MORAL; TERRÓN; SILVA, 2010). Corwin and Lesch (2003) report that attributes with a greater influence on apparent soil electrical conductivity should be identified in each analyzed field. The EC_a can be used to locate the soil sampling points for mapping the chemical and physical soil attributes? Thus, considering the potential application and demand for studying the relationship between the apparent soil electrical conductivity and soil attributes, the objective of this work was to analyze the relations between apparent soil electrical conductivity and soil attributes that are used for managing precision agriculture systems.

MATERIAL AND METHODS

Characterization of the fields used for studying the apparent soil electrical conductivity

This study was performed in two agricultural fields located in the municipalities of Viçosa (Field 1) and Ponte Nova (Field 2), with different physical attributes of soil. According to the Köppen system, the climate of Viçosa and Ponte Nova is characterized as Cwb. This is a mesothermic climate with mild, rainy summers and dry winters. The soil of study area is kandic oxisol, in soft to wavy relief. The predominant vegetation is characterized as a submontane seasonal semideciduous Forest. The geology of the area is characterized by the predominance of banded and migmatite orthogneiss, with the presence of hornblende-biotite and amphibole-biotite.

The Field 1, with an area of 10,703.72 m², presents a sandy clay loam texture and was not being cultivated for four years. The Field 2, with an area of 14,078.68 m², presents a sandy loam texture and was being cultivated with sugarcane around for four years. Table 1 shows the soil chemical and physical characteristics of the study area.

In a preliminary analysis, the apparent soil electrical conductivity was mapped in Field 1 and Field 2. During this analysis, 203 and 163 apparent soil electrical conductivity (EC_a) measurements were made in Field 1 and Field 2, respectively. The measurements were obtained using a portable Landviser® brand model LandMapper® ERM-02, which uses the principle of electrical resistivity through a four-electrode contact probe. The electrodes were configured based on the Wenner matrix, as described by Corwin and Hendrickx (2002) and Corwin and Lesch (2003), to measure the apparent soil electrical conductivity in the 0-0.20 m soil profile. For both fields, the apparent soil electrical conductivity maps, using the kriging methodology, were generated (Figure 1).

From the apparent soil electrical conductivity map of each field, a path through the field was defined for analysis. In both fields, the path was obtained by marking 50 sampling points such that they represented a high variability of the apparent soil electrical conductivity (Figure 1). Each sampling point represented an area of 2×2 m where apparent soil electrical conductivity measurements were conducted on 20 different days. The EC_a determinations were performed over 20 days between October 19th and December 19th, 2012 in the raining season. This procedure permitted the collection of the EC_a under different soil moisture conditions.

The delimitation of the fields and the location of the sampling points in both fields were performed using a ProMark3 model Survey GPS (L1), made by Magellan.

The post-processed differential correction was conducted using the database of the Brazilian Institute of Geography and Statistics (IBGE). The datum used was SIRGAS 2000 and the data correction was performed using the GNSS Solutions™ software supplied by the GPS device maker.

Determination of apparent soil electrical conductivity

Three different four-electrode configurations based on the Wenner matrix (CORWIN; HENDRICKX, 2002; CORWIN; LESCH, 2003) were used. Electrode spacing for the first probe was 0.20 m (designated as EC_a20); the second probe electrode spacing was 0.40 m (designated as EC_a40); and the third electrode spacing was 0.60 m (designated as EC_a60). The probe electrode spacing determined the soil depth of EC_a measurement, thus the three probes measured EC_a from surface to 0.2, 0.4 and 0.6m, respectively. For each daily measurement, the EC_a was measured with the three probes in less than 40 minutes to cover all the 50 sampling points, which was done to avoid the effects of temperature and soil moisture variation during the measurements.

Physical and chemical soil attributes determination

For each of the sampling points (Figure 1), a soil sample composed of four single samples representing the 0-0.20 m layer was obtained. This layer was chosen because it is the most common layer used in precision agriculture management. The samples were collected with a Dutch auger at a distance of up to 2.0 m from the georeferenced sampling point, and after homogenization, a soil sample of approximately 300 g was collected.

The soil moisture content, %, was determined by a thermo-gravimetric Embrapa method (DONAGEMA *et al.*, 2011). From the 50 sampling points, the samples with an even number were chosen for soil moisture determination. Then, the samples were conditioned to avoid the loss of water and sent to the laboratory where they were weighed and dried at 105 °C for twenty-four hours to determine the mass of the dry soil.


Soil particle size analysis was performed using the Embrapa pipette method (DONAGEMA *et al.*, 2011) to determine the fractions of sand, silt and clay present in the

Table 1 - Soil chemical and physical characteristics of Field 1 and Field 2

Variable	Unit	Field 1	Field 2
pH	-	5.67	5.68
P	mg dm ⁻³	3.71	15.40
K	mg dm ⁻³	73.62	52.46
Ca ²⁺	cmolc dm ⁻³	2.08	1.71
Mg ²⁺	cmolc dm ⁻³	0.56	0.59
Al ³⁺	cmolc dm ⁻³	0.07	0.04
H + Al	cmolc dm ⁻³	3.68	2.65
SB	cmolc dm ⁻³	2.83	2.44
CTC t	cmolc dm ⁻³	2.90	2.48
CTC T	cmolc dm ⁻³	6.51	5.09
V	%	43.14	47.88
M	%	3.98	2.02
OM	dag kg ⁻¹	3.30	1.99
P-rem	mg L ⁻¹	31.40	38.30
Sandy	%	53.84	65.76
Silt	%	14.44	13.08
Clay	%	31.72	21.16
Moisture	%	23.84	21.11
Altitude	m	671.21	452.93

pH (1:2,5); P – phosphorus; K – potassium; Ca – calcium; Mg – magnesium; Al – aluminium; H + Al – potential acidity; SB – base sum; CTC t – effective cation exchange capacity; CTC T – total cation exchange capacity; V – base saturation; m – aluminium saturation; OM – organic matter; P-rem – remaining phosphorus. P – K: Mehlich 1 extractor; Ca - Mg - Al – KCl 1 mol/L extractor; H+Al – calcium acetate extractor 0,5 mol/L pH 7,0; Moisture: average of 25 samples in 20 days

Figure 1 - The apparent soil electrical conductivity maps determined for the 0-0.20 m soil layer with the sampling points marked in white for (a) Field 1 and (b) Field 2



soil. A chemical characterization of the soil was completed by determining the active acidity (pH), potential acidity ($H + Al$), calcium (Ca^{2+}), magnesium (Mg^{2+}), potassium (K^+), aluminum (Al^{3+}), phosphorus (P), remaining phosphorus (P-rem), organic matter (OM), sum of bases (SB), effective cation exchange capacity (CEC t), cation exchange capacity of soil at pH 7 (CEC T), aluminum saturation (m) and base saturation (V) using Embrapa method (DONAGEMA *et al.*, 2011).

Analysis of the relationship between apparent soil electrical conductivity and soil physical and chemical attributes

First, an exploratory data analysis was performed. In this analysis, the outliers were removed using the method presented by Hoaglin, Mosteller and Tukey (1992) and Libardi *et al.* (1996). The final decision as to whether a data point needed to be removed was based on the comparison of each outlier candidate and the values of the apparent soil electrical conductivity in the neighboring data points. If the outlier candidate value presented a great discontinuity related to the neighbor points, then this data point was removed. After removing the outliers, the

following statistical parameters were obtained: the mean value, minimum and maximum values, standard deviation and coefficient of variation. The z-test was performed to compare the attributes of the two fields. The Pearson correlation between the apparent soil electrical conductivity and each soil attribute determined by laboratory analysis was obtained.

RESULTS AND DISCUSSION

The results of the descriptive statistical analysis for both fields are presented in Table 2.

Using the z-test, it was found that the mean soil attributes of Field 1 were significantly different from those of Field 2 ($p \leq 0.05$), except for the soil attributes pH, Mg^{2+} and silt. Both fields were characterized by pH values favorable to plant development, and the values of the coefficient of variation for this attribute were lower than 12% (Table 2), which according to the classification proposed by Warrick and Nielsen (1980), is low. According to these authors, this parameter has an average variation of $12\% < CV < 60\%$ and a high variation if $CV > 60\%$. Field 1 presented a high coefficient of variation in the concentration of P, Al^{3+} and m , and a low variation of CEC T, P-rem and sand content (Table 2). The other measured soil attributes from Field 1 had CV values classified as medium (Table 2). Field 2 presented high values of CV for Mg, Al^{3+} , m and silt content; while only the soil attribute of pH presented a low CV value (Table 2). In general, Field 2 showed a greater variability of the 19 soil attributes measured, with 13 showing higher CV values when compared to Field 1. The physical soil attributes tended to have more variation in Field 2 than on Field 1. The main reason for these results was that the Field 2 was being cultivated with sugarcane for four years.

The soil moisture content was measured in 25 of 50 sampling points along the pathway line during 20 different days. The soil moisture average for the 20 days of measurement ranged from 19.81 to 26.07% for Field 1 and from 15.89 to 24.74% for Field 2. This variation occurred because the data was collected during the raining season. The ranges of variation for soil moisture of each point of measurement are shown in Figure 2.

Analysis of the results presented on Figure 2(a) show that Field 1 showed a lower variability of soil moisture. Additionally, Field 1 showed a more uniform soil moisture distribution among the sampling points. Field 2 has a uniform soil moisture distribution up until sampling point 28; however, from this point until the end point, the soil of Field 2 seems to have a different water holding capacity. This variation in soil moisture was directly

related to the soil texture variation observed in the field and confirmed by the texture analysis. Sampling points 30 to 50 had higher percentages of sand (above 70%) and lower clay than the other sampling points. According to


Libardi (2005), the properties of the clay define the soil water holding capacity, and the variation in soil moisture of a field is then related to the spatial variability of clay content.

Table 2 - Summary statistic results for soil attributes of Field 1 and 2

Soil Attributes	Unit	Field 1					Field 2				
		Mean	Min	Max	s ¹	CV ²	Mean	Min	Max	s	CV
pH	-	5.67	4.80	6.20	0.34	5.98	5.68	4.80	6.30	0.41	7.28
P	mg dm ⁻³	3.71	1.20	14.80	2.76	74.52	15.40	5.50	39.70	9.10	59.09
K	mg dm ⁻³	73.62	17.00	189.00	40.46	54.96	52.46	22.00	100.00	21.93	41.81
Ca ²⁺	cmol _c dm ⁻³	2.08	0.60	3.50	0.69	33.10	1.71	0.70	3.00	0.58	33.92
Mg ²⁺	cmol _c dm ⁻³	0.56	0.10	1.00	0.23	40.65	0.59	0.10	1.20	0.38	63.30
Al ³⁺	cmol _c dm ⁻³	0.07	0.00	0.60	0.15	211.92	0.04	0.00	0.30	0.07	182.11
H + Al	cmol _c dm ⁻³	3.68	2.64	5.61	0.82	22.20	2.65	0.17	4.46	0.96	36.39
SB	cmol _c dm ⁻³	2.83	0.86	4.86	0.95	33.57	2.44	0.87	4.15	0.97	39.70
CEC t	cmol _c dm ⁻³	2.90	1.46	4.86	0.85	29.38	2.48	0.97	4.15	0.96	38.83
CEC T	cmol _c dm ⁻³	6.51	5.54	8.00	0.54	8.27	5.09	2.09	8.28	1.76	34.59
V	%	43.14	13.00	64.00	12.83	29.74	47.88	29.00	93.00	10.73	22.40
m	%	3.98	0.00	41.00	9.41	236.44	2.02	0.00	15.00	3.69	182.92
OM	dag kg ⁻¹	3.30	2.40	4.50	0.45	13.66	1.99	1.10	3.00	0.48	24.00
P-rem	mg L ⁻¹	31.40	22.00	36.70	3.57	11.35	38.30	27.30	48.10	5.64	14.73
Sand	%	53.84	49.00	66.00	3.72	6.90	65.76	38.00	82.00	15.93	24.22
Silt	%	14.44	6.00	22.00	4.03	27.92	13.08	3.00	30.00	8.95	68.41
Clay	%	31.72	16.00	41.00	5.05	15.93	21.16	11.00	34.00	7.68	36.29
Soil Moisture	%	23.84	18.52	29.43	2.87	12.03	21.11	8.44	43.57	12.32	58.35

¹s – standard deviation; ²CV – coefficient of variation; pH – hydrogen ion activity (1:2.5); P – phosphorous; K – potassium; Ca – Calcium; Mg – Magnesium; Al – Aluminum; H + Al – Potential acidity; SB – Sum of bases; CEC t – Effective cation exchange capacity; CEC T – Cation exchange capacity at pH 7; V – Base saturation; m – Aluminum saturation; OM – Organic matter Walkley-Black method (organic C × 1.724); P-rem – Remaining phosphorous, P – K: Mehlich-1 Extractor; Ca - Mg - Al: KCl 1 mol/L extractor; H + Al: Calcium acetate 0.5 mol/L pH 7.0 extractor; Soil moisture, mean value of the 25 sampling points in the 20 days of measurement

Figure 2 - The soil moisture variation in the 50 sampling point for the 20 days of measurement in (a) Field 1 and (b) Field 2



The mean values of EC_a for each soil layer in the 50 sampling points in the two Fields are presented in Figure 3.

The average values of EC_a determined in different layers of soil (EC_{a20} , EC_{a40} and EC_{a60}) showed similar behavior within each field. Field 1 presented values of EC_a between 0.5 and 20 $mS\ m^{-1}$ and Field 2 between 1.0 and 24 $mS\ m^{-1}$. These values are similar to those found by other researchers in non-saline soils (AIMRUN *et al.*, 2007; ALCÂNTARA; REIS; QUEIROZ, 2012; FAULIN; MOLIN, 2006; VALENTE *et al.*, 2012). As the EC_a value is directly influenced by the amount of dissolved salts in the soil solution, it is expected that the values found here are lower compared to those of saline soils and those found in temperate climate regions. The source material involved in soil formation, and the climatic conditions that soils have been exposed to, reflect the type, quantity

and quality of clay, and consequently, the cation exchange capacity, the amount of ions available to the soil solution and ultimately the values of EC_a .

When the values of soil moisture to mean values of EC_a in each field are compared (Figures 2 and 3), it was observed that the average EC_a was higher in those points with higher moisture content. This relationship was more expressive for Field 2. To statistically determine this relationship, linear correlation analyses were performed between soil moisture and the EC_a values obtained at different soil depths for both fields and each day of measurement (Table 3).

For Field 1, the correlations between EC_a and soil moisture were not significant for EC_{a60} only on the seventh day of evaluation and for EC_{a40} only on the twentieth day of evaluation (Table 3). For the other correlations,

Figure 3 - The mean values of EC_a (20 days) in the 50 sampling points for soil depths of 0-20 cm (EC_{a20}), 0-40 cm (EC_{a40}), and 0-60 cm (EC_{a60}) for (a) Field 1 and (b) Field 2




Table 3 - Pearson's coefficient of correlation between soil electrical conductivity values and soil moisture evaluated at three soil depths (0-0.20 m EC_a 20, 0-0.40 m EC_a 40 and 0-0.60 m EC_a 60) for Field 1 and Field 2

Day of Evaluation	-----Field 1-----			-----Field 2-----		
	EC _a 20	EC _a 40	EC _a 60	EC _a 20	EC _a 40	EC _a 60
1	0.62	0.58	0.59	0.90	0.90	0.90
2	0.72	0.77	0.72	0.91	0.90	0.90
3	0.71	0.66	0.57	0.93	0.92	0.91
4	0.58	0.55	0.55	0.91	0.90	0.85
5	0.73	0.73	0.69	0.91	0.90	0.85
6	0.63	0.51	0.57	0.91	0.92	0.93
7	0.47	0.46	0.40 ^{ns}	0.90	0.95	0.95
8	0.53	0.55	0.47	0.96	0.93	0.94
9	0.63	0.58	0.45	0.95	0.87	0.83
10	0.56	0.64	0.56	0.87	0.93	0.83
11	0.58	0.59	0.58	0.95	0.97	0.94
12	0.66	0.63	0.52	0.94	0.93	0.88
13	0.66	0.62	0.52	0.91	0.94	0.82
14	0.62	0.66	0.60	0.90	0.91	0.89
15	0.53	0.50	0.49	0.95	0.93	0.97
16	0.56	0.51	0.53	0.92	0.84	0.86
17	0.53	0.53	0.51	0.81	0.76	0.77
18	0.56	0.62	0.51	0.95	0.97	0.96
19	0.56	0.62	0.50	0.95	0.88	0.92
20	0.46	0.38 ^{ns}	0.45	0.94	0.88	0.98

^{ns} – Non-significant (p>0.05). All of the other values were significant at p≤0.05

as well as all sampling points in Field 2, correlation values were statically significant (Table 3), indicating a close relationship between the two soil attributes and that soil moisture positively affects the values of EC_a. The relationship between EC_a and soil moisture has been studied by several researchers (BREVİK; FENTON; LAZARI, 2006; EKWUE; BARTHOLOMEW, 2010; ISLAM *et al.*, 2012; RHOADES, 1996; SERRANO; SHAHIDIAN; SILVA, 2013). In this work, the relationship between EC_a and soil moisture was stronger in Field 2, which is indicated by the higher correlation coefficient values (Table 3). This behavior is justified by a greater soil moisture variability of soil in the field. When a higher soil moisture occurs, there is a greater ability to conduct electrical current through the pore water, which contains dissolved electrolytes. The amount of water does not indicate the presence of loads in the ground, but it plays an important role in the formation of the solution that conducts electric current (CORWIN; LESCH, 2005).

The correlation analysis between the EC_a for different soil depth and the measured soil attributes

showed similar values for all of the data acquisition days. Therefore, only the results of the correlation between the mean values of EC_a of each sampling point and the soil attribute values are presented (Table 4).

In general, there were a large number of significant correlations (p≤0.05) between EC_a and soil attributes, with great similarity between the data obtained for different soil depths (Table 4). In Field 1, there was no correlation between EC_a and organic matter and with the physical attributes of sand and clay contents. The EC_a did not correlate with clay content; however, it presented a significant positive correlation with the remaining phosphorous, which was similar to what was found by Valente *et al.* (2012). In this case, other interactions among soil attributes were more relevant in the change of soil charge, which caused a positive correlation between EC_a and the remaining phosphorus. The largest coefficient was $r = 0.69$, between the EC_a 20 and K content, and this value was also observed for apparent soil electrical conductivity of EC_a 40 and EC_a 60 (Table 4).

Table 4 - Pearson's coefficient of correlation between the mean values of soil apparent electrical conductivity measured at three different depths (0-0.20 m EC_a 20, 0-0.40 m EC_a 40 and 0-0.60 m EC_a 60) and the soil attributes for Field 1 and Field 2

Soil Attributes	-----Field 1-----			-----Field 2-----		
	EC _a 20	EC _a 40	EC _a 60	EC _a 20	EC _a 40	EC _a 60
pH	0.57	0.52	0.53	-0.22 ^{ns}	-0.14 ^{ns}	-0.15 ^{ns}
P	0.65	0.63	0.64	-0.71	-0.66	-0.60
K	0.69	0.67	0.68	0.68	0.71	0.75
Ca ²⁺	0.62	0.58	0.59	0.78	0.78	0.76
Mg ²⁺	0.49	0.44	0.44	0.94	0.95	0.93
Al ³⁺	-0.42	-0.40	-0.39	0.13 ^{ns}	0.12 ^{ns}	0.11 ^{ns}
H + Al	-0.54	-0.50	-0.51	0.80	0.81	0.82
SB	0.64	0.60	0.61	0.87	0.88	0.86
CTC t	0.64	0.60	0.60	0.89	0.89	0.88
CTC T	0.32	0.29	0.30	0.92	0.93	0.92
V	0.61	0.57	0.57	0.09 ^{ns}	0.08 ^{ns}	0.05 ^{ns}
m	-0.40	-0.37	-0.36	-0.09 ^{ns}	-0.10 ^{ns}	-0.10 ^{ns}
OM	0.16 ^{ns}	0.09 ^{ns}	0.08 ^{ns}	0.88	0.88	0.87
P-rem	0.68	0.66	0.65	-0.94	-0.95	-0.93
Sand	-0.21 ^{ns}	-0.22 ^{ns}	-0.18 ^{ns}	-0.96	-0.94	-0.89
Silt	0.54	0.55	0.58	0.92	0.92	0.87
Clay	-0.27 ^{ns}	-0.28 ^{ns}	-0.33 ^{ns}	0.91	0.88	0.82

^{ns} – Non-significant ($p > 0.05$). All of the other values were significant at $p \leq 0.05$. pH – hydrogen ion activity (1:2.5); P – phosphorous; K – potassium; Ca – Calcium; Mg – Magnesium; Al – Aluminum; H + Al – Potential acidity; SB – Sum of bases; CTC t – Effective cation exchange capacity; CTC T – Cation exchange capacity at pH 7; V – Base saturation; m – Aluminum saturation; OM – Organic matter; P-rem – Remaining phosphorous, P – K: Mehlich-1 Extractor; Ca - Mg - Al –KCl 1 mol/L extractor; H + Al – Calcium acetate 0.5 mol/L pH 7.0 extractor; Soil moisture, the mean value of the 25 sampling points in the 20 days of measurement

In Field 2, only the correlations between EC_a and pH, Al, V and *m* soil attributes were not statistically significant. Analyzing the soil texture for Field 2 (different from Field 1), the EC_a had a significant positive correlation with clay ($r = 0.91$ for EC_a 20) and a negative correlation with sand ($r = -0.96$ for EC_a 20), as shown in Table 4. This behavior agrees with those found by Gholizadeh *et al.* (2012) and Souza *et al.* (2004), where the correlations were significant and positive between EC_a and clay content, respectively, and were significant and negative between EC_a and sand content, respectively, with $r = -0.437$ ($p < 0.01$). This occurs because soils with higher clay content present a greater soil water capacity, and consequently, higher electrical conductivity.

Other authors have also found a significant correlation between the apparent soil electrical conductivity and soil physical attributes (ISLAM *et al.*, 2012; LESCH; CORWIN; ROBINSON, 2005; MOLIN; CASTRO, 2008; RODRÍGUEZ-PÉREZ *et al.*, 2011), as was observed in the present work in Field 2 (Table

4). However, in other studies (AIMRUN *et al.*, 2007; MORARI; CASTRIGNANÒ; PAGLIARIN, 2009; SERRANO *et al.*, 2010; VALENTE *et al.*, 2012), there was no correlation or the correlation was low between EC_a and soil particle size, which occurred in Field 1 of this study (Table 4). The dynamics and the interaction among the soil attributes are decisive in the range of variation of the apparent soil electrical conductivity, which can cause positive and negative correlations between EC_a and the other soil attributes. In certain situations, the clay content variability amount is capable of causing high variation in the apparent soil electrical conductivity, which hides the effect of other soil attributes on EC_a. However, when the variability of the clay content is not high enough to cause variation in the apparent soil electrical conductivity, other soil attributes and the interaction between them are the cause of EC_a variability. In this case, the variation of the clay type or the soil chemical attribute variations modify the soil charges, and consequently, the electrical conductivity.

CONCLUSION

The results suggest that it is not feasible to use the apparent soil electrical conductivity to make generalized estimates of soil attributes. However, the apparent soil electrical conductivity can be applied as a tool for mapping the spatial variability of soil attributes. In other words, the EC_a can be used to separate different types of soils presented in a field. Mapping the electrical conductivity, while not a replacement for soil sampling, can be used to define management zones. Management zones could then be used to direct a soil sampling and the application of lime and fertilizers at variable rates.

ACKNOWLEDGMENTS

The authors would like to thank FAPEMIG (The Minas Gerais State Foundation for Research) and CNPq (The Brazilian National Counsel for Research and Development) for their financial support to perform this work.

REFERENCES

- AIMRUN, W. *et al.* Spatial variability of bulk soil electrical conductivity in a Malaysian paddy field: key to soil management. **Paddy Water Environment**, v. 5, n. 2, p. 113-121, 2007.
- ALCÂNTARA, G. R.; REIS, E. F.; QUEIROZ, D. M. Produtividade de culturas correlacionada com condutividade elétrica aparente de um solo sob plantio direto. **Revista Agrotecnologia**, v. 3, n. 2, p. 62-72, 2012.
- BREVIK, E. C.; FENTON, T. E.; LAZARI, A. Soil electrical conductivity as a function of soil water content and implications for soil mapping. **Precision Agriculture**, v. 7, n. 6, p. 393-404, 2006.
- CORWIN, D. L.; HENDRICKX, J. M. H. Electrical resistivity: Wenner array. In: SILVA, J. S. **Methods of soil analysis**: Part 4. Physical methods. Madison, USA: Soil Science Society of America, 2002. p. 1282-1287. (SSSA Book Series, n. 5).
- CORWIN, D. L.; LESCH, S. M. Apparent soil electrical conductivity measurements in agriculture. **Computers and Electronics in Agriculture**, v. 46, n. 1/3, p. 11-43, 2005.
- CORWIN, D. L.; LESCH, S. M. Application of soil electrical conductivity to precision agriculture: theory, principles, and guidelines. **Agronomy Journal**, v. 95, n. 3, p. 455-471, 2003.
- DONAGEMA, G. K. *et al.* **Manual de métodos de análise de solo**. 2. ed. Rio de Janeiro: Embrapa, 2011. 212 p.
- EKWUE, E. I.; BARTHOLOMEW, J. Electrical conductivity of some soils in Trinidad as affected by density, water and peat content. **Biosystems Engineering**, v. 108, n. 2, p. 95-103, 2010.
- FAROOQUE, A. A. *et al.* Delineating management zones for site specific fertilization in wild blueberry fields. **Applied Engineering in Agriculture**, v. 28, n. 1, p. 57-70, 2012.
- FAULIN, G. D. C.; MOLIN, J. P. Amplitude dos valores da umidade e sua influência na mensuração da condutividade elétrica do solo. In: CONGRESSO BRASILEIRO DE AGRICULTURA DE PRECISÃO, 2., 2006, São Pedro, SP. **Anais...** São Pedro, SP: ESALQ: USP, 2006. p. 1-9.
- FRIEDMAN S. P. Soil properties influencing apparent electrical conductivity: a review. **Computers and Electronics in Agriculture**, v. 46, n. 1/3, p. 45-70, 2005.
- GHOLIZADEH, A. *et al.* Relationship between apparent electrical conductivity and soil physical properties in a Malaysian paddy field. **Agronomy and Soil Science**, v. 58, n. 2, p. 155-168, 2012.
- HOAGLIN, D. C.; MOSTELLER, F.; TUKEY, J. W. **Análise exploratória de dados: técnicas robustas: um guia**. Lisboa: Edições Salamandra, 1992. 446 p.
- ISLAM, M. M. *et al.* Comparing apparent electrical conductivity measurements on a paddy field under flooded and drained conditions. **Precision Agriculture**, v. 13, n. 3, p. 384-392, 2012.
- JOHNSON C. K. *et al.* Site-specific management zones based upon soil electrical conductivity in a semiarid cropping system. **Agronomy Journal**, v. 95, n. 2, p. 303-315, 2003.
- LESCH, S. M.; CORWIN, D. L.; ROBINSON, D. A. Apparent soil electrical conductivity mapping as an agricultural management tool in arid zone soils. **Computer and Electronics in Agriculture**, v. 46, n. 1/3, p. 351-378, 2005.
- LIBARDI, P. L. **Dinâmica da água no solo**. São Paulo: Ed. USP, 2005. v. 61, 335 p.
- LIBARDI, P. L. *et al.* Variabilidade da umidade gravimétrica de um solo hidromórfico. **Revista Brasileira de Ciência do Solo**, v. 20, n. 1, p. 1-12, 1996.
- MOLIN, J. P.; CASTRO, C. N. Establishing management zones using soil electrical conductivity and other soil properties by the fuzzy clustering technique. **Scientia Agricola**, v. 65, n. 6, p. 567-573, 2008.
- MOLIN, J. P.; FAULIN, G. D. C. Spatial and temporal variability of soil electrical conductivity related to soil moisture. **Scientia Agricola**, v. 70, n. 1, p. 1-5, 2013.
- MORAL, F. J.; TERRÓN, J. M.; SILVA, J. R. M. D. Delineation of management zones using mobile measurements of soil apparent electrical conductivity and multivariate geostatistical techniques. **Soil and Tillage Research**, v. 106, n. 2, p. 335-343, 2010.
- MORARI, F.; CASTRIGNANÒ, A.; PAGLIARIN, C. Application of multivariate geostatistics in delineating management zones within a gravelly vineyard using geo-electrical sensors. **Computers and Electronics in Agriculture**, v. 68, n. 1, p. 97-107, 2009.
- RHOADES, J. D. Salinity: electrical conductivity and total dissolved solids. In: **Methods of soil analysis**: Part 3.

Chemical methods. Madison, USA: Soil Science Society of America, 1996. p. 417-435. (SSSA Book Series, n. 5).

RODRÍGUEZ-PÉREZ, J. R. *et al.* Using apparent soil electrical conductivity (EC_a) to characterize vineyard soils of high clay content. **Precision Agriculture**, v. 12, n. 6, p. 775-794, 2011.

SERRANO, J. M. *et al.* Medição e mapeamento da condutividade eléctrica aparente do solo em pastagens. **Revista de Ciências Agrárias**, v. 33, n. 2, p. 5-14, 2010.

SERRANO, J. M.; SHAHIDIAN, S.; SILVA, J. R. M. Apparent electrical conductivity in dry versus wet soil conditions in a shallow soil. **Precision Agriculture**, v. 14, n. 1, p. 99-114, 2013.

SOUZA, Z. M. *et al.* Variabilidade espacial da textura de um latossolo vermelho amarelo eutroférico sob cultivo de

cana-de-açúcar. **Engenharia Agrícola**, v. 24, n. 2, p. 309-319, 2004.

SUDDUTH, K. A. *et al.* Relating apparent electrical conductivity to soil properties across the north-central USA. **Computers and Electronics in Agriculture**, v. 46, n. 1/3, p. 263-283, 2005.

TERRÓN, J. M. *et al.* Soil apparent electrical conductivity and geographically weighted regression for mapping soil. **Precision Agriculture**, v. 12, n. 5, p. 750-761, 2011.

VALENTE, D. S. M. *et al.* The relationship between apparent soil electrical conductivity and soil properties. **Revista Ciência Agronômica**, v. 43, n. 4, p. 683-690, 2012.

WARRICK, A. W.; NIELSEN, D. R. Spatial variability of soil physical properties in the field. In: HILLEL, D. **Applications of soil physics**. New York: Academic Press, 1980. p. 319-344.