



Revista Caatinga

ISSN: 0100-316X

ISSN: 1983-2125

Universidade Federal Rural do Semi-Árido

BARBOSA, MARCELA LÚCIA; SILVA, THIERES GEORGE FREIRE DA; ZOLNIER, SERGIO; SILVA, SÉRVULO MERCIER SIQUEIRA E; STEIDLE NETO, ANTONIO JOSÉ  
THE INFLUENCE OF CLADODE MORPHOLOGY ON THE  
CANOPY FORMATION OF FORAGE CACTUS PLANTS<sup>1</sup>

Revista Caatinga, vol. 31, no. 1, January-March, 2018, pp. 180-190  
Universidade Federal Rural do Semi-Árido

DOI: 10.1590/1983-21252018v31n121rc

Available in: <http://www.redalyc.org/articulo.oa?id=237158225021>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in [redalyc.org](http://redalyc.org)

UAEM [redalyc.org](http://redalyc.org)

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

## THE INFLUENCE OF CLADODE MORPHOLOGY ON THE CANOPY FORMATION OF FORAGE CACTUS PLANTS<sup>1</sup>

MARCELA LÚCIA BARBOSA<sup>2</sup>, THIERES GEORGE FREIRE DA SILVA<sup>3\*</sup>, SERGIO ZOLNIER<sup>2</sup>, SÉRVULO MERCIER SIQUEIRA E SILVA<sup>4</sup>, ANTONIO JOSÉ STEIDLE NETO<sup>5</sup>

**ABSTRACT** – Here we aimed to evaluate the influence of cladode morphology on the canopy formation of forage cactus plants. The study was carried out in Serra Talhada, the State of Pernambuco, using the IPA Sertânia (IPA), Miúda (MIU), and Orelha de Elefante Mexicana (OEM) clones, which were submitted to three irrigation depths (2.5, 5.0, or 7.5 mm) and three irrigation intervals (7, 14, or 28 days) from March 2012 to August 2013. Cladode and plant canopy biometric data were obtained during the experimental period. We found that the characteristics of the second and third order cladodes contributed most to the canopy formation of clones of the genus *Nopalea* (IPA and MIU), whereas, for the genus *Opuntia* (OEM), the first and third order cladodes contributed most. Overall, we found that the influence of the cladode variables on the canopy growth of forage cactus is more associated with the characteristics of the genus than to the peculiarities of the clones.

**Keyword:** Statistical analysis. Clones. Biometrics data. *Nopalea* sp.. *Opuntia* sp..

## ASSOCIAÇÃO MORFOLÓGICA DE CLADÓDIOS COM A FORMAÇÃO DO DOSSEL DA PALMA FORRAGEIRA

**RESUMO** - O objetivo foi avaliar a influência das características morfológicas dos cladódios sobre a formação do dossel da palma forrageira. O estudo foi conduzido em Serra Talhada, PE, com os clones IPA Sertânia - IPA, Miúda - MIU e Orelha de Elefante Mexicana - OEM, os quais foram submetidos a três lâminas (2,5; 5,0; e 7,5 mm) e três intervalos de irrigação (7; 14 e 28 dias), no período compreendido entre março 2012 a agosto de 2013. Dados das dimensões dos cladódios e do dossel das plantas foram obtidos ao longo do tempo. A matriz de correlação de Pearson foi elaborada, e análises canônicas e de trilha foram aplicadas para avaliação da associação entre as variáveis. Verificou-se que, as características dos cladódios de 2ª e de 3ª ordens foram as que mais contribuíram na formação do dossel dos clones do gênero *Nopalea* (IPA Sertânia e Miúda), enquanto para a Orelha de Elefante Mexicana do gênero *Opuntia* foram as dos cladódios de primeira 1ª e 3ª ordens. Conclui-se que, a influência das variáveis dos cladódios sobre o crescimento do dossel da palma forrageira esteve mais associada às características do gênero, do que a peculiaridades dos clones.

**Palavras-chave:** Análise estatística. Clones. Dados biométricos. *Nopalea* sp.. *Opuntia* sp..

\*Corresponding author

<sup>1</sup>Received for publication in 06/08/2016; accepted in 03/15/2017.

Paper extracted from the master's dissertation of the first author.

<sup>2</sup>Department of Agricultural Engineering, Universidade Federal de Viçosa, Viçosa, MG, Brazil; marcelalucia.ufvpe@gmail.com, zolnier@ufv.br.

<sup>3</sup>Academic Unit of Serra Talhada, Universidade Federal Rural de Pernambuco, Serra Talhada, PE, Brazil; thieres\_freire@yahoo.com.br.

<sup>4</sup>Experimental Unit of Arcoverde, Instituto Agrônomo de Pernambuco, Arcoverde, PE, Brazil; servulo.mercier@yahoo.com.br.

<sup>5</sup>Universidade Federal de São João Del-Rei, Sete Lagoas, MG, Brazil; steidle@gmail.com.

## INTRODUCTION

As a photosynthetic structure, leaves play a primordial role in plant growth, being in harmony with the other organs and with the edaphoclimatic conditions. Leaves are mostly responsible for the interception of solar radiation, which will promote the metabolic processes of sugar formation and the accumulation of plant mass (GIBSON et al., 2011). The amount of radiation absorbed, as well as water and nutrient supplies, are productive indicators (WEERAKKODY; SURIYAGODA, 2015).

In addition to presenting various physiological adaptations, such as the nocturnal uptake of CO<sub>2</sub> for greater efficiency of water use, Cacti species, such as the cactus forage, display morphological adaptations in their photosynthetic structures. Their leaves are atrophied, with the appearance of thorns in their place, so that photosynthesis is carried out in structures called cladodes (PEÑA-VALDIVIA et al., 2008; SILVA et al., 2010; SILVA et al., 2014a).

Clones of cactus forage of the *Opuntia* and *Nopalea* genera are widely grown in the northeast of Brazil, especially Orelha de Elefante Mexicana, which belongs to the *Opuntia* genus, and Miúda and IPA Sertânia of genus *Nopalea*. These clones are widely used because of their resistance to Carmim cochineal, their major pest, as well as their use as an animal feed source, especially during the dry season (QUEIROZ et al., 2015; SILVA et al., 2015).

The distribution, size, shape, and number of cladodes of the different clones define the canopy architecture of the forage cactus plants (FLORES-HERNÁNDEZ et al., 2004; SILVA et al., 2010; PINHEIRO et al., 2014; SILVA et al., 2015), which will affect the photosynthetic capacity of the crop and, therefore, its yield. However, other factors such as irrigation management, fertilization, spacing, among others, climatic conditions, and growth habit can also influence the architecture of these plants (SILVA et al., 2014a; SILVA; SANTOS, 2006). Thus, the association between the growth of cladodes and the plants allows us to identify which characteristics will contribute most to the formation of the canopy of the forage cactus.

Canonical correlation analysis determines the linear relationship between groups or sets of variables, identifying the existence and the maximum correlation between different groups (BRUM et al., 2011). Track analysis complements this information, revealing whether the relationship between the variables is a cause or an effect or influenced by others, identifying which variables have the greatest weight and contribution. These approaches have been used when testing the interrelationships between characteristics of cladodes, plant, and forage cactus yields conducted under dryland conditions (PEÑA-VALDIVIA et al., 2008; SILVA et al., 2010; NEDER et al., 2013;

PINHEIRO et al., 2014; SILVA et al., 2015).

The objective of this study was to evaluate the influence of the morphological characteristics of the cladodes on the canopy formation of the forage cactus.

## MATERIAL AND METHODS

Clones of forage cactus were planted at the Instituto Agronômico de Pernambuco (7° 56' 20" S; 38° 17' 31" O; 498 m), in Serra Talhada, the state of Pernambuco in the Brazilian semi-arid region, in February 2010. The soil was classified as a yellow-red eutrophic Argisol with sandy loam texture. The spacing adopted in the experiment was 1.6 × 0.2 m, with rows in a counter line. During the first cycle, cultivation was conducted in dry conditions until March 2012, when the harvest was carried out, maintaining only the basal cladode. From this date, the experimental period of the present study was started, characterizing the second production cycle, which finished in August 2013, totaling 532 days.

The forage cactus was arranged in a random block design, in a factorial arrangement of 3 × 3 × 3 + 3 with split-split-plot, plus one control of each clone, and three replications, totaling 90 experimental split-plots. The plots were composed of three fixed water depths for water restoration in the soil (L1: 2.5 mm, L2: 5.0 mm and L3: 7.5 mm, plots), the subplots for three frequencies (F1: 7 days; F2: 14 days, and F3: 28 days, subplots), and the split-split plots by the IPA-Sertânia - IPA (*Napolea cochenillifera* (L.) Salm-Dick.), Miúda - MIU (*Napolea cochenillifera* (L.) Salm-Dick.) and Orelha de Elefante Mexicana - OEM (*Opuntia stricta* (Haw.) Haw.).

In total, the experiment presented 90 split-split-plots, each composed of 4 rows of 20 plants, totaling 80 plants, with an area of 25.6 m<sup>2</sup> and useful area of 11.52 m<sup>2</sup>. The useful area consisted of 32 plants located in the two central rows.

Clones of forage cactus received 756 (L7.5 F7), 672 (L5.0 F7), 622 (L7.5 F14), 586 (L2.5 F7), 579 (L5.0 F14), 555 (L7.5 F28), 536 (L2.5 F14), 535 (L5.0 F28), 514 (L2.5 F28) and 493 (Control) mm year<sup>-1</sup> of water.

Irrigation was carried out using a drip system (drippers spaced at 0.40 m). Crop treatments were carried out to eliminate weeds, such as herbicide application, as well as disease control, whenever necessary. Monthly fertilization was done with the application of 50 kg ha<sup>-1</sup> of NPK formulation 14-00-18, according to the recommendation of the Agronomic Institute of Pernambuco.

The structural characteristics of forage cactus clones were followed-up in 13 monitoring campaigns, according to days after cutting - DAC

(07/24/2012 - DAC 146, 08/22/2012 - DAC 175, 09/19/2012 - DAC 203, 10/27/2012 - DAC 241, 11/24/2012 - DAC 269, 12/22/2012 - DAC 297, 01/26/2013 - DAC 332, 02/23/2013 - DAC 360, 03/23/2013 - DAC 388, 04/27/2013 - DAC 423, 05/25/2013 - DAC 451, 07/06/2013 - DAC 493, 07/27/2013 - DAC 514).

In these campaigns, data on plant growth and cladodes were obtained, and measurements were always performed in the same plants. In the cladode, measurements were made up to the 4<sup>th</sup> order of appearance, from a representative branch of the plant. These measurements included the length of cladodes (BCL, CL1, CL2, CL3 and CL4, where B – base cladode and 1, 2, 3 and 4, orders of cladode emergence), width (BCW, CW1, CW2, CW3 and CW4), the perimeter (BCP, CP1, CP2, CP3 and CP4) and thickness (BCT, CT1, CT2, CT3 and CT4), which were measured with a tape measure and a plastic caliper. In addition, the area values of cladodes were calculated for all orders (BCA, CA1, CA2, CA3 and CA4), using statistical models adjusted by Silva et al. (2014a) for clones, according to data on length and width of cladodes.

The plant measurements consisted of structural characteristics obtained during the biometric monitoring campaigns. In this group, the height of the plant (HP, from the soil surface to the highest cladode) and the width of the plant (PW, at the widest ends) were recorded. These data were obtained using a tape measure, and the number of total cladodes in each plant (TNC) was counted. Finally, a classification was made according to the order of appearance of the cladodes, ranging from the first order (NC1, the first units to appear from the basal cladode) to the fourth order (NC2, NC3, NC4), based on the clone. The cladode area index (CAI) was determined by the ratio between the sum of the cladode areas and the plant spacing ( $1.6 \times 0.2$  m).

The experimental data were subdivided into two groups: A) structural characteristics of the plant: “Plant”; and, B) structural characteristics of the cladodes: “Cladodes”. The elements of the first group were considered as explanatory variables and the second as response variables.

The Plant group was represented by plant height and width (PH, PW); total number of cladodes (TNC); number of cladodes in order, ranging from 1 to 4 on the clone basis (NCOx) and the cladode area index (CAI), which was determined by the ratio between the sum of the area of cladodes and the planting spacing ( $1.6 \times 0.2$  m). The area of the cladodes (CA) was obtained using models proposed by Silva et al. (2014b), which were adjusted for the clones IPA Sertânia, Miúda and Orelha de Elefante Mexicana, based on data of length and width of the cladodes.

The Cladode group consisted of the following variables: cladode length (CL1, CL2, CL3 and CL4), cladode width (CW1, CW2, CW3 and CW4),

cladode thickness (CT1, CT2, CT3 and CT4), cladodes perimeter (CP1, CP2, CP3 and CP4), and cladodes area from first to fourth order (CA1, CA2, CA3 and CA4).

To perform the analyses, the cladodes up to the third order of appearance were included for IPA and OEM clones and up to the 4<sup>th</sup> order for the MIU clone. This procedure was necessary because the number of fourth-order cladodes for IPA and OEM and fifth order clones for MIU was reduced. Thus, to avoid misinterpretation of results, it was decided not to count these cladodes in small numbers.

Also, the grouped data for each clone were mean values of the plant growth and cladode variables, regardless of the water regimes (water depth  $\times$  frequency), that is, the average of each variable was derived from 30 observations, referring to the experimental split-plots with the same clone.

“Plant” (Response) and “Cladodes” (Explanatory) groups were interrelated to evaluate the individual contribution of the morphological characteristics of the cladodes on the formation of the vegetative canopy of the plants.

The experimental data were first submitted to the Lilliefors test to verify their normality at a significance level of 5%.

Then, the response variables were related to the explanatory variables by means of the Pearson Correlation Matrix. In this analysis, the existence, direction, and intensity of the linear relationship between the groups of variables were evaluated by means of Pearson’s linear correlation coefficient ( $r$ ), between -1 a +1. Values closer to 1 indicate the existence of a strong linear relationship, whereas closer to zero is the existence of a weak linear correlation between the variables. The relationship between explanatory variables and responses was obtained by the equation:

$$r_{(XY)} = \frac{\text{Cov}(X, Y)}{\sqrt{V(X) \cdot V(Y)}} = \frac{\sigma(X, Y)}{\sigma(X) \cdot \sigma(Y)} \quad (1)$$

where:  $r_{(XY)}$  is the Pearson correlation coefficient for the variables X (response) and Y (explanatory);  $\sigma(X, Y)$ , the covariance between the variables X and Y; and  $\sigma(X)$  and  $\sigma(Y)$  are the standard deviation of the variables X and Y, respectively.

The classification of the intermediate Pearson correlation coefficients was interpreted according to the one used by Thomaz et al. (2012), which comprised the following ranges: 0 to 0.19, “very weak”; 0.20 to 0.39, “weak”; 0.40 to 0.69, “moderate”; 0.70 to 0.89, “strong” and 0.90 to 1.00, “very strong”. In addition, the correlation signal was analyzed as positive (+1) or negative (-1) and the significance of the coefficients ( $p < 0.01$  and  $p < 0.05$ ) was analyzed by Student’s t-test.

For the multicollinearity analysis, only the data of the explanatory variables correlated with at least one of the response variables were used. In this analysis, a new data screening was performed,

discarding the variables that, within the same group, presented correlation with each other with strong multicollinearity ( $NC > 100$ ) (TOEBE; CARGNELUTTI FILHO, 2013). The presence of strong multicollinearity may lead to misinterpretations of the analyses, which may lead to inconsistent estimates of canonical correlation coefficients, as well as an overestimation of the effects of the explanatory variables on the response variables.

In the canonical correlations, only the variables with weak multicollinearity ( $NC < 100$ ) were used (TOEBE; CARGNELUTTI FILHO, 2013), in which the associations between groups of variables were evaluated, such that the linear correlation between these combinations was maximal. In this analysis, the canonical variables must be orthogonal, that is, linearly independent of each other. The canonical axes were set based on the number of variables of the smallest group, so that the canonical correlations between them were evaluated by means of the Chi-square test, at the level of 1% of probability. The following equation was applied for the determination of the canonical coefficients between the groups of variables (FERREIRA, 2011):

$$\rho_{(U,V)} = \text{Cov}(U, V) = \frac{X^T \sum_{12} Y}{\sqrt{X^T \sum_{11} X} \sqrt{Y^T \sum_{22} Y}} \quad (2)$$

where:  $X^T \sum_{12} Y$  is the covariance among the canonical variables U and V;  $\sqrt{X^T \sum_{11} X}$  is the standard deviation of the canonical variable U and  $\sqrt{Y^T \sum_{22} Y}$  is the standard deviation of the canonical variable V.

In the track analysis, the correlation coefficient was unfolded, allowing us to determine the degree of the effect of an explanatory variable on the response variable, by means of the track coefficient. In this analysis, the coefficient of partial correlation between two variables was calculated, disregarding the effect of the others, by means of the following relationship:

$$r_{(X_i Y, Z)} = \frac{r_{X_i} - r_{X_{iZ}} \cdot r_{YZ}}{\sqrt{(1 - r_{X_{iZ}}^2) \cdot (1 - r_{YZ}^2)}} \quad (3)$$

where i is a response variable,  $r_{X_{iZ}}$  is the correlation coefficient between a response variable and n<sup>th</sup> explanatory variables;  $r_{Y_{iZ}}$  is the correlation coefficient between the response variable and the n<sup>th</sup> explanatory variables. The significance of the partial correlation is identical to that used in the Pearson correlation.

All analyses were carried out in the statistical

software "GENES" (CRUZ, 2006).

## RESULTS AND DISCUSSION

For clones IPA Sertânia (IPA), Miúda (MIU) and Orelha de Elefante Mexicana (OEM), we found that, in the Pearson's correlation between growth characteristics of cladodes and plants, nearly all the characteristics of the cladodes contributed to the formation and architecture of the vegetative canopy (Table 1).

For the IPA clone, the characteristics that represented the plant were the width (PW) and the numbers of cladodes of 1<sup>st</sup> (NC1) and 3<sup>rd</sup> (NC3) orders, which correlated with base cladodes (CB) and 1<sup>st</sup>, 2<sup>nd</sup>, and 3<sup>rd</sup> cladodes (CC1, CC2, CC3, EC1, PC1, ACB), and provided coefficients classified as moderate to very strong (Table 1).

In the MIU, the representative characteristics of plant growth were PW, NC1, NC3, NC4 and cladode area index (CAI), which had a moderate to very strong correlation with the base cladodes and those of the 2<sup>nd</sup>, 3<sup>rd</sup> and 4<sup>th</sup> orders (ECB, EC3, PC2, PC3, AC4).

For OEM, the representative plant characteristics were height (PH), NC1, NC3, and CAI, which correlated with characteristics of base cladodes and those of the 1<sup>st</sup> and 3<sup>rd</sup> order (CC1, ECB, EC1, EC3, PC1, ACB), providing coefficients that were classified as moderate to very strong.

The canonical correlation analysis allowed us to identify the dependence between the group of structural characteristics of the cladodes and the group of the structural characteristics of the plants. For the IPA clone (Table 2), the first canonical axis (0.997\*\*) showed that the plants with the largest widths (PW) and the highest number of 1<sup>st</sup> (NC1) and 3<sup>rd</sup> (NC3) order cladodes were those with higher 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> order cladodes, in addition to the largest basal cladodes.

Canonical correlations were significant for the first and second pair of canonical variables of the MIU clone (Table 3). It was observed in the first canonical axis (0.998 \*\*) that plants with higher amounts of 1<sup>st</sup>, 3<sup>rd</sup> and 4<sup>th</sup> order cladodes and higher CAI were those with higher 2<sup>nd</sup>, 3<sup>rd</sup> and 4<sup>th</sup> cladode orders, as well as the greater thickness of the base and the third order cladodes. In the second canonical axis (0.984 \*), plants with a lower number of 4<sup>th</sup> order cladodes and with lower CAI were those that presented lower thicknesses of the basal and 3<sup>rd</sup> order cladodes, as well as a smaller area of the 4<sup>th</sup> order cladodes.

**Table 1.** Pearson correlation matrices among the variables of the response group “Plant” (structural characteristics of the plants) with the “Cladode” explanatory group (structural characteristics of the cladodes) of forage cactus clones (IPA Sertânia - IPU, Miúda - MIU and *Orelha de Elefante Mexicano* - OEM), under irrigated conditions, in a semi-arid environment.

		CL1	CL2	CL3	CT1	CP1	BCA
IPA	PW	0.613*	0.96**	0.894**	0.867**	0.837**	0.613*
	NC1	0.454	0.898**	0.611*	0.524	0.811**	0.304
	NC3	0.653*	0.803**	0.939**	0.864**	0.706**	0.647*
MIU		BCA	CT3	CP2	CP3	CA4	
	PW	0.507	0.765**	0.617*	0.703**	0.592*	
	NC1	0.201	0.566*	0.879**	0.656*	0.273	
	NC3	0.679*	0.942**	0.817**	0.975**	0.752**	
	NC4	0.775**	0.924**	0.654*	0.884**	0.813**	
	CAI	0.774**	0.964**	0.829**	0.913**	0.878**	
OEM		CLBCA1	BCT	CT1	CT3	CP1	BCA
	PH	0.889**	0.639*	0.744**	0.7**	0.724**	0.519
	NC1	-0.529	-0.245	-0.285	-0.868**	-0.437	-0.066
	NC3	0.635*	0.617*	0.775**	0.289	0.515	0.244
	IAC	0.953**	0.844**	0.913**	0.608*	0.847**	0.595*

Plant Group: PH – plant height, PW – plant width, CAI - cladode area index, NC1 first order cladode number, NC2 – second order cladode number, NC3 – third order cladode number, NC4 – fourth order cladode number.

Cladode number: BCL - basal cladode length, first order cladode length, CL1 – first order cladode length, CL2 – second order cladode length, CL3 – third order cladode length, CW1 - first order cladode width, BCW – basal cladode width, CT1 – first order cladode thickness, CT2 – second order cladode thickness, CT3 – third order cladode thickness, BCP – base cladode perimeter, CP1 – first order cladode perimeter, CP2 – second order cladode perimeter, CP3 – third order cladode perimeter, BCA – basal cladode area, CA4 – fourth order cladode area.

\*\*, \*Significant at 1 and 5%, respectively, by t-test.

**Table 2.** Canonical correlations and canonical pairs between the “Plant” group (structural characteristics of the plants) and the “Cladode” group (structural characteristics of cladodes) of the IPA Sertânia - IPA clone under irrigated conditions in a semi-arid environment.

Groups	Variables	Canonical factors		
		1°	2°	3°
I - Plant	PW	0.998	-0.041	0.039
	NC1	0.836	0.428	-0.343
	NC3	0.901	-0.346	-0.262
II - Cladode	CL1	0.612	-0.237	-0.280
	CL2	0.973	0.223	0.004
	CL3	0.887	-0.390	-0.190
	CT1	0.853	-0.391	0.251
	CP1	0.851	0.220	-0.114
	BCA	0.597	-0.420	0.216
Canonical correlation		0.997**	0.918	0.386
$\chi^2$		49	14	1
Degrees of freedom		18	10	4

Where: Plant group: PW – plant width, NC1 – first order cladode number, NC3 – third order cladode number; Cladode Group: CL1 – first order cladode length, CL2 – second order cladode length, CL3 – third order cladode length, CT1 – first order cladode thickness, CP1 – first order cladode perimeter and BCA – base cladode area. \*\*Significant at 1% by the Chi-square test.

For the OEM clone (Table 4), only the first canonical axis was significant (0.999 \*\*). In this axis, plants with higher CAI, higher heights (PH), and lower NC1 were those with the greatest length, perimeter, and thickness of base cladodes, as well as those of 1<sup>st</sup> and 3<sup>rd</sup> orders.

It was observed that the length of the third-order cladodes was the characteristic that most contributed to the increase in the number of cladodes of the same order, either by direct or indirect effect (via CC1, CC2, EC1, PC1, ACB).

**Table 3.** Canonical correlations and canonical pairs between the “Plant” group (structural characteristics of plants) and the “Cladode” group (structural characteristics of cladodes), of Miúda - MIU clone, under irrigated conditions, in a semi-arid environment.

Groups	Variables	Canonical factors				
		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	5 <sup>th</sup>
I - Plant	PW	0.672	-0.326	0.034	0.642	-0.169
	NC1	0.836	0.217	-0.365	0.115	0.328
	NC3	0.921	-0.338	0.158	0.103	0.041
	NC4	0.756	-0.538	0.197	0.214	0.231
	CAI	0.824	-0.535	-0.170	0.016	0.073
II - Cladode	BCT	0.444	-0.787	0.079	-0.126	0.401
	CT3	0.817	-0.557	0.006	0.151	-0.016
	PC2	0.933	-0.023	-0.358	-0.026	-0.023
	CP3	0.929	-0.323	0.139	-0.111	-0.034
	AC4	0.537	-0.820	-0.052	-0.135	-0.134
Canonical correlation		0.998**	0.984*	0.835	0.394	0.212
$\chi^2$		66	32	9	1	0
Degrees of freedom		25	16	9	4	1

Where: Plant group: PW – plant width, NC1 – first order cladode number, NC3 – third order cladode number, NC4 – fourth order cladode number, IAC – cladode area index; Cladode Group: ECB – basal cladode thickness, CT3 – third order cladode thickness, PC2 – second order cladode perimeter, PC3 – third order cladode perimeter, AC4 – fourth order cladode area. \*\*, \*Significant at 1 and 5%, respectively, by the Chi-square test.

**Table 4.** Canonical correlations and canonical pairs between the “Plant” group (structural characteristics of the plants) and the “Cladode” group (structural characteristics of the cladodes), of *Orelha de Elefante Mexicana* - OEM clone, under irrigated conditions, in a semi - arid environment.

Groups	Variables	Canonical factors			
		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>
I - Plant	PH	0.885	0.186	0.329	-0.273
	CN1	-0.730	0.621	-0.123	-0.257
	CN3	0.465	0.696	0.038	0.546
	CAI	0.869	0.475	-0.021	-0.137
II – Cladode	CT1	0.952	0.286	0.001	0.020
	BCT	0.737	0.439	-0.411	-0.014
	CT1	0.792	0.533	-0.088	0.236
	CT3	0.883	-0.309	0.059	0.320
	CP1	0.841	0.236	-0.327	-0.091
	BCA	0.472	0.309	-0.180	-0.698
Canonical correlation		0.999**	0.944	0.674	0.458
$\chi^2$		62	20	5	2
Degree of freedom		24	15	8	3

Where: Plant Group: PH – plant height, NC1 – number of first order cladodes, NC3 - number of third order cladodes, CAI – cladode area index; Cladode Group: CL1 – first order cladode length, EC1 – first order cladode thickness, EC3 – third order cladode thickness, PC1 – first order cladode perimeter, BCA – base area cladode. \*\*Significant at 1% by the Chi-square test.

In the unfolding of the correlation coefficients between characteristics of cladodes and plant for IPA clone (Table 5), it was verified that the largest plants (PW) and with the highest number of cladodes of the first order (CN1) were those that had higher second order cladodes. These characteristics excelled the others, with direct and indirect effects (via CL1,

CL3, CT1, CP1, BCA). It was observed that the length of the third order cladodes was the characteristic that most contributed to the increase in the number of cladodes of the same order, either by direct or indirect effect (via CL1, CL2, CT1, CP1, BCA).

**Table 5.** Pearson's correlation coefficient on direct and indirect effects among the variables of the response group "Plant" (structural characteristics of plants), with the variables of the explanatory group "Cladode" (structural characteristics of cladodes), IPA Sertânia clone - IPA, under irrigated conditions, in a semi-arid environment.

Variable	Effect	PW	CN1	CN3
CL1	CL1 direct effect	0.037	-	0.127
	via CL2 indirect effect	0.399	-	0.056
	via CL3 indirect effect	0.125	-	0.360
	via CT1 indirect effect	0.085	-	0.098
	via CP1 indirect effect	-0.043	-	0.012
	via BCA indirect effect	0.010	-	0.000
	Total	0.613	-	0.653
CL2	CL2 direct effect	0.699	0.981	0.098
	via CL1 indirect effect	0.021	-0.058	0.072
	via CL3 indirect effect	0.160	0.132	0.463
	via CT1 indirect effect	0.130	-0.304	0.150
	via CP1 indirect effect	-0.071	0.165	0.020
	via BCA indirect effect	0.021	-0.018	0.000
	Total	0.960	0.898	0.803
CL3	CL3 direct effect	0.209	0.172	0.603
	via CL1 indirect effect	0.022	-0.061	0.076
	via CL2 indirect effect	0.536	0.753	0.075
	via CT1 indirect effect	0.148	-0.347	0.171
	via CP1 indirect effect	-0.051	0.120	0.014
	via BCA indirect effect	0.031	-0.027	0.000
	Total	0.894	0.611	0.939
CT1	CT1 direct effect	0.174	-	0.201
	via CL1 indirect effect	0.018	-	0.062
	via CL2 indirect effect	0.522	-	0.073
	via CL3 indirect effect	0.177	-	0.513
	via CP1 indirect effect	-0.056	-	0.016
	via BCA indirect effect	0.032	-	0.000
	Total	0.867	-	0.864
CP1	CP1 direct effect	-0.080	0.186	0.022
	via CL1 indirect effect	0.020	-0.054	0.068
	via CL2 indirect effect	0.621	0.871	0.087
	via CL3 indirect effect	0.134	0.111	0.388
	via CT1 indirect effect	0.122	-0.285	0.141
	via BCA indirect effect	0.020	-0.017	0.000
	Total	0.837	0.811	0.706
BCA	BCA indirect effect	0.044	-	0.001
	via CL1 indirect effect	0.009	-	0.030
	via CL2 indirect effect	0.327	-	0.046
	via CL3 indirect effect	0.144	-	0.416
	via CT1 indirect effect	0.126	-	0.145
	via CP1 indirect effect	-0.036	-	0.010
	Total	0.613	-	0.647
Coefficient of determination		0.992	0.866	0.917

Plant Group: PW – plant width, NC1 - first order cladode number, NC3 – third order cladode number; Cladode Group: CL1 - first order cladode length, CL2 - second order cladode length, CL3 - third order cladode length, CT1 - first order cladode thickness, CP1 - first order cladode perimeter, BCA – basal cladode area. “-” indicates that the variable did not present any correlation of plant group variables.

Similar to IPA, in the MIU clone (Table 6) the characteristics of the 2<sup>nd</sup> and 3<sup>rd</sup> order cladodes were those that most explained the growth and structure of the plant. The perimeter and the thickness of the 3<sup>rd</sup> order cladodes were those that most contributed to the highest PW and the highest number of cladodes of the 3<sup>rd</sup> and 4<sup>th</sup> orders, either directly or indirectly through

other variables (BCT, CP2, CA4). Similarly, the greater the growth of the second-order cladodes (CP2), the greater the stimulus for the development of new first order cladodes (NC1). In addition, it was found that greater thicknesses of the 3<sup>rd</sup> order cladodes provided a greater increase of the CAI.



**Table 6.** Pearson's correlation coefficient unfolding on direct and indirect effects among the variables of the response group "Plant" (structural characteristics of the plants), with the variables of the "Cladode" explanatory group (structural characteristics of the cladodes), of the clone Miúda - MIU, under irrigated conditions, in a semi-arid environment.

Variable	Effect	PW	NC1	NC3	NC4	CAI
BCT	BCT direct effect	-	-	0.014	0.123	0.150
	via CT3 indirect effect	-	-	0.430	0.605	0.313
	via CP2 indirect effect	-	-	-0.020	-0.097	0.151
	via CP3 indirect effect	-	-	0.470	0.270	-0.040
	via CA4 indirect effect	-	-	-0.215	-0.127	0.200
	Total	-	-	0.679	0.775	0.774
CT3	CT3 direct effect	1.242	0.142	0.554	0.779	0.403
	via BCT indirect effect	-0.112	0.060	0.011	0.095	0.117
	via CP2 indirect effect	-0.056	0.739	-0.038	-0.186	0.292
	via CP3 indirect effect	-0.059	0.042	0.639	0.368	-0.055
	via CA4 indirect effect	-0.249	-0.417	-0.224	-0.132	0.208
	Total	0.765	0.566	0.942	0.924	0.964
CP2	CP2 Direct effect	-0.072	0.961	-0.050	-0.242	0.380
	via BCT indirect effect	-0.058	0.031	0.006	0.049	0.060
	via CT3 indirect effect	0.955	0.109	0.426	0.599	0.310
	via CP3 indirect effect	-0.053	0.038	0.573	0.330	-0.049
	via CA4 indirect effect	-0.155	-0.260	-0.139	-0.082	0.129
	Total	0.617	0.879	0.817	0.654	0.829
CP3	CP3 direct effect	-0.064	0.045	0.693	0.399	-0.060
	via BCT indirect effect	-0.098	0.053	0.010	0.083	0.102
	via CT3 indirect effect	1.146	0.131	0.512	0.719	0.372
	via CP2 indirect effect	-0.060	0.796	-0.041	-0.201	0.314
	via CA4 indirect effect	-0.221	-0.369	-0.198	-0.116	0.184
	Total	0.703	0.656	0.975	0.884	0.913
CA4	CA4 Direct effect	-0.285	-	-0.255	-0.150	0.237
	via BCT indirect effect	-0.122	-	0.012	0.104	0.127
	via CT3 indirect effect	1.088	-	0.486	0.683	0.353
	via CP2 indirect effect	-0.039	-	-0.027	-0.132	0.207
	via CP3 indirect effect	-0.049	-	0.537	0.309	-0.046
	Total	0.592	-	0.752	0.813	0.878
Coefficient of determination		0.619	0.841	0.974	0.887	0.974

Where: Plant group: PH – plant height, PT – plant thickness, NC1 - first order cladode number, NC3 – third order cladode number, NC4 – fourth order cladode number, CAI – cladode area index; Cladode group: BCT – basal cladode thickness, CT3 - third order cladode thickness, CP2 – second order cladode perimeter, CP3 – third order cladode perimeter, CA4 – fourth order cladode area “-” indicates that the variable did not present any correlation with one of the variables of the plant group.

The morphogenesis of the cladodes affected canopy formation of the plants, and for the MIU and IPA clones of *Nopalea* genus, the growth of the 2<sup>nd</sup> and 3<sup>rd</sup> order cladodes subsidized plant growth (canopy width, increase of CAI and appearance of

new cladodes of lower or higher orders). This finding can be explained by the location of these cladodes (2<sup>nd</sup> and 3<sup>rd</sup> orders) in the upper part of the plant. Thus, incident solar radiation on them is higher than the incident on lower order cladodes, which are

partially or completely shaded.

According to Nobel (2001), young cladodes display greater stimulus for photosynthesis than those of lower orders, contributing more to plant growth, as well as to the appearance of new cladodes. Acevedo, Badilla, and Nobel (1983)

reported that this occurs because there are differences in the pattern of solar radiation interception by the distinct cladodes over the day, as well as by their side of exposure in relation to the apparent movement of the sun.

**Table 7.** Pearson's correlation coefficient unfolding of direct and indirect effects among the variables of the response group "Plant" (structural characteristics of plants), with the variables of the explanatory group "Cladode" (structural characteristics of cladodes), clone *Orelha de Elefante Mexicana* – OEM under irrigated conditions, in a semi-arid environment.

Variable	Effect	PH	NC1	NC3	CAI
CL1	CL1 direct effect	1.462	-	0.441	0.871
	via ECB	-0.390	-	-0.088	-0.054
	via EC1 indirect effect	-0.113	-	0.899	0.243
	via EC3 indirect effect	-0.012	-	-0.355	-0.127
	via PC1 indirect effect	-0.155	-	-0.169	-0.045
	via BCA indirect effect	0.098	-	-0.093	0.066
	Total	0.889	-	0.635	0.953
BCT	BCT direct effect	-0.456	-	-0.103	-0.064
	via CL1 indirect effect	1.252	-	0.378	0.745
	via CT1 indirect effect	-0.102	-	0.815	0.220
	via CT3 indirect effect	-0.008	-	-0.228	-0.081
	via CP1 indirect effect	-0.142	-	-0.154	-0.041
	via BCA indirect effect	0.095	-	-0.090	0.064
	Total	0.639	-	0.617	0.844
EC1	CT1 direct effect	-0.126	-	1.006	0.272
	via CL1 indirect effect	1.306	-	0.394	0.778
	via BCT indirect effect	-0.369	-	-0.083	-0.051
	via CT3 indirect effect	-0.010	-	-0.292	-0.104
	via CP1 indirect effect	-0.149	-	-0.162	-0.043
	via BCA indirect effect	0.092	-	-0.088	0.062
	Total	0.744	-	0.775	0.913
EC3	CT3 direct effect	-0.016	-0.987	-	-0.169
	via CL1 indirect effect	1.098	-0.225	-	0.654
	via BCT indirect effect	-0.220	0.057	-	-0.031
	via CT1 indirect effect	-0.078	0.443	-	0.168
	via CP1 indirect effect	-0.109	-0.153	-	-0.031
	via BCA indirect effect	0.025	-0.002	-	0.017
	Total	0.700	-0.868	-	0.608
PC1	CP1 direct effect PC1	-0.183	-	-	-0.053
	via CL1 indirect effect	1.245	-	-	0.741
	via BCT indirect effect	-0.354	-	-	-0.049
	via CT1 indirect effect	-0.103	-	-	0.221
	via CT3 indirect effect	-0.010	-	-	-0.100
	via BCA indirect effect	0.128	-	-	0.086
	Total	0.724	-	-	0.847
BCA	BCA direct effect ACB	-	-	-	0.137
	via CC1 indirect effect	-	-	-	0.417
	via ECB indirect effect	-	-	-	-0.030
	via EC1 indirect effect	-	-	-	0.123
	via EC3 indirect effect	-	-	-	-0.021
	via PC1 indirect effect	-	-	-	-0.033
	Total	-	-	-	0.595
Coefficient of determination		0.877	0.897	0.711	0.959

Where: Plant Group: PH – plant height, NC1 – first order cladode number, NC3 – third order cladode number, IAC – cladode area index; Cladode group: CC1 – first order cladode number, ECB – basal cladode thickness, EC1 – first order cladode thickness, EC3 – third order cladode thickness, PC1 – first order cladode perimeter, ACB – basal area cladode. “-” indicates that the variable did not present any correlation with one of the variables of the plant group.

Silva et al. (2015), in a study with clones IPA and Miúda, with data obtained at 754 days after cut, showed that 2<sup>nd</sup> and 3<sup>rd</sup> order cladodes were the largest for these clones. Cunha et al. (2012) reported that there is a correlation between the rate of appearance of new cladodes and the number of cladodes present in the plant because each unit that emerges represents an additional potential for the appearance of new cladodes. These new structural units will perform more photosynthesis and will produce more photoassimilates, which will stimulate plant growth.

Using four years and eight months old MIU clones, Pinheiro et al. (2014) studied the contribution of the growth characteristics of cladodes and plants to the cladode area index, observing the existence of a significant correlation with the height  $\times$  width product and the characteristics of 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> and 4<sup>th</sup> order cladodes. However, the authors pointed out that the characteristics of the fourth-order cladodes were the ones that contributed the most to the IAC clone. The other correlations were of low magnitude.

For the OEM clone (Table 7), higher lengths and thicknesses of the 1<sup>st</sup> order cladodes (CL1 and CT1) contributed to the occurrence of larger plants, with a higher number of 3<sup>rd</sup> order cladodes and higher CAI values, which were verified by means of direct or indirect effects (via BCT, CT3, CP1, BCA). In turn, the lower growth of the 3<sup>rd</sup> order cladodes (CT3) stimulated the development of new 1<sup>st</sup> order cladodes.

The major contribution of the first order cladodes to the OEM clone is because these are more numerous than those of the other orders. Clones of the genus *Opuntia* exhibit fewer cladodes when compared to clones of the genus *Nopalea*, but their cladodes are larger (CAVALCANTE et al., 2014) and play an important role in the growth and storage of water and plant mass.

Silva et al. (2015) report that first order cladodes are the largest in the OEM clone, as the second-order cladodes. The authors highlight that, similarly to the IPA clone; the OEM clone has larger perimeters of basal and first order cladodes when compared to the Miúda clone.

When evaluating the contribution of the morphological characteristics of plants and cladodes to the increase of the cladode area index of the Orelha de Elefante Mexicana clone, Pinheiro et al. (2014) found that the morphological characteristics explained 63.69% of the variation of the cladode area index. The variable that had the greatest weight in this percentage was the total number of cladodes of the plants.

Silva et al. (2010) emphasize that the study of morphological characteristics of clones is important since they will determine the growth habit of the latter, which can be erect or half-open. This habit is directly related to the interception of solar radiation by plants, determined by the architecture of its

canopy.

## CONCLUSIONS

The relationships between the characteristics of the cladodes and the forage cactus plant were more associated with genus peculiarities than with the different clones.

The growth of the 2<sup>nd</sup> and 3<sup>rd</sup> order cladodes was the characteristic that most contributed to the canopy formation of clones of plants of the genus *Nopalea* (IPA and Miúda), whereas for *Orelha de Elefante Mexicana* of the genus *Opuntia*, it was the first and third order cladodes that most contributed.

## ACKNOWLEDGEMENTS

To FACEPE for the financial support (APQ-0215-5.01/10) and to CAPES for scholarship granting.

## REFERENCES

- ACEVEDO, E.; BADILLA, I.; NOBEL, P. S. Water relations, diurnal acidity changes, and productivity of a cultivated cactus, *Opuntia ficus-indica*. **Plant Physiology**, Amsterdam, v. 72, n. 3, p. 775-780, 1983.
- BRUM, B. et al. Correlações canônicas entre variáveis de semente, plântula, planta e produção de grãos em mamoneira. **Ciência Rural**, Santa Maria, v. 41, n. 3, p. 404-411, 2011.
- CAVALCANTE, L. A. D. et al. Respostas de genótipos de palma forrageira a diferentes densidades de cultivo. **Pesquisa Agropecuária Tropical**, Goiânia, v. 44, n. 4, p. 424-433, 2014.
- CRUZ, C. D. **Programa Genes: biometria**. 1. ed. Viçosa, MG: Editora UFV, 2006. 382 p.
- CUNHA, D. N. F. V. et al. Morfometria e acúmulo de biomassa em palma forrageira sob doses de nitrogênio. **Revista Brasileira de Saúde e Produção Animal**, Salvador, v. 13, n. 4, p. 1156-1165, 2012.
- FERREIRA, D. F. **Estatística Multivariada**. 2. ed. Lavras, MG: Editora UFLA, 2011. 675 p.
- FLORES-HERNÁNDEZ, A. et al. Yield and physiological traits of prickly pear cactus 'nopal' (*Opuntia* spp.) cultivars under drip irrigation. **Agricultural Water Management**, Amsterdam, v. 70, n. 1, p. 97-107, 2004.

- GIBSON, et al. Exploiting leaf starch synthesis as a transient sink to elevate photosynthesis, plant productivity and yields. **Plant science**, Amsterdam, v. 181, n. 3, p. 275-281, 2011.
- NEDER, D. G. et al. Correlations and path analysis of morphological and yield traits of cactus pear accessions. **Crop Breeding and Applied Biotechnology**, Amsterdam, v. 13, n. 3, p. 203-207, 2013.
- NOBEL, P. S. Ecophysiology of *Opuntia ficus-indica*. In: MONDRAGÓN-JACOBO, C.; PÉREZ-GONZÁLEZ, S. (Eds.). **Cactus (Opuntia spp.) as forage**. Rome: Food and Agriculture Organization of the United Nations, 2001. p. 13-20. (FAO Plant production and protection paper, 169).
- PEÑA-VALDIVIA, C. B. et al. Morphological characterization of *Opuntia* spp.: a multivariate analysis. **Journal of the Professional Association for Cactus Development**, Chapingo, v. 10, n. 1, p. 1-21, 2008.
- PINHEIRO, K. M. et al. Correlações do índice de área do cladódio com características morfogênicas e produtivas da palma forrageira. **Pesquisa Agropecuária Brasileira**, Brasília, v. 49, n. 12, p. 939-947, 2014.
- QUEIROZ, M. G. et al. Características morfofisiológicas e produtividade da palma forrageira sob diferentes lâminas de irrigação. **Revista Brasileira de Engenharia Agrícola e Ambiental**, Campina Grande, v. 19, n. 10, p. 931-938, 2015.
- SILVA, C. C. F.; SANTOS, L. C. Palma Forrageira (*Opuntia ficus-indica* Mill) como alternativa na alimentação de ruminantes. **Revista Electrónica de Veterinária**, Málaga, v. 7, n. 10, p. 1-13, 2006.
- SILVA, N. G. M. et al. Relação entre características morfológicas e produtivas de clones de palma-forrageira. **Revista Brasileira de Zootecnia**, Viçosa, v. 39, n. 11, p. 2389-2397, 2010.
- SILVA, T. G. F. et al. Indicadores de eficiência do uso da água e de nutrientes de clones de palma forrageira em condições de sequeiro no Semiárido brasileiro. **Bragantia**, Campinas, v. 73, n. 2, p. 184-191, 2014a.
- SILVA, T. G. F. et al. Área do cladódio de clones de palma forrageira: modelagem, análise e aplicabilidade. **Revista Brasileira de Ciências Agrárias**, Recife, v. 9, n. 4, p. 633-641, 2014b.
- SILVA, T. G. F. et al. Crescimento e produtividade de clones de palma forrageira no semiárido e relações com variáveis meteorológicas. **Revista Caatinga**, Mossoró, v. 28, n. 2, p. 10-18, 2015.
- THOMAZ, G. L. et al. Produção do girassol e teor de óleo nos aquênios em função da temperatura do ar, precipitação pluvial e radiação solar. **Ciência Rural**, Santa Maria, v. 42, n. 8, p. 1380-1385, 2012.
- TOEBE, M.; CARGNELUTTI FILHO, A. Não normalidade multivariada e multicolinearidade na análise de trilha em milho. **Pesquisa Agropecuária Brasileira**, Brasília, v. 48, n. 5, p. 466-477, 2013.
- WEERAKKODY, W. A. P.; SURIYAGODA, L. D. B. Estimation of leaf and canopy photosynthesis of pot chrysanthemum and its implication on intensive canopy management. **Scientia Horticulturae**, Amsterdam, v. 192, n. 1, p. 237-243, 2015.