



Pesquisa Agropecuária Tropical

ISSN: 1517-6398

ISSN: 1983-4063

Escola de Agronomia/UFG

Lourenço, Andressa Mirelle Santos; Santos, Carlos
Antonio Fernandes; Silva, Danilo Olegário Matos da
Compatibility and phenotypic variability of guava accessions grafted
on BRS Guaraçá rootstock: vegetative and fruit production traits¹
Pesquisa Agropecuária Tropical, vol. 52, e71259, 2022
Escola de Agronomia/UFG

DOI: <https://doi.org/10.7440/res64.2018.03>

Available in: <https://www.redalyc.org/articulo.oa?id=253070366014>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UFGM  redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Compatibility and phenotypic variability of guava accessions grafted on BRS Guaraçá rootstock: vegetative and fruit production traits¹

Andressa Mirelle Santos Lourenço², Carlos Antonio Fernandes Santos³, Danillo Olegário Matos da Silva⁴

ABSTRACT

One of the main problems faced by guava production is the decline caused by the root-knot nematode (*Meloidogyne enterolobii*), which has a wide distribution in Brazil. This study aimed to evaluate the productive and morphological compatibility and variability of guava accessions grafted on BRS Guaraçá, a rootstock with resistance to the root-knot nematode. The experiment was conducted with 83 accessions from the *Psidium* Germplasm Bank (Petrolina, Pernambuco State, Brazil). The evaluated traits were stem diameter, canopy diameter I and II, plant height, total number of fruits, total fruit weight, mean fruit weight, seed weight, pulp weight, fruit width and length. Except for the stem diameter ratio, all variables showed significant differences ($p < 0.05$), indicating phenotypic variability. The evaluated accessions showed no incompatibility symptoms, e.g., cracks, exudations and differences for stem diameter in the grafted area, indicating high compatibility. The accessions formed vigorous plants in the second harvest cycle, with mean canopy diameter of 2.3 m and mean plant height of 2.2 m, also indicating high compatibility. The accessions Gua64BA, Gua181ES, Gua06MA, Pedro Sato and Gua99AM showed the highest values for pulp and fruit weight (around 160 g), whereas Gua96AM, Gua70RO, Gua192ES, Gua01MA and Gua183ES showed the highest fruit production values (around 20 kg plant⁻¹), being recommended for grafting in BRS Guaraçá or as potential parents for population development.

KEYWORDS: *Psidium guajava*, *Meloidogyne enterolobii*, nematode infection.

RESUMO

Compatibilidade e variabilidade fenotípica de acessos de goiabeira enxertados em porta-enxerto BRS Guaraçá: características vegetativas e de produção de frutos

Um dos principais problemas enfrentados na produção de goiaba é o declínio causado pelo nematoide-das-galhas (*Meloidogyne enterolobii*), que possui ampla distribuição no Brasil. Objetivou-se avaliar a compatibilidade e variabilidade produtiva e morfológica de acessos de goiabeira enxertados em BRS Guaraçá, porta-enxerto com resistência ao nematoide-das-galhas. O experimento foi conduzido com 83 acessos do Banco de Germoplasma de *Psidium* (Petrolina, PE). As características avaliadas foram diâmetro do caule, diâmetro da copa I e II, altura da planta, número total de frutos, peso total do fruto, peso médio do fruto, peso da semente, peso da polpa, largura e comprimento do fruto. Com exceção da razão diâmetro do caule, todas as variáveis apresentaram diferenças significativas ($p < 0,05$), indicando variabilidade fenotípica. Os acessos avaliados não apresentaram sintomas de incompatibilidade, como rachaduras, exsudações e diferenças no diâmetro do caule na área enxertada, indicando alta compatibilidade. Os acessos formaram plantas vigorosas no segundo ciclo de colheita, com diâmetro médio de copa de 2,3 m e altura média de planta de 2,2 m, indicando, também, alta compatibilidade. Os acessos Gua64BA, Gua181ES, Gua06MA, Pedro Sato e Gua99AM apresentaram os maiores valores de polpa e peso de frutos (cerca de 160 g), enquanto Gua96AM, Gua70RO, Gua192ES, Gua01MA e Gua183ES apresentaram os maiores valores de produção de frutos (em torno de 20 kg planta⁻¹), sendo recomendados para enxertia em BRS Guaraçá ou como potenciais genitores para o desenvolvimento populacional.

PALAVRAS-CHAVE: *Psidium guajava*, *Meloidogyne enterolobii*, infecção nematoide.

INTRODUCTION

Guava (*Psidium guajava*) is an important tropical fruit species due to the nutritional quality, aroma and flavor of its fruits (Martins et al. 2020).

This crop contributes significantly to the Brazilian fruit sector, with a total production of 584,223 tons in 2019 that achieved a revenue value close to US\$ 200 million, with the Southeast and Northeast regions of Brazil accounting for 90 % of the national

¹ Received: Dec. 17, 2021. Accepted: Apr. 13, 2022. Published: May 02, 2022. DOI: 10.1590/1983-40632022v5271259.

² Universidade Estadual de Feira de Santana, Feira de Santana, BA, Brasil.

E-mail/ORCID: andressamlourenco@hotmail.com/0000-0001-7692-5307.

³ Empresa Brasileira de Pesquisa Agropecuária (Embrapa Semiárido), Petrolina, PE, Brasil.

E-mail/ORCID: carlos-fernandes.santos@embrapa.br/0000-0002-6932-6805.

⁴ Centro Universitário Maurício de Nassau, Petrolina, PE, Brasil. E-mail/ORCID: danilloolegario@hotmail.com/0000-0002-4874-4880.

guava production (IBGE 2020). Guava fruits may be consumed fresh (*in natura*) or used to produce industrialized products, e.g., sweets, jams and preserves (Cañizares et al. 2003). Moreover, these fruits have significant contents of vitamin C, vitamin A and B-complex vitamins, lycopene and β -carotene (Manica et al. 2000).

The main problems faced by the Brazilian guava production is the disease known as guava decline, initially caused by the root-knot nematode (*Meloidogyne enterolobii*), with an economic impact estimated at US\$ 66 million, in 2008 (Pereira et al. 2009). This complex disease increases the plant sensitivity to environmental stresses and causes the progressive rotting of affected roots, finally resulting in the plant death (Gomes et al. 2010). Early studies aimed at controlling this disease with nematicides, biological control, crop rotation and strawberry guava rootstocks did not show satisfactory results, either due to low control efficacy or incompatibility with grafted guava cultivars (Freitas 2012). Robaina et al. (2015) used *P. cattleyanum* as a rootstock for the guava cultivar Paluma and observed low seedling vigor and incompatibility symptoms, highlighting the limitations of this strategy.

Other research results indicate the BRS Guaraçá hybrid from *P. guajava* \times *P. guineense* as a nematode-resistant rootstock compatible with the guava cultivars Paluma and Pedro Sato, producing 40 t ha⁻¹ year⁻¹ (Souza et al. 2018). However, broader studies evaluating the BRS Guaraçá rootstock with a larger number of guava accessions, as well as other *Psidium* species, have not yet been conducted. Misra (2017) reported an F1 population of *P. molle* \times *P. guajava* with resistance to wilt (*Fusarium oxysporum* and *F. solani*) and serving as a compatible rootstock for any guava variety. Some studies conducted with different species also reported compatibility between rootstock and scions. According to Bowmana & Joubert (2020), most citrus rootstocks are graft-compatible with most citrus scions, with few exceptions. In another study, Gonçalves et al. (2019) evaluated the compatibility of the peach cultivar BRS Libra with 23 clonal rootstocks and identified compatible and incompatible rootstocks. Furthermore, Vršič et al. (2015) evaluated the compatibility of different rootstocks with grapevine varieties, with one of them showing high compatibility with 12 rootstocks.

For Hartmann et al. (1997), graft compatibility is related to the satisfactory union between rootstock and scion and good plant development after grafting. According to Melnyk (2016), a successful grafting begins with the effective adhesion between rootstock and scion, followed by the formation of callus tissue at the graft interface, and ends with the establishment of vascular connections followed by the distribution of nutrients and water throughout the plant body. Conversely, incompatibility may result in increased diameter of the rootstock-scion union area, graft breakage or rupture, premature plant death, leaf yellowing or premature leaf fall (Fachinello et al. 1995). According to He et al. (2018), information on graft incompatibility mechanisms is still limited, and several studies have found that hormones, especially auxin, play significant roles in regulating the development and metabolic traits of higher plants. According to these authors, the content of indole-3-acetic acid (IAA) also increases with compatible graft combination, suggesting that IAA might promote graft compatibility.

In this scenario, the present study evaluated the productive and morphological compatibility and variability of 83 *P. guajava* accessions grafted on BRS Guaraçá, a root-knot nematode-resistant rootstock, aiming to identify limitations and accessions with greater agronomic potential.

MATERIAL AND METHODS

The experiment was conducted with 83 accessions from the *Psidium* Active Germplasm Bank (BAG *Psidium*), located at the Embrapa Semiárido, in Petrolina, Pernambuco State, Brazil (09°08'12.3"S, 40°18'31.6"W and 363 m above the sea level). The guava accessions were collected from different regions of the Brazilian States of Amazonas, Bahia, Espírito Santo, Goiás, Maranhão, Minas Gerais, Pernambuco, Piauí, Roraima, Rio Grande do Sul, Sergipe and Rondônia. The two final letters in the name of the accession identify the State where the material was collected. The soil of the experimental area is an Argissolo Vermelho-Amarelo Distrófico plúntico (Santos et al. 2018) or Ultisol (USDA 2014). The mean temperature is 26 °C, with a hot and dry climate classified as BSh, according to Köppen, and the mean rainfall is 496 mm (Alvares et al. 2013). The transplanting of all plants occurred from April to May of 2018.

The experimental design was randomized blocks, with two replications and two plants per plot, spaced at 4 x 4 m. The root-knot nematode-resistant guava cultivar BRS Guaraçá was used as a rootstock. The rootstock cuttings were propagated in a nursery with suitable conditions for rooting. Seedlings of the BAG *Psidium* accessions, used as scions, were also propagated by seeds in a plant nursery, transplanted to plastic bags and kept under controlled irrigation conditions. The guava accessions were then wedge-grafted in the field at four months after transplanting, and irrigation was performed with a micro-sprinkler system. The area was sprayed to control pests and diseases, and weed control was performed by manual and mechanical hoeing. Finally, the plants were subjected to formative pruning, and the first fruit production pruning was performed in August 2019, with a new production pruning occurring in November 2019.

The evaluation cycle of the 83 guava accessions began in November 2020, by considering the following vegetative traits: plant height (m); stem diameter ratio at 15 cm below and 15 cm above the grafting point; canopy diameter I and II (m). The evaluated production traits were: total number of fruits; mean fruit weight (g); total fruit weight (kg); seed weight (g); pulp weight (g); fruit width and length (mm). Additionally, the accessions were visually evaluated for the presence of cracks and resins in the grafted area of the rootstock.

The data of all variables were subjected to analysis of variance, and the means were compared using the Student-Newman-Keuls test. These analyses were performed using the GLM procedure of the SAS software. The pulp weight and total fruit weight variables were transformed to square root, in order to meet the assumptions of normality. Finally, histograms were constructed using a Microsoft Excel spreadsheet to evaluate the dispersion of the measured traits.

The adopted statistical model used within-plot data, as described by Santos et al. (1995): $Y_{ijk} = \mu + a_i + b_j + e_{ij} + \pi_{ijk}$, where: Y_{ijk} is the value observed in the k-th plant, in the j-th replicate of the i-th accession; μ the overall mean; a_i the effect of the i-th accession, associated with the expected mean square $[E(QM)] \sigma_A^2$, genetic and environmental variance; b_j the effect of block j; e_{ij} the effect of the j-th replicate of the i-th accession, associated with $E(QM) \sigma_E^2$, environmental variance; π_{ijk} the effect

of the k-th plant, in the j-th replicate of the i-th accession, associated with $E(QM) \sigma_A^2$, genetic and environmental variance.

RESULTS AND DISCUSSION

The analyzed variables showed significant statistical differences among the 83 accessions, indicating phenotypic variability, except for the stem diameter ratio at 15 cm below and 15 cm above the grafting point. The statistical significance was $p < 0.01$ for the analyzed variables, except for canopy diameter II ($p < 0.05$). The coefficients of variation ranged from 8.0 to 46.6 % for the analyzed traits, indicating good to limited experimental accuracy in the data measurement (Table 1).

For the studied variables, the variance between plants showed no significant differences by the F-test, indicating no phenotypic variability between plants of a given accession (Table 1). For Pommer & Murakami (2009), guava has a cross-pollination rate of up to 41.3 %, resulting in high phenotypic variability when propagated by seeds. This variability was not observed in the present study, probably due to the low number of plants (four) evaluated per accession.

The grafting success rate was approximately 100 %, although some accessions required re-grafting. This high grafting success rate may be attributed to the fact that the BRS Guaraçá has 50 % of the genome from a guava mother plant, the GUA161PE.

The stem diameter ratio at 15 cm below and 15 cm above the grafting point ranged from 0.73 to 1.14 (Table 1), with most accessions showing values from 0.8 to 1.1, two accessions showing values < 0.8 , and other five accessions with values > 1.1 (Figure 1A), indicating good compatibility in the grafted area for 95 % of the accessions grafted on the BRS Guaraçá rootstock. There were no visual signs of cracks or resins in the grafted area, especially in the accessions with the lowest and highest stem diameter ratio at 15 cm below and 15 cm above the grafting point values (Figure 2). Moreover, the stem in the grafted area had an almost continuous color and brownish spots typical of the *Psidium* genus, indicating compatibility between scion and rootstock.

For Pereira et al. (2014), the presence of necrosis and excessive stem diameter below, above or at the grafted area are associated with graft incompatibility. Grigolo et al. (1997) also reported

Table 1. Mean squares and mean and coefficient of variation (CV) for the traits mean fruit weight (Mfw; g), pulp weight (Pul; g), total number of fruits per plant (Nfp), total fruit weight per plant (Tfw; kg), seed weight per fruit (Swf; g), fruit width (Frw; mm), fruit length (Frl; mm), stem diameter ratio at 15 cm below and 15 cm above the grafting point (Sdr), canopy diameter I (Cd1; m), plant height (Plh; m) and canopy diameter II (Cd2; m) measured in 83 guava accessions grafted on BRS Guaraçá rootstock.

Source of variation	Mean squares										
	Mfw ^{1/}	Pul	Nfp ^{1/}	Tfw ^{1/}	Swf	Frw	Frl	Sdr	Cd1	Plh	Cd2
Block	0.024 ^{ns}	824.14 ^{ns}	1,826.86 ^{**}	115.70 ^{**}	1,018.61 ^{**}	11.92 ^{ns}	32.25 ^{ns}	0.0113 ^{ns}	0.38 [*]	0.032 ^{ns}	0.8789 ^{**}
Accessions	4.39 ^{**}	1,162.50 ^{**}	35.23 ^{**}	1.76 ^{**}	94.51 ^{**}	70.57 ^{**}	105.42 ^{**}	0.0116 ^{ns}	0.15 ^{**}	0.075 ^{**}	0.1269 [*]
Variation between	2.48 ^{ns}	859.41 ^{ns}	25.30 ^{ns}	2.55 ^{ns}	48.54 ^{ns}	7.54 ^{ns}	15.68 ^{ns}	0.023 ^{ns}	0.04 ^{ns}	0.067 ^{ns}	0.0407 ^{ns}
Variation within	1.27	358.67	14.67	0.85	51.12	17.18	33.11	0.0117	0.083	0.037	0.0893
Range	23.0-179.2	19.9-160.0	5-358	0.3-19.3	0.4-45.3	31.8-68.2	31.6-75.4	0.73-1.14	1.6-3.0	1.6-2.9	1.7-2.9
Mean	76.2	61.00	111.58	7.13	15.33	49.72	52.27	0.98	2.33	2.15	2.43
CV(%)	13.1	31.1	43.5	40.2	46.6	8.3	11.0	11.0	12.4	9.0	12.3
	Mfw	Pul	Nfp	Tfw	Swf	Frw	Frl	Rsd	Cd1	Plh	Cd2

Accessions with the five lowest values

Gua73RO Gua73RO Gua89AM Gua89AM Gua147BA Gua17MA Gua17MA Gua82RO Gua33PE Gua33PE Gua22MA
 Gua59SE Gua17MA Gua64BA Gua64BA PedroSato Gua73RO Gua95AM Gua67RO Gua22MA Gua89AM Gua23MA
 Gua02MA Gua59SE Gua181ES Gua145BA Gua23MA Gua02MA Gua01MA Gua70RO Gua23MA Gua145BA Gua89AM
 Gua83AM Gua35PE Gua47PE Gua181ES Gua15MA Gua01MA Gua84AM Gua86AM Gua15MA Gua146BA Gua33PE
 Gua145BA Gua66RO Gua21MA Gua47PE Gua31PI Gua83AM Gua73RO Gua26MA Gua35PE Gua135RR Gua105RS

Accessions with the five highest values

Gua53SE Gua64BA Gua67RO Gua96AM Gua35PE Gua50SE PedroSato Gua99AM Gua83AM Gua132RR PedroSato
 Gua51SE Gua181ES Gua35PE Gua70RO Gua150BA Gua53SE Gua06MA Gua145BA Gua88AM Gua192ES Gua82RO
 PedroSato Gua06MA Gua83AM Gua192ES Gua51SE PedroSato Gua35PE Gua32PE Gua26MA Gua142RR Gua73RO
 Gua06MA PedroSato Gua01MA Gua01MA Gua50SE Gua06MA Gua25MA Gua75RO Gua111GO Gua84AM Gua88AM
 Gua99AM Gua99AM Gua183ES Gua183ES Gua03MA Gua99AM Gua99AM Gua65RO Gua95AM Gua22MA Gua84AM

^{ns}, ^{**} and ^{*}: non-significant and significant at 1 and 5 %, respectively, by the F-test. ^{1/} Data transformed to root square.




Figure 1. Frequency of vegetative traits of guava accessions evaluated for compatibility with the BRS Guaraçá rootstock. A) Stem diameter ratio at 15 cm below and 15 cm above the grafting point; B) canopy diameter I (m); C) canopy diameter II (m); D) plant height (m).

Photos: Carlos Antonio da Silva



Figure 2. Plants without incompatibility symptoms in the field. A and B: plants with lower stem diameter ratio at 15 cm below and 15 cm above the grafting point values; C and D: plants with higher stem diameter ratio at 15 cm below and 15 cm above the grafting point values. The arrows indicate the grafted area.

the absence of incompatibility in grapevine grafted on different rootstocks and highlighted the greater suitability of more vigorous rootstocks, as reported for the BRS Guaraçá (Souza et al. 2018).

Oliveira et al. (2014) studied the affinity between *P. guajava* × *P. guineense* and reported a high genetic proximity between these species when assessing the genetic divergence of *Psidium* accessions using molecular markers. Campos et al. (2017) emphasized that the affinity between scion and rootstock is essential for the grafting success, and, consequently, for the production of grafted seedlings.

Robaina et al. (2015) evaluated the Paluma guava cultivar grafted on strawberry guava (*P. cattleianum*) aiming at nematode resistance and observed incompatibility and low vigor, highlighting the unsuitability of strawberry guava as a rootstock for Paluma.

The present study corroborates the results obtained by Souza et al. (2018), when evaluating the compatibility of guava cultivars grafted on the *P. guajava* × *P. guineense* hybrid, resulting in a high compatibility between rootstock and scion diameter and no exudations or cracks in the stem of grafted plants.

The canopy diameter I ranged from 1.6 to 3.0 m, whereas the canopy diameter II ranged from 1.7 to 2.9 m, with the accessions Gua22MA, Gua23MA and Gua33PE showing the lowest values for canopy diameter I and II (Table 1). The plant height ranged from 1.6 to 2.9 m, with Gua33PE showing the lowest values for canopy diameter I and II and plant height (Table 1).

For canopy diameter, 94 % of the accessions showed values ranging from 1.9 to 2.8 m (Figure 1B). In contrast, for canopy diameter II, 88 % of the accessions showed values ranging from 1.9 to 2.8 m (Figure 1C). For plant height, 87 % showed values from 1.9 to 2.6 m (Figure 1D). Overall, these data also indicate the absence of incompatibility among the accessions grafted on the BRS Guaraçá rootstock and the existence of variability for canopy diameter I/II and plant height among the evaluated accessions. This variability may be exploited to develop new, small-sized cultivars, thus facilitating fruit harvest.

The fruit weight ranged from 31.8 to 179.2 g, with a mean value of 76.2 g (Table 1). For this variable, 64 % of the accessions showed values ranging from 60 to 120 g (Figure 3A). On the other hand, the pulp

weight showed a mean value of 61.0 g and ranged from 19.9 g in the accession Gua59SE to 160 g in Gua73RO (Table 1). For this variable, 78 % of the accessions showed values ranging from 30 to 90 g (Figure 3B). Hussain et al. (2013) emphasized that fruit weight is an essential characteristic for producers and consumers, with medium to large fruits contributing to a greater market acceptance.

The total number of fruits ranged from 5 to 358, whereas the total fruit weight in the plant ranged from 0.3 to 19.7 kg, with the accessions Gua89AM and Gua183ES showing the minimum and maximum values for these variables, respectively (Table 1). Approximately 50 % of the accessions produced less than 60 fruits plant⁻¹, whereas 12 % of the studied accessions produced more than 240 fruits plant⁻¹ (Figure 3C). For total fruit weight, 14 accessions had more than 12 kg of fruits plant⁻¹ (Figure 3D).

Gonçalves et al. (2019) studied the vegetative and productive performance and fruit quality of peach grafted on 23 rootstocks and observed differences in the productive performance. However, the authors emphasized that the production values obtained should not be considered the maximum potential of the rootstocks, since only the first production cycle was evaluated. Likewise, the fruit production values reported in the present study also should not be considered the maximum production potential of the evaluated accessions.

In a study conducted by Souza et al. (2018), the mean values for total number of fruits, mean fruit weight and total fruit weight differed significantly between harvests. The fruit yield of the guava cultivars Paluma and Pedro Sato grafted on the hybrid rootstock was approximately 40 t ha⁻¹, ten times higher than in ungrafted Paluma.

The mean seed weight was 15.3 g, ranging from 0.4 g in Gua147BA to 48.4 g in Gua03MA. Pedro Sato appears among the five accessions with the lowest seed weight values, thus reaffirming one of the traits for this cultivar, which is its small number of seeds (Table 1). For seed weight, 78 % of the accessions showed values ranging from 8 to 24 g, indicating fruits with fewer seeds (Figure 3E).

The fruit width ranged from 9.9 to 69.3 mm, whereas the fruit length ranged from 7.7 to 77.0 mm, with the accessions Gua01MA, Gua17MA and Gua73RO showing the lowest values for both variables. Most accessions showed values ranging from 38 to 62 mm for fruit width (88 %) and fruit




Figure 3. Frequency of fruit production traits of *Psidium guajava* accessions evaluated for compatibility. A) Mean fruit weight (g); B) pulp weight (g); C) total number of fruits; D) total fruit weight (kg); E) seed weight (g); F) fruit width (mm); G) fruit length (mm).

length (84 %) (Figures 3F and 3G). Pedro Sato stood out among the five highest values for mean fruit weight, pulp weight, fruit width, fruit length and canopy length. Among the evaluated accessions, Gua06MA and Gua99AM stood out with the highest values for mean fruit weight, pulp weight, fruit width and fruit length (Table 1).

Vigorous rootstocks translocate water and nutrients more efficiently and show a greater production of growth-promoting substances, thus favoring the canopy development (Pauletto et al. 2001). The present study highlighted the absence of incompatibility symptoms among *P. guajava* accessions grafted on a nematode-resistant rootstock, with plants showing a mean canopy diameter of 2.3 m and a mean plant height of 2.2 m. Gua64BA, Gua181ES, Gua06MA, Pedro Sato and Gua99AM showed the highest values for pulp and fruit weight, whereas Gua96AM, Gua70RO, Gua192ES, Gua01MA and Gua183ES resulted in

plants with greater fruit production, demonstrating their commercial potential for grafting on the BRS Guaraçá rootstock.

CONCLUSIONS

1. There were no incompatibility symptoms, e.g., cracks, exudations and differences in stem diameter in the grafted area, among the 83 guava accessions grafted on the BRS Guaraçá rootstock, indicating a high compatibility after the second harvest cycle;
2. The accessions Gua64BA, Gua181ES, Gua06MA, Pedro Sato and Gua99AM showed the highest values for pulp and fruit weight (around 160 g), whereas Gua96AM, Gua70RO, Gua192ES, Gua01MA and Gua183ES showed a greater fruit production (around 20 kg plant⁻¹), demonstrating their suitability for grafting on the BRS Guaraçá rootstock.

ACKNOWLEDGMENTS

To the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Capes), for the scholarship provided to Andressa M. S. Lourenço; and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), for the scholarship provided to Carlos A. F. Santos.

REFERENCES

- ALVARES, C. A.; STAPE, J. L.; SENTELHAS, P. C.; MORAES-GONÇALVES, J. L.; SPAROVEK, G. Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift*, v. 22, n. 6, p. 711-728, 2013.
- BOWMAN, K. D.; JOUBERT, J. Citrus rootstocks. In: TALON, M.; CARUSO, M.; GMITTER JUNIOR, F. G. (org.). *The genus Citrus*. London: Elsevier, 2020. p. 105-127.
- CAÑIZARES, A.; LAVERDE, D.; PUESME, R. Crecimiento y desarrollo del fruto de guayaba (*Psidium guajava* L.). *Revista Científica UDO Agrícola*, v. 3, n. 1, p. 34-38, 2003.
- FACHINELLO, J. C.; HOFFMANN, A.; NACHTIGAL, J. C.; KERSTEN, E.; FORTES, G. *Propagação de plantas frutíferas de clima temperado*. 2. ed. Pelotas: UFPel, 1995.
- FREITAS, V. M. *Resistência genética de goiabeira e reação de espécies frutíferas visando o manejo de Meloidogyne enterolobii*. Tese (Doutorado em Agronomia) - Faculdade de Agronomia e Medicina Veterinária, Universidade de Brasília, Brasília, DF, 2012.
- GOMES, V. M.; SOUZA, R. M.; MUSSI-DIAS, V.; SILVEIRA, S. F.; DOLINSKI, C. Guava decline: a complex disease involving *Meloidogyne mayaguensis* and *Fusarium solani*. *Journal of Phytopathology*, v. 159, n. 1, p. 45-50, 2010.
- GONÇALVES, E. D.; MONTEIRO, V. F. C.; MAYER, N. A.; MOURA, P. H. A.; ALVARENGA, A. A.; ANTUNES, L. E. C.; TREVISAN, R.; PÁDUA, J. G. Desempenho de pessegueiro 'BRS Libra' autoenraizado e enxertado sobre porta-enxertos clonais em Minas Gerais. *Revista Brasileira de Ciências Agrárias*, v. 14, n. 1, e5610, 2019.
- CAMPOS, G. S.; MARINHO, C. S.; PORTELLA, C. R.; AMARAL, B. D.; CARVALHO, W. S. G. Production of guava mini-grafted on intra or interspecific rootstock. *Revista Brasileira de Fruticultura*, v. 39, n. 1, e635, 2017.
- GRIGOLO, C. R.; CITADIN, I.; FELDBERG, N. P.; SCARIOTTO, S.; PERTILLE, R. H.; HARTMANN, H. T.; KESTER, D. E.; DAVIES, F. T.; GENEVE, R. L. *Plant propagation: principles and practices*. 6. ed. New Jersey: Prentice-Hall, 1997.
- HE, W.; WANG, Y.; CHEN, Q.; SUN, B.; TANG, H. R.; WANG, X. R. Dissection of the mechanism for compatible and incompatible graft combinations of *Citrus grandis* (L.) Osbeck ("Hongmian Miyou"). *International Journal of Molecular Sciences*, v. 19, n. 2, e505, 2018.
- HUSSAIN, S.; CURK, F.; ANJUM, M. A.; PAILLY, O. Performance evaluation of common clementine on various citrus rootstocks. *Scientia Horticulturae*, v. 150, n. 1, p. 278-282, 2013.
- INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATÍSTICA (IBGE). *Sistema IBGE de Recuperação Automática - SIDRA*. 2020. Available at: <https://sidra.ibge.gov.br/tabela/5457>. Access on Mar. 21, 2020.
- MANICA, I.; ICUMA, I. M.; JUNQUEIRA, N. T. V.; SALVADOR, J. O.; MOREIRA, A.; MALAVOLTA, E. *Fruticultura tropical 6: goiaba*. Porto Alegre: Cinco Continentes, 2000.
- MARTINS, A. N.; NARITA, N.; SUGUINO, E.; TAKATA, W. Desempenho de cultivares de goiabeiras em ambientes irrigado e sequeiro. *Colloquium Agrariae*, v. 16, n. 2, p. 82-89, 2020.
- MELNYK, W. C. Plant grafting insights into tissue regeneration. *Regeneration*, v. 4, n. 1, p. 3-14, 2016.
- MISRA, A. K. Progressive steps in understanding and solving guava wilt: a national problem. *Indian Phytopathology*, v. 70, n. 1, p. 1-11, 2017.
- OLIVEIRA, N. N. S.; VIANA, A. P.; QUINTAL, S. S. R.; PAIVA, C. Análise de distância genética entre acessos do gênero *Psidium* via marcadores ISSR. *Revista Brasileira de Fruticultura*, v. 36, n. 4, p. 917-923, 2014.
- PAULETTO, D.; MOURÃO FILHO, F. A. A.; KLUGE, R. A.; SCARPARE FILHO, J. A. Produção e vigor da videira "Niágara Rosada" relacionados com o porta-enxerto. *Pesquisa Agropecuária Brasileira*, v. 36, n. 1, p. 115-121, 2001.
- PEREIRA, F. M.; SOUZA, R. M.; SOUZA, P. M.; DOLINSKI, C.; SANTOS, G. K. Estimativa do impacto econômico e social direto de *Meloidogyne mayaguensis* na cultura da goiaba no Brasil. *Nematologia Brasileira*, v. 33, n. 2, p. 176-181, 2009.
- PEREIRA, I. S.; FACHINELLO, J. C.; ANTUNES, L. E. C.; CAMPOS, A. D.; PINA, A. Incompatibilidade de enxertia em prunus. *Ciencia Rural*, v. 44, n. 9, p. 1519-1526, 2014.
- POMMER, C. V.; MURAKAMI, K. R. Breeding guava (*Psidium guajava* L.). In: JAIN, S. M.; PRIYADARSHAN,

- P. M. (org.). *Breeding plantation tree crops: tropical species*. New York: Springer, 2009. p. 83-120.
- ROBAINA, R. R.; RENATA, G. S.; MARINHO, C. S.; SOUZA, R. M.; BREMENKAMP, C. A. Grafting guava on cattley guava resistant to *Meloidogyne enterolobii*. *Ciência Rural*, v. 45, n. 9, p. 1579-1584, 2015.
- SANTOS, H. G.; JACOMINE, P. K. T.; ANJOS, L. H. C.; OLIVEIRA, V. A.; LUMBRERAS, J. F.; COELHO, M. R.; ALMEIDA, J. A.; CUNHA, T. J. F.; OLIVEIRA, J. B. *Sistema brasileiro de classificação de solos*. 5. ed. Brasília, DF: Embrapa, 2018.
- SANTOS, C. A. F.; REIS, M. S.; CRUZ, C. D.; SEDIYAMA, C. S.; SEDIYAMA, T. Parâmetros genéticos e seleção indireta em progênies F6 de um cruzamento de soja (*Glycine max* (L.) Merrill). *Revista Ceres*, v. 42, n. 240, p. 155-166, 1995.
- SOUZA, R. R. C. de; SANTOS, C. A. F.; COSTA, S. R. da. Field resistance to *Meloidogyne enterolobii* in a *Psidium guajava* × *P. guineense* hybrid and its compatibility as guava rootstock. *Fruits*, v. 73, n. 2, p. 118-124, 2018.
- UNITED STATES DEPARTMENT OF AGRICULTURE (USDA). Soil Survey Staff. *Keys to soil taxonomy*. 12. ed. Washington, DC: USDA, 2014.
- VRŠIČ, S.; PULKO, B.; KOCSIS, L. Factors influencing grafting success and compatibility of grape rootstocks. *Scientia Horticulturae*, v. 181, n. 1, p. 168-173, 2015.