



Pesquisa Agropecuária Tropical

ISSN: 1517-6398

ISSN: 1983-4063

Escola de Agronomia/UFG

Santos, Jônatas Barros dos; Silva, Alberto do Nascimento; Queiroz, Paulo Roberto Martins; Eckstein, Barbara; Monnerat, Rose Gomes
Selection of *Bacillus thuringiensis* strains toxic to *Meloidogyne incognita*
Pesquisa Agropecuária Tropical, vol. 52, e73070, 2022
Escola de Agronomia/UFG

DOI: <https://doi.org/10.1590/1983-40632022v5273070>

Available in: <https://www.redalyc.org/articulo.oa?id=253070366052>

- ▶ How to cite
- ▶ Complete issue
- ▶ More information about this article
- ▶ Journal's webpage in redalyc.org

The logo for Redalyc.org, featuring the text 'redalyc.org' in a stylized font with a red dot above the 'y'.

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Selection of *Bacillus thuringiensis* strains toxic to *Meloidogyne incognita*¹

Jônatas Barros dos Santos², Alberto do Nascimento Silva³,
Paulo Roberto Martins Queiroz⁴, Barbara Eckstein⁵, Rose Gomes Monnerat⁵

ABSTRACT

The *Bacillus thuringiensis* bacterium has demonstrated an effective potential in the control of several agricultural pests, among them nematodes. This study aimed to standardize and establish a methodology of bioassays of *B. thuringiensis* and *Meloidogyne incognita* with the use of resorcinol, select *B. thuringiensis* strains toxic to *M. incognita* *in vitro* and molecularly identify the presence of the *cry6* gene in *B. thuringiensis* strains. Second-stage juveniles were subjected to resorcinol doses, verifying that the concentration of 0.2 % did not cause mortality and provided the Cry6A toxin ingestion. Thereafter, 16 *B. thuringiensis* strains were tested in the presence or absence of resorcinol, resulting in mortality rates of 4-36 %. Among the *B. thuringiensis* strains analyzed for the presence of the *cry6* gene, only one was detected (S1617).

KEYWORDS: Ingestion inducer, *cry6* gene, nematode.

RESUMO

Seleção de estirpes de *Bacillus thuringiensis* tóxicas a *Meloidogyne incognita*

A bactéria *Bacillus thuringiensis* tem demonstrado potencial no controle de diversas pragas agrícolas, dentre elas os nematoides. Objetivou-se padronizar e estabelecer uma metodologia de bioensaios de *B. thuringiensis* e *Meloidogyne incognita* com o uso de resorcinol, selecionar estirpes de *B. thuringiensis* tóxicas a *M. incognita* *in vitro* e identificar molecularmente a presença do gene *cry6* em estirpes de *B. thuringiensis*. Juvenis de segundo estágio foram submetidos a doses de resorcinol, verificando-se que a concentração de 0,2 % não causou mortalidade e proporcionou a ingestão da toxina Cry6A. Após isso, 16 estirpes de *B. thuringiensis* foram testadas na presença ou não de resorcinol, resultando em mortalidades de 4-36 %. Dentre as estirpes de *B. thuringiensis* analisadas quanto à presença do gene *cry6*, detectou-se apenas uma (S1617).

PALAVRAS-CHAVE: Indutor de ingestão, *cry6* gene, nematoide.

INTRODUCTION

Agricultural losses caused by nematodes are increasingly frequent in Brazil. Nematodes of the *Meloidogyne* genus constitute the most important group of phytoparasites in the world, with more than 90 species described (Karssen & Moens 2006) and more than 3,000 host plants, among cultivable plants and weeds, representing a threat to the world agricultural production (Perry et al. 2009, Xiang et al. 2018). In Brazil, *Meloidogyne incognita* (Kofoid and White 1919), Chitwood, 1949 (races 1, 2, 3 and 4), *M. javanica* and *M. arenaria* predominate, being present in any type of soil, mainly in sandy soils and with temperatures above 25 °C (Pinheiro et al. 2010).

The main symptom of *M. incognita* infection is the formation of typical galls in the roots of susceptible host plants (Xiang et al. 2018), resulting in a lower absorption of water and nutrients and presenting weakened plants, of smaller stature, withered at certain times of the day, and with low yield (Priya et al. 2011, Zeng et al. 2018, Sikandar et al. 2019). These nematodes are endophytic, sedentary and have a fast reproduction rate. Therefore, their management depend on crop rotation with non-host plants, what is not interesting from the producer's point of view. Chemical control has been the main control strategy used; however, it presents variable results in the field, high costs and negative impacts on the environment (Engelbrecht et al. 2018) and,

¹ Received: June 09, 2022. Accepted: Sep. 23, 2022. Published: Oct. 27, 2022. DOI: 10.1590/1983-40632022v5273070.

² Universidade de Brasília, Faculdade de Agronomia e Medicina Veterinária, Brasília, DF, Brasil.

E-mail/ORCID: jonatas.bsantos@hotmail.com/0000-0002-3028-2155.

³ Universidade do Estado da Bahia, Barreiras, BA, Brasil. *E-mail/ORCID*: albertosilvaagro@gmail.com/0000-0003-3003-0414.

⁴ Centro Universitário de Brasília, Brasília, DF, Brasil. *E-mail/ORCID*: prqzqueiroz@gmail.com/0000-0002-4672-8123.

⁵ Empresa Brasileira de Pesquisa Agropecuária (Embrapa Recursos Genéticos e Biotecnologia), Brasília, DF, Brasil.

E-mail/ORCID: barbara.eckstein@embrapa.br/0000-0002-9734-6758; rose.monnerat@gmail.com/0000-0001-7822-2525.

as such, many of these products that had a non-tolerable level of toxicity to non-target organisms were withdrawn from use (Oka 2014, Soltanzadeh et al. 2016). In addition, biological control has shown considerable efficacy results by several phytopathogenic nematode control microorganisms and by different forms of action (Engelbrecht et al. 2018, Oliveira et al. 2019, Ramalakshmi et al. 2020).

Among bacteria, *Bacillus* has a great potential in agricultural pest control, since it is a microorganism present in various ecological niches, isolated from various environments, especially the soil, with a direct association with nematodes (Li et al. 2007). In the *Bacillus* genus, *B. thuringiensis* (*Bt*) Berliner, 1915, represents the most successful commercial biopesticide in the biological control market, due to the great diversity of isolated and identified strains. In addition to the spectrum of action that encompasses several agricultural pests, nematodes and public health vectors, *B. thuringiensis* has an easy mass reproduction with the use of different technologies and specific action for its target pests, being harmless to humans, vertebrates, beneficial plants and insects, as well as completely biodegradable (Bravo 2018, Gutierrez et al. 2019).

Cry proteins produced by *B. thuringiensis* are classified into families from Cry1 to Cry78, based on their amino acid sequence identity (Crickmore et al. 2018). Among these known Cry toxin families, Cry5, Cry6, Cry12, Cry13, Cry14 and Cry21 demonstrated nematocidal activity (Guo et al. 2008). Transgenic plants that express the *cry5b* (Li et al. 2008) and *cry6a* (Li et al. 2007) genes significantly reduce the reproductive capacity of *M. incognita* and the number of galls in plant roots. Thus, the isolation and sequencing of *cry* genes should be encouraged in the search for new toxins that can be used in the biological control of pests or in the genetic transformation of plants.

The search for toxic proteins, regardless of the target, undergoes *in vitro* pathogenicity bioassays. In the case of nematodes, it is necessary to ingest the proteins to know their toxicity. In this way, in order to induce the ingestion of proteins, the root extract of plants, phenolic compounds such as resorcinol (Zhang et al. 2012) or carbamoylcholine chloride (Adam et al. 2008) have been used, or tested on model microorganisms such as the free-living bacteriophage nematode *Caenorhabditis elegans* (Montalvão et al. 2018, Verduzco-Rosas et al. 2021).

Resorcinol is a phenolic compound described as a neurostimulant known to induce the impulse of the stylet along with the accumulation of esophageal gland secretions, allowing the absorption of substances present in the solution by the nematode (Jaubert et al. 2002, Williamson & Gleason 2003). Huang et al. (2006) used resorcinol to stimulate dsRNA uptake during the RNAi (interfering RNA) process *in vitro* in second stage juveniles (J2) of *M. incognita*. Zhang et al. (2012), on the other hand, perfected the bioassay protocol involving *B. thuringiensis* and *M. hapla* using resorcinol as an inducer of protein intake to replace root exudates of tomato plants; but the dose used in these assays was not compatible with *M. incognita*. However, adverse effects were reported when stimulants were used to induce the intake of substances, from changes in mobility (Goto et al. 2010) to even the death of nematodes (Adam et al. 2008).

Therefore, this study aimed to standardize a methodology of bioassays of *B. thuringiensis* with *M. incognita* and the use of resorcinol, select strains of *B. thuringiensis* toxic to *M. incognita* in *in vitro* assays and molecularly identify the presence of the *cry6* gene in strains of *B. thuringiensis* with potential to be used in the biocontrol of nematodes.

MATERIAL AND METHODS

The experiments were carried out from 2017 to 2018, at the Embrapa Recursos Genéticos e Biotecnologia (Brasília, DF, Brazil). Sixteen strains of *Bacillus thuringiensis* (*Bt*) belonging to the Invertebrate Bacteria Collection were used: S09, S26, S53, S906, S1295, S1577, S1617, S1615, S1620, S1930, S2493, S2538, S2548, S2557, S2558 and S2560.

The strains were grown using Embrapa medium (Monnerat et al. 2007) at 28 ± 2 °C for 72 h, in a rotary incubator at 200 rpm. After this culture period, the morphology of the strains was analyzed with a phase contrast microscope to observe the presence of spores and protein crystals.

The initial inoculum of *Meloidogyne incognita* race 3 was provided by the Embrapa. The nematodes were multiplied in tomato (*Solanum lycopersicum* L. cv. Santa Clara) for 3 months in a greenhouse. The eggs were obtained using the technique of Hussey & Barker (1973), modified by Boneti & Ferraz (1981). Surface disinfestation of the eggs was performed

in a laminar flow chamber (Zuckerman & Brzeski 1966). The eggs were suspended in 30 mL of 0.12 % chlorhexidine gluconate solution along with antibiotics (10 $\mu\text{g mL}^{-1}$ of erythromycin and 2.5 g L^{-1} of streptomycin) for 30 min. Then, they were centrifuged (3 min; 360 g) and, after that, the supernatant was discarded and the eggs present in the pellet were suspended in sterile distilled water. The procedure was repeated twice in a sterile environment, where the eggs were placed in a modified Baermann funnel (Flegg 1967) for incubation. The *M. incognita* J2 individuals that hatched within 24-48 h were collected for use in the experiments.

The aforementioned strains were used in the *in vitro* pathogenicity test. In 6-cell plates, 1,680 μL of the nematode suspension obtained from the hatching chamber were added, standardized with approximately 45 juveniles J2 of *M. incognita* and 320 μL (16 %) of the bacterial suspensions of *Bt*, totaling 2 mL, with the J2 incubated only in sterile distilled water in the control treatment. The plates were sealed with plastic film and kept in a BOD type chamber at 25 °C, in the absence of light, for 48 h. After the incubation period, the evaluation was performed, adding 10 μL of 1 N NaOH (Xiang & Lawrence 2016) to the solution, in order to help differentiate the paralyzed or immobile nematodes from those that were actually dead. The counts were performed under an optical microscope and Petters slide, determining the percentages of dead J2.

In order to obtain the non-lethal concentration of resorcinol, tests were performed with different doses of the product on the obtained J2, as previously described. For this, different solutions of resorcinol diluted in distilled water (v/v) were prepared in pre-established proportions. Given the regression curve generated by the average of the tests, it was found that the concentration of 0.2 % did not cause toxicity to the nematodes. From this moment on, new assays were carried out to evaluate whether this concentration would provide the ingestion of the Cry6Aa protein, described as a nematicide (Wei et al. 2003, Guo et al. 2008, Palma et al. 2014), obtained from a genetically transformed strain of *Bt*.

To this end, the mortality of nematodes was evaluated after 48 h in contact with the bacterial suspension of *Bt* (Cry6Aa), at a concentration of 16 %, 10 % of resorcinol (0.2 %) and 74 % of nematode suspension containing approximately 45 juveniles J2 per mL. In the control treatment, the J2 were incubated

in sterile distilled water and the material incubated and evaluated after 48 h. After standardization, all *B. thuringiensis* strains were retested in the presence of resorcinol, following this methodology.

To determine the non-toxic dose of resorcinol, regression analysis was performed with increasing doses of the product and mortality rate, using the average of 3 trials and with 4 repetitions for each dose.

The mean mortality data of *M. incognita* with the use of *B. thuringiensis* strains associated or not with resorcinol were submitted to analysis of variance and compared by the Scott-Knott test at the level of 5 % for the formation of groups between the tested treatments.

DNA from the strains was extracted using the methodology described by Bravo et al. (1998), with the bacteria cultured using the Embrapa solid medium (Monnerat et al. 2007) for 16 h, at 30 °C. After this period, for each sample, the bacterial growth was collected with a loop and transferred to a properly identified 1.5-mL polyethylene tube containing 200 μL of sterile ultra-purified water (MiliQ). The samples were then homogenized in a Vortex apparatus and frozen at -80 °C for 20 min, boiled in bain-marie at 100 °C for 10 min, and finally incubated on ice for 2 min. The obtained supernatant was used for reactions, then 5 μL of the DNA from each strain were transferred to a 0.2-mL polyethylene tube containing 0.5 μM of the oligonucleotide, 0.2 mM dNTP, 1X *Taq* buffer and 2.5 U of *Taq* DNA polymerase (5.0 U), totaling a final volume of 25 μL . The characteristics of the initiator used for the reaction are described in Table 1. The PCR program used for the amplification was 94 °C for 5 min, 35 cycles of 94 °C for 1 min, annealing of 48.5 °C for 1 min, 72 °C for 1 min and 30 s, as well as a final extension of 72 °C for 5 min.


For the analysis of the result, 25 μL of the PCR product mixed with 5 μL of 10X running buffer in 1.5 % agarose gel were applied. The electrophoresis run was done in 1X TBE buffer (Tris base; boric acid; 0.5 M EDTA; pH 8.0). After electrophoresis, the gel was stained with ethidium bromide at 1 $\mu\text{g mL}^{-1}$ for 20 min, and decolorized in distilled water for 15 min. The gel was visualized in a transilluminator under UV light and photographed in a photo-documenter.

RESULTS AND DISCUSSION

Figure 1 shows the percentage of resorcinol concentration associated with *M. incognita*,

Table 1. Characteristics of the primers used for the detection of the *cry6* gene in *Bacillus thuringiensis* strains.

Target Gene	Sequence of primers (5' → 3')	Primer name	Fragment size (bp)	Tm (°C)	Genbank number
<i>cry6</i>	TCCACCATTCTGCTGAAGTTT (d)	Cry6AaF	177	48.5	AAA22357
	TAGCCCATTCTGTTTTGGAA (r)	Cry6AaR			AAM46849
					ABH03377
					AAA22358.1

Figure 1. Mean mortality of *Meloidogyne incognita* in three trials, when subjected to resorcinol concentrations. The dotted line indicates 15 % for maximum mortality.

considering a maximum mortality of 15 % in the tests. Mortality values increased as the product concentration increased. Adam et al. (2008) also found that resorcinol doses greater than 0.5 % cause lethality in *M. javanica* J2, what was also observed for *M. incognita* during this assay.

Considering the prescribed mortality limit, a value of 0.2 % was adopted for subsequent validation tests. From this point, it was tested if this selected concentration would maintain the neurostimulant properties of the product. In the following trials, where resorcinol was associated with the bacterial suspension of a genetically modified *B. thuringiensis* strain, expressing only the nematocidal protein Cry6Aa, there was a 3.5-fold higher mortality, when compared to the treatment of the same strain without resorcinol (Table 2). On the other hand, the control treatment with resorcinol alone presented only 4.67 % of mortality, well below the 15 % pre-established as a mortality limit for the control.

The presence of resorcinol stimulated the intake of nematocidal proteins that were in the solution, improving the nematocidal activity, considering that there was a 3.5-fold increase in mortality. Zhang et al. (2012) obtained similar results

Table 2. Mortality rate of *Meloidogyne incognita*, the transformed strain of *Bacillus thuringiensis*, when associated or not with resorcinol.

Treatments	Mortality (%)
Strain with resorcinol	28.93 ± 1.48
Strain without resorcinol	8.18 ± 2.04
Control with resorcinol	4.67 ± 1.76
Control with water	2.63 ± 0.54

for *M. hapla*, verifying an increase in the nematocidal activity with resorcinol, when compared to tomato root exudates.

Of the 16 strains of *B. thuringiensis* tested against *M. incognita* in the presence or absence of resorcinol as an inducer of protein intake (Table 3), it was found that, in the absence of resorcinol, 8 strains (S1615, S1577, S1617, S2538, S1620, S1295, S53 and S2493) were statistically equal to the control treatment, not presenting satisfactory mortality results. Seven other strains presented intermediate results, differing statistically from the control group and presenting an intermediate mortality rate between 8.81 and 17.04 % (S09, S2548, S2557, S2558, S2560, S26 and S906). Finally, the highest mortality obtained without the use of resorcinol was achieved by the S1930 strain, with 35.7 %.

By analyzing the column of the treatments tested with the addition of resorcinol, it is clear that, in general, the mortality rates in the presence of resorcinol were higher than those found in the absence of the product, or statistically equal, except for the S2557 strain, which had its mortality reduced in the presence of the product, probably interfering negatively in some substance produced by the strain.

The S2493 strain, which presents the nematocidal gene *cry6*, which encodes the Cry6 toxin and presented a low mortality rate (2.99 %), stood out in the absence of resorcinol, appearing in the intermediate control group (28.93 %). With the use of resorcinol, taking into account that only the bacterial suspension was used,

Table 3. Mortality results (%) of *Meloidogyne incognita* as a function of strains tested in the presence or absence of resorcinol.

Strain	Resorcinol	
	Without	With
Control	6.21 Ad*	2.99 Ad
S1615	5.38 Bd	21.89 Ab
S1577	4.54 Bd	15.29 Ac
S1617	4.46 Bd	25.11 Ab
S2538	4.59 Ad	8.20 Ad
S09	8.81 Bc	23.06 Ab
S2548	20.28 Ab	20.58 Ab
S2557	9.66 Ac	7.54 Ad
S2558	13.85 Ac	6.28 Bd
S2560	9.75 Ac	6.25 Ad
S1620	3.90 Bd	10.28 Ad
S1295	3.45 Ad	6.04 Ad
S26	12.86 Ac	15.19 Ac
S53	6.19 Ad	8.75 Ad
S2493	2.99 Bd	28.93 Ab
S1930	35.70 Aa	38.87 Aa
S906	17.04 Ab	21.38 Ab

* The same uppercase letter in the row and lowercase letter in the column do not differ from each other by the Scott-Knott test at 5 % of probability.

without any type of concentration and purification technique aiming to obtain only the crystals produced by the strain, it is assumed that the mortality result could be much higher.

Another strain that showed a large increase (greater than 5 times) in mortality was S1617. Subsequently, it was detected that the strain showed a positive result for the PCR product with the *cry6* primer, indicating that this is the reason for the mortality expansion.

The strain that caused the highest percentage of mortality in the *in vitro* tests was S1930, not being influenced by the presence of resorcinol, although its use caused a higher mortality. This strain also did not present the PCR product with the *cry6* primer previously tested. It can be seen, then, in comparison with the strains that presented the nematocidal protein, that, in the presence of resorcinol, there is a large increase in mortality, as observed in the strains S2493 and S1617, indicating that the probable cause of mortality for the strain S1930 was not a Cry protein, but some other toxic metabolite produced by the strain, which was later verified with the sequencing of this strain.

Zheng et al. (2016) demonstrated that 120 spore-forming *Bacillus* strains from different families share virulence factors, what may contribute to their

nematicidal capacity. *B. thuringiensis*, *B. cereus*, *B. subtilis*, *B. pumilus*, *B. firmus*, *B. toyonensis*, *Lysinibacillus sphaericus*, *Brevibacillus laterosporus* and *B. brevis* were highly nematocidal, with the first of them demonstrating the highest activity.

Jouzani et al. (2008) observed *in vitro* that the concentration of spore-crystal mixtures of 22 selected strains provided an average mortality 10 % higher in *M. incognita* with a concentration 2 times the standard. The strains with the best performances presented toxicity of 77 and 81 %, respectively, for the combination of spores and crystals.

Verduzco-Rosas et al. (2021), testing 310 strains of *B. thuringiensis*, using *C. elegans* as a model, found that 10 strains showed toxicity results in spore and crystal concentration. Of these, two strains produced damage to the body and intestine of the nematode. Molecular analysis also revealed the presence of *cry5B*, *cry14A*, *cry21A* and *cyt1A* genes and the new *cry* gene.

The results revealed a very low occurrence of the nematocidal gene *cry6* in the selected strains. As previously mentioned, only the S1617 strain presented the *cry6* gene. According to the methodology used in the present study, only the detection of the *cry6* gene was sought; thus, it cannot be said that the used strains are not producers of other nematocidal proteins.

Of the 70 *Bt* isolates tested by Jouzani et al. (2008), using 12 primers specific for the nematocidal genes *cry5*, *cry6*, *cry12*, *cry13*, *cry14* and *cry21*, the authors found 22 isolates (31.5 %) which contain at least one *cry* gene active against nematodes. Strains containing the *cry6* gene were the most abundant and represent 22.8 % of the isolates.

On the other hand, Bravo et al. (1998), analyzing multiplex PCR-based with general and specific primers, found a total of 496 samples of *B. thuringiensis*, which were isolated from 503 soil samples collected from the five macro-regions of Mexico, of which the proteins are nematocidal and were analyzed in this study (Cry5, Cry12, Cry13 and Cry14), where no strains with the *cry5*, *cry12*, *cry13*, *cry14* or *cry21* genes were found.

Although soil is the main source of most *B. thuringiensis* isolates, and they still have the ability to present several *cry* genes in their molecular profile, their geographical and ecological distribution are very variable, that is, it is not possible to establish a correlation between geographical distribution and frequency of occurrence of *cry* genes of nematocidal activity.

CONCLUSIONS

1. An *in vitro* bioassay methodology was defined using *Bacillus thuringiensis* and resorcinol as inducer of protein intake in *Meloidogyne incognita*;
2. The S1930 strain showed promising results for mortality of *M. incognita*, even at low concentrations;
3. There are strains toxic to the nematode *M. incognita in vitro* and which have their potential increased with the use of resorcinol;
4. The S1617 strain tested positive for the *cry6* gene amplification.

ACKNOWLEDGMENTS

This study was partially financed by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (Capes) (Finance code 001).

REFERENCES

- ADAM, M. A. M.; PHILLIPS, M. S.; JONES, J. T.; BLOK, V. C. Characterisation of the cellulose-binding protein Mj-cbp-1 of the root knot nematode, *Meloidogyne javanica*. *Physiological and Molecular Plant Pathology*, v. 72, n. 1, p. 21-28, 2008.
- BONETI, J. I. S.; FERRAZ, S. Modificação do método de Hussey & Barker para extração de ovos de *Meloidogyne exigua* de raízes de cafeeiros. *Fitopatologia Brasileira*, v. 6, n. 3, p. 553, 1981.
- BRAVO, A. Biodiversity of cry toxins produced by *Bacillus thuringiensis* and evolution of resistance to these toxins in different insect pests. *Toxicon*, v. 149, e98, 2018.
- BRAVO, A.; SARABIA, S.; LOPEZ, L.; ONTIVEROS, H.; ABARCA, C.; ORTIZ, A.; ORTIZ, M.; LINA, L.; VILLALOBOS, F. J.; PEÑA, G.; NUÑEZ-VALDEZ, M. E.; SOBERÓN, M.; QUINTERO, R. Characterization of cry genes in Mexican *Bacillus thuringiensis* strain collection. *Applied and Environmental Microbiology*, v. 64, n. 12, p. 4965-4972, 1998.
- CRICKMORE, N.; BAUM, J.; BRAVO, A.; LERECLUS, D.; NARVA, K.; SAMPSON, K.; SCHNEPF, E.; SUN, M.; ZEIGLER, D. R. *Bacillus thuringiensis* toxin nomenclature. 2018. Available at: <http://www.btnomenclature.info/>. Access on: Sep. 22, 2018.
- ENGELBRECHT, G.; HORAK, I.; RENSBURG, P. J. J.; CLAASSENS, S. Bacillus-based bionematicides: development, modes of action and commercialization. *Biocontrol, Science and Technology*, v. 28, n. 7, p. 629-653, 2018.
- FLEGG, J. J. M. Extraction of *Xiphinema* and *Longidorus* species from soil by a modification of Cobb's decanting and sieving technique. *Annual Applied Biology*, v. 60, n. 3, p. 429-437, 1967.
- GOTO, D. B.; FOSU-NYARKO, J.; SAKUMA, F.; SADLER, J.; FLOTTMAN-REID, M.; UEHARA, T.; KONDO, N.; YAMAGUCHI, J.; JONES, M. G. K. *In planta* observation of live fluorescent plant endoparasitic nematodes during early stages of infection. *Nematological Research*, v. 40, n. 1, p. 15-19, 2010.
- GUO, S.; LIU, M.; PENG, D.; JI, S.; WANG, P.; YU, Z.; SUN, M. New strategy for isolating novel nematicidal crystal protein genes from *Bacillus thuringiensis* strain YBT-1518. *Applied and Environmental Microbiology*, v. 74, n. 22, p. 6997-7001, 2008.
- GUTIERREZ, M. E. M.; CAPALBO, D. M. F.; ARRUDA, R. O.; MORAES, R. O. *Bacillus thuringiensis*. In: SOUZA, B.; VÁZQUEZ, L.; MARUCCI, R. (ed.). *Natural enemies of insect pests in neotropical agroecosystems*. Cham: Springer, 2019.
- HUANG, G.; ALLEN, R.; DAVIS, E. L.; BAUM, T. J.; HUSSEY, R. S. Engineering broad root-knot resistance in transgenic plants by RNAi silencing of a conserved and essential root-knot nematode parasitism gene. *Proceedings of the National Academy of Sciences of the United States of America*, v. 103, n. 39, p. 14302-14306, 2006.
- HUSSEY, R. S.; BARKER, K. R. A comparison of methods of collecting inocula of *Meloidogyne incognita* spp., including a new technique. *Plant Disease Report*, v. 57, n. 12, p. 1025-1028, 1973.
- JAUBERT, S.; LAFFAIRE, J. B.; PIOTTE, C.; ABAD, P.; ROSSO, M.-N.; LEDGER, T. N. Direct identification of stylet secreted proteins from root-knot nematodes by a proteomic approach. *Molecular and Biochemical Parasitology*, v. 121, n. 2, p. 205-211, 2002.
- JOUZANI, G. S.; SEIFINEJAD, A.; SAEEDIZADEH, A.; NAZARIAN, A.; YOUSEFLOO, M.; SOHEILIVAND, S.; MOUSIVAND, M.; JAHANGIRI, R.; YAZDANI, M.; AMIRI, R. M.; AKBARI, S. Molecular detection of nematicidal crystalliferous *Bacillus thuringiensis* strains of Iran and evaluation of their toxicity on free-living and plant-parasitic nematodes. *Canadian Journal of Microbiology*, v. 54, n. 10, p. 812-822, 2008.
- KARSSSEN, G.; MOENS, M. Root-knot nematodes. In: PERRY, R. N.; MOENS, M. (ed.). *Plant nematology*. Wallingford: CABI Publishing, 2006. p. 59-90.
- LI, X. Q.; TAN, A.; VOEGTLIN, M.; BEKELE, S.; CHEN, C. S.; AROIAN, R. V. Expression of cry5B protein from *Bacillus thuringiensis* in plant roots confers resistance to rootknot nematode. *Biological Control*, v. 47, n. 1, p. 97-102, 2008.

- LI, X. Q.; WEI, J.-Z.; TAN, A.; AROIAN, R. V. Resistance to root-knot nematode in tomato roots expressing a nematocidal *Bacillus thuringiensis* crystal protein. *Plant Biotechnology Journal*, v. 5, n. 4, p. 455-464, 2007.
- MONNERAT, R. G.; BATISTA, A. C.; MEDEIROS, P. T.; MARTINS, É. S.; MELATTI, V. M.; PRAÇA, L. B.; DUMAS, V. F.; MORINAGA, C.; DEMO, C.; GOMES, A. C. M.; FALCÃO, R.; SIQUEIRA, C. B.; SILVA-WERNECK, J. O.; BERRY, C. Characterization of Brazilian *Bacillus thuringiensis* strains active against *Spodoptera frugiperda*, *Plutella xylostella* and *Anticarsia gemmatallis*. *Biological Control*, v. 41, n. 3, p. 291-295, 2007.
- MONTALVÃO, S. C. L.; CASTRO, M. T. D.; SOARES, C. M. S.; BLUM, L. E. B.; MONNERAT, R. G. *Caenorhabditis elegans* como indicador de toxicidade de estirpes de *Bacillus thuringiensis* a *Meloidogyne incognita* raça 3. *Ciência Rural*, v. 48, n. 7, e20170712, 2018.
- OKA, Y. Nematicidal activity of fluensulfone against some migratory nematodes under laboratory conditions. *Pest Management Science*, v. 70, n. 12, p. 1850-1858, 2014.
- OLIVEIRA, D. F.; COSTA, V. A.; TERRA, W. C.; CAMPOS, V. P.; PAULA, P. M.; MARTINS, S. J. Impact of phenolic compounds on *Meloidogyne incognita* *in vitro* and in tomato plants. *Experimental Parasitology*, v. 199, n. 1, p. 17-23, 2019.
- PALMA, L.; MUÑOZ, D.; BERRY, C.; MURILLO, J.; CABALLERO, P. *Bacillus thuringiensis* toxins: an overview of their biocidal activity. *Toxins*, v. 6, n. 12, p. 3296-3325, 2014.
- PERRY, R. N.; MOENS, M.; STAR, J. L. *Root-knot nematodes*. Wallingford: CABI, 2009.
- PINHEIRO, J. B.; AMARO, G. B.; PEREIRA, R. B. *Ocorrência e controle de nematoides em hortaliças folhosas*. Brasília, DF: Embrapa Hortaliças, 2010. (Circular técnica, 89).
- PRIYA, D. B.; SOMASEKHAR, N.; PRASAD, J. S.; KIRTI, P. B. Transgenic tobacco plants constitutively expressing *Arabidopsis* NPR1 show enhanced resistance to root-knot nematode, *Meloidogyne incognita*. *BMC Research Notes*, v. 4, n. 1, p. 1-5, 2011.
- RAMALAKSHMI, A.; SHARMILA, R.; INIYAKUMAR, M.; GOMATHI, V. Nematicidal activity of native *Bacillus thuringiensis* against the root knot nematode, *Meloidogyne incognita* (Kofoid and White). *Egyptian Journal of Biological Pest Control*, v. 30, e90, 2020.
- SIKANDAR, A.; ZHANG, M. Y.; ZHU, X. F.; WANG, Y. Y.; AHMED, M.; IQBAL, M. F.; JAVEED, A.; XUAN, Y. H.; FAN, H. Y.; LIU, X. Y.; CHEN, L. J.; DUAN, Y. X. Effects of *Penicillium chrysogenum* strain Snef1216 against root-knot nematodes (*Meloidogyne incognita*) in cucumber (*Cucumis sativus* L.) under greenhouse conditions. *Applied Ecology and Environmental Research*, v. 17, n. 5, p. 12451-12464, 2019.
- SOLTANZADEH, M.; NEJAD, M. S.; BONJAR, G. H. S. Application of soil-borne actinomycetes for biological control against fusarium wilt of chickpea (*Cicer arietinum*) caused by *Fusarium solani* fsp. *lisi*. *Journal of Phytopathology*, v. 164, n. 11-12, p. 967-978, 2016.
- VERDUZCO-ROSAS, L. A.; GARCÍA-SUÁREZ, R.; LÓPEZ-TLACOMULCO, J. J.; IBARRA, J. E. Selection and characterization of two *Bacillus thuringiensis* strains showing nematocidal activity against *Caenorhabditis elegans* and *Meloidogyne incognita*. *FEMS Microbiology Letters*, v. 368, n. 5, efnaa186, 2021.
- WEI, J. Z.; HALE, K.; CARTA, L. *Bacillus thuringiensis* crystal proteins that target nematodes. *Proceedings of the National Academy of Sciences of the United States of America*, v. 100, n. 5, p. 2760-2765, 2003.
- WILLIAMSON, V. M.; GLEASON, C. A. Plant-nematode interactions. *Current Opinion in Plant Biology*, v. 6, n. 4, p. 327-333, 2003.
- XIANG, N.; LAWRENCE, K. S. Optimization of *in vitro* techniques for distinguishing between live and dead second stage juveniles of *Heterodera glycines* and *Meloidogyne incognita*. *Plos One*, v. 11, n. 5, e0154818, 2016.
- XIANG, N.; LAWRENCE, K. S.; DONALD, P. A. Biological control potential of plant growth-promoting rhizobacteria suppression of *Meloidogyne incognita* on cotton and *Heterodera glycines* on soybean: a review. *Journal of Phytopathology*, v. 166, n. 7-8, p. 449-458, 2018.
- ZENG, J.; ZHANG, Z.; LI, M.; WU, X.; ZENG, Y.; LI, Y. Distribution and molecular identification of *Meloidogyne* spp. parasitising flue-cured tobacco in Yunnan, China. *Plant Protection Science*, v. 54, n. 3, p. 183-189, 2018.
- ZHANG, F.; PENG, D.; YE, X.; YU, Z.; HU, Z.; RUAN, L.; SUN, M. *In vitro* uptake of 140 kDa *Bacillus thuringiensis* nematocidal crystal proteins by the second stage juvenile of *Meloidogyne hapla*. *Plos One*, v. 7, n. 6, e38534, 2012.
- ZHENG, Z.; ZHENG, J.; ZHANG, Z.; PENG, D.; SUN, M. Nematicidal spore-forming bacilli share similar virulence factors and mechanisms. *Scientific Reports*, v. 6, n. 1, e31341, 2016.
- ZUCKERMAN, B. M.; BRZESKI, M. W. Methods for the study of plant-parasitic nematodes in gnotobiotic root culture. *Nematologica*, v. 11, n. 4, p. 453-466, 1966.