



Revista mexicana de ciencias agrícolas

ISSN: 2007-0934

Instituto Nacional de Investigaciones Forestales, Agrícolas
y Pecuarias

Hernández-Hernández, Eyra Judith; Hernández-Ríos, Ismael; Almaraz-Suarez, Juan
José; López-López, Aline; Torres-Aquino, Margarita; Morales Flores, Francisco Javier
Caracterización in vitro de rizobacterias y su antagonismo con hongos causantes del damping off en Chile
Revista mexicana de ciencias agrícolas, vol. 9, núm. 3, Abril-Mayo, 2018, pp. 525-537
Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias

DOI: 10.29312/remexca.v9i3.335

Disponible en: <http://www.redalyc.org/articulo.oa?id=263158442003>

- Cómo citar el artículo
- Número completo
- Más información del artículo
- Página de la revista en redalyc.org

UAEM  redalyc.org

Sistema de Información Científica Redalyc

Red de Revistas Científicas de América Latina y el Caribe, España y Portugal

Proyecto académico sin fines de lucro, desarrollado bajo la iniciativa de acceso abierto

***In vitro* characterization of rhizobacteria and their antagonism with fungi that cause damping off in chili**

Eyra Judith Hernández-Hernández¹
 Ismael Hernández-Ríos¹
 Juan José Almaraz-Suarez²
 Aline López-López³
 Margarita Torres-Aquino^{1§}
 Francisco Javier Morales Flores¹

¹Postgraduate College-Campus San Luis Potosí. Agustín de Iturbide num. 73, Salinas de Hidalgo, SLP, Mexico. CP. 78600. Tel. 01 (496) 9630240. (hernandez.eyra@colpos.mx; ismaelhr@colpos.mx; franciscojmf@colpos.mx). ²Microbiology-Colegio de Postgraduados-Campus Montecillo. Mexico- Texcoco Highway km 36.5, Montecillo, Texcoco, Edo. from Mexico. CP. 56230. Tel. 01 (595) 9520200, ext. 1280 and 1269. (jalmaraz@colpos.mx). ³Genetics and Environment Research Center-Autonomous University of Tlaxcala. Freeway Tlaxcala-Tezmelucan km 10.5, Ixtacuixtla, Tlaxcala. CP. 90120. Tel. 01 (248) 4815500. (alopezlo@conacyt.mx).

§Corresponding author: maquino@colpos.mx.

Abstract

The production of chili seedlings (*Capsicum annuum* L.) in seedlings is affected by the disease known as “damping off”. Its prevention and treatment is done regularly by fungicides, whose effects on the fungi that cause this disease are not always effective, but they do represent environmental problems. Given this situation, biological control acquires greater relevance. In the present work three strains of phytopathogenic fungi of the genus *Fusarium* spp. (FP, FN and FA, by the colors purple, orange and yellow developed, respectively). As for bacterial strains, 18 strains of rhizobacteria (B) were selected, which were compared *in vitro* with the three strains of *Fusarium* and one of *Rhizotocnia* (R). Strain B23 was the most efficient in inhibiting the growth of the phytopathogens studied, reducing the growth of FP, FN and FA (50, 44 and 47%, respectively) and R (56%); the production of AIA varied between 1.24 and 11.32 $\mu\text{g mL}^{-1}$, the highest solubilization of inorganic phosphate (104 $\mu\text{g mL}^{-1}$) was obtained with B8. When evaluating the effect of B7, B9, B15 and B23 on the survival of chili seedlings inoculated with *Fusarium* sp. and *Rhizotocnia* sp., it was observed that the B7 and B23 strains were the ones that had the greatest effect on the survival of the seedlings during the first 34 days, with values of between 80 to 100%, while in the control seedlings it was between 30 and 50%. On the other hand, strains B8 and B9 were identified molecularly as *Pseudomonas chlororaphis*; B7 and B14 as *Pseudomonas* sp.; B15 as *Pseudomonas protegens* and B23 as *Bacillus* sp. The presented results show the potential of the use of bacteria as growth promoters and biocontrol agents of fungi causing the damping off in “guajillo” type chili seedlings.

Key words: *Capsicum annuum*, *Fusarium*, *Rhizotocnia*, damping off, rhizobacteria.

Reception date: February 2018

Acceptance date: March 2018

Introduction

In Mexico, the disease known as early drying or “damping off” is considered the main phytosanitary problem in the seedling phase of the pepper crop (Velásquez-Valle *et al.*, 2007) and is attributed to a complex of pathogenic fungi that includes to *Phytophthora capsici*, *Fusarium* spp., *Rhizoctonia solani*, *Sclerotium rolfsii* and *Pythium* sp. (Velásquez-Valle *et al.*, 2013). Seedlings can be affected in two ways: wilting of the plant as a consequence of rotting of its roots and involvement at the base of the stem (neck); both symptoms cause the collapse of the seedlings (Goldberg, 1995). The incidence of diseased seedlings in nurseries can reach up to 29% and constitute a source of inoculum that farmers transfer to the field during the transplant; crop losses in nursery, field and harvest can vary between 10 to 100% (Montero-Tavera *et al.*, 2013).

The use of chemical fungicides for the control of phytopathogens is not always effective (Huang *et al.*, 2012), in addition to causing pollution and harmful effects for human health and the environment (Gomathi and Ambikapathy, 2011). An alternative for the control of diseases caused by these fungi is the use of plant growth promoting rhizobacteria (PGPR) (Kloepper and Schroth 1981), which do not have negative effects on the environment (Mehetre and Kale *et al.*, 2011).

The PGPR can act as: a) biocontrol agents of diseases, through competition for space or food, or through the production of antibiotics and induction of plant defense systems (Matar *et al.*, 2009), decreasing or preventing disease (Saravanakumar *et al.*, 2008); b) growth promoters, through the production of growth regulators, as in the case of auxin-3-acetic acid (AIA) (Goswami *et al.*, 2015), synthesized by approximately 80% of the rhizobacteria (Ahmad *et al.*, 2005), having as an effect a greater nutrient uptake and consequently promoting the growth and yield of the crops; and c) inorganic phosphate solubilizers through the secretion of phosphatases and organic acids that release the unavailable mineral or organic P, estimating that 40% of the PGPR present this capacity (Richardson *et al.*, 2011).

The objective of this study was to characterize *in vitro* 18 bacterial isolates based on their ability to inhibit the growth of four fungal wilt fungi in chili pepper seedlings, their ability to produce indoleacetic acid and the solubilization of inorganic phosphate.

Materials and methods

Microbial isolations. Sampling of rhizospheric maize soil (*Zea mays* L.) established in an agroforestry module (22° 43' 08.4" north latitude, 101° 48' 17.7" west longitude) located in The Palma Pegada, municipality of Salinas, SLP was carried out, with an elevation of 2 222 meters above sea level, where the previous crop was chili. Isolation of the strains was performed as described by Kavitha *et al.* (2005), but the culture medium used was yeast-glucose-agar peptone (YPGA) (De Leon *et al.*, 2008), incubating at 28 °C for 24 h. Subsequently, the morphological and cultural characterization was carried out in a period of 24 h, where the color, shape, elevation and edge of the colony was considered.

On the other hand, in seedlings of “guajillo” chili seedlings located in La Dulce Grande (23° 00' 00.1" north latitude 102° 10' 29.1" west longitude), municipality of Villa of Ramos, SLP, a sampling of seedlings that had symptoms of “damping off”. In the laboratory of the Postgraduate College, Campus San Luis Potosi, the roots and neck of diseased seedlings were washed; they were then dissected in small portions (2 mm length) and placed in a culture medium potato-dextrose-agar (PDA, DIBICO) at 28 °C for seven days, after which the purification of the fungal colonies was carried out, obtaining three strains belonging to the genus *Fusarium* spp. (F). Additionally, a strain of *Rhizoctonia* spp. (R) provided by the Edaphology Laboratory of the Postgraduate College, Campus Montecillo. The strains were conserved in PDA.

Determination of *in vitro* microbial antagonism. Reactivation of bacterial and fungal isolates was performed in YPGA and PDA media, respectively. To evaluate the antifungal activity of 18 bacterial isolates against four pathogenic fungi, the dual culture method was used (Coskuntuna and Özer, 2008). The one-day-old bacterial isolates were placed 2 cm from the periphery of the Petri dish, each in triplicate, and incubating at 28 °C for 24 h; after that period, a 5 mm diameter disc with fungal mycelium was placed at a distance of 4 cm from the bacteria. The cultures were incubated at 28 °C, until the control (without bacteria) filled the box. The results were expressed as the percentage of inhibition corresponding to each of the bacterial strains. The inhibition percentage was calculated according to Mehetre and Kale (2011).

Production of indoleacetic acid (AIA). The Luria-Bertani liquid medium (Bric *et al.*, 1991) modified according Goswami *et al.* (2014), which was incubated at 25 ±2 °C in rotating agitation (New Brunswick Scientific Edison) for five days. The 2 ml of each isolate was taken and centrifuged at 8 000 rpm; the supernatant of each culture was mixed with the Salkowski reagent (Balota *et al.*, 1998) in a ratio of 2:1 and incubated in the dark for 30 min, to show the characteristic color development (pink-violet). Finally, the absorbance of the colored complex was read at 530 nm in a spectrophotometer (Bio Tek Laboratory Instrument, Synergy 2, USA). The concentration of AIA was determined by means of a standard curve in a range of 0 to 300 µg mL⁻¹. For each isolation, three replications were evaluated using as reference the culture medium without bacterial inoculum.

Solubilization of inorganic phosphorus. The 18 bacterial strains were inoculated in flasks containing 20 ml of liquid culture medium Pikovskaya (Pikovskaya, 1948), which had insoluble tricalcium phosphate [Ca₃(PO₄)₂] as the sole source of phosphorus, and incubated at 25 ±2 °C in constant agitation for 7 days. The 2 ml of culture were taken from each vial and centrifuged at 8 000 rpm. The soluble phosphorus of the supernatant was estimated by the blue colorimetric method of molybdenum (Jackson, 1967) at 480 nm. The corresponding amount of soluble phosphate was calculated at 7 days of the growth of the culture through a standard curve of soluble phosphate in a range of 0 to 300 µg mL⁻¹ (Banerjee *et al.*, 2010).

Seedling survival. This part of the research was conducted under greenhouse conditions. The guajillo chili seed was seeded in a mixture of peat moss substrates and vermiculite (1:1) previously sterilized twice at 121° for 1 h, with an interval of 3 days. Seven days after the emergence, the seedlings were transplanted to uncel trays. A period of 5 days was given for the acclimation of the seedlings and the inoculation was carried out with 1 mL of bacterial culture containing 10⁹ cells mL⁻¹ of each bacterial strain (B7, B9, B15 and B23) selected from the *in vitro* study.

The bacteria were applied with a micropipette at the base of the stem of the seedlings. An interval of 4 days was given to perform the inoculation of the pathogenic fungus (Hassan *et al.*, 2015), the fungal mycelium came from a 7-day-old culture and was placed where the seedling roots began to develop. The survival of the seedlings was recorded and in the end the treatment with the greatest number of surviving plants was obtained, which was attributed to the effect of the inoculated antagonist bacteria against each phytopathogenic fungus.

Molecular identification of rhizobacteria and fungi. The isolation of genomic DNA from rhizobacteria B7, B8, B9, B14, B15 and B23 and fungi *F. purple*, *F. orange* and *F. yellow* from bacterial culture on YPGA nutritional agar for rhizobacteria and PDA for fungi was performed, using the extraction protocol CTAB and sodium acetate (LaMontagne *et al.*, 2002). Amplification and sequencing of genomic DNA was carried out at the Potosino Institute for Scientific and Technological Research (IPICYT). A 16S rRNA gene fragment was amplified using oligonucleotides 533F (Weisburg *et al.*, 1991) and 1391R (Turner *et al.*, 1999) in a Verity thermocycler for endpoint PCR (Applied Biosystems). The products obtained were sequenced with the dideoxynucleotide method labeled in the 3130 Genetic Analyzer sequencer (Applied Biosystems). The percent identity of the sequences of the 16S rRNA gene fragment with the type strains was determined using BLASTN from the GenBank database.

Statistical analysis. The behavior of the bacteria in the inhibition of fungal growth, the production of indoleacetic acid, as well as the solubilization of phosphorus as a reflection of the activity to reduce the “damping off” were analyzed identifying differences (at 95% reliability.) After to identify the relevant sources of variation using InfoStat (Di Rienzo, 2014).

Results and discussion

Microbial isolations. A total of 50 strains of rhizobacteria were isolated, of which 18 were selected taking their growth rate as a criterion in a period of 24 h. On the other hand, three fungal strains were isolated from chili seedlings, which belong to the genus *Fusarium*, based on their pigmentation of the fungal mycelium and the culture medium where they were grown, they were assigned the nomenclature FP, FN and FA (*Fusarium purple*, *F. orange* and *F. yellow*, respectively).

Determination of *in vitro* microbial antagonism. The results obtained in dual culture showed that strain B23 was statistically the most efficient, inhibiting the mycelial growth of FP (50.02%), FN (44.05%), FA (46.67%) and *Rhizoctonia* spp. (R; 55.56%) (Figure 1). These results are similar to those reported by Ríos-Velasco *et al.* (2016), who report a percentage of growth inhibition of *F. oxysporum* of 42 and 51.5% due to the effect of *B. methylotrophicus* and *B. amyloliquefaciens*, respectively, as well as those found by Mojica-Morin *et al.* (2009), when confronting *F. oxysporum* with the strains of *B. thuringiensis* GM-23 (43.02%) and HD-121 (42.04%) and observed higher percentages of inhibition for *R. solani* due to the effect of the GM-64 strains. (66.66%) and HD-203 (65.99%). On the other hand, Djordjevic *et al.* (2011), when evaluating 16 bacterial isolates as antagonists of *F. oxysporum*, report a higher inhibition percentage (70.98%) on the growth of this fungus with strain Ab23, with results similar to those found in the present study with the isolates Ab9 (59.74%), Ab17 (57.7%) and Ab1 (56.6%).

In Figure 1A it can be seen that strains B6; B13; B11; B22; B15; B9; B17 and B7 exert a statistically similar antagonistic effect on *Fusarium* purple (FP), but different from the rest of the bacterial strains studied; B23 inhibited FP in 50%. When contrasting the 18 bacterial strains with *Fusarium* orange (FO) it was found that strains B23 and B9 showed a percentage of inhibition of 44.05 and 42.86%, respectively, which was higher than the rest of the treatments (Figure 1B).

While strains B7 and B15 inhibited fungal growth of *Fusarium* yellow (FY) by 30%, and with strain B23 almost reached 50% (Figure 1C). On the other hand, percentages of *Rhizoctonia* growth inhibition of 47.7, 42.2 and 35.5% were found when comparing strains B7, B15 and B14 respectively (Figure 1D). Finally, strains B3, B4, B5, B10, B20, B24 had no inhibitory effect on the four phytopathogens studied, which indicates that these strains were not able to produce a sufficient amount of substances that can inhibit the growth of pathogens, so that a threshold that allows them to exhibit an antagonistic response is not reached (Figuroa-López *et al.*, 2016).




Figure 1. Inhibitory effect of 18 bacterial strains on the mycelial growth of four phytopathogenic fungi. A) *Fusarium* Purple, B) *Fusarium* Orange, C) *Fusarium* Yellow and D) *Rhizoctonia*. Negative values indicate growth stimulation.

It is well documented that the antagonistic effect of rhizobacteria on the inhibition of the growth of phytopathogens is due to the synthesis of secondary metabolites (Khannous *et al.*, 2014). Trivedi *et al.* (2008) showed that volatile metabolites produced by *Pseudomonas corrugata* play a predominant role in the inhibition of fungi *F. oxysporum* and *A. alternata*, while diffusible

metabolites play only a secondary role. On the other hand, it has been shown that some species of *Pseudomonas* produce lytic enzymes, which can break the cell membranes of some phytopathogenic fungi; the best known are: chitinases, cellulases, proteases and 1,3-glucanases (Compant *et al.*, 2005).

Production of indoleacetic acid (AIA). The production of AIA is a property of some rhizobacteria to stimulate and facilitate plant growth (Mohite, 2013). This auxin is a product derived from the metabolism of L-tryptophan produced by several soil microorganisms, including PGPR (Goswami *et al.*, 2015). In the present work, when analyzing the AIA production of 18 bacterial isolates from the maize rhizosphere, a variation of 1.3 to 11.2 $\mu\text{g mL}^{-1}$ was found. The production of AIA significantly higher ($p \leq 0.05$) corresponded to B14 and the lowest to B4 (Figure 2).

These results agree with those reported in several investigations, where the production of AIA in bacterial strains coming from the maize rhizosphere was evaluated. Noh *et al.* (2014) mention a production of 3.3 $\mu\text{g mL}^{-1}$ of AIA. On the other hand, Josic *et al.* (2012) point out that when adding 5mM of tryptophan to the King B liquid medium, the concentration of AIA increased from 8.6 to 26 $\mu\text{g mL}^{-1}$ in strain Q4m (*Pseudomonas* sp.). It has been observed that the production of AIA by rhizobacteria varies between different species and strains; which is influenced by the growing conditions, growth stage and substrate availability (Mirza *et al.*, 2001).




Figure 2. Production of indoleacetic acid from 18 bacterial isolates from the corn rhizosphere. Averages with different letters indicate significant differences (Tukey 0.05).

According to the production of AIA *in vitro*, the rhizobacteria can be divided into three groups: low (1 to 10 $\mu\text{g mL}^{-1}$), medium (11 to 20 $\mu\text{g mL}^{-1}$) and high producing (21 to 30 $\mu\text{g mL}^{-1}$) of AIA (Rojas *et al.*, 2010). According to this classification, most of the strains analyzed in the present study are classified as low producers of indoleacetic acid. However, not all bacteria that produce a high amount of AIA stimulate plant growth (Noh *et al.*, 2014). For example, *Agrobacterium tumefaciens* produces up to 78 $\mu\text{g mL}^{-1}$ of AIA, which induces the development of the disease known as “crown gall” (Lara *et al.*, 2011), while lower concentrations (1.1 to 12.1 $\mu\text{g mL}^{-1}$) increased the growth of wheat plants *in vitro* (Khalid *et al.*, 2004).

Ramírez-Malagon and Ochoa-Alejo (1996), in an in vitro study showed that the AIA has an effect on the induction and growth of adventitious buds of the pepper culture, being its concentration the one that determines the type of response. Peña *et al.* (2016) report an AIA production, between 4.0 to 24.3 $\mu\text{g mL}^{-1}$ in five *Bacillus* species, which is sufficient to promote the growth of *Capsicum annuum* L. Therefore, in the present study eight strains can be considered promising for promote the growth of the chili crop.

Solubilization of inorganic phosphorus. The amount of inorganic phosphate (Pi) dissolved in the culture medium varied among the different isolates (Figure 3). Strain B8 was the one that solubilized the highest amount of Pi (100.4 $\mu\text{g mL}^{-1}$), followed by strain B14, with which 80.92 $\mu\text{g mL}^{-1}$ and then B17 and B21 were recorded (60.67 and 56.39 $\mu\text{g mL}^{-1}$, respectively). The amount of phosphate solubilized by the rest of the strains, with the exception of B10 and B9 (null effect), was found in the range of 2.21 to 49.12 $\mu\text{g mL}^{-1}$. The results recorded in strains B8, B14 and B17 are superior to those reported by Nautiyal (1999), who found a variation of 8 to 35 $\mu\text{g mL}^{-1}$ in five species of *Pseudomonas* and from 8 to 17 $\mu\text{g mL}^{-1}$ in three *Bacillus* species.

It has been pointed out that bacteria solubilize Pi through the production of CO_2 , organic acids, proton excretion and NH_4^+ assimilation. The most important mechanism of solubilization of calcium phosphate is acidification through the biosynthesis and secretion of organic acids (Paredes and Espinosa, 2009). Such organic acids can directly dissolve the Pi as a result of the exchange of phosphate anions by an acid anion or by chelating the Fe, Al and Ca ions associated with P (Omar, 1998), which consequently increases the bioavailability of P, which can be exploited by the crops or the wild flora present.




Figure 3. Phosphate solubilized by bacterial isolates. Columns with different letters indicate significant differences (Tukey 0.05).

Seedling survival. When evaluating the effect of strains B7, B9, B15 and B23 on the survival of chili seedlings inoculated with *Fusarium* spp. (FA, FP and FN) and *Rhizotocnia* sp. (R), we observe in Figure 4 (a, b and d) 100% survival in seedlings inoculated with B23 and a survival in control plants of FP, FA and R of 80, 77.8 and 83%, respectively (only inoculated with the

pathogen) at eight days. In this same period, 100% survival was also observed when inoculating B9 and B15 for the case of FP (4a) and B15 for the case of *Rhizoctonia* (Figure 4d). It is observed that from 34 to 60 days, there is a decrease in the survival of the control seedlings in comparison with the seedlings inoculated with rhizobacteria. On the other hand, only strains B7 and B23 have an effect on the survival of FA infected seedlings up to 55 days (Figure 4b), this survival coincides with that reported by Abdel-Monaim (2013) who obtained 72% and 76.34% of survival by *Bacillus megaterium* in bean plants in two different seasons, whose plants were infested with the pathogens *R. solani*, *F. solani*, *F. oxysporum* and *M. phaseolina* causing damping off to the plant.




Figure 4. Effect of four antagonistic bacteria on the survival of “guajillo” chili seedlings inoculated with different phytopathogenic fungi. a) *Fusarium Purple*; b) *F. yellow*; c) *F. orange*; and d) *Rhizoctonia*.

With respect to the fungus FN (Figure 4c), both in control seedlings and in the seedlings inoculated with B7 and B23, 100% survival was recorded. This same trend is maintained until 34 days for the case of control seedlings and those inoculated with B7. The behavior of the control seedlings could possibly be due to the fact that the amount of fungal mycelium was not enough for this fungus to express virulence in the first 34 days or that its effect to cause the disease is later than the other pathogens used in the present study.

Finally, for the case of *Rhizotocnia* spp., Strain B7 had an effect on the survival of the seedlings, because it was 80% until 55 days, while the control seedlings were 41.4% (Figure 4d). In this regard, Toppo and Tiwari (2015), report a variation in the survival of tomato seedlings (of 30

days) infected with *Rhizoctonia* which varied between 93.33 to 46.67.6% when inoculating *Pseudomonas alcaligenes* PKJ25 and *Pseudomonas* PKS10. In the present study, at 34 days, the highest survival was 83.3 and the lowest was 46.2% when inoculating B7 and B23, respectively.

Molecular identification of rhizobacteria and fungi. Isolates B8 and B9 were identified as *Pseudomonas chlororaphis*; B7 and B14 as *Pseudomonas* sp. and B15 as *Pseudomonas protegens*. For its part, B23 was identified as *Bacillus* sp. All presented an identity 99%, with a coverage of 100%. The genus *Pseudomonas* is composed of gram-negative bacteria, which belong to the γ -Proteobacteria, are distributed in a wide variety of environments and are known for their metabolic versatility in the use of organic compounds as carbon and energy sources (Lessie and Phibbs, 1984; Wu *et al.*, 2011) and in the production of various secondary metabolites (Gross and Loper, 2009). *Pseudomonas* is one of the most complex bacterial groups, with more than 100 species described so far, in which there is the presence of groups and subgroups.

The group of *Pseudomonas fluorescens* contains several species (*P. brassicacearum*, *P. protegens*, *P. chlororaphis* and *P. fluorescence*), described as PGPR for its ability to suppress disease-causing pathogens (Raajimakers *et al.*, 2009) and produce antibiotics or secondary metabolites (Gross and Loper, 2009) that make them very viable for biotechnological applications (Saravanakumar *et al.*, 2008). Bacterial identification through the 16S ribosomal gene has been widely used; however, it is well known that nearby bacterial species cannot be differentiated by the use of this gene due to its high conservation rate (Fox *et al.*, 1992; Kämpfer and Glaeser, 2012), so the use of the analysis of multilocus sequences by the use of other genes as molecular markers: *rpoB*, *rpoD*, *gyrB* (Yamamoto *et al.*, 2000; Ait Toyeb *et al.*, 2005), allowing identification up to the species level.

In this work, by using the 16S rRNA gene as the only molecular marker, it was possible to designate the obtained isolates, which belong to *Pseudomonas* and *Bacillus*, up to gender level. Members of these genera have been reported as PGPR, with different mechanisms such as production of phytohormones, suppression of pathogens or producers of secondary metabolites of biotechnological interest.

On the other hand, FP and FN fungal isolates correspond to *Fusarium* sp.; the first with 93% identity and coverage 84% and the second with 86% identity and 80 coverage. *F. yellow* corresponds to a non-cultivated fungus, with 94% identity and 33% coverage. *Fusarium* sp belongs to the family *Nectriaceae*. The genus *Fusarium* is a group of filamentous fungi widely distributed in soil and plants. Due to their ability to grow at 37 °C, they are considered opportunistic (Tapia and Amaro, 2014). Santos (2010) mentions *Fusarium* sp. as one of the main causal agents of the wilt of the chili that diminishes the yield of the crop.

Conclusions

Strain B23 was found to be the most efficient in inhibiting the growth of the four pathogenic fungi evaluated, *Rhizoctonia* (56%), *F. purple* (50%), *F. yellow* (47%) and *F. orange* (44%), whereas strain B14 was the most outstanding in the production of IAA (11.2 $\mu\text{g mL}^{-1}$). On the other hand,

strain B8 showed the highest solubilization of phosphate ($104.4 \mu\text{g mL}^{-1}$). The Bacteria isolated from the rhizosphere of maize, in a plot with previous cultivation of chili, have antagonistic capacity against four strains of phytopathogenic fungi, in addition to showing growth promoting activity (production of AIA and solubilization of inorganic phosphorus).

On the survival of chili seedlings, strain B7 maintained the highest survival of seedlings in time, while those inoculated with B15 were those that died in greater proportion and in less time. The results of this study suggest that diseases caused by *Fusarium* or *Rhizoctonia* can be effectively managed through the use of antagonistic bacteria such as B23, or the promotion of plant growth through the use of bacterial strains such as B14 and B8, decreasing damage to the environment and health. In addition, the use of phosphate solubilizing microorganisms as an inoculant in seeds and seedlings can represent a biotechnological alternative to increase the availability of phosphorus for crops.

Cited literature

- Abdel, M. M. F. 2013. Improvement of biocontrol of damping-off and root rot/wilt of Faba bean by salicylic acid and hydrogen peroxide. *Mycobiology*. 41(1):47-55.
- Ahmad, F.; Ahmad, I. and Khan, M. S. 2005. Indole acetic acid production by the indigenous isolated of *Azotobacter* and fluorescent *Pseudomonas* in the presence and absence of tryptophan. *Turk. J. Biol.* 29:29-34.
- Ait, T. L.; Ageron, E.; Grimont, F. and Grimont P. A. 2005. Molecular phylogeny of the genus *Pseudomonas* based on *rpoB* sequences and application for the identification of isolates. *Res Microbiology*. 156(5-6):763-773.
- Balota, E. L.; Colozzi, F. A.; Andrade, D. S. e Hungria, M. 1998. Biomassa microbiana e sua atividade em solos sob diferentes sistemas de preparo e sucessã ode culturas. *Rev. Bras. Ciencia do Solo*. 22:641-649.
- Banerjee, S.; Palit, R.; Sengupta, C. and Standing, D. 2010. Stress induced phosphate solubilization by *Arthrobacter* sp. and *Bacillus* sp. isolated from tomato rhizosphere. *Austr. J. Crop Sci.* 4(6):378-383.
- Bric, J. M.; Bostock, R. M. and Silverstone S. E. 1991. Rapid in situ assay for indoleacetic acid production by bacteria immobilized on a nitrocellulose membrane. *Appl. Environ. Microbiol.* 57(2):535-538.
- Compant, S.; Duffy B.; J. Nowak, C. Clément and Barka E. A. 2005. Use of plant growth-promoting bacteria for biocontrol of plant diseases: principles, mechanisms of action, and future prospects. *Appl. Environ. Microbiol.* 71:4951-4959.
- Coskuntuna, A. and Özer, N. 2008. Biological control of onion basal rot disease using *Trichoderma harzianum* and induction of antifungal compounds in onion set following seed treatment. *Crop Prot.* 27:330-336.
- De León, L.; Rodríguez, A.; López, M. M. and Siverio, F. 2008. Evaluation of the efficacy of immunomagnetic separation for the detection of *Clavibacter michiganensis* subsp. *michiganensis* in tomato seeds. *J. Appl. Microbiol.* 104:776-786. doi:10.1111/j.1365-2672.2007.03595.x.
- Di Rienzo, J. A.; Casanoves, R. F. M. G.; Balzarini, B.; González, M. L.; Tablada T. M. y Robledo S. C. W. 2014. InfoStat versión 2014. Grupo InfoStat, FCA. Universidad Nacional de Córdoba, Argentina. <http://www.infostat.com.ar>.

- Djordjevic, M.; Ugrinovic, M.; Sevic, M.; Djordjevic, R. and Mijatovic, M. 2011. Antagonistic effect of soil bacteria against fusarium wilt of pepper *in vitro*. *Acta Agriculturae Serbica*. 16:19-31.
- Figueroa, L. A. M.; Cordero, R. J. D; Martínez, A. J. C; López, M. M; Lizárraga, S. G. J; Félix, G. R; Castro, M. C. and Maldonado, M. I. E. 2016. Rhizospheric bacteria of maize with potential for biocontrol of *Fusarium verticillioides*. Springer Plus. DOI: 10.1186/s40064-016-1780-x.
- Fox, G. E.; Wisotzkey, J. D. and Jurtshuk, P. Jr. 1992. How close is close: 16S rRNA sequence identity may not be sufficient to guarantee species identity. *Inter. J. System. Bacteriol.* 42(1):166-170.
- Goldberg, N. P. 1995. Chile pepper diseases. Agricultural Experiment Station. College of Agriculture and Home Economics. New Mexico State University. Circular 549. Las Cruces, NM, USA. 20 p.
- Gomathi, S. and V. Ambikapathy. 2011. *Adv App Sci Res.* 2:291-297.
- Goswami, D., Dhandhukia P., Patel P. and Thakker J. N. 2014. Screening of PGPR from saline desert of Kutch: Growth promotion in *Arachis hypogea* by *Bacillus licheniformis* A2. *Microbiol. Res.* 169:66-75.
- Goswami, D.; Thakker, J. N. and Dhandhukia, P. C. 2015. Simultaneous detection and quantification of indole-3-acetic acid (IAA) and indole-3-butyric acid (IBA) produced by rhizobacteria from L-tryptophan (Trp) using HPTLC. *J. Microbiol. Methods.* 110:7-14.
- Gross, H. and Loper, J. E. 2009. Genomics of secondary metabolite productions by *Pseudomonas* spp. *Natural Proceedings Reports.* 26(11):1408-46
- Hassan, D. Gh.; Hassan, M. G.; Rashid, H.; Ali, D. W. and Majeed, M. 2015. Evaluation of microbial antagonists for the management of wilt/root rot and damping-off diseases in chilli (*Capsicum annuum*). *Rev. Vegetos.* 28(4):102-110.
- Huang, X.; Zhang, N.; Yong, X.; Yang, X. and Shen, Q. 2012. Biocontrol of *Rhizoctonia solani* damping-off disease in cucumber with *Bacillus pumilus* SQR-N43. *Microbiol. Res.* 167:135-143.
- Jackson, M. L. 1967. Soil chemical analysis. (Ed). Prentic Hall, Inc., Engle Wood Cliff. USA.
- Josic, D.; Delic, D.; Rasulic, N.; Stajkovic, O.; Kuzmanovic, D.; Stanojkovic, A. and Pivic, R. 2012. Indigenous pseudomonads from rhizosphere of maize grown on pseudogley soil in Serbia. *Bulgarian J. Agric. Sci.* 18(12):197-206.
- Kavitha, K.; Mathiyazhagan, S.; Senthilvel, V.; Nakkeeran, S. and Chandrasekar, G. 2005. Development of bioformulations of antagonistic bacteria for the management of damping off of Chilli (*Capsicum annuum* L.). DOI: 10.1080/0323540040000838.
- Kämpfer, P. and Glaeser, S. P. 2012. Prokaryotic taxonomy inn the sequencing era-the poluphasic approach revisited. *Environ. Microbiol.* 14(2):291-317.
- Khalid, A.; Arshad M. and Zahir, Z. A. 2004. Screening plant growth-promoting rhizobacteria for improving growth and yield of wheat. *J. Appl. Microbiol.* 96:473-480.
- Khannous, L.; Mouna, J.; Mouna, D.; Ramzi, M.; Nour, C.; Bassem, K.; Néji G. and Imen, F. 2014. Isolation of a novel amylase and lipase-producing *Pseudomonas luteola* strain: study of amylase production conditions. *Lipids in Health and Disease.* 13:9.
- Kloepper, J. W. and Schroth M. N. 1981. Plant growth-promoting rhizobacteria and plant growth under gnotobiotic conditions. *Phytopatology.* 71:642-644.
- LaMontagne, M. G.; Michel, Jr. F. C.; Holden, P. A. and Reddy, C. A. 2002. Evaluation of extraction and purification methods for obtaining PCR-amplifiable DNA from compost for microbial community analysis. *J. Microbiol. Methods.* 49(3):255-264.

- Lessie, T. G. and Phibbs, P. V. Jr. 1984. Alternative pathways of carbohydrate utilization in Pseudomonads. Annual Rev. Microbiol. 38:359-388.
- Lara, M. C.; Oviedo Z. L. y Betancur H. C. A. 2011. Bacterias nativas con potencial en la producción de ácido indolacético para mejorar los pastos. Zootec. Trop. 29:187-194.
- Matar, S. M.; El-Kazzaz, S. A.; Wagih, E. E.; El-Diwan, A. I.; Moustafa, H. E.; Abo-Zaid, G. A.; Abd-Elsalam, H. E. and Hafez, E. E. 2009. Antagonistic and inhibitory effect of *Bacillus subtilis* against certain plant pathogenic fungi, I. Biotechnology. 8(1):53-61.
- Matar, S. M.; El-Kazzaz, S. A.; Wagih, E. E.; El-Diwan, A. I.; Moustafa, H. E.; Abo-Zaid, G. A.; Abd-Elsalam, H. E. and Hafez E. E. 2009. Antagonistic and inhibitory effect of *Bacillus subtilis* against certain plant pathogenic fungi, I. Biotechnology. 8(1):53-61.
- Mojica, M. V; Luna, O. H. A; Sandoval, C. C. F.; Pereyra, A. B; Morales, R. L. H.; González, A. N. A.; Hernández, L. C. E. y Alvarado, G. O. G. 2009. Control biológico de la marchitez del chile (*Capsicum annuum* L.) por *Bacillus thuringiensis*. Phytom. 78(2):105-110.
- Mehetre, S. T. and Kale, S. P. 2011. Comparative efficacy of thermophilic bacterium, *Bacillus licheniformis* (NR1005) and antagonistic fungi, *Trichoderma harzianum* to control *Pythium aphanidermatum*-induced damping off in chilli (*Capsicum annuum* L.). Arch. Phytopathol. Plant Protec. 44(11):1068-1074.
- Mirza, M. S.; Ahmad, W.; Latif, F.; Haurat, J.; Bally, R.; Normand P. and Malik, K. A. 2001. Isolation, partial characterization, and the effect of plant growth-promoting bacteria (PGPB) on micro-propagated sugarcane *in vitro*. Plant Soil 237:47-54.
- Mohite, B. 2013. Isolation and characterization of indole acetic acid (IAA) producing bacteria from rhizospheric soil and its effect on plant growth. J. Soil Sci. Plant Nutr. 13(3):638-649.
- Montero, T. V.; Guerrero, A. B. Z.; Anaya, L. J. L.; Martínez, M. T.O.; Guevara, O. L. y González, C. M. M. 2013. Diversidad genética de aislados de *Rhizoctonia solani* (Kuhn) de Chile en México. Rev. Mex. Cienc. Agríc. 4(7):1043-1054.
- Nautiyal, C. S. 1999. An efficient microbiological growth medium for screening phosphate solubilizing microorganisms. FEMS Microbiology Letters. 170:265-270.
- Noh, M. J.; Zúñiga, A. J. J.; Borges, G. L.; Yam, C. C. y Godoy, H. G. 2014. Aislados bacterianos con potencial biofertilizante para plántulas de tomate. Terra Latinoam. 32(4):273-281
- Omar, S. A. 1998. The role of rock phosphate solubilizing fungi and vesicular arbuscular mycorrhiza (VAM) in growth of wheat plants fertilized with rock phosphate. World J. Microbiol. Biotechnol. 14:211-219.
- Paredes, M. M. y Espinosa, V. D. 2009. Ácidos orgánicos producidos por rizobacterias que solubilizan fosfato: una revisión crítica. Terra Latinoam. 28(1):61-70.
- Peña, Y. L. P.; Ruíz, S. E.; Barbosa, C. J. E. and Reyes, R. A. 2016. Isolation of Mexican *Bacillus* species and their effects in promoting growth of chili pepper (*Capsicum annuum* L. cv. Jalapeño). Indian J. Microbiol. 56(3):375-378. DOI 10.1007/s12088-016-0582-8.
- Pikovskaya, R. I. 1948. Mobilization of phosphorus in soil in connection with the vital activity of some microbial species. Mikrobiologiya. 17:362-370.
- Ramírez, M. R. and Ochoa, A. N. 1996. An improved and reliable chili pepper (*Capsicum annuum* L.) plant regeneration method. Plant Cell Rep. 16:226-231.
- Raajimakers, J. M.; Paulitz, T.C.; Steinberg, C.; Alabouvette, C. and Moënne, L. Y. 2009. The rhizosphere: a playground and battlefield for soilborne pathogens and beneficial microorganisms. Plant Soil. 321(1-2):20.

- Richardson, A. E.; Lynch, J. P.; Ryan, P. R.; Delhaize, M.; Smith, F. A.; Smith, S. E.; Harvey, P. R.; Ryan, M. H.; Veneklaas, E. J.; Lambers, H.; Oberson, A.; Culvenor, R. A. and Simpson, R. J. 2011. Plant and microbial strategies to improve the phosphorus efficiency of agriculture. *Plant Soil* 349:121-156. DOI 10.1007/s11104-011-0950-4.
- Rios, V. C.; Caro, C. J. N.; Berlanga, R. D. I.; Ruíz, C. M. F.; Ornelas, P. J. J.; Salas, M. M. A.; Villalobos, P. E. and Guerrero, P. V. M. 2016. Identification and antagonistic activity in vitro of *Bacillus* spp. and *Trichoderma* spp. isolates against common phytopathogenic fungi. *Rev. Mex. Fitopatol.* 34:84-99.
- Rojas, C. A.; Rodríguez, D. A. M.; Montes, V. S. y Pérez J. S. 2010. Evaluación de la promoción de crecimiento de *Cynodon dactylon* L. por rizobacterias productoras de fitohormonas aisladas de un suelo contaminado con hidrocarburos derivados del petróleo. *Rev. Polibotánica.* 29:1-147.
- Santos, J. P. 2010. Estrategias para el control de *Phytophthora capsici* Leo. y *Fusarium solani* Mart. en el cultivo de chile (*Capsicum annum* L.). Tesis de maestría. Colegio de Postgraduados.
- Saravanakumar, D.; Lavanya, N.; Muthumeena, B.; Raguchander, T.; Suresh S. and Samiyappan R. 2008. *Pseudomonas fluorescens* enhances resistance and natural enemy population in rice plants against leaf folder pest. *J. Appl. Entomol.* 132:469-479.
- Tapia, C. y Amaro J. 2014. Género *Fusarium*. *Rev Chilena Infectol.* 31(1):85-86.
- Trivedi, P.; Pandey, A. and Palni, L. M. S. 2008. In vitro evaluation of antagonistic properties of *Pseudomonas corrugata*. *Microbiol. Res.* 163:329-336.
- Turner, S.; Pryer, K. M.; Miao, V. P. y Palmer, J. D. 1999. Investigating deep phylogenetic relationships among cyanobacteria and plastids by small subunit rRNA sequence analysis. *J. Eukaryotic Microbiol.* 46(4):327-338.
- Toppo, S. R. and Tiwari, P. 2015. Biocontrol potentialities of native *Pseudomonas* isolates against plant pathogenic fungi *Rhizoctonia* spp., *Fusarium* spp. and *Colletotricum* spp. in tomato rhizosphere under greenhouse condition. *The Bioscan.* 10(1):373-377.
- Velásquez, V. V. R.; Amador R. M. D.; Medina, A. M. M. y Lara V. F. 2007. Presencia de Patógenos en Almácigos y Semilla de Chile (*Capsicum annum* L.) en Aguascalientes y Zacatecas, México. *Rev. Mex. Fitopatol.* 25(1):75-79.
- Velásquez, V. V. R.; Reveles, T. L. R. y Reveles, H. M. 2013. Manejo de las principales enfermedades del chile para secado en el norte centro de México. *Campo Experimental Zacatecas. CIRNOC-INIFAP. Folleto técnico* núm. 50. 57 p.
- Weisburg, W. G.; Barns, S. M.; Pelletier, D. A. and Lane, D. J. 1991. 16S ribosomal DNA amplification for phylogenetic study. *J. Bacteriol.* 173(2):697-703.
- Wu, X.; Monchy, S.; Taghavi, S.; Zhu, W.; Ramos, J. and van der Lelie, D. 2011. Comparative genomics and functional analysis of niche-specific adaptation in *Pseudomonas putida*. *FEMS microbiology Reviews.* 35(2):299-323.
- Yamamoto, S.; Kasai, H.; Arnold, D. L.; Jackson, R. W.; Vivian, A. y Harayamas, S. 2000. Phylogeny of the genus *Pseudomonas*: intergenic structure reconstructed from nucleotide sequences of *gyrB* and *rpoD* genes. *Microbiology.* 146:2385-2394.