



Surgical & Cosmetic Dermatology

ISSN: 1984-8773

Sociedade Brasileira de Dermatologia

Dehó, Isabella Zurita; Oliveira Filho, Jayme de; Afonso,
Irina Andrea Pires; Oliveira, Anna Rita Ferrante Mitidieri de
Unexpected riziform bodies
Surgical & Cosmetic Dermatology, vol. 14, e20220057, 2022
Sociedade Brasileira de Dermatologia

DOI: <https://doi.org/10.5935/scd1984-8773.2022140057>

Available in: <https://www.redalyc.org/articulo.oa?id=265570220005>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in [redalyc.org](https://www.redalyc.org)

redalyc.org

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative



Unexpected riziform bodies

Corpos riziformes em localização inusitada

DOI: <http://www.dx.doi.org/10.5935/scd1984-8773.2022140057>

ABSTRACT

Riziform bodies are structures in the synovial fluid or attached to the synovium, which macroscopically resemble rice grains. They are common in patients with rheumatoid arthritis and result from a chronic inflammation. They are found in 25% of aspiration procedures or joint surgery. However, there are still no publications reporting its presence in the subcutaneous tissue. We report for the first time the occurrence of riziform bodies in the subcutaneous plane of the palmar region in a patient with rheumatoid arthritis, confirmed by sonographic and anatomopathological examination, successfully treated with surgical drainage, without recurrences after two years of follow-up.

Keywords: Abscess; Rheumatoid arthritis; Dermatopathies; Hand dermatoses; Outpatient Surgical Procedures; Surgical Diagnostic Techniques

RESUMO

Corpos riziformes são estruturas que podem ser encontradas no líquido sinovial ou aderidas à sinóvia, que se assemelham macroscopicamente a grãos de arroz. São frequentes em pacientes com artrite reumatoide e decorrem de resposta inflamatória crônica. São encontrados em 25% dos procedimentos de aspiração ou cirurgias das articulações. Porém, ainda não existem publicações relatando sua presença no subcutâneo. Relatamos, pela primeira vez, a ocorrência de corpos riziformes no subcutâneo da região palmar em paciente com artrite reumatoide, confirmados por exames ultrassonográfico e anatomopatológico, tratados com sucesso com drenagem cirúrgica, sem recidivas após dois anos de seguimento.

Palavras-chave: Abscesso; Artrite reumatoide; Dermatopatias; Dermatoses da mão; Procedimentos cirúrgicos ambulatoriais; Técnicas de diagnóstico por cirurgia

Case Report

Authors:

Isabella Zurita Dehó¹
Jayme de Oliveira Filho²
Irina Andrea Pires Afonso²
Anna Rita Ferrante Mitidieri de Oliveira¹

- ¹ Hospital Sírio Libanês, Department of Dermatology, São Paulo (SP), Brasil.
- ² Tez Dermatologia, Department of Dermatology, São Paulo (SP), Brasil.

Correspondence:

Isabella Zurita Dehó
Email: isazdeho@gmail.com / isazdeho@gmail.com

Financial support: None.

Conflict of interest: None.

Submitted on: 27/05/2021

Approved on: 04/06/2021

How to cite this article: Dehó IZ, Oliveira Filho J, Afonso IAP, Oliveira ARFM. Unexpected riziform bodies. Surg Cosmet Dermatol. 2022;14:e20220057.



INTRODUCTION

Riziform bodies are structures found in the synovial fluid or adhered to the synovia. They receive this name because they macroscopically resemble rice grains.¹ These agents occur due to a chronic and nonspecific synovial inflammatory response.² They are composed of an acidophilic center of collagen types I, II, and IV, encapsulated in the periphery by fibrin.^{3,4}

These structures were described for the first time in 1895 in a patient with tuberculous arthritis.¹ They were later described in association with other diseases, such as rheumatoid arthritis (RA), juvenile idiopathic arthritis (JIA), osteoarthritis, tenosynovitis associated with brucellosis, joint infection by mycobacteria and fungi, or even idiopathically, in patients without comorbidities.^{1,3}

They are currently more frequently associated with rheumatoid arthritis than seronegative arthritis. Its removal ensures clinical improvement for patients, reduction of synovitis, and the possibility of investigating infectious causes.⁵

Riziform bodies are found in 25% of aspiration procedures or joint surgeries.^{3,4} However, there are still no publications reporting their presence in the subcutaneous plane, even in patients with rheumatoid arthritis. In this article, we report, for the first time, the occurrence of riziform bodies in the subcutaneous tissue of the palmar region in a patient with rheumatoid arthritis, confirmed by ultrasound and pathological examinations. They were successfully surgically drained, with no recurrences after a two-year follow-up.

CASE REPORT

A 67-year-old woman, skin phototype II, born and resident in São Paulo, sought medical care for three weeks complaining of pain and difficulty opening her right hand (Figure 1). Dermatological examination revealed an erythematous tumor, poorly delimited, measuring 2.7 cm in its longest axis, surmounted by hyperkeratotic areas in the right palmar region, at the level of the metacarpophalangeal joints, between the second and fourth fingers (Figures 2A and 2B).

She had a personal history of hepatitis C, *diabetes mellitus*, sickle cell anemia, rheumatoid arthritis, and kidney transplantation 26 years ago. The patient was in use of ribavirin, NPH human insulin, azathioprine, folic acid, B complex, alpha epoetin, and prednisone 5 mg/d.

The differential diagnoses included cutaneous abscess, tenosynovitis, epidermal inclusion cyst, adnexal tumor, and Dupuytren's contracture.

Given that the clinical characteristics suggested the diagnosis of abscess, we chose treatment with oral ciprofloxacin 500 mg 12/12h for ten days and requested exams, including imaging. After a week, the patient returned with the results of the exams, and the lesion still looked the same.

The ultrasound of the right hand showed tendons with an extensive circumscribed collection, walls, and hypoechoic content associated with thick suspension debris surrounding the tendons (Figure 3).



FIGURE 1: Contracture present in the first consultation




FIGURE 2: **A** - Dermatological examination: erythematous tumor, poorly delimited, surmounted by hyperkeratotic areas in the region of the metacarpophalangeal joints of the right hand. **B** - Oblique view: erythematous tumor surmounted by hyperkeratotic crust. **C** - Surgical drainage: visualization of multiple granular structures, pearlescent, with fibroelastic consistency and smooth surface. **D** - Drainage of the entire content of the lesion




FIGURE 3: **A** - Ultrasonography of the right palmar region: perpendicular section showing a circumscribed hypoechoic collection. **B** - Ultrasonography of the right palmar region: tendons with extensive circumscribed collection of hypoechoic content associated with debris




FIGURE 4: **A** - Surgical drainage: visualization of multiple granular structures, pearlescent, with fibroelastic consistency and smooth surface. **B** - Drainage of the entire content of the lesion

We conducted the surgical drainage of the lesion. During the procedure, we observed the presence of a fibrous nodule adhered to the deep planes, associated with the emergence of multiple granular, cystic structures, with fibroelastic consistency and smooth surface (Figures 4A and 4B).

The material was sent to the anatomopathological exam (Figures 5A, 5B, 5C, and 5D), which showed granular material represented by hyalinized collagen, intermingled with cells with elongated or ovoid nuclei and rare mononuclear elements, and presence of foci of dystrophic calcification. In the periphery of the lesions, we observed the presence of fibrin and macrophages,




FIGURE 5: **A** - Material sent for anatomopathological study.

B - Anatomopathological: general view. **C** - Image with microscopic magnification of 40 times, showing hyalinized collagen intermingled with cells with elongated or ovoid nuclei and rare mononuclear elements. Presence of foci of dystrophic calcification. In the periphery of the lesions, there is the presence of fibrin and macrophages.

D - 70-fold magnification on optical microscopy, showing hyalinized collagen intermingled with cells with elongated or ovoid nuclei and rare mononuclear elements. Presence of dystrophic calcification foci. In the periphery of the lesions, there is the presence of fibrin and macrophages




FIGURE 6: A - Follow-up three weeks after the procedure. B - Follow-up two months after surgical drainage. C - Image after two years of follow-up, showing no recurrence in the site or in other joints

some of them phagocytizing a light brown pigment with the characteristics of hemosiderin. There was an absence of signs of malignancy. These features confirmed the diagnosis of “riziform bodies”.

One month after the drainage, the patient returned for consultation, presenting clinical improvement; however, she still had difficulty flexing her fingers (Figure 6A).

Two months after the drainage, the patient returned with significant clinical improvement, including the ability to flex her fingers, which was previously difficult to perform (Figure 6B).

The patient evolved with complete wound healing and no recurrence after a two-year follow-up (Figure 6C).

This article presents the complete clinical picture and the follow-up of the lesions within one month, two months, and two years after the procedure.

DISCUSSION

Riziform bodies are prevalent in patients with rheumatoid arthritis. Previous studies have already demonstrated its presence in 72% of the joint effusions of these patients.⁵ On average, they appear in rheumatoid arthritis with six to nine years of evolution. Still, they can occur both at the onset of the condition and in later stages, without association with the disease severity.⁵

Its pathogenesis remains uncertain. It is believed that these bodies are the end product of an inflammatory and proliferative process in the synovia, associated with ischemia with subsequent degeneration and desquamation of the same, encapsulated by fibrin.⁶

In this article, the patient had a previous diagnosis of rheumatoid arthritis. On the other hand, other authors reported the presence of riziform bodies before the diagnosis of inflammatory arthritis, highlighting the importance of following up on these patients for at least one to two years for etiological investigation.²

Riziform bodies have already been isolated in chronic infectious arthritis of the hands, mainly caused by *Mycobacterium marinum* and *M. avium*.⁷ However, it is the first article to report their occurrence in this location, associated with rheumatoid arthritis.

Furthermore, during surgical drainage, we observed that they originated from the subcutaneous tissue of the patient's palmar region, unlike what is expected in rheumatoid arthritis, in which they predominantly occur within the bursa, with the subacromial being the most affected. This subcutaneous occurrence was previously reported only in tenosynovitis caused by mycobacteria.⁸

In patients with rheumatoid arthritis, consideration of concomitant infection is essential, especially in patients using immunosuppressants and corticosteroids.⁷

As shown the particles are characterized, macroscopically, by a smooth, light, yellow surface. However, it has already been demonstrated that riziform bodies can present several morphological variations: color (between white and yellow), consistency (smooth or rough), and size (from less than 2 mm to greater than 7mm).⁵

In this case, the patient had complete remission of the lesion after surgical drainage without the need for additional treatments. Other authors have reported similar results after treatment but with a shorter follow-up time than described in this article.² The improvement with drainage seems to be due to the removal of fibrin deposits that make up the riziform bodies since this substance presents the ability to perpetuate the inflammatory stimulus in the joints.⁵ Thus, in addition to being curative, surgical drainage reduces the recurrence risk, as observed during the two-year follow-up of our patient.⁹

Another measure associated with surgical treatment is the clinical control of underlying arthritis with specific pharmacological therapy, and, in cases of associated infection, antimicrobial therapy should be introduced.

CONCLUSION

What most caught our attention in this patient was the rarity we came across similar cases.

The diagnosis was made through physical examination combined with ultrasound examination and confirmed by pathological examination. Along with the unique characteristics of the surgical finding, we concluded this unusual and rare diagnosis.

In this case, we believe that the etiology is rheumatoid arthritis. As the clinical evolution has been satisfactory, we did not conduct other tests, nor the proposal in the literature for the excision of the affected tendon sheath. However, we are closely following the evolution of the clinical picture, which continues without recurrences and with good recovery of joint movements after two years of follow-up. ●

REFERENCES:

1. Campos LR, Sztajnbock FCN, Galvão S, Lessa MA, Aymoré IL, et al. Presence of riziform bodies in a patient with juvenile idiopathic arthritis: case report and literature review. *Rev Bras Reumatol Engl Ed*. 2017;57(6):610-2.
2. Subramaniam R, Tan JW, Chau CY, Lee KT. Subacromial bursitis with giant rice bodies as initial presentation of rheumatoid arthritis. *J Clin Rheumatol*. 2012;18(7):352-5.
3. Reda FM, Talal G, Moncef B, Reda-Allah B, Omar LM, Saleh BM. Mass of the thenar eminence hiding idiopathic massive rice bodies formation with a compression of the median nerve: case report and review of the literature. *Int J Surg Case Rep*. 2018;50:28-31.
4. Cheung HS, Ryan LM, Kozin F, McCarty DJ. Synovial origins of Rice bodies in joint fluid. *Arthritis Rheum*. 1980;23(1):72-6.
5. Popert AJ, Scott DL, Wainwright AC, Walton KW, Williamson N, Chapman JH. Frequency of occurrence, mode of development, and significance of rice bodies in rheumatoid joints. *Ann Rheum Dis*. 1982;41(2):109-17.
6. Karaoğlu S, Karaaslan F, Mermerkaya MU. A mechanically locked knee joint due to free-floating flake-shaped rice bodies: a case report. *Acta Orthop Traumatol Turc*. 2015;49(5):565-7.
7. Matcuk Jr GR, Patel DB, Lefebvre RE. Horseshoe abscess of the hand with rice bodies secondary to mycobacterium avium intracellulare infection. *Clin Imaging*. 2020;63:24-9.
8. Saraya T, Fukuoka K, Maruno H, Komagata Y, Fujiwara M, Kaname S. Tenosynovitis with rice body formation due to mycobacterium intracellulare infection after initiation of infliximab Therapy. *Am J Case Rep*. 2018;19:656-62.
9. Cegarra-Escolano M, Jaloux C, Camuzard O. Rice-body formation without rheumatic disease or tuberculosis in a "sausage" ring finger. *Hand Surg Rehabil*. 2018;S2468-1229(18)30067-7.

AUTHORS' CONTRIBUTION:

Isabella Zurita Dehó  ORCID 0000-0002-2884-7073

Preparation and writing of the manuscript.

Jayme de Oliveira Filho  ORCID 0000-0003-0239-0981

Approval of the final version of the manuscript; active participation in research orientation; intellectual participation in propaedeutical and/or therapeutic conduct of studied cases; critical revision of the manuscript.

Irina Andrea Pires Afonso  ORCID 0-000-0001-6787-8529

Author's contribution: Approval of the final version of the manuscript; preparation and writing of the manuscript.

Anna Rita Ferrante Mitidieri de Oliveira  ORCID 0000-0002-5551-7440

Approval of the final version of the manuscript.