



Pastos y Forrajes

ISSN: 0864-0394

ISSN: 2078-8452

Estación Experimental de Pastos y Forrajes Indio Hatuey

Trejo-Arista, Laura Karen; Cortés-Díaz, Enrique; Martínez-Hernández, Pedro Arturo; Huerta-Bravo, Maximino  
Sistemas Silvopastoriles: Una estrategia para la resiliencia ambiental en empresas ganaderas  
Pastos y Forrajes, vol. 46, e19, 2023, January-December  
Estación Experimental de Pastos y Forrajes Indio Hatuey

Available in: <https://www.redalyc.org/articulo.oa?id=269176991018>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UDEM  redalyc.org

Scientific Information System Redalyc  
Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

## Silvopastoral systems: A strategy for environmental resilience in livestock enterprises

Laura Karen Trejo-Arista<sup>1</sup> <https://orcid.org/0009-0008-6812-0531>, Enrique Cortés-Díaz<sup>1\*</sup> <https://orcid.org/0000-0003-1676-0402>, Pedro Arturo Martínez-Hernández<sup>1</sup> <https://orcid.org/0000-0003-2197-3736> and Maximino Huerta-Bravo<sup>1</sup> <https://orcid.org/0000-0002-9150-8971>

<sup>1</sup>Universidad Autónoma Chapingo. Carretera México-Texcoco km 38.5, Chapingo, 56230, Estado de México.

E-mail: laurakaren00@hotmail.com, ecodia@yahoo.com.mx\*, pedroarturo@correo.chapingo.mx, mhuertab@correo.chapingo.mx.

\*Corresponding author

### Abstract

**Objective:** To provide an information bank on the management and benefits of silvopastoral systems to enable the creation of environmentally resilient livestock enterprises.

**Materials and Methods:** Through a search for information, scientific papers on the impact of the conformation and management of silvopastoral systems on indicators of environmental resilience were compiled. For this purpose, Google Scholar, Springer, SciELO, Scopus, Redalyc and ScienceDirect were reviewed. Inclusion and exclusion criteria were established and only 63 documents were selected for inclusion in the information bank.

**Results:** The benefits of silvopastoral systems were explained for four indicators: preservation of soil health, mitigation of greenhouse gas emissions, conservation of biodiversity and promotion of animal welfare. Their effect on the improvement of soil physical, chemical and microbiological properties, reduction of CO<sub>2</sub> and CH<sub>4</sub> emissions, generation of food resources, as well as their performance as a source of refuge for wild animals and a comfort zone and welfare for livestock were discussed.

**Conclusions:** Silvopastoral systems promote environmental resilience, which is achieved by improving soil physical, chemical and microbiological properties, conserving biodiversity, promoting animal welfare and contributing to the mitigation of greenhouse gas emissions.

**Keywords:** biodiversity, mitigation, soil

### Introduction

The management of pastoral animal husbandry enterprises, based on an environment poor in plant species and rich in the application of external inputs to eradicate plant and animal, non-livestock species, and promote the maximum possible yield of a few relevant plant species for livestock feeding, has been associated with the generation of fragile ecosystems, of poor environmental and economic resilience (Gómez-Villalva *et al.*, 2019; Pérez-Sánchez *et al.*, 2021). Environmental fragility originates in the scarce plant and animal biodiversity, which allows soil degradation, the extinction of plant biomass under extreme climatic conditions, as well as the interruption of mutualistic symbiont chains and the increase of the water footprint, among other consequences (Rodríguez-Moreno *et al.*, 2020; Yalta *et al.*, 2021). In turn, economic fragility stems from a production cost that is dependent on fluctuations in the cost of external inputs. The latter are increasingly necessary to maintain or improve the production level of the

animal husbandry enterprise (Rodríguez-Moreno *et al.*, 2020; Aguilar-Jiménez *et al.*, 2023).

El-Bilali *et al.* (2019) define sustainability as development that makes it possible to meet present needs without compromising the ability of future generations to meet their needs. Areas of broad plant biodiversity are environmentally and economically resilient ecosystems, because plant biodiversity favors mutualistic relationships that, in turn, promote healthy soils, a fundamental premise for sustainable plant production (Gutiérrez-Bermúdez *et al.*, 2020; Lehmann *et al.*, 2020). Pastoral or grazing areas with planned plant biodiversity, with the incorporation of herbaceous, semi-ligneous and ligneous species of medium to tall size, have been referred to as silvopastoral systems (SPS) and have been identified as an option in the design and implementation of resilient animal husbandry enterprises (Díaz-Lezcano *et al.*, 2020; Peña-Domene *et al.*, 2022).

SPSs can be an alternative in the management of natural resources to reverse damage to

Received: May 08, 2023

Accepted: September 29, 2023

How to cite a paper: Trejo-Arista, Laura Karen; Cortés-Díaz, Enrique; Martínez-Hernández, Pedro Arturo & Huerta-Bravo, Maximino. Silvopastoral systems: A strategy for environmental resilience in livestock enterprises. *Pastures and Forages*, 46:e19, 2023.

This is an open access article distributed in Attribution NonCommercial 4.0 International (CC BY-NC4.0) <https://creativecommons.org/licenses/by-nc/4.0/> The use, distribution or reproduction is allowed citing the original source and authors.

ecosystems disturbed by human activity, in which ecosystems with low plant and animal biodiversity were established in the search for maximum yield (González-Valdivia *et al.*, 2018). In addition to obtaining direct anthropocentric consumption and use satisfiers, agricultural enterprises must pay attention to the provision of ecosystem services: clean water, healthy soil formation, microclimate due to aerial vegetation cover, pollination, among others (Gómez-Villalva *et al.*, 2019; Peña-Domene *et al.*, 2022).

The adequate planning of the plant components of a SPS promotes healthy soils, capable of producing higher yields due to their abundant biology, extensive plant and animal biodiversity, in addition to intervening in the mitigation of greenhouse gas emissions and contributing to animal welfare and the resilience of animal husbandry enterprises (González-Valdivia *et al.*, 2018; Vásquez *et al.*, 2020; Deniz, 2021; Oyelami, 2022). However, it is necessary to compile and critically analyze the scientific evidence about the shaping and management of SPSs to achieve positive environmental impacts and resilient and sustainable animal husbandry enterprises (Améndola *et al.*, 2016; Hanisch *et al.*, 2019). The objective of this scientific literature review was to provide a bank of information about the influence of the conformation and management of SPSs as grazing areas on indicators associated with environmentally resilient animal husbandry enterprises.

## Materials and Methods

Through a search for information, scientific papers about the impact of SPS formation and management on indicators of environmental resilience were compiled. These included soil health, mitigation of greenhouse gas (GHG) emissions, biodiversity and animal welfare. The strategy followed for the search was based on the selection of keywords to obtain information on each indicator. The keywords that were connected with the Boolean operators AND, OR and NOT, according to the meaning of the phrase were: silvopastoral system, soil health, soil organic matter, soil bulk density, soil fauna, soil minerals, carbon sequestration, GHG mitigation, methane, climate change, biodiversity, animal welfare, animal production, sustainability, ecological resilience. The search was conducted in Google Scholar, Springer, SciELO, Scopus, Redalyc and ScienceDirect. Scientific and literature review papers were considered, as well as books and theses.

The search was prioritized with the filter for information published as of 2019. The following criteria were used to select the information.

### *Inclusion criteria:*

- Documents that clearly and accurately describe the impact of SPSs or traditional animal husbandry systems, and both, on soil physical, chemical and microbiological properties, carbon sequestration, GHG emissions, biodiversity and animal welfare indicators.
- Documents detailing the impact of SPSs or traditional animal husbandry systems, and of both, on environmental sustainability and resilience.
- Papers that consider SPSs based on native tree species.

### *Exclusion criteria:*

- Papers that do not clearly and concisely describe the effect of SPSs or traditional animal husbandry systems, and of both, on the environmental resilience indicators considered in this work.
- Documents that consider SPSs based on exotic tree species.

Eighty-six documents were identified, which could be included in the review. After reading the title, 9 were discarded. Seventy-seven were reviewed in full, of which 14 were rejected, i.e. only 63 were selected for inclusion in this review.

## Results and Discussion

*Soil health in silvopastoral systems.* Sustainable soil management is based on the maintenance and improvement of soil health over time, so as to avoid compromising the support services it provides to the ecosystem (FAO *et al.*, 2015). Soil health can be evaluated through the state of its physical, chemical and microbiological properties (Cabrera-Dávila *et al.*, 2021; Castillo-Valdez *et al.*, 2021), since the management of this resource in production systems alters them significantly, mainly when energy inputs are lower than outputs (Castillo-Valdez *et al.*, 2021). Compaction, salinization, acidification, loss of nutrients or organic matter (OM), and of both, are considered types of soil degradation and refer to the alteration of its properties (FAO *et al.*, 2015). Silva-Olaya *et al.* (2022) refer that, in soils with degraded conditions, SPSs help to better use the land, as the integration of multiple-use trees favors nutrient recycling and improves environmental conditions. Several studies have been conducted to determine the impact of these systems on soil quality, in terms of chemical, physical and microbiological properties.

Regarding physical properties, Contreras-Santos *et al.* (2020) found in a SPS, lower bulk density (BD) and mechanical resistance to root penetration (MRRP), up to 25 and 37 %, respectively, compared with a soil from a system without trees (SWT) (table 1). Oliva *et al.* (2016) referred a similar trend, with values up to 31 and 65 % lower in the soil of a SPS with regards to that of a SWT (table 1). Romero-Delgado *et al.* (2021) pointed out that ligneous species show higher root development than herbaceous species, which has a direct effect on improving soil aggregation and structuring levels and decreasing BD, compaction and MRRP.

Table 1 also shows the studies conducted by Bugarín *et al.* (2010), who after 14 months of evaluation of a SPS, recorded that the BD was maintained, despite sheep trampling. Escobar *et al.* (2020) found no differences in the MRRP when comparing the areas under the trees of a SPS and the open areas, which had 54 % higher soil depth. In the SPS under tree canopies, temperatures are usually lower than in open areas, which generates a more suitable microclimate for animals, so they tend to rest in these areas at various times of the day (Deniz *et al.*, 2020). As a result, trampling in these areas increases, which can lead to higher soil compaction (Escobar *et al.*, 2020; Romero-Delgado *et al.*, 2021). However, it is encouraging that indicators such as BD and MRRP are maintained, despite this animal behavior.

Regarding chemical properties, the contributions of macronutrients and micronutrients to the soil via leaf litter and biological nitrogen fixation by systems with leguminous trees are shown (Bueno-López and Camargo-García, 2015; Oliva *et al.*, 2016), which proves that SPSs have a positive effect on improving soil fertility. This was found by Escobar *et al.* (2020), who refer a saturation of bases (K, Ca and Mg) up to 54 %, higher in SPSs with 30 years of exploitation compared with open areas and under eight-year-old trees. Toledo (2016) and Vargas-Batis *et al.* (2020) consider base saturation as one of the properties that determine soil fertility. The above allows to infer that the longer the soil is maintained under silvopastoral management, its fertility will improve.

Another important aspect in determining soil health is the status of OM. Most soil ecosystem functions depend on it, which are essential for C sequestration, N mineralization, aggregation, promotion of plant health, and water and nutrient retention (Cruz-Macias *et al.*, 2020; Hoffland *et al.*, 2020;

Núñez-Ravelo *et al.*, 2021). It has been pointed out that soil OM comes from the microbial decomposition of plant (litter, exudates and root residues) and animal biomass residues, so a greater contribution of these residues is essential for its regeneration and increase (Crespo and Fraga, 2002; Haddix *et al.*, 2020; Núñez-Ravelo *et al.*, 2021). The table above shows the benefits of SPS in terms of the amount of soil OM when compared with SWT, and with regards to the increase in OM content over time, an issue that has been addressed by several authors (Hernández-Chavez *et al.*, 2008; Bugarín *et al.*, 2010; Oliva *et al.*, 2016; Escobar *et al.*, 2020).

Regarding soil organic carbon (OC) content, increases of up to 70, 12 and 47 %, respectively, have been reported in soils under silvopastoral management compared with SWT soils (Oliva *et al.*, 2016; Contreras-Santos *et al.*, 2020; Escobar *et al.*, 2020). Likewise, Contreras-Santos *et al.* (2020) estimated at least 18 times more fine root biomass in soils under silvopasture when compared with SWT. The above-explained fact is in agreement with the report by Sanderman *et al.* (2017), who found that, in soils with higher plant diversity and biomass productivity, OC storage increased compared with conventionally managed areas. Haddix *et al.* (2020) reported that lower OC storage may undermine the long-term sustainability of food production and contribute to climate change. Lok *et al.* (2013) state that carbon stored in the soil can serve as an indicator of the efficiency, stability and functioning of forage systems, as it combines the effects of management, soil type, vegetation present and biological activity. Therefore, due to their positive effect on OC storage, SPSs contribute to sustainability.

Regarding microbiological properties, Gutiérrez-Bermúdez *et al.* (2020) and Cabrera-Dávila *et al.* (2021) indicated that a fundamental factor to achieve healthy soils is their biology, formed by microorganisms and macroorganisms, which exert the mining, movement, retention and recirculation of minerals in the soil-plant-animal complex. However, in turn, soil functioning and prosperity depend on the quantity and diversity of the root mass, for which soil biology demands broad plant biodiversity to create a soil profile with abundant and diverse root mass. In this regard, several studies (Hernández-Chavez *et al.*, 2008; Escobar-Montenegro *et al.*, 2017; Caicedo-Rosero *et al.*, 2018; Escobar *et al.*, 2020) have estimated higher diversity or abundance of edaphic microfauna and

Table 1. Contributions of silvopastoral systems to the soil.

| Study                               | Components of SPS                                                                  |                                                                                              | Density of the tree species                 | Contributions to soil properties                                         |                                                                                                                                                                                                                                                                                                                                              |                                                                                         | Treatments and comparisons                                                                |
|-------------------------------------|------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------|---------------------------------------------|--------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------|
|                                     | Trees                                                                              | Herbaceous plants                                                                            |                                             | Physical                                                                 | Chemical                                                                                                                                                                                                                                                                                                                                     | Microbiological                                                                         |                                                                                           |
| Bueno-López & Camargo-García (2015) | <i>Leucaena leucocephala</i> (Lam.) de Wit                                         | NS                                                                                           | 10 000 plants/ha.                           |                                                                          | Contributed in seven months 250 kg N/ha.                                                                                                                                                                                                                                                                                                     |                                                                                         | It was not compared with SWT                                                              |
| Bugarín et al. (2010)               | <i>L. leucocephala</i> (Ll) and <i>Leucaena glauca</i> (L.) Benth. (Lg)            | <i>Urochloa brizantha</i> (A. Rich.) R. D. Webster (Ub) and <i>Clitoria ternatea</i> L. (Ct) | 2 500 plants/ha in all the SPSs.            | BD (1,33 g/cm <sup>3</sup> ) was maintained in spite of sheep trampling. | In T1, T2 y T3, at 14 months of establishment of the SPS, increase of -OM from 1,68 to 1,94 %<br>-pH from 6,3 to 6,7 as average.                                                                                                                                                                                                             |                                                                                         | Four SPS:<br>T1 (Lj+Ub)<br>T2 (Lg+Ub)<br>T3 (Lj+Ub+Ct)<br>T4 (Lg+Ub+Ct)<br>1 SWT: T5 (Ub) |
| Caicedo-Rosero et al. (2018)        | 2 SPS;<br>T1: <i>Alnus acuminata</i> Kunth<br>T2: <i>Acacia melanoxylon</i> R. Br. | Forage mix of grasses in both SPS                                                            | NS                                          |                                                                          | 9 taxa in each SPS and 10 in T3.<br>Haptotaxida with higher relative abundance in the three treatments (> 66,5 %).<br>According to Shannon – Wiener index, T2 had higher diversity (1,161), followed by T3 (0,924) and finally T1 (0,902).<br>According to Simpson index, T2 was the highest (0,531), followed by T1 (0,396) and T3 (0,394). |                                                                                         | T3: SWT with forage mixture of grasses.                                                   |
| Camero-Rey & Rodríguez-Díaz (2015)  | <i>Erythrina berteiroana</i> Urb.                                                  | <i>U. brizantha</i> cv. Toledo CIAT 26110                                                    | Tree biomass production: 2724 kg DM/ha/year |                                                                          | Higher OM quantity in the SPS vs SWT (1,70 vs 1,20 %)                                                                                                                                                                                                                                                                                        | Higher quantity of earthworms in the SPS vs SWT (315 vs 262 earthworms/m <sup>2</sup> ) | SWT of <i>U. brizantha</i> cv. Toledo CIAT 26110                                          |

Table 1. Contributions of silvopastoral systems to the soil.

| Study                                 | Components of SPS                                                                     |                                                                                           | Density of the tree species | Contributions to soil properties                                                                                                                                                                             |                                                                                                                                                                                                   |                                                                                                                                                                                                   | Treatments and comparisons      |
|---------------------------------------|---------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------|-----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------|
|                                       | Trees                                                                                 | Herbaceous plants                                                                         |                             | Physical                                                                                                                                                                                                     | Chemical                                                                                                                                                                                          | Microbiological                                                                                                                                                                                   |                                 |
| Contreras-Santos <i>et al.</i> (2020) | Four SPS (T1, T2, T3 and T4) with combinations of forage and timber shrubs and trees. | <i>Megathyrus maximus</i> (Jacq.) B. K. Simon & S. W. L. Jacobs cv. Mombasa in the 4 SPSs | 39 trees/ha in all the SPSs | Lower BD in the four SPS at 20 cm of depth (1,16-1,22 g/cm3) than in T5 (1,54 g/cm3).<br>Lower MRRP at 20 cm of depth in the four SPS (288,1-1864,7 kg/ha) than in T5 (1,93-2,26 MPa) than in T5 (3,07 MPa). | Higher accumulation of OC in the four SPSs (60,6-65,1 t/ha) than T5 (38,3 t/ha).<br>Higher biomass of fine roots at 20 cm of depth in the four SPS (288,1-1864,7 kg/ha) than in T5 (15,37 kg/ha). | T1: pasture + forage shrubs<br>T2: pasture + forage trees<br>T3: pasture + forage shrubs + forage trees<br>T4: pasture + forage shrubs + forage trees + timber trees<br>T5: pasture without trees |                                 |
|                                       |                                                                                       |                                                                                           |                             |                                                                                                                                                                                                              |                                                                                                                                                                                                   |                                                                                                                                                                                                   |                                 |
| Crespo y Fraga (2002)                 | <i>Cajanus cajan</i> (L.) Millsp.                                                     | <i>Cynodon nlemfuensis</i> Vanderyst                                                      | 3 300 plants/ha             | In five months it contributed via litter:<br>973 g DM/shrub = 63, 10, 44, 30 and 6 kg/ha of N, P, K, Ca and Mg, respectively                                                                                 |                                                                                                                                                                                                   |                                                                                                                                                                                                   | They were not compared with SWT |
|                                       | <i>Albizia lebbeck</i> (L.) Benth.                                                    | <i>C. nlemfuensis</i>                                                                     | 100 plants/ha               | In four months it contributed via litter:<br>2 871 g DM/tree = 11, 1, 7, 9, and 1 kg/ha of N, P, K, Ca and Mg, respectively                                                                                  |                                                                                                                                                                                                   |                                                                                                                                                                                                   |                                 |

Table 1. Contributions of silvopastoral systems to the soil.

| Study                                   | Components of SPS                                                                                                                                                                                                                                 |                                                         | Contributions to soil properties                                |                                                                                                                                                     |                                                                                                                                                                                                                                                                                    | Treatments and comparisons                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-----------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------|-----------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                         | Trees                                                                                                                                                                                                                                             | Herbaceous plants                                       | Density of the tree species                                     | Physical                                                                                                                                            | Chemical                                                                                                                                                                                                                                                                           | Microbiological                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Escobar-Montenegro <i>et al.</i> (2017) | SPS (T1) with <i>Guazuma ulmifolia</i> Lam., <i>Vachellia pinnatula</i> (Schltdl. & Cham.) Seigler & Ebinger, <i>Psidium guajava</i> L., <i>Gliricidia sepium</i> (Jacq.) Kunth ex Walp. and <i>Caesalpinia velutina</i> (Britton & Rose) Standl. | <i>Andropogon gayanus</i> Kunth                         | Spatial distribution of scattered trees and live fences.        |                                                                                                                                                     | Similar values in the three treatments of:<br><br>-pH (6,16-6,36)<br>-OM (4,12-5,27 %)<br>-N (0,2-0,26 %)<br>-P (6,01-6,41 ppm)<br>-CEC (36,9-47,2 Meq/100 g soil)                                                                                                                 | T1 with higher abundance of macrofauna (4 231 individuals) and richness (24 orders) than T2 (2 664 individuals and 22 orders) and T3 (2 985 individuals and 23 orders).<br><br>T2: SWT of traditional enclosed pasture ( <i>Paspalum notatum</i> Flügge, <i>A. gayanus</i> and undergrowth).<br><br>T3: latifoliate forest (large proportion of deciduous xerophyte species).                                                                |
|                                         |                                                                                                                                                                                                                                                   |                                                         |                                                                 |                                                                                                                                                     |                                                                                                                                                                                                                                                                                    |                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Escobar <i>et al.</i> (2020)            | Eight- and 30-year old <i>A. acuminata</i> trees                                                                                                                                                                                                  | <i>Cenchrus clandestinus</i> (Hochst ex Chiov.) Morrone | 35 % of cover in arrangement of scattered trees and live fences | Lower depth of the soil in the areas under trees (22 - 25 cm) than in the open areas (34 cm).<br><br>There were no differences among areas in MRRP. | Areas under the trees with higher quantity of OM (9,5-9,7 %) and OC (5,5-5,6 %) than the open ones (8,6 % OM and 5 % OC).<br><br>Under 30-year old trees higher saturation of bases (K, Ca and Mg: 34,15 %) than in open areas (25,47 %) and under eight-year old trees (22,13 %). | In areas under the trees, higher quantity of:<br><br>-Scarabaeidae: 11-26/m2<br><br>-Actinomycetes: 17-18 x105 CFU/g<br><br>-PSM: 31-160 x104 CFU/g<br><br>In open areas:<br><br>-Scarabaeidae 3/m2<br><br>-Actinomycetes: 13 x105 CFU/g<br><br>-PSM 6 x104 CFU/g<br><br>In the silvopastoral system the areas under the 8 and 30 year-old trees, and the open areas, were evaluated separately. Implementation of agroecological practices. |
|                                         |                                                                                                                                                                                                                                                   |                                                         |                                                                 |                                                                                                                                                     |                                                                                                                                                                                                                                                                                    |                                                                                                                                                                                                                                                                                                                                                                                                                                              |

Table 1. Contributions of silvopastoral systems to the soil.

| Study                          | Components of SPS                                 |                                                                                  | Density of the tree species | Contributions to soil properties                                                                                                                             |                                                                                                                                                                                                                                  |                                                              | Treatments and comparisons            |
|--------------------------------|---------------------------------------------------|----------------------------------------------------------------------------------|-----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------|---------------------------------------|
|                                | Trees                                             | Herbaceous plants                                                                |                             | Physical                                                                                                                                                     | Chemical                                                                                                                                                                                                                         | Microbiological                                              |                                       |
| Hernández-Chavez et al. (2008) | 2 SPS (T1 y T2): both with <i>L. leucocephala</i> | T1: <i>M. maximus</i> .<br>T2: <i>Cynodon plechtostachyus</i> K. (Schum) Pilger. |                             |                                                                                                                                                              | Higher quantity of OM in T1 and T2 (4,4 % both) than T3 (3,0 %).                                                                                                                                                                 | Higher quantity of taxa in T1 (6) and T2 (7) than in T3 (3). | T3: SWT of natural pasture            |
| Oliva et al. (2016)            | 2 SPS:                                            |                                                                                  |                             |                                                                                                                                                              | Higher levels of P in SWT in open field.                                                                                                                                                                                         |                                                              |                                       |
|                                | T1: <i>Pinus patula</i> Schl. et Cham.            | NS                                                                               | NS                          | Lower compaction in T4 (61,4 psi), followed by T1 (136,9 psi) and T2 (141,5 psi), higher in T3 (395 psi).                                                    | T4: forest. T1 (5,83 ppm) and T2 (5,81 ppm) than in T3 (4,22 ppm) and T4 (5,45 ppm).                                                                                                                                             |                                                              | T3: SWT in open field.<br>T4: forest. |
|                                | T2: <i>A. acuminata</i>                           |                                                                                  |                             | Lower BD in T4 (0,55 g/cm <sup>3</sup> ), followed by T1 (0,69 g/cm <sup>3</sup> ) and T2 (0,64 g/cm <sup>3</sup> ), higher in T3 (0,93 g/cm <sup>3</sup> ). | Higher quantity of OC (5,56 %), OM (9,58 %) and N (0,48 %) in T4 followed by T1 (OC: 5,34 %, OM: 9,2 % and N: 0,47 %) and T2 (OC: 4,76 %, OM: 8,2 % and N: 0,41 %), lower quantity in T3 (OC: 3,64 %, OM: 5,92 % and N: 0,31 %). |                                                              |                                       |

CEC-cation exchange capacity, OC-organic carbon, BD-bulk density, OM-organic matter, PSM-phosphate-solubilizing microorganisms, NS- not specified, MRRP- mechanical resistance to root penetration, SWT— system without trees, SPS- silvopastoral system, T- treatment.

mesofauna, and both, in soils with SPS with regards to soils with SWT (table 1). The improvement in the microbiological part has a positive effect on the sustainability of the system, since soil biology plays an important role in the processing of organic residues, resulting in the storage of OM and OC (Sanderman *et al.*, 2017; Núñez-Ravelo *et al.*, 2021).

*Mitigation of greenhouse gas emissions in silvopastoral systems.* According to IPCC (2018), the increase in the concentration of GHGs, such as CO<sub>2</sub> and CH<sub>4</sub>, in the atmosphere is related to the warming of the earth's surface, which can cause drastic changes in the climate pattern. Several studies (table 2) have found the potential of SPSs to carry out carbon sequestration and thereby contribute to mitigating the effects of climate change (Lok *et al.*, 2013; Oliva *et al.*, 2017; Vásquez *et al.*, 2020). The inclusion of tree species in these systems increases their productivity and, in turn, nutrient recycling (Oliva *et al.*, 2018; Azuara-Morales *et al.*, 2020), which can translate into higher carbon sequestration compared with SWT. According to a paper by Oliva *et al.* (2017), the tree component of the SPS contributed 8,9 % of the carbon sequestered by the system; while the herbaceous one only contributed 1,6 %. Other authors (Landholm *et al.*, 2019; Vásquez *et al.*, 2020; Contreras-Santos *et al.*, 2023) showed, at the system level, that the treatments with SPS showed higher carbon captures by 56, 321 and 37 %, respectively, compared with the evaluated SWT.

As discussed in the previous section, soil represents a storehouse for carbon recycled by plants. In this regard, Contreras-Santos *et al.* (2020) mentioned that SPSs have the capacity to capture atmospheric carbon and immobilize it in the soil, so they are of great importance in the recovery of degraded soils and GHG mitigation. This can be corroborated in the studies conducted by Pérez-Atehortúa *et al.* (2019) and Contreras-Santos *et al.* (2023), who found higher carbon sequestration (up to 23 and 31 %, respectively) in the soil of SPSs, compared with SWTs. According to these authors (table 2), this difference was associated with the increase in root biomass in the SPS, due to the development of the trees, since they observed that carbon sequestration in the soil increased proportionally with the increase in this biomass. This highlights the role of roots in the carbon cycle and the importance of considering the inclusion of trees in the design of agroecosystems due to their capacity for root development.

One of the risks of continuing to develop conventional production systems is the contribution they make to GHG generation. Lok *et al.* (2013) estimated that, after three years, 16,7 t C/ha were lost in the soil of a monoculture pasture system; while 10,9 t C/ha were stored in the soil of a SPS. Landholm *et al.* (2019) reported 76 % lower CO<sub>2</sub> emission in a SPS compared with one with degraded pasture. De Bernardi *et al.* (2020) determined that a SPS can capture up to 374 % more CH<sub>4</sub> than a conventionally managed agricultural land. In this regard, the authors commented that the CH<sub>4</sub> uptake capacity of the soil is related to CH<sub>4</sub> diffusivity. Therefore, it is influenced by BD, water content and soil temperature, properties that are favorably influenced by SPS.

Commonly used tree species in SPSs have potential to reduce CH<sub>4</sub> emissions due to their nutritive value and secondary metabolite content. Berhanu *et al.* (2019) estimated higher CH<sub>4</sub> concentration in the gas produced *in vitro* by *C. cajan* (52,2 %), corn stubble (31,7 %) and *Sesbania sesban* (L.) Merr. (31,0 %) with regards to *L. leucocephala* (25,4 %), *Moringa stenopetala* (Baker fil.) Cufod. (24,1 %), *Crotalaria juncea* L. (23,4 %) and *Lablab purpureus* (L.) Sweet (16,2 %). The above-cited authors point out that the concentration of CH<sub>4</sub> in the gas produced *in vitro* by the above-mentioned species is inversely related to their ability to reduce CH<sub>4</sub> production and to their crude protein content. Irawan *et al.* (2020) observed in an *in vitro* study that, in a cattle diet consisting of 25 % concentrate feed and 75 % forage of *Cenchrus purpureus* (Schumach.) Morrone and *L. leucocephala*, the addition of this tree (500 and 750 g/kg DM of the diet) positively influenced ruminal fermentation and reduced CH<sub>4</sub> production by up to 54,2 %. This reduction was associated with the existence of tannins in the tree, which contributes directly to the suppression of methanogens and indirectly through inhibition of protozoan growth (Wanapat *et al.*, 2015). In turn, it can be ascribed to the improvement in the total digestibility of the diet, since there was a tendency that the higher the proportion of *L. leucocephala* in the diet, the lower the NDF contents (without *L. leucocephala* in the diet: 540,2 g/kg DM vs. *L. leucocephala* in the diet: 540,2 g/kg DM vs. with *L. leucocephala* (750 g/kg DM: 328,1 g/kg DM) and the ADF contents (without *L. leucocephala* in the diet: 312,9 g/kg DM vs. with *L. leucocephala* (750 g/kg DM: 198,0 g/kg DM) of the diet. Likewise, non-fiber carbohydrates were higher (without

Table 2. Contribution of silvopastoral systems to the reduction of greenhouse gas emissions.

| Study                                | Evaluated treatments                                                                                                                                                                                                                                                                                   | Tree density,<br>trees/ha | Sequestration, t/ha                           |                 | CH <sub>4</sub>                            | Storage                         | Time of<br>evaluation | Observations                                             |
|--------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------|-----------------------------------------------|-----------------|--------------------------------------------|---------------------------------|-----------------------|----------------------------------------------------------|
|                                      |                                                                                                                                                                                                                                                                                                        |                           | C                                             | CO <sub>2</sub> |                                            |                                 |                       |                                                          |
| Conteras-Santos <i>et al.</i> (2023) | SPS of <i>Erythrina fusca</i> Lour., <i>Pachira quinata</i> (Jacq.) W.S.Alverson, <i>Enterolobium cyclocarpum</i> (Jacq.) Griseb., <i>G. ulmifolia</i> , <i>Inga edulis</i> Mart., <i>Tabebuia rosea</i> (Bertol.) DC., <i>Samanea saman</i> (Jacq.) Merr and <i>M. maximus</i> cv. Sabanera Agrosavia | 19,0                      | AB: 2,18 RB: 0,63 LB: 2,51 S: 28,05 To: 33,21 |                 |                                            | Plant tissue, soil and system   | Two years             | Higher accumulation of C in AB and S                     |
|                                      | SPS of <i>E. cyclocarpum</i> , <i>E. fusca</i> , <i>Gmelina arborea</i> Roxb., <i>S. saman</i> , <i>G. sepium</i> , <i>Ficus</i> sp and <i>M. maximus</i> cv. Mombasa                                                                                                                                  | 81,0                      | AB: 4,51 RB: 1,16 LB: 3,09 S: 28,98 To: 37,70 |                 |                                            | Plant tissue, soil and system   | Two years             | Higher accumulation of C in AB, RB, LB, S and To         |
|                                      | SWT of <i>Bothriochloa pertusa</i> (L.) A. Camus                                                                                                                                                                                                                                                       |                           | AB: 0,19 RB: 0,08 LB: 0,30 S: 23,54 To: 24,12 |                 |                                            | Plant tissue, soil and system   | Two years             | Control. Lower accumulation of C in AB, RB, LB, S and To |
| De Bernardi <i>et al.</i> (2020)     | Natural pastureland                                                                                                                                                                                                                                                                                    |                           |                                               |                 | -9,5 ngCH <sub>4</sub> /m <sup>2</sup> /s  | Flow of CH <sub>4</sub> in soil | NS                    | Control                                                  |
|                                      | Agricultural land with <i>Hordeum vulgare</i> L. and <i>Glycine max</i> (L.) Merr.                                                                                                                                                                                                                     |                           |                                               |                 | -4,5 ngCH <sub>4</sub> /m <sup>2</sup> /s  | Flow of CH <sub>4</sub> in soil | NS                    | Lower capture of CH <sub>4</sub>                         |
|                                      | SPS of <i>Pinus radiata</i> D.Don and <i>Dactylis glomerata</i> L.                                                                                                                                                                                                                                     | 727,0                     |                                               |                 | -21,3 ngCH <sub>4</sub> /m <sup>2</sup> /s | Flow of CH <sub>4</sub> in soil | NS                    | Higher capture of CH <sub>4</sub>                        |
|                                      | Forestation of <i>P. radiata</i>                                                                                                                                                                                                                                                                       | 977,0                     |                                               |                 | -22,5 ngCH <sub>4</sub> /m <sup>2</sup> /s | Flow of CH <sub>4</sub> in soil | NS                    | Higher capture of CH <sub>4</sub>                        |

Table 2. Contribution of silvopastoral systems to the reduction of greenhouse gas emissions.

| Study                               | Evaluated treatments                                                                                                                                               | Tree density,<br>trees/ha | Sequestration,<br>t/ha |                 | CH <sub>4</sub> | Storage      | Time of evalua-<br>tion | Observations                                                        |
|-------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------|------------------------|-----------------|-----------------|--------------|-------------------------|---------------------------------------------------------------------|
|                                     |                                                                                                                                                                    |                           | C                      | CO <sub>2</sub> |                 |              |                         |                                                                     |
| Landholm<br><i>et al.</i><br>(2019) | SWT of degraded pastures<br>( <i>Urochloa spp</i> )                                                                                                                |                           |                        | 3,4 (1)         |                 | System       | One year                | Control (the C sequestration was<br>not evaluated)                  |
|                                     | SWT of cultivated pastures:<br><i>M. maximus</i> , <i>Urochloa humi-<br/>dicola</i> (Rendle) Morrone &<br>Zuloaga y <i>Axonopus scoparius</i><br>(Flügge) Kuhlman. |                           | 0,38                   | 1,3 (1)         |                 | System       | One year                |                                                                     |
|                                     | Forage bank and cultivated<br>pastures                                                                                                                             |                           | 0,66                   | 2,2 (1)         |                 | System       | One year                |                                                                     |
|                                     | SPS of <i>Calycopteryx spruce-<br/>anum</i> (Benth.) Hook.f. ex<br>K. Schum., <i>G. arborea</i> , cultiva-<br>ted pastures and forage bank                         | 104,0                     | 1,6                    | 0,8 (1)         |                 | System       | One year                | Higher sequestration of C and<br>lower emissions of CO <sub>2</sub> |
| Lok <i>et al.</i><br>(2013)         | SPS of <i>M. maximus</i> and <i>L.<br/>leucocephala</i>                                                                                                            | 9,014,0                   | 10,9                   |                 |                 | Soil         | Three years             |                                                                     |
|                                     | Grasses and mixture of cree-<br>ping legumes                                                                                                                       |                           | 9,8                    |                 |                 | Soil         | Three years             |                                                                     |
|                                     | SWT of <i>M. maximus</i>                                                                                                                                           |                           | -16,7                  |                 |                 | Soil         | Three years             | C in the soil was reduced                                           |
|                                     | Trees of <i>P. patula</i> associated<br>with native herbaceous plants                                                                                              | 625,0                     | 8,2                    | 30,2            |                 | Plant tissue | NS                      | Capacity of C capture of 42,7 %                                     |
| Oliva <i>et al.</i><br>(2017)       | Native herbaceous plants                                                                                                                                           |                           | 1,4                    | 5,4             |                 | Plant tissue | NS                      | (15 to 20 days of age)                                              |
|                                     | Litter                                                                                                                                                             |                           | 1,1                    | 4,1             |                 | Plant tissue | NS                      |                                                                     |
|                                     | Soil                                                                                                                                                               |                           | 81,2                   | 297,3           |                 | Soil         | NS                      |                                                                     |
|                                     | SPS as a whole                                                                                                                                                     |                           | 92,1                   | 337,2           |                 | System       | NS                      |                                                                     |

Table 2. Contribution of silvopastoral systems to the reduction of greenhouse gas emissions.

| Study                                   | Evaluated treatments                                                                         | Tree density, trees/ha | Sequestration, t/ha |                 | CH <sub>4</sub> | Storage                                | Time of evaluation | Observations                                                      |
|-----------------------------------------|----------------------------------------------------------------------------------------------|------------------------|---------------------|-----------------|-----------------|----------------------------------------|--------------------|-------------------------------------------------------------------|
|                                         |                                                                                              |                        | C                   | CO <sub>2</sub> |                 |                                        |                    |                                                                   |
| Pérez-Atehortúa<br><i>et al.</i> (2019) | SPS of <i>A. acuminata</i> ,<br><i>Acacia decurrens</i> Willd.<br>and <i>C. clandestinus</i> | 285,0                  | AB: 1,9<br>RB: 2,1  |                 |                 | Plant tissue of <i>C. clandestinus</i> | One year           | Yields<br>AB: 3,9 t/ha RB: 4,3 t/ha.<br>Higher capture of C in RB |
|                                         | SWT of <i>C. clandestinus</i>                                                                |                        | AB: 2,0<br>RB: 1,6  |                 |                 | Plant tissue                           | One year           | Yields<br>AB: 4,2 t/ha RB: 3,3 t/ha                               |
|                                         | SPS of <i>A. acuminata</i> ,<br>pastures and weeds                                           | Strips                 | B: 6,8<br>S: 101,3  | 396,4           |                 | Plant tissue and<br>soil               | Seven months       |                                                                   |
|                                         | SPS of <i>P. patula</i> , pastures<br>and weeds                                              | Strips                 | B: 11,7<br>S: 49,0  | 589,3           |                 | Plant tissue and<br>soil               | Seven months       |                                                                   |
| Vásquez <i>et al.</i><br>(2020)         | SPS of <i>Cupressus macrocarpa</i> Hartw. ex Gord.,<br>pastures and weeds                    | Live fence             | B: 32,8<br>S: 17,2  | 550,2           |                 | Plant tissue and<br>soil               | Seven months       |                                                                   |
|                                         | SPS of <i>Ceroxylon quin-diense</i> (H.Karst.)<br>H.Wendl., pastures and<br>weeds            | Scattered<br>trees     | B: 57,8<br>S: 121,6 | 658,0           |                 | Plant tissue and<br>soil               | Seven months       | Higher retention of C and CO <sub>2</sub>                         |
|                                         | SWT of pastures and<br>weeds                                                                 |                        | B: 1,7<br>S: 29,6   | 481,4           |                 | Plant tissue and<br>soil               | Seven months       |                                                                   |

(I)-emissions, B-biomass, AB-aerial biomass, LB-leaf litter biomass, RB-root biomass, NS-not specified, S-soil, To-total.

*L. leucocephala* in the diet: 15,3 g/kg DM vs with *L. leucocephala* (750 g/kg DM: 36,3 g/kg DM). The existence of secondary metabolites in tree species used in SPS, which can contribute to the reduction of CH<sub>4</sub> emissions, has been documented by Sandoval-Pelcastre *et al.* (2020), who compiled information on the content of these compounds in 20 species.

From the information discussed in this section, it is confirmed that SPSs contribute to the sustainability of animal husbandry systems by mitigating GHG emissions; while they could represent a benefit for social actors, since in many countries in the region (Oliva *et al.*, 2017) carbon sequestration is translated as an environmental service; while gaining access to other products for household use, such as timber and firewood.

*Biodiversity in silvopastoral systems.* The integration of biological variability at all levels, from genetic resources to species, ecosystems and landscapes, defines the term biodiversity. Oropesa-Casanova *et al.* (2020) pointed out that biodiversity is a reservoir of genetic resources that provides diverse ecosystem services and makes it possible to reduce the use of external inputs in agroecosystems. Appropriate management of biodiversity is a key aspect in the design of agricultural production systems, because it is the basis for life in the land and sustainability in agroecosystems (Sarandón and Flores, 2014). Its conservation depends on maintaining the functional integrity of the latter, as this is the best way to reduce the loss of species; it is also necessary to prioritize biodiversity conservation with emphasis on the definition of functional groups of organisms, identified on the basis of ecosystemic processes that have lower redundancy (Walker, 1992; Pascual *et al.*, 2021).

González-Valdivia *et al.* (2018), in Estelí, Nicaragua, documented the existing diversity of malacofauna in two traditional SPSs: dry, shrub and deciduous forest (BS) and oak-pine forest (BP). The studied systems contained 47 of the 216 species of terrestrial mollusks recorded for Nicaragua, 41 of which were found in the BS and 22 in the BP, with 16 in common. Among the species identified, 13 may be potential vectors of livestock diseases or agricultural pests. The authors also identified four species categorized as endemic and two with potential for biological control, predators of other snails. In both systems, very rare species prevailed (24 and 11 for BS and BP, respectively), followed in descending order: rare, occasional and frequent.

These conditions showed that the diversity of mollusks depends on the existence of a tree stratum. Therefore, in pastures and meadows it is convenient to have this component, as it can ensure the existence of niches that support the diversity of mollusks. In turn, when conserving biodiversity, trophic chains are present that can favor the presence and vigor of vegetation that feeds livestock, and others that can cause damage. However, the existence of functional groups of organisms with higher diversity favors the regulation of the functions that species fulfill in ecosystems and promotes their conservation (Walker, 1992).

In a commercial dairy farm with high plant diversity (grasses, *Tithonia diversifolia* (Hemsl.), *C. clandestinus* and *A. acuminata*), beneficial food chains for the growth of forage species were established by controlling the intensity of attack by the pests *Collaria* sp, *Draeculacephala* sp. and *Aeneolamia* sp. while improving the diet of grazing cows, which reduced the amount of canoe supplements and farm production costs (Lopera *et al.*, 2017).

Regarding diurnal avifauna, Ordoñez-Solarte *et al.* (2015) compared three production systems (I-*Coffea arabica* L. in monoculture, II-*C. arabica* in agroforestry arrangement with *I. edulis*, III-SSP with *C. clandestinus* and *P. guajava* in monoculture) and in system I the lowest values were found in the Shannon and Simpson indices, number of species and abundance of individuals, with respective values of 2,16; 0,88; nine species and 202 individuals, respectively; while the other two silvopastoral arrangements recorded an average of 2,75; 0,93; 15,5 species and 426 individuals. These authors concluded that a plant canopy of greater botanical complexity promotes more niches for feeding, shelter and refuge for birds and, consequently, brings with it abundant biodiversity.

Marinero and Grau (2015), in the Argentine Chaco, recorded higher variability in the Simpson index for birds and mammals in an animal husbandry landscape dominated by introduced grasses, compared with naturalized grassland and SPS (native trees-introduced grasses) livestock landscapes. From these findings, these authors indicated that botanically more complex landscapes, with the combination of plant, herbaceous and ligneous strata, promote greater homogeneity in the distribution of feeding, shelter and refuge niches and, therefore, better distribution of animal biodiversity. Therefore, they propose that animal husbandry

landscapes should be complex in their botanical conformation so that pastoral animal husbandry to be a factor of biodiversity, with homogeneous distribution across the area they occupy, so as to reverse the environmental degradation associated with pastoral animal husbandry. In an analysis of indicator species, the aforementioned authors identified *Chunga burmeisteri* Hartlaub among the species associated with the SPS, which only nests in trees, so its presence was associated with the characteristic structure of these systems.

The information discussed here allows us to conclude that the use of SPSs avoids the simplification of animal husbandry systems, since functional groups of organisms are created and established, aimed at covering livestock production and maintain the functional integrity of the ecosystem. These systems resemble natural ecosystems, which ultimately has positive effects on biodiversity conservation.

*Animal welfare in silvopastoral systems.* Animal welfare contributes to the economic sustainability of animal husbandry systems in the face of growing consumer concern for purchasing products that incorporate practices that preserve animal health. This improves the marketing status of products that incorporate these measures in their production (Molina, 2021). Blanco-Penedo and Cantalapiedra (2020) and Szorobura *et al.* (2022) refer that the implementation of practices that reduce animal stress and contribute to their welfare will facilitate their adaptation to environmental changes caused by climate change, which will improve the resilience and sustainability of the system.

Social interactions, classified as affiliative and agonistic, develop among grazing cattle. The former are considered sociopositive to high productive behavior and the latter, detrimental. Améndola *et al.* (2016) reported that the frequency of affiliative interactions was 40 % lower among cows grazing a canopy of *C. nlemfuensis*, compared with those grazing a SPS of *M. maximus* and *C. nlemfuensis*, associated with planted *L. leucocephala* and scattered trees. These authors conclude that the silvopastoral environment promotes a more positive social interaction among animals and this leads to a higher degree of welfare.

Deniz (2021) when recording the microclimate characteristics of shaded and unshaded grazing areas, determined that shaded areas showed a microclimate with lower ambient temperature, wind speed and radiant heat load, but higher relative

humidity than unshaded areas. This determined that during the hours and days with the highest heat load, cows sought the shaded areas. The authors conclude that planned silvopastoral arrangements or with the presence of trees can offer favorable microclimate conditions that translate into higher animal comfort, especially under conditions of high heat load environments.

Zúñiga-López *et al.* (2020) compared animal welfare indicators in cows grazing in an *A. acuminata*-*C. clandestinus* SPS and a *C. clandestinus* SWT. They observed that the number of *Haematobia irritans* L. flies per animal was higher in the SWT (56) than in the SPS (44). The average coat temperature was higher in SWT animals (37,4 °C) than in SPS (33,8 °C). Also, the level of soiling (0 clean animal - 3 dirty animal) and escape distance were lower in SPS (1,5 and 0,7 m) than in SWT (2,0 and 1,3 m). The differences were associated with the diversification of the SPS, given by the inclusion of trees and the subsequent generation of better distributed comfort spaces, which avoid the overcrowding of the animals in the moments of rest and rumination. In another study, Galloso-Hernández (2021) proved that water buffalo grazing on a SPS of *M. maximus* cv. Likoni and *L. leucocephala* cv. Cunningham spent more time grazing under the trees than in a SWT, where they spent more time in the thermoregulatory bath. In turn, this author determined that it is possible to dispense with these baths or wallows, and both, when managing buffalo under shaded conditions in a SPS.

The above-cited works prove that the generation of a microclimate by means of shade in SPSs has a positive effect on animal welfare, which can benefit productive performance by facilitating grazing in areas that reduce the animals' thermal stress. According to what was discussed in this section, it was found that SPSs provide comfort areas for the animals, which reduces their stress and improves their coexistence, which has a positive impact on their welfare.

## Conclusions

Silvopastoral systems promote environmental resilience, which can be achieved by improving the physical, chemical and microbiological properties of the soil, conserving biodiversity, promoting animal welfare and contributing to the mitigation of greenhouse gas emissions.

## Acknowledgments

The authors thank the Universidad Autónoma Chapingo for providing the space and digital media to carry out this review.

## Conflict of interest

The authors declare that there is no conflict of interest.

## Authors' contribution

- Laura Karen Trejo-Arista. Information search and analysis and paper writing.
- Enrique Cortés-Díaz. Information search and analysis and critical revision of the content.
- Pedro Arturo Martínez-Hernández. Information analysis and critical revision of the content.
- Maximino Huerta-Bravo. Information analysis and critical revision of the content.

## Bibliographic references

- Aguilar-Jiménez, J. R.; Aguilar-Jiménez, C. E.; Guevara-Hernández, F.; Galdámez-Galdámez, J.; Martínez-Aguilar, F.; La-O-Arias, M. A. *et al.* Clasificación y caracterización de los sistemas familiares de producción bovina de la región Frailesca de Chiapas, Mexico, con base en el aporte de la ganadería al ingreso familiar. *Trop. Subtrop. Agroecosyst.* 26:017, 2023. DOI: <http://doi.org/10.56369/tsaes.4131>.
- Améndola, L.; Solorio, F. J.; Ku-Vera, J. C.; Améndola-Massiotti, R. D.; Zarza, H. & Galindo, F. Social behaviour of cattle in tropical silvopastoral and monoculture systems. *Anim. Nutr.* 10(5):863-867, 2016. DOI: <https://doi.org/10.1017/S1751731115002475>.
- Azuara-Morales, I.; López-Ortiz, Silvia; Jarillo-Rodríguez, J.; Pérez-Hernández, P.; Ortega-Jiménez, E. & Castillo-Gallego, E. Forage availability in a silvopastoral system having different densities of *Leucaena leucocephala* under Voisin grazing management. *Agroforest Syst.* 94:1701-1711, 2020. DOI: <https://doi.org/10.1007/s10457-020-00487-5>.
- Berhanu, Y.; Olav, L.; Nurfeta, A.; Abdeta, A. & Aune, J. Methane emissions from ruminant livestock in Ethiopia. Promising forage species to reduce CH<sub>4</sub> emissions. *Agriculture.* 9:130, 2019. DOI: <https://doi.org/10.1080/10549811.2020.1745653>.
- Blanco-Penedo, Isabel; Cantalapiedra, J. & Llonch, P. Impacto del cambio climático sobre el bienestar animal en los sistemas ganaderos. *ITEA-Inf. Tec. Econ. Agrar.* 116 (5):424-443, [https://www.aida-itea.org/aida-itea/files/itea/revistas/2020/116-5/\(424-443\)%20ITEA%20116-5%20EXTRA.pdf](https://www.aida-itea.org/aida-itea/files/itea/revistas/2020/116-5/(424-443)%20ITEA%20116-5%20EXTRA.pdf), 2020.
- Bueno-López, Liliana & Camargo-García, J. C. Nitrógeno edáfico y nodulación de *Leucaena leucocephala* (Lam.) de Wit en sistemas silvopastoriles. *Acta Agron., Palmira.* 64 (4):349-354, 2015. DOI: <https://doi.org/10.15446/acag.v64n4.45362>.
- Bugarín, J.; Bojórquez, J. I.; Lemus, C.; Murray, R. M.; Ontiveros, H.; Aguirre, J. *et al.* Comportamiento de algunas propiedades físico-químicas del suelo con diferente sistema silvopastoril en la llanura norte de Nayarit. *Cultivos Tropicales.* 31 (2):48-55, [http://scielo.sld.cu/scielo.php?script=sci\\_arttext&pid=S0258-59362010000200007&lng=es&tlng=es](http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S0258-59362010000200007&lng=es&tlng=es), 2010.
- Cabrera-Dávila, Grisel de la C.; Sánchez-Rendón, J. A. & Ponce-de-León-Lima, D. Macrofauna edáfica: composición, variación y utilización como bioindicador según el impacto del uso y calidad del suelo. *Acta Botánica Cubana.* 221:1-21, <https://revistasgeotech.com/index.php/abc/article/view/404>, 2021.
- Caicedo-Rosero, D. M.; Benavides-Rosales, H. R.; Carvajal-Pérez, L. A. & Ortega-Hernández, Jessica P. Población de macrofauna en sistemas silvopastoriles dedicados a la producción lechera: análisis preliminar. *La Granja.* 27 (1):77-85, 2018. DOI: <https://doi.org/10.17163/lgr.n27.2018.06>.
- Camero-Rey, A. & Rodríguez-Díaz, H. Características químicas del suelo, producción forrajera y densidad poblacional de lombrices en un sistema silvopastoril en la zona Huetar Norte de Costa Rica. *Tecnología en Marcha.* 28 (1):91-104, <https://dialnet.unirioja.es/servlet/articulo?codigo=5051542>, 2015.
- Castillo-Valdez, Ximena; Etchevers, J. D.; Hidalgo-Moreno, Claudia Ma. I. & Aguirre-Gómez, A. Evaluación de la calidad de suelo: generación e interpretación de indicadores. *Terra Latinoam.* 39:e698, 2021. DOI: <https://doi.org/10.28940/terra.v39i0.698>.
- Contreras-Santos, J. L.; Falla-Guzmán, Cindy K.; Rodríguez, J. L.; Fernando-Garrido, J.; Martínez-Atencia, Judith & Aguayo-Ulloa, Lorena. Reserva de carbono en sistemas silvopastoriles: Un estudio en el Medio Sinú, Colombia. *Agron. Mesoam.* 34 (1):49138, 2023. DOI: <https://dx.doi.org/10.15517/am.v34i1.49138>.
- Contreras-Santos, J. L.; Martínez-Atencia, Judith; Cadena-Torres, J. & Falla-Guzmán, Cindy K. Evaluación del carbono acumulado en suelo en sistemas silvopastoriles del Caribe colombiano. *Agron. Costarricense.* 44 (1):29-41, 2020. DOI: <https://dx.doi.org/10.15517/rac.v44i1.39999>.
- Crespo, G. & Fraga, S. Nota técnica acerca del aporte de hojarasca y nutrientes al suelo por las especies *Cajanus cajan* (L.) Millsp y *Albizia lebbbeck* (L.) Benth en sistemas silvopastoriles. *Rev. cubana Cienc. agric.* 36 (4):397-402. <https://www.redalyc.org/pdf/1930/193018080014.pdf>, 2002.
- Cruz-Macias, W. O.; Rodríguez-Larramendi, L. A.; Salas-Marina, M. Á.; Hernández-García, V.; Campos-Saldaña, R. A.; Chávez-Hernández, M. H. *et al.* Efecto de la materia orgánica y la capacidad de intercambio catiónico en la acidez de suelos cultivados con maíz en dos regiones de Chiapas, México. *Terra Latinoam.* 38 (3):475-

- 480, 2020. DOI: <https://doi.org/10.28940/terra.v38i3.506>.
- De Bernardi, María; Priano, María E.; Fusé, Victoria S.; Fernández, María E.; Gyenge, J.; Guzmán, S. A. *et al.* High methane uptake from soils of low and high density radiata pine afforestations compared to herbaceous systems. *J. Sustain. For.* 40 (1):99-109, 2020. DOI: <https://doi.org/10.1080/10549811.2020.1745653>.
- Deniz, M.; Sousa, Karoline T. de; Moro, M. F.; Vale, M. M. do; Dittrich, J. R.; Pinheiro-Machado Filho, L. C. *et al.* Social hierarchy influences dairy cows' use of shade in a silvopastoral system under intensive rotational grazing. *Applied Anim. Behaviour Sci.* 244:105467, 2021. DOI: <https://doi.org/10.1016/j.applanim.2021.105467>.
- Deniz, M.; Schmitt-Filho, A. L.; Hötzel, M. J.; Sousa, K. T. de; Pinheiro-Machado Filho, L. C. & Sinisgalli, P. A. Microclimate and pasture area preferences by dairy cows under high biodiversity silvopastoral system in Southern Brazil. *Int. J. Biometeorol.* 64:1877-1887, 2020. DOI: <https://doi.org/10.1007/s00484-020-01975-0>.
- Díaz-Lezcano, Maura I.; Gamarra-Lezcano, Cynthia C.; Ruiz-Díaz, Susana & Vera-de-Ortiz, Mirtha. Contenido de materia orgánica en suelos de sistemas silvopastoriles establecidos en el Chaco Central paraguayo. *Rev. Soc. cient., Parag.* 25 (2):131-143, 2020. DOI: <https://doi.org/10.32480/rscsp.2020.25.2.131>.
- El-Bilali, H.; Callenius, Carolin; Strassner, Carola & Probst, L. Food and nutrition security and sustainability transitions in food systems. *Food Energy Security.* 8 (2):e00154, 2019. DOI: <https://doi.org/10.1002/fes3.154>.
- Escobar-Montenegro, Alexa del C.; Bartolomé-Filella, J. & González-Valdivia, N. A. Estudio comparativo macrofauna del suelo en sistema agroforestal, potrero tradicional y bosque latifoliado en microcuenca del trópico seco, Tomabú, Nicaragua. *Rev. Cient. FAREM-Estelí.* 6 (22):39-49, <https://repositorio.unan.edu.ni/5829/1/320-1168-1-PB.pdf>, 2017.
- Escobar, María I.; Navas-Panadero, A.; Medina, Claudia A.; Corrales-Álvarez, J. D.; Tenjo, Ana I. & Borrás-Sandoval, L. M. Efecto de prácticas agroecológicas sobre características del suelo en un sistema de lechería especializada del trópico alto colombiano. *LRRD.* 32 (4), <http://www.lrrd.org/lrrd32/4/maria.es32058.html>, 2020.
- FAO; ITPS & SWSR. *Main Report. Food and Agriculture Organization of the United Nations and Intergovernmental Technical Panel on Soils.* Rome: FAO, Intergovernmental Technical Panel on Soils, Status of the World's Soil Resources. <http://www.fao.org/3/a-i5199e.pdf>, 2015.
- Gallos-Hernández, M. A. *Potencial de los sistemas silvopastoriles para la producción bufalina en ambientes tropicales.* Tesis para optar por el grado de Doctor en Veterinaria, Sección Biociencias y Ciencias Agroalimentarias. Córdoba, España: Departamento de Producción Animal, Universidad de Córdoba. <https://helvia.uco.es/bitstream/handle/10396/22085/2021000002348.pdf?sequence=1&isAllowed=y>, 2021.
- Gómez-Villalva, J.; Cobos-Mora, F. & Hasang-Moran, E. Sostenibilidad de los sistemas de producción de ganadería extensiva. *Revista Ciencia e Investigación.* 4:180-195, 2019. DOI: <https://doi.org/10.5281/zenodo.3594078>.
- González-Valdivia, N. A.; Martínez-Puc, J. F. & Echavarría-Góngora, E. de J. Malacofauna en dos sistemas silvopastoriles en Estelí, Nicaragua. *J. Selva Andina Anim. Sci.* 5 (1):3-13, 2018. DOI: <https://doi.org/10.36610/j.jsaas.2018.050100003>.
- Gutiérrez-Bermúdez, C. del C.; Mendieta-Araica, B. G. & Noguera-Talavera, Á. J. Composición trófica de la macrofauna edáfica en sistemas ganaderos en el Corredor Seco de Nicaragua. *Pastos y Forrajes.* 43 (1):32-40, <https://www.redalyc.org/articulo.oa?id=269163399005>, 2020.
- Haddix, Michelle L.; Gregorich, E. G.; Helgason, B. L.; Janzen, H.; Ellert, B. H. & Cotrufo, M. Francesca. Climate, carbon content, and soil texture control the independent formation and persistence of particulate and mineral-associated organic matter in soil. *Geoderma.* 363:114160, 2020. DOI: <https://doi.org/10.1016/j.geoderma.2019.114160>.
- Hanisch, Ana L.; Negrelle, Raquel R. B.; Bonatto, R. A.; Nimmo, Evelyn R. & Lacerda, A. E. B. Evaluating sustainability in traditional silvopastoral systems (caívas): Looking beyond the impact of animals on biodiversity. *Sustainability.* 11 (11):3098, 2019. DOI: <https://doi.org/10.3390/sul1113098>.
- Hernández-Chavez, Marta; Sánchez-Cárdenas, Saray & Simón-Guelmes, L. Efecto de los sistemas silvopastoriles en la fertilidad edáfica. *Zootecnia Trop.* 26 (3):319-321, [http://ve.scielo.org/scielo.php?script=sci\\_arttext&pid=S0798-72692008000300035&lng=es&tlng=es](http://ve.scielo.org/scielo.php?script=sci_arttext&pid=S0798-72692008000300035&lng=es&tlng=es), 2008.
- Hoffland, E.; Kuyper, T. W.; Comans, R. N. J. & Creamer, Rachel E. Eco-functionality of organic matter in soils. *Plant Soil.* 455 (1):1-22, 2020. DOI: <https://doi.org/10.1007/s11104-020-04651-9>.
- IPCC. *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty.* Cambridge, UK; New

- York, USA.: Cambridge University Press, 2018. DOI: <https://doi.org/10.1017/9781009157940.001>.
- Irawan, Adiwinarta; Noviani, C. T.; Kustantinah; Widyobroto, B. P.; Astuti, A. & Ates, S. Effect of *Leucaena leucocephala* and corn oil on ruminal fermentation, methane production and fatty acid profile: an *in vitro* study. *Anim. Prod. Sci.* 61 (5):459-469, 2020. DOI: <https://doi.org/10.1071/AN20003>.
- Landholm, D. M.; Pradhan, P.; Wegmann, P.; Romero-Sánchez, M. A.; Suárez-Salazar, J. C. & Kropp, J. P. Reducing deforestation and improving livestock productivity: greenhouse gas mitigation potential of silvopastoral systems in Caquetá. *Environ. Res. Lett.* 14:11400755, 2019. DOI: <https://doi.org/10.1088/1748-9326/ab3db6>.
- Lehmann, J.; Bossio, D. A.; Kögel-Knabner, Ingrid & Rilling, M. C. The concept and future prospects of soil health. *Nat. Rev. Earth Environ.* 1:544-553, 2020. DOI: <https://doi.org/10.1038/s43017-020-0080-8>.
- Lok, Sandra; Fraga, S.; Noda, Aida & García, M. Almacenamiento de carbono en el suelo de tres sistemas ganaderos tropicales en explotación con ganado vacuno. *Rev. cubana Cienc. agríc.* 47 (1):75-82, <http://www.redalyc.org/articulo.oa?id=193028545014>, 2013.
- Lopera, J. J.; Márquez, Sara M.; Ochoa, D. E.; Calle, Zoraida; Sossa, Claudia P. & Murgueitio, E. Producción agroecológica de leche en el trópico de altura: sinergia entre restauración ecológica y sistemas silvopastoriles. *Agroecología*. 10 (1):79-85, <https://revistas.um.es/agroecologia/article/view/300761>, 2017.
- Marinero, Sofia & Grau, R. H. Comparison of animal biodiversity in three livestock systems of open environments of the semi-arid Chaco of Argentina. *Rangeland J.* 37 (5):497-505, 2015. DOI: <https://doi.org/10.1071/RJ15052>.
- Molina, N. D. P. Hacia una nueva normalidad: la Responsabilidad Social Corporativa y el bienestar animal como una oportunidad para las empresas. En: Lorena Miranda-Navarro, M. Á. Santinelli-Ramos y Diana Molina, coords. *Responsabilidad social y sostenibilidad: disrupción e innovación ante el cambio de época*. México: Universidad Anáhuac. p. 474-492. <http://pegaso.anahuac.mx/accesoabierto/publicaciones.php?Accion=Informacion&Pub=153>, 2021.
- Núñez-Ravelo, F.; Ugas-Pérez, María; Calderón-Castellanos, Rossany & Rivas-Meriño, F. Cuantificación del carbono orgánico y materia orgánica en suelos no rizosféricos o cubiertos por *Avicennia germinans* (L.) y *Conocarpus erectus* (L.) emplazados en Boca de Uchire, laguna de Unare, Estado de Anzoátegui, Venezuela. *RGE*. 66 (1):371-398, 2021. DOI: <https://doi.org/10.15359/rgac.66-1.13>.
- Oliva, M.; Culqui, M. L.; Leiva, S.; Collazos, R.; Salas, R.; Vásquez, H. V. *et al.* Reserva de carbono en un sistema silvopastoril compuesto de *Pinus patula* y herbáceas nativas. *Sci. Agropecu.* 8 (2):149-157, 2017. DOI: <https://doi.org/10.17268/sci.agropecu.2017.02.07>.
- Oliva, M.; Valqui, L.; Meléndez, J.; Milla, M.; Leiva, S.; Collazos, R. *et al.* Influencia de especies arbóreas nativas en sistemas silvopastoriles sobre el rendimiento y valor nutricional de *Lolium multiflorum* y *Trifolium repens*. *Sci. Agropecu.* 9 (4):579-583, 2018. DOI: <https://doi.org/10.17268/sci.agropecu.2018.04.14>.
- Oliva, M.; Vásquez-Pérez, H. V. & Vigo-Mestanza, Carmen N. Efecto de los sistemas de producción sobre las características físico-químicas de los suelos del distrito de Molinopampa, provincia de Chachapoyas, región Amazonas. *Revista Científica UNTRM*. 2 (1):44-52, 2016. DOI: <https://doi.org/10.25127/indes.20142.63>.
- Ordoñez-Solarte, Griseli; Erazo-Gómez, R. E. & Navia-Estrada, J. F. Evaluación de la biodiversidad en aves diurnas bajo diferentes sistemas de producción en el municipio de La Unión, departamento de Nariño, Colombia. *FAGROPEC*. 7 (1):3-10, <https://editorial.uniamazonia.edu.co/index.php/fagropec/article/view/318>, 2015.
- Oropesa-Casanova, Katherine; Pentón-Fernández, Gertrudis; Lezcano-Fleires, J. C.; Miranda-Tortoló, Taymer & Núñez-García, N. F. Biodiversidad y manejo de los residuos agropecuarios en una finca del municipio de Perico, Matanzas. *Pastos y Forrajes*. 43 (2):112-119, [http://scielo.sld.cu/scielo.php?script=sci\\_arttext&pid=S0864-03942020000200112&lng=es&tlng=es](http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S0864-03942020000200112&lng=es&tlng=es), 2020.
- Oyelami, B. A. & Osikabor, B. Adoption of silvopastoral agroforestry system for a sustainable cattle production in Nigeria. *J. Applied Sci. Environ. Manag.* 26 (8):1397-1402, 2022. DOI: <https://doi.org/10.4314/jasem.v26i8.12>.
- Pascual, U.; Adams, W. M.; Díaz, Sandra; Sharachandra, L.; Mace, G. M. & Turnhout, Esther. Biodiversity and the challenge of pluralism. *Nat. Sustain.* 4:567-572, 2021. DOI: <https://doi.org/10.1038/s41893-021-00694-7>.
- Peña-Domene, Marín de la; Ayestarán-Hernández, Luz M.; Márquez-Torres, J. F.; Martínez-Monroy, F.; Rivas-Alonso, Edith; Carrasco-Carballido, Patricia V. *et al.* Sistemas silvopastoriles enriquecidos: una propuesta para integrar la conservación en la producción ganadera en comunidades rurales de Los Tuxtlas, México. *Acta Botanica Mexicana*. 129:e1925, 2022. DOI: <https://doi.org/10.21829/abm129.2022.1925>.
- Pérez-Atehortúa, Maritza; Medina-Aguirre, María F.; Hurtado-Granada, A.; Arboleda-Zapata, E. M.

- & Medina-Sierra, Marisol. Reservas de carbono del pasto *Cenchrus clandestinus* (Poaceae) en los sistemas de manejo tradicional y silvopastoril, en diferentes relieves. *Rev. biol. trop.* 67 (4):769-783, 2019. DOI: <http://dx.doi.org/10.15517/rbt.v67i4.34529>.
- Pérez-Sánchez, E.; Hernández-Hernández, Erika; Jiménez-Trujillo, J. A.; Betanzos-Simón, J. E.; Casasola-Coto, F.; Martínez-Salinas, Alejandra *et al.* Reconversión de ganadería convencional a silvopastoril: Estudio de caso rancho El Once en Campeche, México. *ALA*. 25 (3):174-175, 2021. DOI: <https://doi.org/10.53897/RevAIA.21.25.46>.
- Rodríguez-Moreno, Orquidia G.; Nahed-Toral, J.; Guevara-Hernández, F.; Alayón-Gamboa, J. A. & Grande-Cano, J. D. Historia y caracterización técnica y socioeconómica de la ganadería bovina en la costa de Chiapas, México. *Trop. Subtrop. Agroecosyst.* 23:55, <https://www.revista.ccba.uady.mx/ojs/index.php/TSA/article/view/3058>, 2020.
- Romero-Delgado, G.; Trillo-Zárate, F.; Orellana-Chirinos, J.; Quiroga-Espilco, Pamela; Gamarra-Bojorquez, J.; Rojas-Egoavil, D. *et al.* Efecto de *Acacia macracantha* en las propiedades físicas y químicas del suelo en un sistema silvopastoril. *Rev. investig. vet. Perú.* 32 (3):e20389, 2021. DOI: <https://doi.org/10.15381/rivep.v32i3.20389>.
- Sanderman, J.; Creamer, Courtney.; Baisden, W. T.; Farrell, M. & Fallon, S. Greater soil carbon stocks and faster turnover rates with increasing agricultural productivity. *Soil Sci.* 3 (1):1-16, 2017. DOI: <https://doi.org/10.5194/soil-3-1-2017>.
- Sandoval-Pelcastre, A. A.; Ramírez-Mella, Mónica; Rodríguez-Ávila, Norma L. & Candelaria-Martínez, B. Árboles y arbustos tropicales con potencial para disminuir la producción de metano en rumiantes. *Trop. Subtrop. Agroecosyst.* 23:33, <https://www.revista.ccba.uady.mx/ojs/index.php/TSA/article/view/3061>, 2020.
- Sarandón, S. J. & Flores, Claudia C. *Agroecología: Bases teóricas para el diseño y manejo de agroecosistemas sustentables*. La Plata, Argentina: Universidad Nacional de La Plata, 2014.
- Silva-Olaya, A. M.; Olaya-Montes, A.; Polanía-Hincapié, K. L.; Cherubin, M. R.; Duran-Bautista, E. H. & Ortiz-Morea, F. A. Silvopastoral systems enhance soil health in the Amazon region. *Sustainability*. 14:320, 2022. DOI: <https://doi.org/10.3390/su14010320>.
- Szorobura, Florencia A.; Lynch, Gloria M.; Simonetti, Laura; Ghibaudo, Mercedes; Cormick, Mercedes Mc & Arioni, J. M. Bienestar animal: estrés al destete en ovinos. *RCYTAAA*. 9 (1), <http://servicios.ingenieria.unlz.edu.ar:8080/ojs/index.php/agrarias/article/view/90>, 2022.
- Toledo, M. *Manejo de suelos ácidos de las zonas altas de Honduras: Conceptos y métodos*. Tegucigalpa: IICA. <https://repositorio.ica.int/handle/11324/3108>, 2016.
- Vargas-Batis, B.; Escobar-Perea, Y.; Rodríguez-Fonseca, R.; Ramos-García, Y. M.; Rodríguez-Suárez, E. J. & Fuente-Miranda, O. Propiedades químicas del suelo en cuatro fincas de la agricultura suburbana en Santiago de Cuba. *Agrosost.* 26 (3):1-10, 2020. DOI: <https://doi.org/10.5281/zenodo.7433751>.
- Vásquez, H. V.; Valqui, L.; Alegre, J. C.; Gómez, C. & Maicelo, J. Análisis de cuatro sistemas silvopastoriles en Perú: Caracterización física y nutricional de pasturas, composición florística, reserva de carbono y CO<sub>2</sub>. *Sci. Agropecu.* 11 (2):167-176, 2020. DOI: <https://doi.org/10.17268/sci.agropecu.2020.02.03>.
- Walker, B. H. Biodiversity and ecological redundancy. *Conserv. Biol.* 6 (1):18-23, 1992. DOI: <https://doi.org/10.1046/j.1523-1739.1992.610018.x>.
- Wanapat, M.; Cherdthong, A.; Phesatcha, K. & Kang, S. Dietary sources and their effects on animal production and environmental sustainability. *Anim. Nutr.* 1 (3):96-103, 2015. DOI: <https://doi.org/10.1016/j.aninu.2015.07.004>.
- Yalta, J.; Ríos, N.; Valqui, L.; Bobadilla, Leidy G.; Vigo, Carmen N. & Vásquez, H. V. Huella hídrica de la producción lechera en la cuenca ganadera Pomacochas, Perú. *LRRD*. 33:125, <http://www.lrrd.org/lrrd33/10/33125hvasq.html>, 2021.
- Zúñiga-López, A.; Rodríguez-Serrano, A. C.; Benavides-Cruz, J. C.; Medrano-Galarza, Catalina & García-Castro, F. E. Indicadores de bienestar animal en vacas lecheras en un sistema silvopastoril del trópico alto colombiano. *Rev. investig. vet. Perú.* 31 (4):e16871, 2020. DOI: <http://dx.doi.org/10.15381/rivep.v31i4.16871>.