



Revista Latino-Americana de Enfermagem
ISSN: 1518-8345
Escola de Enfermagem de Ribeirão Preto /
Universidade de São Paulo

Occipital Neuralgia: a noninvasive therapeutic approach

López-Soto, Pablo Jesús; Bretones-García, José Miguel; Arroyo-García, Verónica; García-Ruiz, Margarita; Sánchez-Ossorio, Eduardo; Rodríguez-Borrego, María Aurora
Occipital Neuralgia: a noninvasive therapeutic approach
Revista Latino-Americana de Enfermagem, vol. 26, 2018
Escola de Enfermagem de Ribeirão Preto / Universidade de São Paulo
Available in: <http://www.redalyc.org/articulo.oa?id=281458425030>
DOI: 10.1590/1518-8345.2621.3067

Occipital Neuralgia: a noninvasive therapeutic approach

Neuralgia Occipital: un abordaje terapéutico no invasivo
Neuralgia Occipital: uma abordagem terapêutica não invasiva

Pablo Jesús López-Soto¹²³

Instituto Maimonides de Investigación Biomédica de Córdoba, Spain

ORCID: <http://orcid.org/0000-0002-1046-6686>

José Miguel Bretones-García⁴

Centro de Ergodinámica Córdoba, Spain

Verónica Arroyo-García⁴

Centro de Ergodinámica Córdoba, Spain

Margarita García-Ruiz³

Hospital Universitario Reina Sofía, Spain

Eduardo Sánchez-Ossorio⁵

Centro Ergodinámica Barcelona, Spain

María Aurora Rodríguez-Borrego¹²³

Instituto Maimonides de Investigación Biomédica de Córdoba, Spain

Revista Latino-Americana de
Enfermagem, vol. 26, 2018

Escola de Enfermagem de Ribeirão
Preto / Universidade de São Paulo

Received: 05 February 2018

Accepted: 13 August 2018

DOI: 10.1590/1518-8345.2621.3067

© 2018 Revista Latino-Americana de
Enfermagem
CC BY

ABSTRACT

Objective: to evaluate the application of a noninvasive intervention consisting of a postural modification using personalized models and osteopathy in people with occipital neuralgia.

Method: retrospective study of the intervention performed in adult population with occipital neuralgia, consisting of postural modification using personalized plantar orthoses and osteopathy, in a study period of four years. The observed variables were: persistence of headache, alignment of the axes, plantar support, center of gravity and center of mass; medical interview data, visual analogue scale, Win-Track gait analysis system and Kinovea software for video analysis (clinical assessment instruments used).

Results: a total of 34 records of people with occipital neuralgia were studied. A fraction of 58.8% of the patients reported improvement after the intervention. The visual analogue scale data were provided for 64.7% of the records and significant differences ($p < 0.001$) between the means before (8.4 ± 1.7) and after the intervention (2.6 ± 2.7) were found.

Conclusion: postural modification using personalized orthoses and osteopathy substantially improves the symptomatology of patients with occipital neuralgia.

Descriptors: Neuralgia++ Therapeutics++ Foot Orthoses++ Gait++ Pain++ Pain Management.

Resumen: Objetivo: evaluar la aplicación de una intervención no invasiva consistente en una modificación postural empleando plantillas personalizadas y osteopatía en personas con neuralgia occipital. Método: estudio retrospectivo de la intervención llevada a cabo en población adulta con neuralgia occipital, de modificación postural empleando órtesis plantares personalizadas y osteopatía, en un período de estudio de cuatro años. Las variables observadas fueron: persistencia de cefalea, alineación de los ejes, apoyo plantar, centro de gravedad y centro de masa; datos de la entrevista médica, la escala visual análoga, el sistema de análisis de la marcha Win-Track y el software Kinovea

para análisis de vídeo (instrumentos de valoración clínica utilizados). Resultados: un total de 34 registros de personas con neuralgia occipital fueron estudiados. Un 58,8% refería presentar mejoría tras la intervención. Los datos de la escala visual análoga eran proporcionados un 64,7% de los registros, encontrándose diferencias significativas (p menor que 0,001) entre las medias previas a la intervención ($8,4 \pm 1,7$) y aquellas posteriores ($2,6 \pm 2,7$). Conclusión: la modificación postural empleando órtesis personalizadas y osteopatía mejora sustancialmente la sintomatología de los pacientes con neuralgia occipital.

Descriptores: Neuralgia, Terapéutica, Órtesis Plantares, Marcha, Dolor, Manejo del Dolor.

Resumo: Objetivo: avaliar a aplicação de uma intervenção não invasiva que consiste em uma modificação postural usando modelos personalizados e osteopatia em pessoas com neuralgia occipital. Método: estudo retrospectivo da intervenção realizada em população adulta com neuralgia occipital que consiste de modificação postural empregando órteses plantares personalizadas e osteopatia, em um período de estudo de quatro anos. As variáveis observadas foram: persistência de cefaleia, alinhamento dos eixos, apoio plantar, centro de gravidade e centro de massa; dados da entrevista médica, escala visual analógica, sistema de análise de marcha Win-Track e o software Kinovea para análise de vídeo (instrumentos de avaliação clínica utilizados). Resultados: foram estudados no total 34 registros de pessoas com neuralgia occipital. Uma fração de 58,8% dos pacientes informou apresentar melhoria após a intervenção. Os dados da escala visual analógica foram fornecidos para 64,7% dos registros, encontrando diferenças significativas (p menor que 0,001) entre as médias antes ($8,4 \pm 1,7$) e depois da intervenção ($2,6 \pm 2,7$). Conclusão: a modificação postural empregando órteses personalizadas e osteopatia melhora substancialmente a sintomatologia dos pacientes com neuralgia occipital.

Descritores: Neuralgia, Terapêutica, Órtoses do Pé, Marcha, Dor, Manejo da Dor.

Introduction

Neuralgia is considered the most frequent neuropathy. It was described by Beruto and Lentijo and Ramos in 1821, being defined as a disabling alteration characterized by recurrent headaches located in the occipital region¹. According to the classification by the International Headache Society (IHS), occipital neuralgia (ON) is defined as “unilateral or bilateral paroxysmal pain, of lancinating or acute nature that is located in the posterior part of the scalp in the distribution of the major, minor and third occipital nerves, which on certain occasions is accompanied by reduced sensitivity or dysesthesia in the affected area and is usually associated to hypersensitivity of the affected nerve or nerves”². This type of neuralgia is classified into a subset that includes post-traumatic pain, lash, cervical spine deformity, tension headache, chronic daily headache and migraine³⁻⁴. In addition to occipital neuralgia, the scientific community employs other terms for its definition, such as C2 neuralgia, Arnold neuralgia, or occipital neuritis⁵⁻⁶. Other characteristics are the presence of pain that recovers the ocipucius that diminishes after the use of anesthetic blocks of the affected nerves^{2,4}.

ON is a multiple-cause clinical condition. In its manifestation, several factors⁷ may be involved: traumatic factors (fracture, hematoma, iatrogenic), anatomical factors (Chiari malformation, pinching, compression sensitivity), fibromyalgia, tumors (osteochondroma, neuroma, multiple myeloma, atlantoaxial lateral masses) infections

(pyomyositis, neurosyphilis, pachymeningitis) and degenerative changes (atlantoaxial lateral osteoarthritis, C1-C2 arthrosis syndrome).

The therapeutic approach is very broad. Regarding treatment, the conservative methods were: use of antiepileptics, antidepressants, nervous block, non-steroidal analgesics, opioids, neuromodulators, and transcutaneous electrical nerve stimulation, as well as cervical orthoses. However, the scientific community also approves the use of surgery as the last therapeutic option: neurolysis and decompression, neurectomy, rhizotomy and ganglionectomy, C1-C2 fusion, radiofrequency ablation, and peripheral nerve stimulation⁷. Nevertheless, the use of different therapeutic options does not always result in the remission of the symptomatology, together with the fact that they are painful. Therefore, it is considered necessary to propose therapeutic alternatives that lead to more effective therapeutic results.

The starting hypothesis of the present study was the fact that the appearance of the symptoms associated with occipital neuralgia was of biomechanical origin, considering the cause of the occipital neuralgia a postural alteration that has as consequence the nervous/venous compression of the root of the C2 vertebra⁸. In this sense, we describe the results that the research group obtained with the application of customized insoles in patients with ON, obtaining an improvement in the symptomatology of ON with the postural correction of standing/walking position, as well as osteopathy. Therefore, the objective of the study is to evaluate the application of a noninvasive intervention consisting of a postural modification using personalized insoles and osteopathy in people with occipital neuralgia (ON).

Methods

This is a retrospective observational study of pre/post-intervention records, which includes three periods of clinical evaluation (after 15 days of intervention, after 40 weeks and after one year). The objects of study were the clinical records of people older than 14 years who voluntarily sought a biomechanical clinical unit in southern Spain in the period between May 2012 and May 2016, with signs of ON.

We included records related to occipital neuralgia (previously diagnosed by a specialist), of persons older than 14 years and corresponding to the study period, in which there was a noninvasive intervention on the symptomatology.

In the clinical evaluation, the following variables were collected: socio-demographic characteristics (age and gender), clinical characteristics (years of persistence of neuralgia) and clinical-assistance characteristics. The latter variables were determined by the ergodynamic gait study [axis alignment, plantar support, center of gravity and center of mass (N/cm²)], posturology study [center of gravity and center of mass (N/cm²) and dysmetria of the lower limbs, as well as the level of pain (using the Visual Analogue Scale - VAS).

The noninvasive therapeutic intervention consisted in performing a postural modification of the affected person using plantar orthoses and osteopathy.

Subsequent to the application of the treatment, data were collected on the persistence of headache and whether some type of complementary medication was necessary, and it was advised, according to clinical criteria, some modification of the plantar orthoses. The treatment follow-up was performed in three observations (consultations at 15 days, 40 weeks and approximately one year from the start of treatment), in which the following tests were repeated in order to assess the efficacy of the intervention: ergodynamic gait study, posturology study and lower limb dysmetria; as well as post-intervention pain level.

As indicated, in addition to the medical interview for data collection, VAS was used to determine pain level, and baropodographies (N/cm^2) were used to determine the different pressures exerted by the foot, using the NOVEL-EMED-Model AT[®] device. As a gait analysis system, the Win-Track device (Medicapteurs, France-USA)⁹⁾ was used.

Finally, to carefully measure the postural axes, computer software Kinovea video analysis was used. This software draws a line from the external auditory canal to the caudal line through a normal axis. This line must coincide with distinct anatomical points (acromion-clavicular, trochanter and external-malleolus). The software measures the separation with each of certain points of the medial line, noting if there is any abnormal axis (kyphotic or lordotic).

The clinical-assistance variables were described by numerical values and percentages, and the differences in mean values and percentages were calculated by the Pearson test χ^2 and Fisher's test, according to the type of variable. For comparison of means of independent groups, the Student's t-test or the Mann Whitney U-test were used, according to normal or non-normal parameters. According to the type of variable, statistical tests of hypothesis contrast were performed, assuming a statistical significance of $p < 0.5$ and confidence intervals with 95% confidence.

The study was carried out in agreement with the institution's ethical guidelines and with the Helsinki Declaration, and with approval of the Research Ethics Committee (Registration No. 3281), abiding to the Organic Law 15/1999 on Personal Data Protection of the Spanish State.

Results

Thirty-four adult individuals with episodes of ON were treated with the noninvasive intervention. The mean age of people whose records were included in the study ($N = 31$) was 34.1 ± 10.9 years. There was a greater proportion of women (67.6%) than men. An important percentage (35.5%) had suffered for more than 10 years with neuralgia; and in 10 patients (29.4%) there was no record of this variable. In the other registries, it was documented that the duration of the condition was 5-6 years (11.8%), 2-3 years (11.8%), and from months to one year, only

three people (8.8%). Of the 34 patients studied, 17 (50%) reported taking previous medications to treat neuralgia, with anti-inflammatories being the most used type of medication (64.7%).

The ergodynamic gait study, posturology study and lower limb dysmetria evaluation were performed in all patients. Nevertheless, there are records of only one third of the subjects included in the study (N = 11, 32.3%).

Based on the reference points (acromioclavicular, major trochanter and peroneal malleolus), the established vertical did not correspond in 5 of the 11 subjects (45.4%). Besides the approximation to the reference points, the alignment between these points is also important, being the same the percentage of people (45.4%) who did not comply with this criterion, as shown in Figure 1.



Figure 1

Pre-intervention posturology study. Cordoba, Andalusia, Spain, 2012-2016

Plantar orthoses were prescribed to all patients in the study. However, osteopathy was performed in only 15 subjects (44.1%). The type of prescribed orthosis depended on the person's type of plantar support, center of gravity and center of mass, as shown in Figures 2 and 3.




Figure 2

Static baropodography prior to intervention. Cordoba, Andalusia, Spain, 2012-2016




Figure 3

Dynamic baropodography of left and right foot before the intervention. Cordoba, Andalusia, Spain, 2012-2016

As previously indicated, 2-3 follow-up visits were performed to know the evolution of the patients and potential improvement of their condition. The first control was performed in a time frame with a median after the intervention of 64.0 days (interquartile range - RI: 47.0-72.75), in 24 subjects (82.3%). At this visit, a high percentage of patients presented a favorable evolution (79.1% of the patients evaluated, only one of them was asymptomatic). In the other patients, it was necessary to manipulate with osteopathy or to place a loop in one of the limbs due to dysmetria. The next consultation ($N = 27$), scheduled for the sixth month after the intervention, was performed in a median time frame of 230.0 days (RI: 201.0-284.0). The symptomatology at this visit had disappeared

in five of the subjects studied (18.5% of those who attended) and was favorable in the same percentage. In this second consultation, in 44.1% of the total patients studied, no data was obtained. On the other hand, four patients (14.8%) required changes in the insoles; three decided not to continue with the insoles and presented recurrence of symptoms; and in two other subjects, the discomfort continued. The proposed consultation at 18 months (N=19) occurred after a period whose median was 608 days (RI: 546-885). Six of the patients reported being asymptomatic (31.5%). Although five of those who participated in the consultation (26.3%) reported that symptomatology had appeared, then the orthoses were modified. Another third of those who attended (36.8%) reported stability in symptomatology and one patient reported not agreeing with the treatment provided in the approach. At the last visit, with a wide variation in time, since many subjects did not require a high number of medical visits (N = 29), the period was a median of 1,312 days (RI: 409-1608.5) (43.1 month/3.6 years).

After the three follow-up visits, a large percentage of patients (N=13; 38.2%) reported no persistence of occipital neuralgia; 20.6% reported improvement and neuralgia was less persistent. However, five patients (14.7%) indicated that the symptomatology persisted and from 23.5% data was not obtained. Concerning VAS, only 64.7% (N=22) provided information on the records, with significant differences ($p < 0.001$) between means previous (2.6 ± 2.7) and after (8.4 ± 1.7) the intervention.

At the end of the study period, 20 of the 34 subjects studied (58.8%) remained with plantar orthoses and four of them did not use them (11.7%); of the other subjects, no data was obtained. On the other hand, 35.2% of the patients did not take any medication at the end of the study; 11.7% only took it when they had a crisis and 14.7% took anti-inflammatory drugs; regarding the other subjects studied, there is no data available.

Regarding the posturology study (N = 9), including the plantar orthosis, all the subjects presented improvement in the alignment of the axis, as well as the distance to said axis, according to Figure 4. Moreover, baropodographies with insoles showed improvement in the person's plantar support, center of gravity and center of mass, according to Figure 5.



Figure 4
Posturology study using plantar orthoses. Cordoba, Andalusia, Spain, 2012-2016




Figure 5
Static baropodography employing plantar orthoses. Left image: Left foot; Right: Right foot. Cordoba, Andalusia, Spain, 2012-2016

Discussion

ON has been a well-established entity for many years, but there is no consensus on the intervention to be performed as treatment. Usually, most are invasive conservative interventions and noninvasive measures are poorly considered. Nevertheless, the present study provides

information about the effectiveness of postural correction as an ON treatment with plantar orthosis, and when the clinician considered it appropriate, osteopathy.

The most common therapeutic measure is the infiltration of local anesthetic agents with and without steroids, a technique that alleviates pain in some cases (15% -36%)¹⁰⁻¹². In this sense, a high percentage of patients (58.8%) experienced pain reduction after orthoses implantation. In addition to infiltration with or without steroids, botulinum toxin A was used for its inhibitory effect on the motor plaque as a muscle relaxant, although its possible analgesic action was indirect¹³⁻¹⁴, reducing only the associated acute and pungent ON pain, not constant and only pain for several months¹⁵. The postural correction intervention showed continuous improvement in the three follow-up visits, a fact that strengthens the application of these measures in future studies.

On the other hand, some studies show the efficacy of pulsed radiofrequency in the treatment of ON. Nevertheless, the known studies are observational without controls and in the same way as in the previous case, only short and medium-term pain control is observed¹⁶⁻¹⁸.

Surgical techniques are also used for the treatment of ON, such as occipital neurolysis and stimulation of the occipital nerve. However, it has been documented that the use of these techniques increases the possibilities of developing entities of worse therapeutic control than neuralgia, for example, neuroma or regional pain syndrome¹⁹.

However, the present study has several limitations. Despite the fact that there were statistically significant differences after postural correction in ON symptomatology, the retrospective nature and consequently, follow-up losses, does not allow to determine exclusively the effectiveness of the treatment, due to the fact that there are no control groups. Moreover, the VAS scale determines the perception of pain, nevertheless, the control of neuropathic pain requires evaluation of the psychological aspects²⁰, a fact that is not usually considered in interventional studies in this neuralgia.

Conclusion

The application of customized orthoses, and in some cases osteopathy, substantially improves the postural alignment (acromion-clavicular, trochanter and external malleolus) and as a consequence, the symptomatology of ON. It is possible to conclude that after the noninvasive intervention, the level of neuropathic pain decreased significantly.

This therapeutic alternative, according to our knowledge, was not approached by the scientific community and could be considered as a first approach in the treatment of ON. On the other hand, clinicians should consider that invasive and/or surgical techniques may trigger less controllable clinical conditions than the underlying entity. In this sense, a

consensus is needed in the scientific community to establish an adequate therapeutic algorithm.

Despite the interesting results of our study, the data obtained should be considered with caution due to its limitations. Future prospective studies should be conducted that include control groups and evaluation of psychological variables that may influence the perception of pain.

Acknowledgements

The authors gratefully acknowledge the assistance of all the nurses and physicians who participated in the study and the relevant institutions involved.

References

- Perelson HN. Occipital nerve tenderness; a sign of headache. *South Med J*. [Internet]. 1947 [cited Dec 1, 2017];40(8):653-6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20252372>
- Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders. 3ed. *Cephalgia*. 2013;33(9):629-808. doi: 10.1177/0333102413485658
- Hecht JS. Occipital nerve blocks in postconcussive headaches: a retrospective review and report of ten patients. *J Head Trauma Rehabil*. 2004;19(1):58-71. doi: 10.1097/00001199-200401000-00006
- Ducic I, Hartmann EC, Larson EE. Indications and outcomes for surgical treatment of patients with chronic migraine headaches caused by occipital neuralgia. *Plast Reconstr Surg*. 2009;123(5):1453-61. doi: 10.1097/PRS.0b013e3181a0720e
- Ballesteros-Del Rio B, Ares-Luque A, Tejada-Garcia J, Muela-Molinero A. Occipital (Arnold) neuralgia secondary to greater occipital nerve schwannoma. *Headache*. 2003;43(7):804-7. doi: 10.1046/j.1526-4610.2003.03142.x
- Slavin KV, Nersesyan H, Wess C. Peripheral neurostimulation for treatment of intractable occipital neuralgia. *Neurosurgery*. 2006;58(1):112-9; discussion -9. doi: 10.1227/01.NEU.0000192163.55428.62
- Cesmebasi A, Loukas M, Hogan E, Kralovic S, Tubbs RS, Cohen-Gadol AA. The Chiari malformations: a review with emphasis on anatomical traits. *Clin Anat*. 2015;28(2):184-94. doi: 10.1002/ca.22442
- Lucas G, Laudanna A, Chopard R, Raffaelli E. Anatomy of the lesser occipital nerve in relation to cervicogenic headache. *Clin Anatomy*. 1994;7(2):6. doi: 10.1002/ca.980070207
- Ramachandra P, Maiya AG, Kumar P. Test-retest reliability of the Win-Track platform in analyzing the gait parameters and plantar pressures during barefoot walking in healthy adults. *Foot Ankle Spec*. 2012;5(5):306-12. doi: 10.1177/1938640012457680
- Hammond SR, Danta G. Occipital neuralgia. *Clin Exp Neurol*. [Internet]. 1978 [cited Dec 5, 2017];15:258-70. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/756019>

- Kuhn WF, Kuhn SC, Gilberstadt H. Occipital neuralgias: clinical recognition of a complicated headache. A case series and literature review. *J Orofac Pain*. [internet]. 1997 [cited Dec 5, 2017];11(2):158-65. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/10332322>
- Anthony M. Headache and the greater occipital nerve. *Clin Neurol Neurosurg*. [Internet]. 1992 [cited Dec 1, 2017];94(4):297-301. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/1335856>
- Cui M, Khanijou S, Rubino J, Aoki KR. Subcutaneous administration of botulinum toxin A reduces formalin-induced pain. *Pain*. 2004;107(1-2):125-33. doi: /10.1016/j.pain.2003.10.008
- Durham PL, Cady R. Regulation of calcitonin gene-related peptide secretion from trigeminal nerve cells by botulinum toxin type A: implications for migraine therapy. *Headache*. 2004;44(1):35-42; discussion -3. doi: 10.1111/j.1526-4610.2004.04007.x
- Taylor M, Silva S, Cottrell C. Botulinum toxin type-A (BOTOX) in the treatment of occipital neuralgia: a pilot study. *Headache*. 2008;48(10):1476-81. doi: 10.1111/j.1526-4610.2008.01089.x
- Choi HJ, Oh IH, Choi SK, Lim YJ. Clinical outcomes of pulsed radiofrequency neuromodulation for the treatment of occipital neuralgia. *J Korean Neurosurg Soc*. 2012;51(5):281-5. doi: 10.3340/jkns.2012.51.5.281
- Huang JH, Galvagno SM, Hameed M, Wilkinson I, Erdek MA, Patel A, et al. Occipital nerve pulsed radiofrequency treatment: a multi-center study evaluating predictors of outcome. *Pain Med*. 2012;13(4):489-97. doi: 10.1111/j.1526-4637.2012.01348.x
- Vanelderen P, Rouwette T, De Vooght P, Puylaert M, Heylen R, Vissers K, et al. Pulsed radiofrequency for the treatment of occipital neuralgia: a prospective study with 6 months of follow-up. *Reg Anesth Pain Med*. 2010;35(2):148-51. doi: 10.3410/f.2835957.2502064
- Slavin KV, Colpan ME, Munawar N, Wess C, Nersesyan H. Trigeminal and occipital peripheral nerve stimulation for craniofacial pain: a single-institution experience and review of the literature. *Neurosurg Focus*. [Internet]. 2006 [cited Dec 10, 2017];21(6):E5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/17341049>
- Choi I, Jeon SR. Neuralgias of the Head: Occipital Neuralgia. *J Korean Med Sci*. 2016;31(4):479-88. doi: 10.3346/jkms.2016.31.4.479

Author notes

Corresponding Author:

Pablo Jesús López-Soto E-mail: pablo.lopez@imibic.org