



Colombia Medica
ISSN: 0120-8322
ISSN: 1657-9534
Universidad del Valle

Rengifo, Aura Caterine; Durán, María Alexandra; Ortiz,
Yamileth; Rodriguez, Jorge Martín; Ospina, Martha Lucía
Reflections on Afro-descendant origin and the outcome of dengue fever cases in Colombia
Colombia Medica, vol. 48, no. 2, 2017, April-June, pp. 98-100
Universidad del Valle

Available in: <https://www.redalyc.org/articulo.oa?id=28354693014>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal
Project academic non-profit, developed under the open access initiative

Letters to editor

Reflections on Afro-descendant origin and the outcome of dengue fever cases in Colombia.

Reflexiones sobre el origen de los afrodescendientes y los casos de dengue en Colombia

Dear editors, cordial greetings

Article ref: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5091271/>

As a group of researchers and professionals signing this letter to the editors, respectfully, we want to make a series of observations and reflections regarding the article “Afro-Colombian ethnicity, a paradoxical protective factor against dengue”¹, published in your prestigious journal.

We considered that the article in question lightly concludes that the Afro-Colombian population present a lower risk of complications compared to the non-Afro-Colombian population. This situation, besides of not being completely based on data and facts that it describes and analyzes, is an assertion that can be a source of deeply disorientation for the Colombian health services, upon a time these conclusions get disseminated, and it can also mistakenly indicate to Afro-Colombian people that the risk of complications is lower in the presence of a virus causing a dengue disease.

In our point of view, several of the article’s affirmations can be object of xenophobic and exclusive evaluations towards the Afro-Colombian populations, mostly perceived when the authors conclude that in zones or neighborhoods with high number of Afro-Colombian there is a greater incidence of dengue among this population. The authors did not show data or measurements of this condition to identify whether (or not) Afro-descendants were infected at the site of evaluation and whether (or not) they developed the disease, so that these findings should have been assessed more rigorously before making this kind of affirmations and “risky” conclusion.

Although there are characteristics of Afro-descendants that may give to them less susceptibility to severe dengue manifestations, it is already known about the multifactorial pathogenesis of this disease; in which, variables such as environmental factors², Age³, gender⁴, different blood types AB versus types A, O and B⁵, allele polymorphisms of human leukocyte antigens⁶, and factors associated with leakage Vascular...among others, play an important role.

The conditions related to host genetics and environmental factors might predispose to severe forms of the disease, but the important thing is to bear in mind that most of them have been the result of conclusions obtained by studies of association. In addition,

recent meta-analysis studies on factors that may influence dengue’s manifestations indicate that malnutrition has an inverse association with dengue shock syndrome and dengue hemorrhagic fever^{7,8}. We might then, paradoxically, think that “malnutrition offers a kind of protection against the development of the severe form of the disease”. Therefore, being a person of black race does not mean that he/she is protected against some forms of dengue, neither it indicates that it is a risk factor for other ethnic groups⁹. It is also possible that the article is failing to measure or has bias in misclassification, regarding to the auto-recognition for each patient’s race¹⁰. A dichotomous classification as Afro-Colombian or non-Afro-Colombian can generate measurement errors; particularly, considering that an Afro descendant patient (mulatto, zambo, mestizo, and others) can be classified as non-Afro-Colombian. Additionally, genetics of their predecessors makes it much vulnerable to possible complications of external origin infectious events. It is possible that of the 402/431 cases of severe dengue cases with complications in non-Afro-Colombians reported by the authors, several of them have genetic elements and traits of Afro-Colombian population, generating a measurement error in which it would be needed to perform Genetic and biological measurements that were not considered.

On the other hand, it is confusing the fact that when comparing with the conclusions obtained by Rojas et al.¹ in relation to the epidemiological reports made by the National Institute of Health (NIH) in 2013, it differs substantially. For example, the NIH reported that the most prevalent provincial o territories (in order of magnitude) with cases of classical dengue and severe dengue are Tolima, Valle, Santander, Norte de Santander and Cundinamarca. In these territories/provinces, (with the exemption of Valle) there are low black race population¹¹.

Although the authors state that there is no risk of a possible ecological fallacy, it is obvious that most conclusions were obtained with data analysis of population-level records, which exceed individual conclusions¹⁰. To avoid this type of situation, multi-level or multi-step method validations are required to ensure consistency between the two measurement levels.

Finally, it should be considered that dengue is a complex disease in which mechanisms and severe forms of infection have not been fully elucidated, requiring robust and ethical studies to establish new theories about viral pathogenesis, as to avoid making wrong assumptions for the individuals under the study.

Authors:

Aura Caterine Rengifo¹, María Alexandra Durán², Yamileth Ortiz³, Jorge Martín Rodríguez⁴, Martha Lucía Ospina⁵

¹ Grupo de Morfología Celular. Instituto Nacional de Salud, Bogotá, Colombia.

² Red Nacional de laboratorios. Instituto Nacional de Salud. Bogotá, Colombia.

³ Subdirección de Investigación, Científica y Tecnológica. Instituto nacional de Salud. Bogotá, Colombia.

⁴ Director de Investigación en Salud Pública. Instituto Nacional de Salud. Bogotá, Colombia.

⁵ Directora del Instituto Nacional de Salud de Colombia. Bogotá, Colombia.

Conflicts of interest:

None to disclaim

References

1. Rojas-Palacios JH, Alzate A, Martinez-Romero HJ, Concha-Eastman AI. Afro-Colombian ethnicity, a paradoxical protective factor against Dengue. *Colomb Med (Cali)* 2016;47(3):133–141.
2. Thammapalo S, Chongsuvivatwong V, Geater A, Dueravee M. Environmental factors and incidence of dengue fever and dengue haemorrhagic fever in an urban area, Southern Thailand. *Epidemiol Infect.* 2008;136(1):135–143.
3. Were F. The dengue situation in Africa. *Paediatr Int Child Health.* 2012;32(s1):18–21.
4. Shekhar KC, Huat OL. Epidemiology of dengue/dengue hemorrhagic fever in Malaysia A retrospective epidemiological study 1973-1987. Part I: dengue hemorrhagic fever. *Asia Pac J Publ Health* 1992 -6(2):15–25.
5. Kalayanaroj S, Gibbons RV, Vaughn D, Green S, Nisalak A, Jarman RG. Blood group AB is associated with increased risk for severe dengue disease in secondary infections. *J Infect Dis.* 2007;195:1014–1017.
6. Coffey L, Mertens E, Brehin A, FernándezGarcía M, Amara A, Despres PH. Human genetic determinants of dengue virus susceptibility. *Microbes Infect.* 2009;11:143–156.
7. Trang NT, Long NP, Hue TT, Hung le P, Trung TD, Dinh DN, *et al.* Association between nutritional status and dengue infection: a systematic review and meta-analysis. *BMC Infect Dis.* 2016;16:172–172.
8. Huy NT, Van Giang T, Thuy DHD, Kikuchi M, Hien TT, Zamora J. Factors Associated with Dengue Shock Syndrome A Systematic Review and Meta-Analysis. *PLoS Negl Trop Dis.* 2013;7(9):e2412
9. Whitehorn J, Farrar J. Dengue. *Br Med Bull.* 2010;95(1):161–173.
10. Szklo M, Nieto J. *Epidemiología Intermedia: Conceptos y Aplicaciones.* Madrid: Diaz de Santos; 2000. pp. 17–151.
11. Sivigila Situación del dengue en Colombia hasta el 16 de febrero de 2013. [Nov. 2016]. Disponible en <http://www.ins.gov.co/noticias/paginas/situaci%C3%B3n-del-dengue-en-colombia-hasta-el-16-de-febrero-de-2013.aspx#.WKPSbPnhCM9>.

Authors Response: Reflections on Afro-descendant origin and the outcome of dengue fever cases in Colombia.

Article ref: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5091271/>

Answer letter to the Editor:

We have carefully read Rengifo *et al.*¹, views on our research on dengue and ethnicity². Most of their observations point to the classification of ethnicity in the study design, in which we recognize a great difficulty to establish the category of Afro-descendant to each one of the individuals. Some proposals to classify that ethnic group in Latin American populations, based on genetic and biological tests, have shown a great miscegenation, which makes classification difficult³. The method of self-recognition of the ethnicity is imperfect as we see in the methodology of our study; but it is still valid for the scientific community at this time, while other more efficient, reliable and robust methods arise^{4,5}.

We disagree with the letter correspondents in considering xenophobic any element of this research. The study design considered any risk of discrimination on the grounds that it directly affected the outcome, and that it safeguarded the principle of equity in the selection and evaluation of variables. Significant results were verified by adjusted models that in some cases refuted the original findings (cumulative incidence by age and ethnic origin). Our interpretations of the results also warn of a silent phase of dissemination of the epidemic in communities with a larger Afro-Colombian population, which may require sanitary adjustments to modify the case numbers of dengue and severe dengue.

Finally, we agree on the complexity of the disease and the limitations of ecological studies; but these two arguments do not detract from the value of the conclusions of the study beyond the limitations we recognize in the discussion. Public health research can contribute to better policy decisions in a region by allowing decision-makers to better understand a problem or situation. But defining what decisions authorities should take in health is a responsibility only of decision-makers, in which they have to take into account other aspects, such as social, economic, technical and reliance on other sciences⁶. In this way, our publication does not encompass defining moral and politically convenient decisions.

Author:

Jorge Humberto Rojas Palacios^{1,2}, Alberto Alzate³, Héctor Jairo Martínez Romero^{2,4}, Alberto Ignacio Concha-Eastman⁵

¹ Grupo de Epidemiología y Salud Pública, Secretaría de Salud Pública Municipal de Cali, Alcaldía Municipal de Santiago de Cali, Cali Colombia

² Grupo de Investigación en modelos y métodos matemáticos para el control y vigilancia del dengue, Cali, Colombia

³ Grupo de Investigación en Epidemiología y Servicios (GRIEPIS). Postgrado de Epidemiología, Universidad Libre, Cali, Colombia

⁴ Departamento de Matemáticas, Universidad del Valle. Cali, Colombia

⁵ Alcaldía Municipal de Santiago de Cali, Cali, Colombia

Corresponding author:

Jorge Humberto Rojas Palacios. Secretaría de Salud Municipal de Cali, Alcaldía Municipal de Santiago de Cali, Colombia. Mobile: 316 5252701. E-mail: jorge.rojas@cali.gov.co

References:

1. Rengifo AC, Durán MA, Ortiz Y, Rodriguez JM, Ospina ML. Reflections on Afro-descendant origin and the outcome of dengue fever cases in Colombia. *Colomb Med (Cali)*. 2017; 48(2): 98-9.
2. Rojas-Palacios JH, Alzate A, Martinez-Romero HJ, Concha-Eastman AI. Afro-Colombian ethnicity, a paradoxical protective factor against Dengue. *Colomb Med (Cali)*. 2016; 47(3): 133-41.
3. Wang S, Ray N, Rojas W, Parra MV, Bedoya G, Gallo C, et al. Geographic patterns of genome admixture in Latin American mestizos. *PLoS Genet*. 2008;4(3): e1000037.
4. Agudelo-Suárez AA, Martínez-Herrera E, Posada-López A, Rocha-Buelvas A. Ethnicity and health in Colombia: what do self-perceived health indicators tell us? *Ethn Dis*. 26(2): 147-56.
5. Larifla L, Beaney KE, Foucan L, Bangou J, Michel CT, Martino J, et al. Influence of genetic risk factors on coronary heart disease occurrence in Afro-Caribbeans. *Can J Cardiol*. 2016; 32(8): 978-85.
6. Feldhay R. The fragile boundary between the political and the academic. *Israel Studies Review*. 2013; 28(1):1-7.