



International Journal of Psychological Research

ISSN: 2011-2084

ISSN: 2011-7922

Facultad de Psicología. Universidad de San Buenaventura,
Medellín

Mento, Carmela; Rizzo, Amelia; Muscatello, Maria
Rosaria Anna; Zoccali, Rocco Antonio; Bruno, Antonio
Negative Emotions in Skin Disorders: A Systematic Review
International Journal of Psychological Research, vol. 13, no. 1, 2020, January-July, pp. 71-86
Facultad de Psicología. Universidad de San Buenaventura, Medellín

DOI: <https://doi.org/10.21500/20112084.4078>


Available in: <https://www.redalyc.org/articulo.oa?id=299067956008>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative



Vol 13, N° 1

<https://revistas.usb.edu.co/index.php/IJPR>

ISSN 2011-2084

E-ISSN 2011-7922

 OPEN ACCESS

Editor: Jorge Mauricio Cuartas Arias,
Universidad de San Buenaventura,
Medellín, Colombia

Manuscript received: 30-04-2019

Revised: 15-08-2019

Accepted: 23-09-2019

***Corresponding author:**

Amelia Rizzo

Email: amrizzo@unime.it

Copyright: ©2020. International Journal of Psychological Research provides open access to all its contents under the terms of the license [creative commons](https://creativecommons.org/licenses/by-nc-nd/4.0/) Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0)

Declaration of data availability: All relevant data are within the article, as well as the information support files.

Conflict of interests: The authors have declared that there is no conflict of interest.

Negative Emotions in Skin Disorders: A Systematic Review

Emociones negativas en enfermedades de la piel: una revisión sistemática

Carmela Mento¹, Amelia Rizzo^{2*}, Maria Rosaria Anna Muscatello², Rocco Antonio Zoccali², Antonio Bruno²

¹ *Department of Cognitive Sciences, Psychological, Educational and Cultural Studies, University of Messina, Italy.*

² *Department of Biomedical and Dental Sciences and Morphofunctional Imaging, University of Messina, Italy.*

Abstract.

The main purpose of this study is to describe how negative emotions were investigated in the sphere of dermatological diseases, in order (1) to summarize literature trends about skin disorders and emotions, (2) to highlight any imbalances between the most studied and neglected emotions, (3) and to offer directions for future research. A computerized literature search provided 41 relevant and potentially eligible studies. Results showed that the study of emotions in skin disease is limited to Sadness/depression and Fear/anxiety. The emotions of Anger and Disgust have been poorly explored in empirical studies, despite they could be theoretically considered a vulnerability factor for the development of skin disorders and the dermatological extreme consequences, as negative emotionality toward self and the pathological skin condition. The bibliometric qualitative analysis with VOSViewer software revealed that the majority of the studies have been focused on the relationships between vitiligo and Sadness/depression, dermatitis and Fear/anxiety, psoriasis, and Anger, suggesting the need of future research exploring Disgust and, in general, a wider emotional spectrum.

Resumen.

El objetivo principal de este estudio fue describir cómo se han investigado las emociones negativas en el ámbito de las enfermedades dermatológicas, con el fin de (1) resumir las tendencias de la literatura sobre las enfermedades de la piel y las emociones, (2) para resaltar las diferencias entre las emociones más estudiadas y las más descuidadas y (3) ofrecer direcciones para futuras investigaciones. Una búsqueda bibliográfica computarizada proporcionó 41 estudios relevantes y potencialmente elegibles. Los resultados mostraron que el estudio de las emociones en la enfermedad de la piel se limita a Tristeza/depresión y Miedo/ansiedad. Las emociones de la ira y el asco han sido poco exploradas en estudios empíricos, a pesar de que teóricamente podrían considerarse un factor de vulnerabilidad para el desarrollo de enfermedades de la piel y las consecuencias dermatológicas extremas, como la emocionalidad negativa hacia uno mismo y la condición patológica de la piel. El análisis cualitativo bibliométrico con el software VOSViewer reveló que la mayoría de los estudios se han centrado en las relaciones entre vitiligo y Tristeza/depresión, dermatitis y Miedo/ansiedad, psoriasis e Ira, lo que sugiere la necesidad de futuras investigaciones que exploren asco y, en general, un espectro emocional más amplio.

Keywords.

negative emotions, skin disorders, depression, anxiety, anger, disgust.

Palabras Clave.

emociones negativas, enfermedades de la piel, depresión, ansiedad, ira, asco.

1. Introduction

From the physiological point of view, the skin is an organ strictly linked to emotional activities: redness, pallor, sweating, itching can be consequences of somatic activation, expressing a range of affective feeling-states. The relationship between psychological distress and skin alterations has always been a topic of great interest for researchers and clinicians. Unlike many other organs in the body, the human epidermis has an immediate reaction to mental stress, so that several authors aimed to demonstrate the so-called “brain-skin connection” (Paus, Theoharides, & Arck, 2006; Arck et al., 2010). It was found that mental, physical, and emotional stress definitely affects the skin and this may depend on several reasons. In stressful conditions, in fact, hormones relapse encourages inflammation and the decreasing blood flow to the skin (Dhabhar, 2000), irritating the skin nerves and increasing inflammation or allergic reactions, with systemic alterations of neuroendocrine and immunological parameters (Pavlovic et al., 2008). Furthermore, skin recruits the immune system to fight, dysregulating the innate immune response and causing inflammations such as Rosacea, Acne, and Psoriasis (Yamasaki & Gallo, 2009). As a consequence, triggering a vicious circle, natural moisturizing factors, and plumping lipids production decline and skin healing, repair, and restoration processes are delayed (Eming, Krieg, & Davidson, 2007).

The connection between skin and mind has deep roots and is the object of a specific clinic and research field called Psychodermatology. Research hypothesis has arisen spontaneously on the basis of careful observation of clinical data in medical settings and active collaboration between dermatologists, psychiatrists and psychologists. In the last twenty years, a number of dermatological studies have shown that people affected by a skin disorder often have a related psychological problem. For example, Gupta and Gupta (2001) have observed that patients with dermatological diseases show high rates of comorbid Major Depressive Disorder (MDD), Obsessive Compulsive Disorder (OCD), Body Dysmorphic Disorder, Social Phobia, and Post-Traumatic Stress Disorder (PTSD), which required prescriptions for anxiolytic and antidepressant medications. Picardi, Abeni, Melchi, Puddu, and Pasquini (2000) showed that the prevalence of emotional disorders was even 25.2% in about 2500 outpatients of a dermatological clinic. More specifically, a prevalence of more than 30% was found in patients with acne, pruritus, urticaria and alopecia.

In the study carried on by Gieler, Niemeier, Kupfer, Brosig, and Schill (2001), a survey was sent to 69 dermatology hospitals. In nearly all dermatoses, with the exception of hyperhidrosis and seborrheic eczema, the authors found that the extent of emotional influences was increased, compared to a similar study performed ten years before. However, the most interesting result

was the discovery of a subpopulation of about 23% of dermatology patients with psychosomatic disorders.

Fritzsche et al. (2001), using the ICD-10 criteria for the assessment of mental disorders, found mental, emotional, and behavioral disorders –of which the most frequent were mood and anxiety disorders– in 46% of 86 dermatological patients. Depression and anxiety, in fact, are mostly risk factors for several skin diseases, as can be seen in empirical studies on psoriasis vulgaris (Devrimci-Ozguven, Kundakci, Kumbasar, & Boyvat, 2000).

Furthermore, there are some review studies that have sought to establish relationships between psychiatric disorders and skin disorders. Buljan, Buljan, and Situm (2005), arguing that the treatment of psychodermatologic disorders is almost impossible without a holistic team approach involving psychiatrist, dermatologist and psychologist, proposed a review about psychodermatologic disorders, the psychosomatic disorders, primary psychiatric disorders, and secondary psychiatric disorders. The review conclusion is that psychopharmacologic treatment with anxiolytics, antidepressants, antipsychotics, and mood stabilizers can be prescribed by the dermatologist after consulting the psychiatrist.

Even Yadav, Narang, and Kumaran (2013) presented a comprehensive review of salient features and treatment updates in primary psychiatric dermatoses and also discussed the secondary psychiatric morbidity. The authors’ aim was to allow the dermatologist to be able to initiate basic pharmacotherapy, to know about various non-pharmacological treatments, such as psychotherapy, and the right time to refer the patient to the psychiatrist.

Again, Tareen and Tareen (2015) described how major affective disorders (such as major depression and bipolar disorders), anxiety spectrum disorders (such as social phobia and obsessive-compulsive disorder), and some commonly encountered personality disorders are frequently encountered in patients with skin conditions, often complicating the treatment plan and creating a vicious cycle where both disease processes keep perpetuating each other.

More recently, Krooks, Weatherall, and Holland (2017) discussed the etiology, epidemiology, clinical presentation, diagnosis, and first-line treatment of specific primary psychiatric causes of dermatologic conditions, writing a review, useful as a guide for dermatologists to use while prescribing atypical antipsychotics, selective serotonin reuptake inhibitors, and tricyclic antidepressants and/or cognitive behavioral therapy.

However, this literature review highlights some critical points. First of all, the common denominator is that these studies are psychiatric/medical guidelines, focused on the dermatological clinical practice and management, and not on psychological features or to the comprehension of such mechanisms. As stated by Marshall, Taylor,

and Bewley (2016) “patients who present to a psychodermatology clinic usually believe they have a primary skin problem (though this is not always the case)” (p. 31). Hence, there is the need of a study taking into account not only the nosographic aspects, or specific skin disorders, but also the different types of skin disorders and affective components, by including the entire range of negative emotions, that is, sadness, fear, anger, and disgust.

In the comprehension of patient suffering it should be considered the link between skin, skin pathologies, emotions, emotion expression, and emotion pathology. Among skin disorders there is a psychosomatic quote. Several psychodynamics theories suggest that the affected skin represents the expression of a negative emotion that is difficult for the individual to differentiate and describe. However, the role played by emotions is still unclear. It has not been yet established if negative emotions can be considered a reaction to the pathological condition or if, as stable response patterns, they can increase the vulnerability for the development of skin diseases. To reach this purpose a literature review is necessary, in order to understand how negative emotions have been studied hereinbefore, and to plan future studies answering to this issue and helping clinicians to identify the most effective treatments, both dermatological and psychological, for these patients.

From these premises, the main aim of this study is to describe how negative emotions were investigated in the sphere of dermatological diseases, in order (1) to summarize literature trends about skin disorders and emotions, (2) to highlight any imbalances between the most studied and the most neglected emotions, (3) and to offer directions for future research.

2. Methods

A computerized literature searches of each item contained in the ICD-10 classification for skin diseases –from code L00 to L99– has been conducted on three scientific article databases: Psyc-INFO, Web of Science, and PubMed; moreover, in order to obtain even the most obscure information, an electronic search on Google Scholar has been carried on. For many skin disorders, the search did not produce any relevant results to the field of emotions. Figure 1 shows some of the terms inserted on the literature search, representing the most popular results. Anyway, at the beginning of the research, each skin disorder has been associated with each emotion.

Search limits restricted the results to: 1) research articles published in English between 2000 and 2017 and identified as empirical, epidemiological or longitudinal studies; 2) researches conducted in a clinical sample; 3) measuring negative emotions such as sadness, disgust, anger and fear. As regards exclusion criteria, the search did not include: a) Studies on Quality of Life (681); b)

Acne		Anxiety OR Fear
Rosacea		
Vitiligio		Depression OR Sadness
Psoriasis		
Pruritus	AND	Anger OR Aggression
Dermatitis		
Urticaria		Disgust
Alopecia		

Figure 1. Literature search terms.

Studies on caregivers of skin disordered patients (315); c) Studies on children or adolescents (287); d) Dermatological review studies, book chapters, dissertation, and thesis (92). Following this second stage in the screening process, 41 studies provided relevant and potentially eligible information for coding. According to the recommendation of Moher and the Prisma Group (2009), Figure 2 shows the detailed flow chart with all research steps and selection criteria.

For each publication, data regarding authors name, year of publication, country of the study (population, instruments language), the skin disorders observed, the psychological instruments used, and the main results, were entered in a spreadsheet (MS Excel 5.0) to allow the construction of synthetic tables. To understand the relationship between skin and emotions we have divided the studies on the basis of the primary emotions considered: 1) sadness; 2) fear; 3) anger; 4) disgust. However, despite anxiety and depression are separate syndromes, they often co-occur. Hence, data of the studies that considered both the diseases have been grouped, to avoid duplicate, resulting in two tables. Finally, the qualitative data analysis was conducted using VOSviewer 1.6.1, a software tool for constructing and visualizing bibliometric networks based on data downloaded from bibliographic databases (Van Eck & Waltman, 2014). It produces a bubble map through the analyses of words reported in publications’ titles and abstracts. Each bubble represents a keyword or a phrase. The bubble size and color, and inter-Buble distance indicate keyword’s frequency, citation per publication (CPP) count, and frequency of co-occurrence or corresponding terms in publications, respectively (Van Eck & Waltman, 2014).

3. Results

3.1 Skin disorders, Sadness/Depression and Fear/Anxiety

From a clinical point of view, sadness is the typical emotion of loss, disappointment, separation, mourning. The intensity of sadness depends on the value that the indi-




Figure 2. Systematic Review Flow Chart.

vidual attributes to the loss, and the purposes which the loss incurred compromise. The more it is considered indispensable, irreplaceable and irretrievable what it lacks, the more intense and prolonged will be the emotional response of sadness (Lewis, Haviland-Jones, & Barrett, 2010). It is necessary to distinguish depression by “moments of sadness”, which may be present in the life of every individual and usually have fairly obvious causes and are characterized by a limited duration. Instead, depression and the states of melancholy mood usually are long lasting and accompanied by: a) low self-esteem; b) loss of interest and/or pleasure in activities that are normally pleasurable for the subject (anhedonia); c) lack of motivation. In fact, depression is contemplated in the international classifications of mood disorders (Wakefield & Demazeux, 2016). Conceptually, it is impossible to study the emotion of sadness as a primary emotion, as the existing literature was mainly interested and focused on the pathological expression of sadness, i.e., depression. Moreover, in the case of dermatological disorders there may be a moment of physiological sadness linked to the diagnosis, but the more interesting object is the pathological, prolonged, and pervasive sadness, because of the increased level of subjective suffering and psychological pain. To date, too little is known about the role played by this emotion. It is consolidated that the two conditions co-occur, but it is not clear whether the depression is the result of the failure to accept the diagnosis (i.e., a secondary disease) or if the depression, as pre-existing condition, could increase the vulnerability to the development of the somatic disorder.

A similar discussion regards the emotion of fear, an immediate response to danger and undoubtedly a close

relative of anxiety. Both are the response to a “threat”. However, even though anxiety and fear are similar in their physiological manifestation (through physical symptoms such as rapid heartbeat, difficulty in breathing, sweating, tightness, etc.), they differ substantially: fear is an emotional reaction to a *real* danger, while anxiety is an emotional reaction to a *perceived* danger (Perusini & Fanselow, 2015). The studies analyzed refer to the latter category, that is, to the pathological fear activation (anxiety). Up to date, we can only hypothesize that anxiety may worsen the course of dermatological pathology or may negatively influence the outcome and management of the disease, which is often chronic. Moreover, is not established if the anxiety trait can predispose to the development of dermatological pathologies, although many of them have been found in correlation.

Table 1 represents the characteristics of the studies that explored Sadness/Depression and Fear/Anxiety in Skin Disorders. As stated, depression and anxiety often co-occur. We found 29 studies (only 8 measuring exclusively depression), which explored depression and anxiety in skin disordered patients. Studies increased in the last years, but are homogeneous in different continents such as Europe, Asia, and America, showing how the incidence of depression and anxiety are similar in different cultural and ethnic groups and also in the comparison between different skin disorders. However, it can be observed that the majority of the studies were focused on vitiligo and psoriasis (65.5%), and often the two groups are compared, with almost all studies reporting higher levels of depression and anxiety in favor of the psoriatic patients.

Table 1

Characteristics of the researches studying Sadness/Depression and Fear/Anxiety in Skin Disorders

Author	Country	Skin disorder	Sample	Measure	Results
Mattoo, Handa, Kaur, Gupta, and Malhotra (2001)	India	Vitiligo and Psoriasis	113 patients with vitiligo (30.11 ± 12.49) and 103 with psoriasis (40.91 ± 14.26)	ICD-10 psychiatric diagnoses and General Health Questionnaire (GHQ)	Positive cases of depressive episode were 22% vs 29% in vitiligo and psoriasis, respectively. Depression, anxiety, and total psychopathology levels were similar in the two GHQ positive subgroups
Sharma, Koranne, and Singh (2001)	India	Vitiligo and Psoriasis	30 untreated psoriasis and vitiligo patients aged 18–60 yrs	General Health Questionnaire (GHQ-H)	The prevalences of depression were 23.3% and 10% in psoriasis and vitiligo respectively and anxiety was observed in 3.3% of each group.
Esposito, Saraceno, Giunta, Mac-carone, and Chimenti (2006)	Italy	Psoriasis	2.391 patients	Center for Epidemiological Studies-Depression Scale (CES-D) questionnaire Depressive symptomatology was observed in 1.482/2.391 patients (62% overall)	
Engin, Uguz, Yilmaz, Özdemir, and Mevlitoglu (2008)	Turkey	Chronic idiopathic urticaria (CIU)	73 patients with CIU, and 34 healthy subjects matched for age and sex	Beck Depression Inventory (BDI), the Beck Anxiety Inventory (BAI)	When compared with healthy controls, patients with CIU had significantly higher BDI and BAI scores
Ahmed, Ahmed, and Nasreen (2007)	Pakistan	Vitiligo	100 patients 15-60 years with a mean age of 24.6 years	Psychiatric Assessment Schedule (PAS)	Major depressive illness (15), was the most frequent psychiatric illness followed by generalized anxiety (10), mixed anxiety and depression, social phobia, agoraphobia and sexual dysfunction.
Tabolli et al. (2008)	Italy	Pemphigus vulgaris and pemphigus foliaceus	58 patients of whom 47% < 50 years 53% >50 years	36-item short form health survey (SF-36); ASQ anxiety scale questionnaire; Clinical Depression Questionnaire	Patients with higher depression and anxiety had a more severe degree of mucocutaneous involvement
Ehsani, Toosi, Shahshahani, Arbabi, and Noor-mohammadpour (2009)	Iran	Neurotic excoriation, trichotillomania, delusion of parasitosis and dermatitis artefacta	178 patients with psychocutaneous disorder (40.5 ± 15.6 years)	DSM-IV diagnostic criteria	Mood and anxiety disorders were common in patients with dermatitis artefacta, as patients with neurotic excoriation

Dieris-Hirche, Gieler, Kupfer, and Milch (2009)	Germany	Atopic Dermatitis (AD)	62 adult atopic dermatitis patients and 62 healthy controls	HADS-D	A significantly higher level of suicidal ideation, anxiety and depression was shown among patients with atopic dermatitis
Abram, Silm, Maaros, and Oona (2009)	Estonia	Rosacea	70 consecutive patients (seekers) and 56 subjects with rosacea from population (non-seekers)	Mood Scale questionnaire and visual analogue scale (VAS)	Higher mean VAS scores were not related to severity of rosacea, but were associated with the presence of depressive symptoms among seekers.
Layegh, Arshadi, Shahriari, Pezeshkpour, and Nahidi (2010)	Iran	Psoriasis, acne, alopecia and vitiligo	300 patients with a mean age of 26.55 ± 10.81 years.	Beck depression questionnaire	The prevalence rate of clinical depression was found to be 47.4% in patients with acne; 69.4% in those with psoriasis; 70.1% in those suffering from vitiligo; 50% in diffuse alopecia areata; 60% in universalis alopecia areata; 100% in ophiasis alopecia areata and 68.3% in patients with localized alopecia areata.
Uhlenhake, Yentzer, and Feldman (2010)	North Carolina	Acne	Acne patients aged 18 and over	Total Resource Utilization Benchmarks ^{OTM}	Depression was two to three times more prevalent in acne patients than in the general population, with a reported 8.8% of acne patients having clinical depression.
Ghajarzadeh, Ghiasi, and Kheirkhah (2012)	Iran	psoriasis, vitiligo, and alopecia areata	300 patients (100 with alopecia areata, 100 with psoriasis and 100 with vitiligo)	Beck Depression Inventory (BDI), SF-36 and Dermatology Life Quality Index (DLQI)	Significant correlation was found between DLQI and BDI in all disease groups ($r = 0.44, P < 0.001$). BDI scores were the highest in psoriasis group but this difference was not significant ($P = 0.2$).
Rashid et al. (2011)	Bangladesh	Vitiligo and Psoriasis	50 patients with psoriasis 35.32 ± 10.05 years and same number of patients with vitiligo 33.50 ± 9.99 years and were compared	Diagnostic and Statistical Manual for Mental and Behavioural Disorders-Fourth Edition)DSM-IV	The proportion of major depressive disorders (MDD) was higher among psoriasis patients 15(30.0%) compared to vitiligo patients 04(8.0%). However, no statistical difference was found between two groups of patients in terms of anxiety disorders (8% in psoriasis and 12% in vitiligo patients).
Chan et al. (2012)	Singapore	Vitiligo	145 vitiligo patients aged 21+	Strucured questionnaire	Among the patients, 17.2% ($n = 25$) had been identified as depressed.
Alfani et al. (2012)	Italy	Alopecia Areata	73 patients and 73 controls 35.2 ± 9.2 and 35.1 ± 9.1 years, respectively	Minnesota Multiphasic Personality Inventory (MMPI-2)	Some scales (i.e. Depression, Anxiety, Family relationships) were higher for patients with alopecia areata than for controls.

Al-Harbi (2013)	Saudi Arabia	Vitiligo	308 vitiligo patients	Beck Depression Inventory Scale	According to the Beck Depression Scale, 54.5% were found to be depressed, most with mild depression.
Settineri, Guarneri, Saitta, Mento, and Cannavò (2013)	Italy	Seborrheic dermatitis and Psoriasis	Respectively: 33 with SD (39.85 ± 15.45) and 36 psoriatic (41.11 ± 13.72) years	Profile of Mood States (POMS) and the Toronto Alexithymia Scale TAS-20	Two different depressive profiles emerged: psoriatic showed symptoms of major depression, SD patients to minor depression.
Sellami et al. (2014)	Tunisia	Alopecia areata (AA)	50 patients-mean age 32.92 years	Hospital Anxiety and Depression scale questionnaire, Toronto Alexithymia scale 20, and Severity of Alopecia Tool.	Depression and anxiety were detected respectively in 38% and 62% of patients.
Duman, Ozdemir, Yucel, and Akin (2014)	Turkey	Pilonidal sinus	205 patients: 107 operated with primary closure (26.4 ± 4.5 years) and 98 Limberg flap reconstruction (28.3 ± 4.8 years)	Beck Depression Inventory; Beck Anxiety Inventory; Short Form 36	Mean Beck Depression Inventory (19 ± 6.13 vs. 16 ± 4.90 $p < 0.001$) and Beck Anxiety Inventory (19 ± 6.27 vs. 16 ± 4.90 $p < 0.001$)
Dalgard et al. (2015)	13 countries	Infections of the skin and subcutaneous tissue	3635 patients aged 47.2 ± 17.9 years	Hospital Anxiety and Depression Scale	Clinical depression = 10.1% Clinical Anxiety = 17.2% Suicidal Ideation = 12.7%
Pärna, Aluoja, and Kingo (2015)	Estonia	Dermatitis and Eczema	40 patients with psoriasis, 40 with eczema, 40 with acne, 15 with seborrheic dermatitis and 40 healthy controls ($M = 38.9 \pm 14.5$)	Emotional State Questionnaire (EST-Q); Dermatology Life Quality Index (DLQI); RAND-36-item HRQoL survey	Depression was similar in all dermatological groups. Anxiety was significantly higher in dermatological patients, compared to healthy controls ($p = .005$).
Kim et al. (2015)	Korea	Atopic Dermatitis (AD)	1517 (19.8 ± 1.0 years)	Korean Military Multiphasic Personality Inventory (KMPI)	The adjusted odds ratios for depression, anxiety, and somatization were significantly greater for individuals with AD compared with those without AD.

Cheng et al. (2015)	Taiwan	Atopic Dermatitis (AD)	8208 (32.60 ± 16.06 years)	ICD-9 Diagnosis	Higher incidence of depressive disorder With AD=273(4.32) Without AD=48(0.74) <0.001 and Anxiety disorders With AD=180(2.83) Without AD=45(0.70) <0.001.
Konda et al. (2015)	India	Prurigo Nodularis (PN)	39 patients with PN (27.87 ± 17.6) and 39 age and gender matched healthy controls	Pruritus Grading Scale; Pittsburgh Sleep Quality Index; Hamilton rating scale for depression (HDRS)	Severity of depressive symptoms was significantly higher in PN patients ($U = 350.5; p = 0.01$).
Khatttri, Bist, Arun, and Mehta (2015)	India	Vitiligo	50 patients with vitiligo and 50 healthy controls, aged between 20 and 70 years old	DSM-IV Axis I Disorders (SCID-I), Hospital Anxiety Depression Scale (HADS), Rosenberg Self-Esteem Scale (RSES), Liebowitz Social Anxiety Scale (LSAS) and Sheehan Disability Scale (SDS)	In comparison to healthy controls, the rate of depression and anxiety was found to be higher and mean self-esteem score was found to be lower in the vitiligo group. There was no significant difference between groups in terms of social anxiety.
Egeberg, Hansen, Gislason, and Thyssen (2016)	Denmark	Rosacea	30.725 and 24.712 patients with mild and moderate-to-severe rosacea > 18 years	Data on 4.632.341 Danish citizens	Mild and moderate- to-severe rosacea increased the risk of both depression [IRR 1.89 (95% CI 1.82–1.96) and IRR 2.04 (95% CI 1.96–2.12)] and anxiety disorders [IRR 1.80 (95% CI 1.75–1.86) and IRR 1.98 (95% CI 1.91–2.05)].
Halioua, Cribier, Frey, and Tan (2017)	UK, France, Germany and US	Rosacea	807 (31.3 ± 14.5 years)	Online survey of the general population over 18 years of age	Those with FS were more likely to avoid social situations (54.2% vs. 2.0%, $p < 1.00E - 10$) and had a higher rate of depression (36.7% vs. 21.1%, $p < 1.00E - 10$).
Wojtyna, Łakuta, Marcinkiewicz, Bergler-Czop, and Brzezińska-Wcisło (2017)	Poland	Psoriasis	219 patients with psoriasis, aged 18–70 years	Beck Depression Inventory, the Appearance Schemas Inventory-Revised, the Berlin Social Support Scales, and the Distress Thermometer	The main contributors to depression were: female gender, beliefs about appearance and its salience to one's self-worth, greater psychological distress, and lower levels of emotional social support.
Vernwal (2017)	India	Vitiligo	100 vitiligo patients and 100 controls	Hospital Anxiety and Depression Scale	In comparison to healthy controls, the psychiatric morbidity was found to be significantly higher in the vitiligo group (62% v/s 25%). 37%, 18%, and 7% vitiligo patients suffered from Mixed anxiety and depressive disorder, Depressive disorder and Generalized anxiety disorder respectively.

3.2 Skin disorders and Anger/Aggression

For most of the theories, anger is the typical reaction to both physical and psychological frustration and constriction. The causal relationship between frustration and anger is quite complex. Other factors seem in fact to be involved in the origins of anger emotion as, for example, the responsibility and the awareness attributed to the person or the situation that causes frustration or constriction (Averill, 2012). The emotion of anger can be defined as the reaction that follows a precise sequence of events: a) a state of need; b) an object (living or nonliving) that is opposed to the realization of the need; c) the attribution of intentionality to such object of opposing; d) a lack of fear toward the frustrating object; e) the strong intention to remove and attack the frustrating object; f) an act of aggression implemented through the attack (Boswell, 2016). Therefore, there are different kinds of anger and specifically three: anger directed toward others, self-directed or denied (Livingstone, Shepherd, Spears, & Manstead, 2016).

Table 2 shows the studies related to Anger/Aggression in Skin Disorders. The studies that have found the emotion of anger, in association with skin diseases, is quite low. Only 11 out of 41 studies (29%) addressed this emotion, finding that this is one of the emotions mostly experienced by people with psoriasis and atopic dermatitis, followed by urticaria and erythema, chronic idiopathic urticaria, acne and vitiligo.

In particular, Conrad et al. (2008) measured alexithymia (TAS-20), emotional distress (SCL-90-R), and anger (STAXI) in skin-disordered patients, founding that that chronic idiopathic urticaria (CIU) and psoriasis are associated with personality-based difficulties in emotional regulation, particularly with regard to the feeling of anger. However, it remains unclear if anger is directed toward others, self-directed or denied. The high prevalence of alexithymia seems connected with difficulties in communicating anger and in both skin disorders, the relationship between pruritus as a major symptom of negative affectivity could explain anger towards self and others.

3.3 Skin Disorders and Disgust

Disgust is an emotion characterized by a feeling of nausea, revulsion, especially in respect of odors and flavors, and, figuratively speaking, moral repulsion, intolerance, hate with a strong psychological value. It is manifested by a behavior tending to remove something or someone, and from a physiological reaction, such as nausea and vomiting in response to dirt and contamination sensations (Lateiner & Spatharas, 2016).

Over the years, not many authors have given their contribution to the empirical study of the emotion of disgust, but several studies show that disgust has the function to keep us away from situations or substances which could be dangerous, or from anything that is

abhorrent and dirty, including values, thoughts, people, and, in some cases, even the self (Yoder, Widen, & Russell, 2016). Although it can be assumed that disgust plays a key role in the perception of body image and dermatological disease, the studies on this topic are quite low.

Unfortunately, only one study addressed the emotion of disgust in patients with psoriasis. Lahousen et al. (2016), in fact, investigated the differences psoriasis patients and skin-healthy controls concerning appraisal of touching, shame and disgust in one hundred and seventy-one patients with psoriasis and 171 skin healthy controls who completed the Touch-Shame-Disgust-Questionnaire (TSD-Q), obtaining that skin-related disgust and shame were significantly higher in psoriasis patients. The study of disgust is one of the most neglected, not only in the field of psychology and psychiatry, but also in its applications in the dermatological context, presenting a significantly low incidence even in the field of psychodermatology.

4. Qualitative Analysis

Thanks to VOSviewer has been possible to analyze a text that contained the main information of a paper i.e. the disease and the associated primary emotion studied. On the basis of data collected in the review of the forty-one studies, a list has been formed. Figure 3 shows the bibliographic landscapes of the studies about skin disorders and emotions. The more the colors are close to red, the higher their occurrence and the closer the words, the higher the co-occurrence.

The qualitative analysis provides visual information on: 1) the proximity of vitiligo and sadness/depression; 2) the proximity of fear/anxiety and atopic dermatitis; 3) the proximity of psoriasis and anger; 4) the apparent independence of disgust.

This visual analysis remarks on the findings discussed: literature on emotions and skin diseases has been focused on specific emotions, neglecting others. In particular, the map highlights the imbalance between studies that addressed depression and anxiety and studies that considered anger and/or disgust. In fact, still little is known about the involvement of anger and its role as a predisposing factor and also as a secondary problem towards the dermatological pathology. Furthermore, the emotion of disgust is under-represented: in comparison to other emotions, disgust is the most neglected, despite it is undoubtedly involved. The individual can feel disgust against his/her body (self-disgust) or can believe that the other perceives him/her as disgusting.

5. Discussion

The present review study was aimed to describe in what way negative emotions were investigated in the sphere

Table 2

Characteristics of the researches studying Anger/Aggression in Skin Disorders

Author	Country	Skin disorder	Sample	Measure	Results
Sayar, Ugurad, Kural, and Acar (2000)	Turkey	Acne	31 acne vulgaris patients and 25 controls	Beck Depression Inventory, State-Trait Anxiety Inventory, Rosenberg Self-Esteem Scale, and Spielberger State-Trait Anger Expression Inventory.	No significant difference was detected on anger-related subscales between the acne and control groups.
Conrad et al. (2008)	Germany	Chronic idiopathic urticaria (CIU) and psoriasis	41 CIU patients 44 psoriasis patients and 49 healthy controls	Alexithymia (TAS-20), emotional distress (SCL-90-R) and anger (STAXI).	Anger was the only significant predictor of pruritus severity in CIU.
Linder et al. (2009)	Italy	Psoriasis	323 patients (aged 18–85 years; mean –SD age 51.2 – 28.7 years)	Interview and a questionnaire developed by clinicians and researchers	Psoriasis elicited anger, annoyance at the inconvenience of the disease, and irritation in approximately 50% of the patients
Kossakowska, Poland Cieś- cińska, Jaszewska, and Placek (2010)		Vitiligo and Psoriasis	60 patients with psoriasis ($n = 30$) and vitiligo ($n = 30$) as well as healthy persons ($n = 60$) matched	Watson and Greer's Courtauld Emotional Control Scale and Kossakowska's Chronic Patients Questionnaire.	Psoriasis patients control negative emotions more intensively than healthy people. Vitiligo patients do not differ in the control of negative emotions compared with healthy subjects.
Sampogna, Tabolli, and Abeni (2012)	Italy	Psoriasias	936 patients	Skindex-29 questionnaire	The emotions most frequently experienced were: shame, anger, Worry.
Takaki and Ishii (2013)	Japan	Atopic Dermatitis (AD)	43 adult patients with AD, 32 adults with remission from AD, and 63 adults without AD 20–47 years	Anger expression trait scale (AX)	Results indicated a significant impact of anger suppression on depression, in patients with AD.

Altınöz, Taşkın-tuna, Altınöz, and Ceran (2014)	Turkey	Chronic spontaneous urticaria (CSU) and alopecia areata (AA)	30 CSU 30 AA 30 controls	Hospital Anxiety and Depression Scale and Multi-Dimensional Anger Inventory	More of the CSU patients were observed to respond with excessive anger to most situations, to have high levels of anxiety anger and passive aggressive interpersonal Relationships.
Sarkar, Sarkar, Saha, and Sarkar (2014)	India	Psoriasis	Forty-eight patients of psoriasis and equal number of healthy controls	Self-reporting questionnaire-24 (SRQ-24) and skindex (A 61-item survey questionnaire)	the most common psychiatric morbidity in psoriasis patients was anger (58.3%)
Altunay and Demirci (2015)	Turkey	Pruritus	40 patients with a mean age 46.55 ± 13.20 years	State-Trait Anger Expression Index and the Mini International Neuropsychiatric Interview (M.I.N.I)	Pruritus duration was correlated with anger trait and anger-in subscale scores in all patients.
Aydin et al. (2017)	Turkey	Psoriasis	Eighty-five patients with psoriasis and 86 healthy controls	State-Trait Expression Inventory for Anger (STAXI) and Roserberg Self-esteem Scale (RSES)	Reduced self-esteem and increased anger levels are remarkable in psoriasis patients.
(Coneo, Thompson, & Lavda, 2017)	England	Dermatology patients	ninety-one participant	Optimism (LOT-R), perceptions of social support (SFSSQ) social acceptance, fear of negative Evaluation (FNE), appearance concern (CAR-VAL/CARSAL), appearance discrepancy PADQ), social comparison (INCOMM) and wellbeing (HADS) on aggression levels (RAQ)	Anxiety was also found to have a significant positive relationship with aggression ($\beta = 0.356, t = 2.564, p = 0.01$).




Figure 3. Bibliometric landscape of primary emotions and skin diseases.

of dermatological diseases, in order to highlight any possible imbalances between the most studied and the most neglected emotions, to identify a possible association model between the type of dermatological condition and the emotion studied, and to offer cues for future research.

Clinical observation suggested that dermatological patients live an intense emotional activation. From the moment of the received diagnosis they have to face psychological difficulties and have to manage disease acceptance and therapy management. This can raise different reactions in the individual: from depression for the loss of health, to anxiety for symptoms worsening, to anger (why me?), until disgust towards pustule, desquamation, rash, in other terms, against his/her own skin and body. The loss of a healthy and attractive image can severely impair the relationships with the body image and with others. Studies on quality of life, despite interesting and useful, do not cover or explain the aspects more related to the inner emotional experience. To reach this purpose, i.e., to understand how negative emotions are declined among skin disordered patients, a review was necessary as a first step in the comprehension of the psychological functioning.

The present review suggests –despite taking into account that studies are published from one to two years after their actual implementation– an increasing interest, over the last five years, by authors who explored the role of emotions in skin diseases, or more likely their emotional consequences. It is increasingly evident the impact of the psychosomatic medicine, the collaboration

between dermatologists, psychiatrists, and psychologists, and the vision of the disorder in the psyche-soma unit (Muscatello, Bruno, Scimeca, Pandolfo, & Zoccali, 2014; Rizzo et al., 2018). However, the interest on negative emotions in skin disorders reveals some contradiction and imbalance. For example, we observed that the exploration of the emotional aspects has been more successful for some diseases such as psoriasis, dermatitis and vitiligo. This may depend on the epidemiology of the disease and thus on the rarity or representativeness of the observations, or from a possible greater interest for the diseases in which the psychological implications are more observable. For example, the incidence of psoriasis in adults varies from 78.9/100,000 person-years in the United States to 230/100,000 person-years in Italy. The occurrence of psoriasis varies also according to age and geographic region, being more frequent in countries more distant from the equator and in the adult age range (Parisi, Symmons, Griffiths, & Ashcroft, 2013) while the worldwide incidence of pemphigus is 0.75-5 cases/1,000,000 per year. Most cases of PV in North America, Europe and Asia are sporadic, with a higher incidence among Ashkenazi Jews, with an estimate of 1.6 per 100,000 populations per year in Jerusalem (Femiano, 2007).

The imbalance of the epidemiology of skin diseases fails to explain the imbalance in the study of primary emotions, which was found rather inhomogeneous.

Most of the studies have shown consistent results. For example, we have found that: a) sadness was studied

in terms of depression and, therefore, in its pathological manifestation and not physiological; b) most of the studies addressed the fear in terms of anxiety, or a reaction to a perceived danger; c) many of the studies that have studied depression have found an association with anxiety, as already consolidated in the literature.

While other studies raise some controversial issues. Results show that the emotion of disgust is considered involved in dermatological disorders by different authors, but is not sufficiently represented on the side of empirical exploration. In other words, the authors argue that disgust is undoubtedly associated with, for example, papulosquamous or bullous diseases as a theoretical assumption, but empirical studies on clinical samples and plausibly appropriate tools for measuring the emotion of disgust are almost completely unavailable (Settineri & Mento, 2014).

The study of the emotion of anger, on the other hand, resulted very limited. It becomes difficult to determine whether or not this emotion is involved in skin diseases, as it would suggest the clinical observation. The collected data must take into account a possible bias: when the emotion of anger appears in the studies cited, could be the result of variability due to chance (e.g., an upstream selection of the variable in the research design, a lack of publication of studies that disconfirm the involvement of anger in skin diseases, etc.). For these reasons, future studies on the role of anger and disgust in dermatological diseases would be advantageous.

6. Limitations

Of course, this review has also several limitations. Having included some reviews, it is possible that the number of studies dealing with depression and anxiety has been underestimated. However, this could slightly modify results, since depression and anxiety resulted as the most studied among skin problems.

On the contrary, some of the studies cited also analyzed the relationship between skin disorders and the feelings of shame or guilt. These emotions have been excluded because should need specific space of discussion and scientific deepening. In clinical practice, we observe that the patient is immersed in a mix of different emotions with regard to the dermatological problem, because it has an impact on the self-image and the social and relational sphere. Every dermatological disease has its particular features, some of them affect exposed body areas, such as face or neck, and can generate intense negative emotions not only towards self, but also against other leading to avoidance of exposure, fear of judgement, fear of intimacy, etc.

Besides, due to the disparity between the studies, but also to the heterogeneity of the dermatological disorders treated, it was not possible to conduct a meta-analysis. Unfortunately, statistical conclusions and comparisons

are not applicable to this research. Hence, a problem emerges with respect to the comparison of results. This study has not mitigated the effects of measuring negative emotions through different tools, which span from structured interview to self-report scales or questionnaires vs. DSM-IV or ICD-10 diagnostic criteria. This makes the discussion a narrative reflection of a qualitative nature, supported by the visual analysis, as a bibliographic landscape.

Finally, the failure to consult of search engines (e.g., Scopus) most appropriate for this kind of study, may result in the exclusion of manuscripts useful for the development of the revision theme.

7. Conclusion

We believe the most innovative points are two. The first is that, to our knowledge, the present review considers the whole negative emotional spectrum, i.e., not only depression and anxiety, which are the most studied, but also anger and disgust. The second point is the application of an innovative method of bibliographic data analysis software (VOS), resulting in a graphical representation of the studies on the topic. This method allows to create a sort of map, a landscape that immediately suggests to the reader the “state of art”, arising new issues and future research cues.

Dermatology is an interesting research field, a preferential point of observation to understand the somatization mechanisms and the mind-body relationship. The link between emotional reactions and skin responses is the background of the success of techniques such as biofeedback, which allows visualizing the skin reactions to a series of stimuli (Shenefelt, 2010). Since the emotional reaction steps away from the skin (Damasio, 2003), the visualization of one's emotional reactions can help the individual to recognize the importance of these reactions and to manage them more effectively.

In conclusion, as well as in many other fields of medicine, in which emotions intervene (Settineri, Rizzo, Liotta, & Mento, 2017; Mento, Le Donne, Crisafulli, Rizzo, & Settineri, 2017; Mento et al., 2015), the deepening of the neglected aspects can foster a better understanding of the skin disorder and can be the basis of therapeutic interventions increasingly targeted, such as biofeedback technique and psychotherapy based on emotion regulation and mindfulness based stress reduction.

8. Contributors

R.A. and B.A. conducted literature searches, analysis and wrote the manuscript.

Z.R.A. and M.M.R.A. contributed to and have approved the final manuscript.

M.C. conceived the study and wrote the hypothesis.

9. Compliance with Ethical Standards

Funding. This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Conflict of Interest. The authors declare that there is no financial, general, and institutional conflict of interest regarding the publication of this article.

Ethical approval. This article does not contain any studies with human participants performed by any of the authors.

References

- Abram, K., Silm, H., Maaroos, H., & Oona, M. (2009). Subjective disease perception and symptoms of depression in relation to healthcare-seeking behaviour in patients with rosacea. *Acta dermato-venereologica*, 89(5), 488–491.
- Ahmed, I., Ahmed, S., & Nasreen, S. (2007). Frequency and pattern of psychiatric disorders in patients with vitiligo. *J Ayub Med Coll Abbottabad*, 19(3), 19–21.
- Alfani, S., Antinone, V., Mozzetta, A., Pietro, C., Mazzanti, C., Stella, P., & Abeni, D. (2012). Psychological status of patients with alopecia areata. *Acta dermato-venereologica*, 92(3), 304–306.
- Al-Harbi, M. (2013). Prevalence of depression in vitiligo patients. *Skinmed*, 11(6), 327–330.
- Altınöz, A., Taşkıntuna, N., Altınöz, S., & Ceran, S. (2014). A cohort study of the relationship between anger and chronic spontaneous urticaria. *Advances in therapy*, 31(9), 1000–1007.
- Altunay, I., & Demirci, G. (2015). The relationship of functional pruritus with anger and associated psychiatric disorders. *Turkderm*, 49(A00101s1), 28–32.
- Arck, P., Handjiski, B., Hagen, E., Pincus, M., Bruenahl, C., Bienenstock, J., & Paus, R. (2010). Is there a ‘gut-brain-skin axis’? *Experimental dermatology*, 19(5), 401–405.
- Averill, J. R. (2012). *Anger and aggression: An essay on emotion*. Springer Science & Business Media.
- Aydin, E., Atis, G., Bolu, A., Aydin, C., Karabacak, E., Dogan, B., & Ates, M. (2017). Identification of anger and self-esteem in psoriasis patients in a consultation-liaison psychiatry setting: a case control study. *Psychiatry and Clinical Psychopharmacology*, 27(3), 222–226.
- Boswell, J. (2016). Recognizing anger in clinical research and practice. *Clinical Psychology: Science and Practice*, 23(1), 86–89.
- Buljan, D., Buljan, M., & Situm, M. (2005). Psychodermatology: a brief review for clinicians. *Psychiatria Danubina*, 17(1–2), 76–83.
- Chan, M., Chua, T., Goh, B., Aw, C., Thng, T., & Lee, S. (2012). Investigating factors associated with depression of vitiligo patients in Singapore. *Journal of clinical nursing*, 21(11–12), 1614–1621.
- Cheng, C., Hsu, J., Huang, K., Bai, Y., Su, T., Li, C., & Chen, M. (2015). Risk of developing major depressive disorder and anxiety disorders among adolescents and adults with atopic dermatitis: A nationwide longitudinal study. *Journal of affective disorders*, 178, 60–65.
- Coneo, A., Thompson, A., & Lavda, A. (2017). The influence of optimism, social support and anxiety on aggression in a sample of dermatology patients: an analysis of cross-sectional data. *British Journal of Dermatology*, 176(5), 1187–1194.
- Conrad, R., Geiser, F., Haidl, G., Hutmacher, M., Liedtke, R., & Wermter, F. (2008). Relationship between anger and pruritus perception in patients with chronic idiopathic urticaria and psoriasis. *Journal of the European Academy of Dermatology and Venereology*, 22(9), 1062–1069.
- Dalgard, F., Gieler, U., Tomas-Aragones, L., Lien, L., Poot, F., Jemec, G., & Evers, A. (2015). The psychological burden of skin diseases: a cross-sectional multicenter study among dermatological out-patients in 13 european countries. *Journal of Investigative Dermatology*, 135(4), 984–991.
- Damasio, A. (2003). Feelings of emotion and the self. *Annals of the New York Academy of Sciences*, 1001(1), 253–261.
- Devrimci-Ozguven, H., Kundakci, N., Kumbasar, H., & Boyvat, A. (2000). The depression, anxiety, life satisfaction and affective expression levels in psoriasis patients. *Journal of the European Academy of dermatology and venereology*, 14(4), 267–271.
- Dhabhar, F. (2000). Acute stress enhances while chronic stress suppresses skin immunity: the role of stress hormones and leukocyte trafficking. *Annals of the New York Academy of Sciences*, 917(1), 876–893.
- Dieris-Hirche, J., Gieler, U., Kupfer, J., & Milch, W. (2009). Suicidal ideation, anxiety and depression in adult patients with atopic dermatitis. *Der Hautarzt; Zeitschrift fur Dermatologie, Venerologie, und verwandte Gebiete*, 60(8), 641–646.
- Duman, K., Ozdemir, Y., Yucel, E., & Akin, M. (2014). Comparison of depression, anxiety and long-term quality of health in patients with a history of either primary closure or limberg flap reconstruction for pilonidal sinus. *Clinics*, 69(6), 384–387.
- Egeberg, A., Hansen, P., Gislason, G., & Thyssen, J. (2016). Patients with rosacea have increased risk of depression and anxiety disorders: a Danish nationwide cohort study. *Dermatology*, 232(2), 208–213.
- Ehsani, A., Toosi, S., Shahshahani, M., Arbabi, M., & Noormohammadpour, P. (2009). Psycho-cutaneous disorders: an epidemiologic study. *Journal of the*

- European Academy of Dermatology and Venereology*, 23(8), 945–947.
- Eming, S., Krieg, T., & Davidson, J. (2007). Inflammation in wound repair: molecular and cellular mechanisms. *Journal of Investigative Dermatology*, 127(3), 514–525.
- Engin, B., Uguz, F., Yilmaz, E., Özdemir, M., & Mevlitoglu, I. (2008). The levels of depression, anxiety and quality of life in patients with chronic idiopathic urticaria. *Journal of the European Academy of Dermatology and Venereology*, 22(1), 36–40.
- Esposito, M., Saraceno, R., Giunta, A., Maccarone, M., & Chimenti, S. (2006). An italian study on psoriasis and depression. *Dermatology*, 212(2), 123–127.
- Femiano, F. (2007). Pemphigus vulgaris: recent advances in our understanding of its pathogenesis. *Minerva stomatologica*, 56(4), 215–223.
- Fritzsche, K., Ott, J., Zschocke, I., Scheib, P., Burger, T., & Augustin, M. (2001). Psychosomatic liaison service in dermatology. *Dermatology*, 203(1), 27–31.
- Ghajarzadeh, M., Ghiasi, M., & Kheirkhah, S. (2012). Associations between skin diseases and quality of life: a comparison of psoriasis, vitiligo, and alopecia areata. *Acta Medica Iranica*, 50(7), 511.
- Gieler, U., Niemeier, V., Kupfer, J., Brosig, B., & Schill, W. (2001). Psychosomatic dermatology in germany: a survey of 69 dermatologic clinics. *Der Hautarzt; Zeitschrift für Dermatologie, Venerologie, und verwandte Gebiete*, 52(2), 104–110.
- Gupta, M., & Gupta, A. (2001). The use of antidepressant drugs in dermatology. *Journal of the European Academy of Dermatology and Venereology*, 15(6), 512–518.
- Halioua, B., Cribier, B., Frey, M., & Tan, J. (2017). Feelings of stigmatization in patients with rosacea. *Journal of the European Academy of Dermatology and Venereology*, 31(1), 163–168.
- Khatttri, S., Bist, J., Arun, A., & Mehta, A. (2015). Clinical correlates of vitiligo with depression and anxiety: A comparative study in patients and their caregivers. *International Journal*, 3(1), 200–205.
- Kim, S., Hur, J., Jang, J., Park, H., Hong, C., Son, S., & Chang, K. (2015). Psychological Distress in Young Adult Males with Atopic Dermatitis: A Cross-Sectional Study. *Medicine*, 94(23).
- Konda, D., Chandrashekar, L., Rajappa, M., Kattimani, S., Thappa, D., & Ananthanarayanan, P. (2015). Serotonin and interleukin-6: Association with pruritus severity, sleep quality and depression severity in prurigo nodularis. *Asian journal of psychiatry*, 17, 24–28.
- Kossakowska, M., Cieścińska, C., Jaszewska, J., & Placek, W. (2010). Control of negative emotions and its implication for illness perception among psoriasis and vitiligo patients. *Journal of the European Academy of Dermatology and Venereology*, 24(4), 429–433.
- Krooks, J., Weatherall, A., & Holland, P. (2017). Review of epidemiology, clinical presentation, diagnosis, and treatment of common primary psychiatric causes of cutaneous disease. *Journal of Dermatological Treatment*, (just-accepted), 1–45.
- Lahousen, T., Kupfer, J., Gieler, U., Hofer, A., Linder, M., & Schut, C. (2016). Differences Between Psoriasis Patients and Skin-healthy Controls Concerning Appraisal of Touching, Shame and Disgust. *Acta dermato-venereologica*, 96(217), 78–82.
- Lateiner, D., & Spatharas, D. (2016). *The ancient emotion of disgust*. Oxford University Press.
- Layegh, P., Arshadi, H., Shahriari, S., Pezeshkpour, F., & Nahidi, Y. (2010). A comparative study on the prevalence of depression and suicidal ideation in dermatology patients suffering from psoriasis, Acne, Alopecia areata and Vitiligo. *Iranian Journal of Dermatology*, 13(4), 106–111.
- Lewis, M., Haviland-Jones, J., & Barrett, L. E. (2010). *Handbook of emotions*. Guilford Press.
- Linder, D., Dall’Olio, E., Gisondi, P., Berardesca, E., De Gennaro, E., Pennella, A., & Girolomoni, G. (2009). Perception of disease and doctor-patient relationship experienced by patients with psoriasis. *American journal of clinical dermatology*, 10(5), 325–330.
- Livingstone, A. G., Shepherd, L., Spears, R., & Manstead, A. S. (2016). “fury, us”: Anger as a basis for new group self-categories. *Cognition and Emotion*, 30(1), 183–192.
- Marshall, C., Taylor, R., & Bewley, A. (2016). Psychodermatology in Clinical Practice: Main Principles. *Acta dermato-venereologica*, 96(217), 30–34.
- Mattoo, S., Handa, S., Kaur, I., Gupta, N., & Malhotra, R. (2001). Psychiatric morbidity in vitiligo and psoriasis: a comparative study from india. *The Journal of dermatology*, 28(8), 424–432.
- Mento, C., Le Donne, M., Crisafulli, S., Rizzo, A., & Settineri, S. (2017). Bmi at early puerperium: body image, eating attitudes and mood states. *Journal of Obstetrics and Gynaecology*, 37(4), 428–434.
- Mento, C., Piraino, B., Rizzo, A., Vento, R., Rigoli, L., Moschella, E., & Settineri, S. (2015). Affective control and life satisfaction in thalassemics. *International Journal of Psychological Research*, 8(1), 90–97.
- Muscattello, M., Bruno, A., Scimeca, G., Pandolfo, G., & Zoccali, R. (2014). Role of negative affects in pathophysiology and clinical expression of irritable bowel syndrome. *World Journal of Gastroenterology: WJG*, 20(24), 7570.
- Parisi, R., Symmons, D., Griffiths, C., & Ashcroft, D. (2013). Global epidemiology of psoriasis: a systematic review of incidence and prevalence. *Journal*

- of *Investigative Dermatology*, 133(2), 377–385.
- Paus, R., Theoharides, T., & Arck, P. (2006). Neuroimmunoenocrine circuitry of the 'brain-skin connection'. *Trends in immunology*, 27(1), 32–39.
- Pavlovic, S., Danilchenko, M., Tobin, D., Hagen, E., Hunt, S., Klapp, B., & Peters, E. (2008). Further exploring the brain-skin connection: stress worsens dermatitis via substance P-dependent neurogenic inflammation in mice. *Journal of Investigative Dermatology*, 128(2), 434–446.
- Perusini, J., & Fanselow, M. (2015). Neurobehavioral perspectives on the distinction between fear and anxiety. *Learning & Memory*, 22(9), 417–425.
- Picardi, A., Abeni, D., Melchi, C., Puddu, P., & Pasquini, P. (2000). Psychiatric morbidity in dermatological outpatients: an issue to be recognized. *British Journal of dermatology*, 143(5), 983–991.
- Pärna, E., Aluoja, A., & Kingo, K. (2015). Quality of life and emotional state in chronic skin disease. *Acta Derm Venereol*, 95, 312–316.
- Rashid, M., Mullick, M., Jaigirdar, M., Ali, R., Nirola, D., Salam, M., & Ahsan, M. (2011). Psychiatric Morbidity in Psoriasis and Vitiligo in Two Tertiary Hospitals in Bangladesh. *Bangabandhu Sheikh Mujib Medical University Journal*, 4(2), 88–93.
- Rizzo, A., Muscatello, M., Autunno, M., Borgese, C., Pandolfo, G., Zoccali, R., & Bruno, A. (2018). Negative emotions in headache patients. *Recenti progressi in medicina*, 109(7), 393–397.
- Sampogna, F., Tabolli, S., & Abeni, D. (2012). Living with psoriasis: prevalence of shame, anger, worry, and problems in daily activities and social life. *Acta dermato-venereologica*, 92(3), 299–303.
- Sarkar, S., Sarkar, A., Saha, R., & Sarkar, T. (2014). Psoriasis and psychiatric morbidity: a profile from a tertiary care centre of Eastern India. *Journal of family medicine and primary care*, 3(1), 29.
- Sayar, K., Ugurad, I., Kural, Y., & Acar, B. (2000). The psychometric assessment of acne vulgaris patients. *Dermatology and Psychosomatics/Dermatologie und Psychosomatik*, 1(2), 62–65.
- Sellami, R., Masmoudi, J., Ouali, U., Mnif, L., Amouri, M., Turki, H., & Jaoua, A. (2014). The relationship between alopecia areata and alexithymia, anxiety and depression: a case-control study. *Indian journal of dermatology*, 59(4), 421.
- Settineri, S., Guarneri, F., Saitta, A., Mento, C., & Cannavò, S. (2013). Depression profiles in skin disorders. *Open Journal of Psychiatry*, 3(1).
- Settineri, S., & Mento, C. (2014). Questionnaire of Disgust. *Mediterranean Journal of Clinical Psychology*, 2(1).
- Settineri, S., Rizzo, A., Liotta, M., & Mento, C. (2017). Clinical Psychology of Oral Health: The Link Between Teeth and Emotions. *SAGE Open*, 7(3), 2158244017728319.
- Sharma, N., Koranne, R., & Singh, R. (2001). Psychiatric morbidity in psoriasis and vitiligo: a comparative study. *The Journal of dermatology*, 28(8), 419–423.
- Shenefelt, P. (2010). Psychological interventions in the management of common skin conditions. *Psychology research and behavior management*, 3(51).
- Tabolli, S., Mozzetta, A., Antinone, V., Alfani, S., Cianchini, G., & Abeni, D. (2008). The health impact of pemphigus vulgaris and pemphigus foliaceus assessed using the Medical Outcomes Study 36-item short form health survey questionnaire. *British Journal of Dermatology*, 158(5), 1029–1034.
- Takaki, H., & Ishii, Y. (2013). Sense of coherence, depression, and anger among adults with atopic dermatitis. *Psychology, health & medicine*, 18(6), 725–734.
- Tareen, R., & Tareen, A. (2015). Dermatology practice and psychiatric disorders. *Journal of Alternative Medicine Research*, 7(1), 47.
- Uhlenhake, E., Yentzer, B., & Feldman, S. (2010). Acne vulgaris and depression: a retrospective examination. *Journal of cosmetic dermatology*, 9(1), 59–63.
- Van Eck, N., & Waltman, L. (2014). Visualizing bibliometric networks. In Y. Ding, R. Rousseau, & D. Wolfram (Eds.), *Measuring Scholarly Impact: Methods and practice*. New York, NY: Springer International Publishing, pp. 285–320.
- Vernwal, D. (2017). A study of anxiety and depression in Vitiligo patients: New challenges to treat. *European Psychiatry*, 41, S321.
- Wakefield, J., & Demazeux, S. (2016). *Sadness or Depression? International Perspectives on the Depression Epidemic and Its Meaning*. Springer.
- Wojtyna, E., Łakuta, P., Marcinkiewicz, K., Bergler-Czop, B., & Brzezińska-Wcisło, L. (2017). Gender, body image and social support: Biopsychosocial determinants of depression among patients with psoriasis. *Acta dermato-venereologica*, 97(1), 91–97.
- Yadav, S., Narang, T., & Kumaran, M. (2013). Psychodermatology: A comprehensive review. *Indian Journal of Dermatology, Venereology, and Leprolology*, 79(2), 176.
- Yamasaki, K., & Gallo, R. (2009). The molecular pathology of rosacea. *Journal of dermatological science*, 55(2), 77–81.
- Yoder, A., Widen, S., & Russell, J. (2016). The word disgust may refer to more than one emotion. *Emotion*, 16(3), 301.