



Acta Scientiarum. Agronomy

ISSN: 1679-9275

ISSN: 1807-8621

Editora da Universidade Estadual de Maringá - EDUEM

Tavone, Luciana Alves da Silva; Nascimento, Kauyze Matos; Fachina, Yasmin Jaqueline;
Madrona, Grasielle Scaramal; Bergamasco, Rita de Cássia; Scapim, Mônica Regina da Silva

Mathematical modeling and effect of thin-layer drying and lyophilization on
antioxidant compounds from ultrasonic-assisted extracted *Muntingia calabura* peels

Acta Scientiarum. Agronomy, vol. 43, e50301, 2021, January-December

Editora da Universidade Estadual de Maringá - EDUEM

DOI: <https://doi.org/10.4025/actasciagron.v43i1.50301>

Available in: <https://www.redalyc.org/articulo.oa?id=303067924034>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

EDUEM
redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative



Mathematical modeling and effect of thin-layer drying and lyophilization on antioxidant compounds from ultrasonic-assisted extracted *Muntingia calabura* peels

Luciana Alves da Silva Tavone^{*}, Kauyse Matos Nascimento, Yasmin Jaqueline Fachina, Grasielle Scaramal Madrona, Rita de Cássia Bergamasco and Mônica Regina da Silva Scapim

Universidade Estadual de Maringá, Av. Colombo, 5790, 87020-900, Maringá, Paraná, Brazil. *Author for correspondence. E-mail: luciana.alves.engali@gmail.com

ABSTRACT. *Muntingia calabura* fruits are rich in bioactive compounds such as antioxidants, and the consumption of these compounds is associated with cancer prevention and aging. In this study, mathematical models were used to fit the experimental data of the *Muntingia calabura* peel drying kinetics, and the effective diffusion coefficient, activation energy and thermodynamic properties of the process were determined. Then, the effect of the drying temperature on the antioxidant activity and phenolic compounds of fruit peels was examined using conventional extraction and ultrasonication. Among the analyzed models, the logarithmic model was selected to represent the drying phenomenon of the calabura peel kinetics. The effective diffusion coefficient decreased by 74% as the temperature increased from 40 to 60°C, and the activation energy for liquid diffusion during drying was 23.96 kJ mol⁻¹. The enthalpy and entropy decreased with increasing temperature, while the Gibbs free energy increased by 5% for each 10°C increase in temperature. Regarding the content of phenolic compounds and the antioxidant activity of the calabura peel, it was observed that an increase in the drying temperature had a positive effect on the conservation of the bioactive compounds, making it possible to conclude that drying at 60°C and ultrasound extraction are the most suitable approach to conduct the process.

Keywords: effective diffusion coefficient; lyophilized; bioactive compounds.

Received on October 10, 2019.

Accepted on March 13, 2020.

Introduction

Muntingia calabura L. is a fruit grown in Southeast Asia and Central and South America (Buhian, Rubio, & Martin-Puzon, 2017). Known popularly as both calabura or cherry jamaica, the plant belongs to the family Elaeocarpaceae, the only species in the genus *Muntingiaceae* (Gomathi, Anusuya, & Manian, 2013).

Scientific studies on calabura have shown that stems, flowers, roots and fruits of the plant have antinociceptive properties (Amiruddin et al., 2016), anticarcinogenic activity (Nasir et al., 2017), antidiabetic activity (Sridhar, Thirupathi, Chaitanya, Kumar, & Mohan, 2011), antimicrobial activity (Patrick et al., 2017; Sufian, Ramasamy, Ahmat, Zakaria, & Yusof, 2013) and quinone reductase activity (Su et al., 2003).

Natural antioxidants, found in fruits and vegetables, cooperate in endogenous elimination of free radicals, and this effect is helpful to prevent diseases such as cancer and to aid in delaying cellular aging. In fruits, the retention rates of bioactive compounds in peels and seeds are higher than those in pulps (Morais et al., 2015)

The use of drying for fruits, seeds, peels and other foods is the oldest conservation technique that exists. In this way, the kinetics of thin-layer drying have been applied in plant material, such as pears (Silva, Figueiredo, Costa, & Guiné, 2014), apples (Vega-Galvez et al., 2012), passion fruit (Nascimento, Mulet, Ascheri, Carvalho, & Cárcel, 2016), rosemary leaves (Mghazli et al., 2017), tomatoes (Azeez, Adebisi, Oyedeji, Adetoro, & Tijani, 2019; Workneh & Oke, 2013), avocado (Avhad & Marchetti, 2016), hawthorn fruits (Aral & Bese, 2016), and grape seed (Roberts, Kidd, & Padilla-Zakour, 2008).

From the literature, there are no reports on the drying kinetics of *Muntingia calabura* or how drying affects the bioactive compounds of the fruit peel. The objective of the study was to select the best mathematical model to describe the drying curves, calculate the effective diffusion and thermodynamic properties, and evaluate the effects of forced convection drying and lyophilization on the antioxidant compounds from the peel using the ultrasound extraction method.

Material and methods

Plant material

Calabura (*Muntingia calabura*) mature fruits were harvested at the headquarters campus of the State University of Maringá (UEM) in the city of Maringá, Paraná State, Brazil (latitude: 23° 25' 31" S; Longitude: 51° 56' 19" W). The fruits were selected and sanitized with sodium hypochlorite solution at 200 ppm for 15 minutes at 25°C. The peels were divided into two parts: one part frozen at $-18 \pm 1^\circ\text{C}$ to be used in the lyophilization process and the other part used to conduct the drying kinetics analysis.

Forced convection drying process and lyophilization

The forced convection drying process of the calabura peel was determined in triplicate at temperatures of 40, 50, and $60 \pm 2^\circ\text{C}$ in a greenhouse with recirculation of air at a constant speed of 1 m s^{-1} . The reading intervals were 15 minutes in the first hour, 30 minutes during the first six hours and then every 60 minutes until the fruit peels reached a constant mass with a final moisture content lower than $0.10 \pm 0.01 \text{ g water g}^{-1}$ peel (Aral & Bese, 2016). The dry samples in the drying kinetics experiment were coded as follows: dry sample at 40°C = CD40, at 50°C = CD50, and at 60°C = CD60. The samples of frozen peel at $-18 \pm 1^\circ\text{C}$ were subjected to lyophilization with the aid of equipment (CHRIST Mark - Model Alpha 1-2 LDplus) for 24 hours.

Mathematical Models

To determine the values of the moisture content of the calabura (MR) fruit peel at any time, Equation 1 was used. Nonlinear regression analysis was performed by using the Gauss-Newton method to obtain the constants of the mathematical models selected. To evaluate each model, the coefficient of determination (R^2) (Equation 2), Chi-square test (X^2) (Equation 3), and square root mean error (RMSE) (Equation 4) were calculated.

$$MR = \frac{(M_x - M_{x0})}{(M_{xi} - M_{x0})} \quad (1)$$

$$R^2 = \sum_{i=1}^N \frac{MR_{est} - \bar{MR}_{obs}}{(MR_{obs} - \bar{MR}_{obs})^2} \quad (2)$$

$$X^2 = \sum_{i=1}^n \frac{(MR_{obs} - MR_{est})}{GLR} \quad (3)$$

$$RMSE = \sqrt{\left(\frac{\sum_{i=1}^n (MR_{obs} - MR_{est})^2}{GLR} \right)} \quad (4)$$

where: MR is the water content ratio (dimensionless value), M_x is the water content of the product represented on a dry basis (b.s), M_{x0} is the equilibrium water content of the product (b.s.), and M_{xi} is the initial water content of the product (b.s). MR_{obs} is the value observed experimentally; MR_{est} is the value calculated by the model; and GLR is the degrees of freedom of the model (observations minus the number of parameters of the model).

To adjust the experimental data collected during the drying process of the *Muntingia calabura* peel, the most commonly used mathematical models for plant products were selected (Table 1).

Determination of the effective diffusion coefficient and Thermodynamic Properties

To obtain the thickness (L) of the calabura peel, a digital micrometer was used; the average thickness was calculated based on 90 repetitions, 30 fruit peels were evaluated, and in each peel, the thickness was evaluated at three different points. The average value found for the thickness of the peel was 0.176 mm. The shape of the calabura peel was geometrically similar to a flat plate, and the geometric form of the product is considered to approximate eight terms based on Fick's second law (Equation 5) (Darvishi et al., 2014). The thickness of the shell was considered the value for the thin layer. The activation energy was calculated by the Arrhenius equation (Equation 6).

$$MR = \frac{(M_x - M_{x0})}{(M_{xi} - M_{x0})} = \frac{8}{\pi^2} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} \exp \left[-(2n+1)^2 \pi^2 Di \left(\frac{t}{4L} \right)^2 \right] \quad (5)$$

$$D_{eff} = D_0 \exp \frac{-E_a}{RT} \quad (6)$$

where: MR is the ratio of the water content (dimensionless value); "n" is the number of terms in the model; Di denotes the effective diffusion coefficient ($\text{m}^2 \text{ s}^{-1}$); "t" is the drying time (s); and L is the thickness of the

product (m). Do is the pre-exponential factor; Ea is the activation energy (kJ mol⁻¹); R is the universal gas constant (8,134 kJ kmol⁻¹ K⁻¹); and T is the absolute temperature (K).

Table 1. Mathematical models used in *Muntingia calabura* drying kinetics.

Name of model	Mathematical expression.	References
Difussion Approach	$MR = a \exp(-kt) + (1-a) \exp(-kbt)$	(Akpınar, 2006)
Two Term	$MR = a \exp(-k_1t) + b \exp(-k_2t)$	(Sharaf-Eldeen, Blaisdell, & Hamdy, 1980)
Henderson & Pabis	$MR = a \exp(-kt)$	(Henderson, 1974)
Logaritmic	$MR = a \exp(-kt) + c$	(Kingsly, Goyal, Manikantan, & Ilyas, 2007)
Midilli et.al. (2002)	$MR = a \exp(-kt^n) + bt$	(Midilli et al., 2002)
Newton	$MR = \exp(-kt)$	(Lewis, 1921)
Page	$MR = \exp(-kt^n)$	(Page, 1949)
Thompson	$MR = \exp(-a - (a^2 + 4bt)^{0.5} / (2b))$	(Thompson, Peart, & Foster, 1968)
Valcam	$MR = a + bt + ct^{1.5} + dt^2$	(Madamba, Driscoll, & Buckle, 1996)
Verma	$MR = a \exp(-k_1t) + (1-a) \exp(-k_2t)$	(Lalit, Verma, Bucklin, Endan, & Wratten, 1985)

The thermodynamic properties associated with the drying process were determined according to the method proposed by Jideani and Mpotokwana (2009). Equations 7, 8, and 9 were used to calculate the specific enthalpy, specific entropy and Gibbs free energy, respectively.

$$\Delta H = E_a - RT_a \quad (7)$$

$$\Delta S = R \left(\ln D_0 - \ln \frac{K_B}{h_p} - \ln T_a \right) \quad (8)$$

$$\Delta G = \Delta H - T_a \Delta S \quad (9)$$

where: ΔH is the specific enthalpy (J mol⁻¹); ΔS is the specific entropy (J mol⁻¹ K⁻¹); ΔG is the Gibbs free energy (J mol⁻¹); KB is the Boltzmann constant (1.38 x 10⁻²³ J K⁻¹); and hp is Planck's constant (6.626 x 10⁻³⁴ m² J⁻¹ s⁻¹).

Ultrasonic and conventional assisted extraction

The extraction of the bioactive compounds from the calabura peel was performed in an ultrasonic bath in indirect contact (Model USC - 1400) with an ultrasonic frequency of 40 kHz and ultrasonic power of 135 watts RMS and by the method of extraction by maceration (conventional). The extraction conditions (time 45 minutes, temperature 40°C and 25% aqueous ethanol) for both methodologies were identical for comparison purposes. The extracts were obtained with modifications of the method of Singh, Chidambara Murthy, and Jayaprakasha (2002); samples of peel at a ratio of 1:25 mg mL⁻¹ (peel/solvent) were placed in an Erlenmeyer flask and homogenized for 5 minutes in a homogenizer at a temperature of 25°C.

Analyses to quantify bioactive compounds

The ABTS radical assay was performed according to Rufino et al. (2007), and the results are expressed in μM Trolox g⁻¹ sample. The iron reduction power (FRAP) method was determined according to Benzie and Strain (1996), and the antioxidant potential of the extracts was determined in triplicate based on the calibration curve drawn using ferrous sulfate (FeSO₄) (2 mmol L⁻¹), expressed in μM FeSO₄ mg⁻¹ sample. The content of total phenolic compounds (TPC) was determined by the Folin-Ciocalteu (FC) spectrophotometric method, according to Singleton and Rossi (1965), expressed in mg gallic acid 100 g⁻¹ of sample, using the standard curve of gallic acid.

Statistical analysis

The experimental data were recorded as the mean \pm standard deviation, and three replicates were made for each sample. The results were analyzed by the analysis of variance (ANOVA) method, followed by the Tukey test with 95% confidence interval, using Statistic 8.0 software.

Results and discussion

Drying kinetics and thermodynamic properties

The peel of *Muntingia calabura* fruits submitted to drying at temperatures of 40 to 60°C presented an initial mean moisture content of 3.00 ± 0.495 g of water 100 g⁻¹ of dry matter (decimal b.s.). The time required to dry

the peels to equilibrium moisture content values of 0.0123 ± 0.001 , 0.0139 ± 0.001 , and 0.0135 ± 0.001 (b.s.) was 990, 690, and 510 minutes at 40, 50, and 60°C, respectively. The time needed to reach the equilibrium moisture of the sample decreased with increasing temperature; at 40°C, it took 30% longer than at 50°C, 26% longer at 50°C than at 60°C, and 48% longer at 40°C than at 60°C (48%).

This phenomenon was also observed by Kaleta and Górnicki (2010) in drying apple peels, by Oliveira, Nascimento, Borges, Ribeiro, and Ruback (2002) for pear slices, and by Zielinska and Michalska (2016) with blueberry fruits. It is also worth mentioning that during the initial 60 minutes of drying, a higher reduction in humidity was observed for the higher temperature, and at 60°C, the humidity was 22.8% lower than at the temperatures of 40°C and 14.03% lower than at 50°C.

According to Fratianni et al. (2017), the temperature used in drying is related to the internal diffusion of the sample, which is responsible for the loss of water in the process; as the internal resistance for the conduction of water in the liquid state increases, the speed of drying decreases.

To evaluate the quality of the adjustments, the highest values of R^2 and the lowest values of X^2 and RMSE were used as the criteria for comparison. Table 2 indicates all the values found for each model, and all models presented values greater than 0.99 for R^2 and less than 0.0004 for X^2 , and 0.01 for RMSE.

Regarding the comparison criteria, the models that presented the best adjustments for the three drying temperatures (40, 50, and 60°C) were the Midilli models, followed by the Valcam and Logarithmic models. In food-drying studies, similar results were found in slices of tomato, strawberry, gooseberry, hawthorn, and apple (Aral & Bese, 2016; Azeez et al., 2019; Junqueira, Luiz, & Corr, 2017; Kaleta & Górnicki, 2010; Méndez-Lagunas, Rodríguez-Ramírez, Cruz-Gracida, Sandoval-Torres, & Barriada-Bernal, 2017).

The models of Midilli and Valcam present four coefficients (a, k, n, and b; a, b, c, and d, respectively), while the logarithmic model presents only three coefficients (a, c, and k). Thus, the logarithmic model was chosen as the best model to represent the drying phenomenon of *Muntingia calabura* peel considering its simplicity in relation to the other models.

Table 2. Coefficients of determination (R^2 , decimal), Chi-square test (X^2 , decimal) and square root mean error (RMSE, decimal) for the ten mathematical models used to describe the drying process of *Muntingia calabura* peel at 40, 50, and 60°C.

Model Name	Drying temperatures								
	40°C			50°C			60°C		
	R^2	X^2	RMSE	R^2	X^2	RMSE	R^2	X^2	RMSE
Diffusion Approach	0.9989	0.00023	0.01503	0.9995	0.00012	0.01117	0.9979	0.00046	0.02155
Two Term	0.9996	0.00010	0.00989	0.9995	0.00012	0.02203	0.9985	0.00036	0.01897
Henderson e Pabis	0.9996	0.00009	0.00950	0.9995	0.00011	0.01043	0.9985	0.00031	0.01774
Logaritmic	0.9996	0.00009	0.00938	0.9997	0.00007	0.00863	0.9989	0.00025	0.01588
Midilli, Kucuk, and Yapar (2002)	0.9999	0.00003	0.00542	0.9998	0.00005	0.00696	0.9993	0.00017	0.01314
Newton	0.9989	0.00021	0.01442	0.9994	0.00012	0.01075	0.9979	0.00041	0.02024
Page	0.9996	0.00009	0.00927	0.9995	0.00011	0.01067	0.9983	0.00035	0.01879
Thompson	0.9992	0.00017	0.01292	0.9995	0.00012	0.01099	0.9980	0.00043	0.02079
Valcam	0.9998	0.00004	0.00650	0.9998	0.00004	0.00628	0.9994	0.00015	0.01205
Verma	0.9989	0.00023	0.01501	0.9995	0.00012	0.01117	0.9979	0.00046	0.02155

The increase in temperature directly affected the effective diffusion of the sample since as observed in Table 3, there was a decrease of 74% in Def compared to that at temperatures of 40 and 60°C. Similar values were found in the literature for fruits and vegetables under similar temperature conditions, with Def values of $15.30 \times 10^{-9} \text{ m}^2 \text{ s}^{-1}$ for apple (Vega-Galvez et al., 2012). In the drying procedures, the lower the activation energy is, the greater the water diffusivity is in the product.

The activation energy was calculated through the Arrhenius plot, and the average value for the three drying temperatures (40, 50, and 60°C) employed in the calabura peel was $23.96 \text{ kJ mol}^{-1}$. Galaz et al. (2017) found a value of 58.9 kJ mol^{-1} for pomegranate peel, and Nascimento et al. (2016) found a value of $43.86 \text{ kJ mol}^{-1}$ for passion fruit peel; regarding the activation energy found for the calabura peel, it is stated that the structure of the material interferes directly with the barrier that must be broken so that effective diffusion occurs in the sample.

When analyzing the values of the thermodynamic properties shown in Table 3, it can be seen that the specific enthalpy (ΔH) decreased as the temperature used in the drying kinetics experiments increased (40, 50, and 60°C), confirming that the higher the temperature used, the less energy that will be spent during the drying process, thus explaining why the time to reach RX at 60°C is half that used at 40°C. On the other hand,

the specific entropy (ΔS) and Gibbs free energy had a reverse behavior to that of the enthalpy, with values increasing as the temperatures increased. The small entropy change from -0.3399 to -0.3404 is related to the low variation of the temperatures used (10°C), and the negative values are usually related to the changes in the structure of the material.

Table 3. Thermodynamic properties of *Muntingia calabura* peel: specific enthalpy (ΔH), specific entropy (ΔS), Gibbs free energy (ΔG), and effective diffusion (Def).

Propriedades termodinâmicas				
Temperature (K)	ΔH (kJ mol ⁻¹)	ΔS (kJ mol ⁻¹ K ⁻¹)	ΔG (kJ mol ⁻¹)	Def (m ² s ⁻¹)
313.16	21.3607	-0.3399	127.4250	1.9164×10^{-11}
323.16	21.2775	-0.3402	131.1610	2.5447×10^{-11}
333.16	21.1944	-0.3404	134.8996	3.3302×10^{-11}

Effects of drying temperatures on antioxidant compounds

Quantification of the phenolic compounds and antioxidant activity are presented in Table 4. Of the four samples analyzed (CD40, CD50, CD60, and lyophilized), those with the best bioactive compound values were the CD60 and lyophilized samples, and the type of extraction used also influenced these results.

With the dry samples that underwent the forced convection drying process, the phenolic content was not significantly different between the CD40, CD50, and CD60 samples when compared to that from conventional extraction. However, in the extraction method performed with ultrasonication, the phenolic content of the CD60 sample showed a significant difference in relation to that of samples CD40 and CD50. Chen et al. (2015) explain that higher temperatures favor the extraction process since they increase both the diffusion coefficient and the solubility of the sample; however, the maximum temperature that the sample can be subjected to must be considered because very high temperatures end up denaturing the phenolic compounds.

Méndez-Lagunas et al. (2017) studied the drying effects with conventional extraction of bioactive compounds from strawberry residues and reported values ranging from 1,230.1 to 848 mg of gallic acid 100 g⁻¹ for dry samples at 50 and 60°C. Siqueira, Rosa, Fustinoni, Sant'Ana, and Arruda (2013) extracted conventional antioxidants from Brazilian Cerrado fruits and reported values of 580 ± 143 and $1,095 \pm 159$ mg of gallic acid 100 g⁻¹ in ethanoic and aqueous solvents, respectively, from *Annona crassiflora* Mart. and 651 ± 61 and $1,015 \pm 62$ mg of gallic acid 100 g⁻¹ from *Genipa americana* L. under extraction conditions of 30°C for 60 minutes. It is possible to observe that the results reported in the literature are similar to those found for calabura peel.

According to Hani, Torkamani, Zainul Abidin, Mahmood, and Juliano (2017), ultrasonic waves facilitate the penetration of solvents into the cellular matrix because they cause amplification of diffusion into the medium and allow the use of milder temperatures in the extraction process, resulting in higher yields.

Regarding the antioxidant activity, the values from the FRAP and ABTS methods showed that when comparing the type of extraction (ultrasound or conventional), there was a significant difference among all the dry samples by forced convection. When comparing the drying kinetics samples with each other, CD60 obtained the best results, and when the drying methods were compared, there was, in large part, no significant difference between the CD60 and lyophilized samples.

Chen et al. (2015) explained that the higher the temperature is, the higher the diffusion coefficient and the better the solubility of the sample. However, caution must be taken because very high temperatures will denature the phenolic compounds to be studied. In general, with increasing drying temperature, the phenolic compound and antioxidant activity indexes increased significantly, positive impacting these values.

Table 4. Effect of drying temperatures on phenolic compounds and antioxidant activity of *Muntingia calabura* peels. Mean values for TPC (mg gallic acid 100 g⁻¹), antioxidant activity ABTS (μM Trolox g⁻¹), and FRAP (μM FeSO₄ mg⁻¹).

Extraction		Sample			
		CD40	CD50	CD60	Lyophilized
Conventional	TPC	$1,102.83 \pm 87.37^{\text{bB}}$	$1,086.17 \pm 5.77^{\text{bA}}$	$1,092.83 \pm 81.29^{\text{bB}}$	$1,544.5 \pm 207.85^{\text{aA}}$
	ABTS	$22.56 \pm 2.07^{\text{bB}}$	$29.94 \pm 1.99^{\text{bB}}$	$34.9 \pm 0.94^{\text{abB}}$	$45.31 \pm 1.5^{\text{aA}}$
	FRAP	$482.32 \pm 39.81^{\text{cB}}$	$582.46 \pm 12.96^{\text{cB}}$	$782.74 \pm 45.4^{\text{bB}}$	$857.6 \pm 95.34^{\text{aA}}$
Ultrassound	TPC	$1,322.83 \pm 135.31^{\text{bCA}}$	$1,246.17 \pm 60.48^{\text{cA}}$	$1,579.5 \pm 43.3^{\text{aA}}$	$1,531.17 \pm 87.8^{\text{abA}}$
	ABTS	$48.61 \pm 1.1^{\text{bA}}$	$48.51 \pm 9.6^{\text{bA}}$	$62.71 \pm 10.22^{\text{aA}}$	$51.43 \pm 4.39^{\text{abA}}$
	FRAP	$587.81 \pm 27.53^{\text{cA}}$	$739.96 \pm 65.74^{\text{bA}}$	$888.22 \pm 33.71^{\text{aA}}$	$1,159.96 \pm 6.68^{\text{abB}}$

Expressed results with mean \pm standard deviation (triplicate). ^{abcd} The values with lowercase letters on the same line are not significantly different (Tukey test, $p < 0.05$). ^{ABCD} Values with uppercase letters in the same column and of the same chemical analysis are not significantly different (Tukey test, $p < 0.05$).

Conclusion

The logarithmic model best fit the data, and it can be used to represent the kinetics of drying for the peels. The lyophilization method can be replaced by convective drying at 60°C since the former is an expensive method. In relation to conventional extraction and ultrasonication, ultrasound-assisted techniques presented better results for phenolic compound amounts and antioxidant activity values. This study may provide useful information for future work as it provides data on the drying behavior and thermodynamic properties of calabura peel and the results of unconventional extraction methods.

References

- Akpınar, E. K. (2006). Mathematical modelling of thin layer drying process under open sun of some aromatic plants. *Journal of Food Engineering*, 77(4), 864-870. DOI: 10.1016/j.jfoodeng.2005.08.014
- Amiruddin, Z., Hijaz, M., Sani, M., Abdul, A., Lay, T., & Zaki, M. (2016). Antinociceptive effect of semi-purified petroleum ether partition of *Muntingia calabura* leaves. *Revista Brasileira de Farmacognosia*, 26(4), 408-419. DOI: 10.1016/j.bjp.2015.12.007
- Aral, S., & Bese, A. V. (2016). Convective drying of hawthorn fruit (*Crataegus* spp.): Effect of experimental parameters on drying kinetics, color, shrinkage, and rehydration capacity. *Food Chemistry*, 210, 577-584. DOI: 10.1016/j.foodchem.2016.04.128
- Avhad, M. R., & Marchetti, J. M. (2016). Mathematical modelling of the drying kinetics of Hass avocado seeds. *Industrial Crops and Products*, 91, 76-87. DOI: 10.1016/j.indcrop.2016.06.035
- Azeez, L., Adebisi, S. A., Oyediji, A. O., Adetoro, R. O., & Tijani, K. O. (2019). Bioactive compounds' contents, drying kinetics and mathematical modelling of tomato slices influenced by drying temperatures and time. *Journal of the Saudi Society of Agricultural Sciences*, 18(2), 120-126. DOI: 10.1016/j.jssas.2017.03.002
- Benzie, I. F. F., & Strain, J. J. (1996). The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": The FRAP assay. *Analytical Biochemistry*, 239(1), 70-76. DOI: 10.1006/abio.1996.0292
- Buhian, W. P. C., Rubio, R. O., & Martin-Puzon, J. J. (2017). Chromatographic fingerprinting and free-radical scavenging activity of ethanol extracts of *Muntingia calabura* L. leaves and stems. *Asian Pacific Journal of Tropical Biomedicine*, 7(2), 139-143. DOI: 10.1016/j.apjtb.2016.11.016
- Buhian, W. P. C., Rubio, R. O., Valle Jr., D. L., & Martin-Puzon, J. J. (2016). Bioactive metabolite profiles and antimicrobial activity of ethanolic extracts from *Muntingia calabura* L. leaves and stems. *Asian Pacific journal of tropical biomedicine*, 6(8), 682-685. DOI: 10.1016/j.apjtb.2016.06.006
- Chen, Q., Bi, J., Wu, X., Yi, J., Zhou, L., & Zhou, Y. (2015). Drying kinetics and quality attributes of jujube (*Zizyphus jujuba* Miller) slices dried by hot-air and short- and medium-wave infrared radiation. *LWT - Food Science and Technology*, 64(2), 759-766. DOI: 10.1016/j.lwt.2015.06.071
- Darvishi, H., Asl, A. R., Ali Asghari, Azadbakht, M., Najafi, G., & Khodaei, J. (2014). Study of the drying kinetics of pepper. *Journal of the Saudi Society of Agricultural Sciences*, 13(2), 130-138. DOI: 10.1016/J.JSSAS.2013.03.002
- Fратиanni, A., Niro, S., Messina, M. C., Cinquanta, L., Panfili, G., Albanese, D., & Di Matteo, M. (2017). Kinetics of carotenoids degradation and furosine formation in dried apricots (*Prunus armeniaca* L.). *Food Research International*, 99(Part 2), 862-867. DOI: 10.1016/j.foodres.2016.12.009
- Galaz, P., Valdenegro, M., Ramírez, C., Nuñez, H., Almonacid, S., & Simpson, R. (2017). Effect of drum drying temperature on drying kinetic and polyphenol contents in pomegranate peel. *Journal of Food Engineering*, 208, 19-27. DOI: 10.1016/j.jfoodeng.2017.04.002
- Gomathi, R., Anusuya, N., & Manian, S. (2013). A dietary antioxidant supplementation of Jamaican cherries (*Muntingia calabura* L.) attenuates inflammatory related disorders. *Food Science and Biotechnology*, 22(3), 787-794. DOI: 10.1007/s10068-013-0146-1
- Hani, N. M., Torkamani, A. E., Zainul Abidin, S., Mahmood, W. A. K., & Juliano, P. (2017). The effects of ultrasound assisted extraction on antioxidative activity of polyphenolics obtained from *Momordica charantia* fruit using response surface approach. *Food Bioscience*, 17, 7-16. DOI: 10.1016/j.fbio.2016.11.002
- Henderson, S. M. (1974). Progress in developing the thin layer drying equation. *Transactions of the ASAE*, 17(6), 1167-1168. DOI: 10.13031/2013.37052
- Jideani, V. A., & Mpotokwana, S. M. (2009). Modeling of water absorption of Botswana bambara varieties using Peleg's equation. *Journal of Food Engineering*, 92(2), 182-188. DOI: 10.1016/j.jfoodeng.2008.10.040

- Junqueira, D. J., Luiz, J., & Corr, G. (2017). Convective drying of cape gooseberry fruits : Effect of pretreatments on kinetics and quality parameters. *LWT - Food Science and Technology*, 82, 404-410. DOI: 10.1016/j.lwt.2017.04.072
- Kaleta, A., & Górnicki, K. (2010). Evaluation of drying models of apple (var. McIntosh) dried in a convective dryer. *International Journal of Food Science & Technology*, 45(5), 891-898. DOI: 10.1111/j.1365-2621.2010.02230.x
- Kingsly, R. P., Goyal, R. K., Manikantan, M. R., & Ilyas, S. M. (2007). Effects of pretreatments and drying air temperature on drying behaviour of peach slice. *International Journal of Food Science & Technology*, 42(1), 65-69. DOI: 10.1111/j.1365-2621.2006.01210.x
- Lalit, R., Verma, L. R., Bucklin, R. A., Endan, J. B., & Wratten, F. T. (1985). Effects of drying air parameters on rice drying models. *Transactions of the ASAE*, 28(1), 296-301. DOI: 10.13031/2013.32245
- Lewis, W. K. (1921). The rate of drying of solid materials. *Journal of Industrial & Engineering Chemistry*, 13(5), 427-432. DOI: 10.1021/ie50137a021
- Madamba, P. S., Driscoll, R. H., & Buckle, K. A. (1996). The thin-layer drying characteristics of garlic slices. *Journal of Food Engineering*, 29(1), 75-97. DOI: 10.1016/0260-8774(95)00062-3
- Méndez-Lagunas, L., Rodríguez-Ramírez, J., Cruz-Gracida, M., Sandoval-Torres, S., & Barriada-Bernal, G. (2017). Convective drying kinetics of strawberry (*Fragaria ananassa*): Effects on antioxidant activity, anthocyanins and total phenolic content. *Food Chemistry*, 230, 174-181. DOI: 10.1016/j.foodchem.2017.03.010
- Mghazli, S., Ouhammou, M., Hidar, N., Lahnine, L., Idlimam, A., & Mahrouz, M. (2017). Drying characteristics and kinetics solar drying of Moroccan rosemary leaves. *Renewable Energy*, 108, 303-310. DOI: 10.1016/j.renene.2017.02.022
- Midilli, A., Kucuk, H., & Yapar, Z. (2002). A new model for single-layer drying. *Drying Technology*, 20(7), 1503-1513. DOI: 10.1081/DRT-120005864
- Morais, D. R., Rotta, E. M., Sargi, S. C., Schmidt, E. M., Bonafe, E. G., Eberlin, M. N., ... Visentainer, J. V. (2015). Antioxidant activity, phenolics and UPLC-ESI(-)-MS of extracts from different tropical fruits parts and processed peels. *Food Research International*, 77(Part 3), 392-399. DOI: 10.1016/j.foodres.2015.08.036
- Nascimento, E. M. G. C., Mulet, A., Ascheri, J. L. R., Carvalho, C. W. P., & Cárcel, J. A. (2016). Effects of high-intensity ultrasound on drying kinetics and antioxidant properties of passion fruit peel. *Journal of Food Engineering*, 170, 108-118. DOI: 10.1016/j.jfoodeng.2015.09.015
- Nasir, N. L. Md., Kamsani, N. E., Mohtarrudin, N., Othman, F., Tohid, S. F. Md., & Zakaria, Z. A. (2017). Anticarcinogenic activity of *Muntingia calabura* leaves methanol extract against the azoxymethane-induced colon cancer in rats involved modulation of the colonic antioxidant system partly by flavonoids. *Pharmaceutical Biology*, 55(1), 2102-2109. DOI: 10.1080/13880209.2017.1371769
- Oliveira, L. F., Nascimento, M. R. F., Borges, S. V., Ribeiro, P. C. N., & Ruback, V. R. (2002). Aproveitamento alternativo da casca do maracujá-amarelo (*Passiflora edulis* F. Flavicarpa) para produção de doce em calda. *Food Science and Technology*, 22(3), 259-262. DOI: 10.1590/S0101-20612002000300011
- Page, G. E. (1949). *Factors influencing the maximum rates of air drying shelled corn in thin layers*. Retrieved on Aug. 8, 2019 from <https://docs.lib.purdue.edu/dissertations/AAI1300089>
- Roberts, J. S., Kidd, D. R., & Padilla-Zakour, O. (2008). Drying kinetics of grape seeds. *Journal of Food Engineering*, 89, 460-465. DOI: 10.1016/j.jfoodeng.2008.05.030
- Rufino, M. S. M., Alves, R. E., Brito, E. S., Morais, S. M., Sampaio, C. G., Pérez-Jiménez, J., & Calixto, F. D. S. (2007). *Metodologia científica: determinação da atividade antioxidante total em frutas pela captura do radical livre DPPH*. Fortaleza, CE: Embrapa. (Comunicado Técnico, 127).
- Sharaf-Eldeen, Y. I., Blaisdell, J. L., & Hamdy, M. Y. (1980). A model for ear corn drying. *Transactions of the ASAE*, 23(5), 1261-1265. DOI: 10.13031/2013.34757
- Silva, V., Figueiredo, A. R., Costa, J. J., & Guiné, R. P. F. (2014). Experimental and mathematical study of the discontinuous drying kinetics of pears. *Journal of Food Engineering*, 134, 30-36. DOI: 10.1016/j.jfoodeng.2014.02.022
- Singh, R. P., Chidambara Murthy, K. N., & Jayaprakasha, G. K. (2002). Studies on the antioxidant activity of pomegranate (*Punica granatum*) peel and seed extracts using in vitro models. *Journal of Agricultural and Food Chemistry*, 50(1), 81-86. DOI: 10.1021/jf010865b
- Singleton, V. L. & Rossi, J. A. (1965). Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture*, 16(3), 144-158.

- Siqueira, E. M. A., Rosa, F. R., Fustinoni, A. M., Sant'Ana, L. P., & Arruda, S. F. (2013). Brazilian savanna fruits contain higher bioactive compounds content and higher antioxidant activity relative to the conventional Red Delicious apple. *PLoS ONE*, 8(8), 1-7. DOI: 10.1371/journal.pone.0072826
- Sridhar, M., Thirupathi, K., Chaitanya, G., Kumar, B. R., & Mohan, G. K. (2011). Antidiabetic effect of leaves of *Muntingia calabura* L., in normal and alloxan-induced diabetic rats. *Pharmacologyonline*, 2, 626-632.
- Su, B. N., Jung Park, E., Vigo, J. S., Graham, J. G., Cabieses, F., Fong, H. H. S., ... Kinghorn, A. D. (2003). Activity-guided isolation of the chemical constituents of *Muntingia calabura* using a quinone reductase induction assay. *Phytochemistry*, 63(3), 335-341. DOI: 10.1016/S0031-9422(03)00112-2
- Sufian, A. S., Ramasamy, K., Ahmat, N., Zakaria, Z. A., & Yusof, M. I. M. (2013). Isolation and identification of antibacterial and cytotoxic compounds from the leaves of *Muntingia calabura* L. *Journal of Ethnopharmacology*, 146(1), 198-204. DOI: 10.1016/j.jep.2012.12.032
- Thompson, T. L., Peart, R. M., & Foster, G. H. (1968). Mathematical simulation of corn drying — A new model. *Transactions of the ASAE*, 11(4), 582-586. DOI: 10.13031/2013.39473
- Vega-Galvez, A., Ah-Hen, K., Chacana, M., Vergara, J., Martinez-Monzo, J., Garcia-Segovia, P., ... Di Scala, K. (2012). Effect of temperature and air velocity on drying kinetics, antioxidant capacity, total phenolic content, colour, texture and microstructure of apple (var. Granny Smith) slices. *Food Chemistry*, 132(1), 51-59. DOI: 10.1016/j.foodchem.2011.10.029
- Workneh, T. S., & Oke, M. O. (2013). Thin layer modelling of microwave-convective drying of tomato slices. *International Journal of Food Engineering*, 9(1), 75-90. DOI: 10.1515/ijfe-2012-0205
- Zielinska, M., & Michalska, A. (2016). Microwave-assisted drying of blueberry (*Vaccinium corymbosum* L.) fruits: Drying kinetics, polyphenols, anthocyanins, antioxidant capacity, colour and texture. *Food Chemistry*, 212, 671-680. DOI: 10.1016/j.foodchem.2016.06.003