



Acta Scientiarum. Agronomy

ISSN: 1807-8621

Editora da Universidade Estadual de Maringá - EDUEM

Ehrhardt-Brocardo, Natalia Carolina Moraes; Coelho,  
Cileide Maria Medeiros; Souza, Clovis Arruda  
Storage protein composition during germination and its  
association with physiological seed quality in common bean  
Acta Scientiarum. Agronomy, vol. 44, e53434, 2022  
Editora da Universidade Estadual de Maringá - EDUEM

DOI: <https://doi.org/10.4025/actasciagron.v44i1.53434>

Available in: <https://www.redalyc.org/articulo.oa?id=303071489012>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc  
Network of Scientific Journals from Latin America and the Caribbean, Spain and  
Portugal

Project academic non-profit, developed under the open access initiative



# Storage protein composition during germination and its association with physiological seed quality in common bean

Natalia Carolina Moraes Ehrhardt-Brocardo, Cileide Maria Medeiros Coelho\* and Clovis Arruda Souza

Programa de Pós-Graduação em Produção Vegetal, Departamento de Agronomia, Universidade do Estado de Santa Catarina, Av. Luiz de Camões, 2090, 88520-000, Lages, Santa Catarina, Brazil. \*Author for correspondence. E-mail: cileidecoelho@yahoo.com.br

**ABSTRACT.** Seed reserve mobilization is considered a post-germination process; however, seed storage protein mobilization occurs during germination. Thus, the knowledge of seed protein composition is important to understand various processes during germination, and it be associated with seed vigor. Therefore, this study aimed to characterize landrace genotypes of common bean (*Phaseolus vulgaris*) through fractionation of seed storage proteins during germination and to verify the association between seed protein composition and seed vigor. Genotypes of the highest (55 and 81) and lowest (23 and 50) physiological quality were selected. Protein content analysis, profiling, and characterization were performed by combining different hydration times. Mass spectrometry involving in-gel digestion and MALDI-ToF analysis was employed for the qualitative identification of proteins. Glutelin extraction detected the enzyme lipoxygenase in genotypes 55 and 81 alone. In plants, this enzyme may be involved in diverse physiological processes, including growth and development, pest resistance, senescence, and response to wounding. In the process of germination, lipoxygenase removes reactive oxygen species during reserve mobilization; therefore, lipoxygenase may be a candidate biochemical marker for high-vigor genotypes.

**Keywords:** *Phaseolus vulgaris*; lipoxygenase; glutelin; 1D SDS-PAGE; mass spectrometry.

Received on April 29, 2020.

Accepted on July 22, 2020.

## Introduction

Germination involves various physiological processes that are sequentially initiated by water uptake (imbibition; D'Hooghe et al., 2019). Seed hydration increases respiration and metabolism, thus allowing the mobilization of carbon and nitrogen reserves necessary to support germination and post-germination growth until autotrophy is reached (Anzala, Paven, Fournier, Rondeau, & Limami, 2006). Therefore, seed germination is a physiological and biochemical process that involves different cascades of signal transduction and gene expression regulation.

The elucidation and subsequent availability of bean genome have enabled the rapid advancement of seed proteomics (Rossi, Valentim-Neto, Blank, Faria, & Arisi, 2017). Many studies of bean seed proteomics have demonstrated the efficiency of different protein extraction methods to optimize the quality of separation and subsequent identification of proteins (Martínez et al., 2012).

In 1924, Osborne developed the first classification of seed storage proteins (SSPs), in which SSPs are grouped according to their solubilities in water (albumins), dilute salt solutions (globulins), aqueous alcohols (prolamins), and weakly acidic or alkaline or dilute sodium dodecyl sulfate (SDS) solution (glutelins; Chéreau et al., 2016). Biochemical reactions during seed germination are carried out by different proteins, and the analysis of seed protein profile may be a highly precise approach to clarify this physiological process (Bojórquez-Velázquez, Barrera-Pacheco, Espitia-Rangel, Herrera-Estrella, & Rosa, 2019). With the availability of massive genomic data and development of mass spectrometric techniques, proteomics has enabled the analysis of dynamic and diverse biological processes (Gu et al., 2019). Most studies involving the characterization, bioavailability, and diversity of specific proteins in bean seeds have focused on phaseolin, a type of globulin (Martínez et al., 2012; Carović-Stanko et al., 2017).

According to Rajjou et al. (2012), further research into germination and seed vigor would greatly help in unraveling the key biochemical and molecular mechanisms underlying these complex traits, which are unique to plants. Characterization of biomarkers for seed vigor to improve seeds for breeding programs as

well as other biotechnological approaches will allow for the production of seeds of the highest possible quality, with the goal of improving crop yield, particularly under environmental stresses. Thus, studies including landrace genotypes can be advantageous, because these genotypes have an extensive genetic base and can adapt to different environmental conditions relatively easily, as opposed to commercial cultivars that have a restricted genetic base (Michels, Souza, Coelho, & Zilio, 2014).

To this end, the present study aimed to characterize the SSP composition of landrace genotypes of common bean (*Phaseolus vulgaris*) during germination and examine its association with physiological seed quality.

## Material and methods

### Seed material

This study was conducted using seeds of four landrace genotypes of common bean (23, 50, 55, and 81) acquired from the Bean Active Germplasm Bank of the Santa Catarina State University. These genotypes had undergone successive selfing selection that favored greater homogeneity and led to higher stability within the population, and they were previously characterized and studied during eight consecutive seasons. Seeds were produced through a field experiment conducted in a randomized complete block design during the agricultural season of 2012-2013 by conventional farming in Campos Novos, Santa Catarina. The harvested seeds were stored in a dry chamber at  $10 \pm 2^\circ\text{C}$  under  $50 \pm 5\%$  relative humidity until further analysis. In a previous study, the physiological seed quality of these genotypes was determined using germination and vigor (accelerated aging) tests. Genotypes 55 and 81 showed the highest physiological quality (germination rate: 93 and 94%; vigor: 83 and 87%), whereas genotypes 23 and 50 showed the lowest physiological quality (germination rate: 85 and 83%; vigor: 79 and 78%) (Ehrhardt-Brocardo & Coelho, 2016).

### Cotyledons extraction and quantification

Protein content determination, profile analysis, and characterization were performed by combining different hydration times (0, 9, 12, 15, 18, and 21 hours) for each genotype to form a gene pool. Protein extraction followed the sequential extraction method modified by Landry, Delhay, and Damerval (2000) using 1 mL of extraction buffer. Globulins were extracted in 500 mM sodium chloride (NaCl; pH 2.4) at  $4^\circ\text{C}$ , and glutelins were extracted from the previous pellet in 0.5% SDS (pH 10; 25 mM sodium borate buffer) containing 0.6% 2-mercaptoethanol at room temperature.

The extraction procedure was repeated twice for each protein fraction. Three replicates of 0.1 g of fine cotyledons powder (lyophilized) were vortexed for 1 min. (AP56; Phoenix), shaken for 30 min. on an orbital shaker table (Oxy304; Oxylab), and then centrifuged at  $1,500 \times g$  for 7 min. (CE01-B1; Kacil); the supernatants were stored for further analysis. The soluble protein content was determined as described by Bradford (1976) using bovine serum albumin as a standard; absorbance was measured at 595 nm using a spectrophotometer (UV/VIS Spectro800S; Mars).

### One-dimensional gel electrophoresis of proteins

Electrophoresis was performed under denaturing conditions (SDS-PAGE) in a mini-vertical apparatus (K33-10V; Kasvi) with 12 resolving gel and 4% stacking gel. Each previously extracted protein fraction was loaded onto the gel, as described by Ma and Bliss (1978). After quantification, soluble proteins samples (15 for globulins and 9  $\mu\text{g}$  for glutelins) were dissolved in a sample buffer containing Tris-HCl (pH 6.7), glycerol, 10% SDS, 0.5% bromophenol blue and 2-mercaptoethanol at a ratio of 1:1, and distilled water. The electrophoresis system containing the samples was placed in a 10% buffer solution (25 mM Tris, 192 mM glycine, and 1% SDS; pH 8.3) at 400 mA for 3h at room temperature. After electrophoresis, the gels were washed with distilled water to completely remove the buffer and then stained for 3h with Coomassie Brilliant Blue R-250. The sample protein profiles were compared with a molecular range marker (Precision Plus Protein™, Bio-Rad, CA, USA). Reproducibility was confirmed by a minimum of two repetitions of each gel electrophoresis run under similar conditions and using samples from two independent protein extractions. Finally, digital documentation was performed in a white light transilluminator (DNR Bio-Imaging Systems Mini Bis Pro, Jerusalem - Israel) with Gel Capture.

### In-gel digestion and matrix-assisted laser desorption/ionization (MALDI) time-of-flight (ToF) MS analysis of proteins

Bands of each fraction were selected based on differential intensity between high- (55 and 81) and low-vigor (23 and 50) genotypes using GelAnalyzer 2010a (one-dimensional gel electrophoresis image analysis freeware) with an automatic band detection tool. After image analysis, each intensely Coomassie-stained lane of the gel was divided into approximately 2.5 mm slices, and the slices were then subjected to in-gel digestion with trypsin (EC 3.4.21.4).

The stained protein bands were individually excised from the gel and subjected to destaining twice with a solution containing acetonitrile (50%) and 25 mM ammonium bicarbonate ( $\text{NH}_4\text{HCO}_3$ ) for 30 min. at 25°C. The solution was replaced by pure acetonitrile and dried under vacuum (10 min.). The excised bands were reduced and alkylated before digestion. The bands were incubated with 20 ng  $\mu\text{L}^{-1}$  trypsin (Trypsin Gold V5280 Promega) and 25 mM  $\text{NH}_4\text{HCO}_3$  on ice for 30 min. and then at 37°C for 14 hours. After digestion and extraction (Shevchenko, Tomas, Havli, Olsen, & Mann, 2006), peptides were mixed with the MALDI Matrix Solution (5  $\mu\text{g}\cdot\text{mL}^{-1}$   $\alpha$ -cyano-4-hydroxycinnamic acid, 50% acetonitrile, and 0.1% trifluoroacetic acid) at a 1:1 ratio, placed on an MTP 384 target plate (Bruker Daltonics, Germany), and dried at room temperature. MALDI-ToF MS was employed for protein identification using trypsin-digested bands on an Autoflex III MALDI-TOF Mass Spectrometer (Bruker Daltonics) with the full automatic mode using flexControl 3.3 (Bruker Daltonics), and the peak lists were created using flexAnalysis 3.0 (Bruker Daltonics).

Database search was performed with Mascot (Matrix Science; <http://www.matrixscience.com>) using the *Phaseolus vulgaris* Database (downloaded from Phytozome – phaseolus\_20160609) and Viridiplantae Database (SwissProt 2016\_06). The following search parameters were used: (1) type of search: peptide mass fingerprint; fixed modifications: carbamidomethylation of cysteine; variable modifications: methionine oxidation; mass values: monoisotopic; protein mass: unrestricted; peptide mass tolerance:  $\pm 200$  ppm; max missed cleavages: 1; and (2) type of search: MS/MS ion search; fixed modifications: carbamidomethylation of cysteine; variable modifications: methionine oxidation; mass values: monoisotopic; protein mass: unrestricted; peptide mass tolerance:  $\pm 2$  Da; fragment mass tolerance:  $\pm 0.7$  Da; max missed cleavages: 1. The Universal Protein Resource Knowledgebase (UniProt Consortium, 2017) was used to assess the relevant biological functions and descriptions of proteins.

### Statistical analysis

The experiment was conducted using a randomized complete block design, and analysis of variance (*F* test) was performed. Averages were compared using Tukey test with Assistat (Silva & Azevedo, 2016). Cotyledons proteins of landrace genotypes of common bean identified by MS were converted into a binary matrix by grouping the genotypes into hierarchical clustering, generated through the Paleontological Statistics Software Package for Education and Data Analysis (PAST) software, using the Jaccard similarity coefficient (Hammer, Harper, & Ryan, 2001).

## Results and discussion

### Cotyledons protein analysis

Majority of the proteins were salt-soluble (globulins). Genotypes 23 and 55 showed higher protein content throughout germination (pool) than the other genotypes (Table 1).

**Table 1.** Soluble cotyledons protein content of landrace genotypes of common bean during germination.

| Genotype | Globulins               | Glutelins               |
|----------|-------------------------|-------------------------|
|          | mg 100 mg <sup>-1</sup> | mg 100 mg <sup>-1</sup> |
| 23       | 4.41Ab                  | 0.40b                   |
| 50       | 4.33B                   | 0.51a                   |
| 55       | 4.82A                   | 0.25c                   |
| 81       | 4.35b                   | 0.40b                   |
| CV (%)   | 3.89                    | 4.76                    |

Data are expressed as mg:100 mg<sup>-1</sup> of cotyledons dry matter. The values are means of three repetitions. Mean values followed by the same letter do not differ significantly (Tukey test, 5%).

### One-dimensional gel electrophoresis


In electrophoretic profiles of globulin, the tested landrace genotypes showed bands ranging from 15 to 150 kDa, with higher band intensity in molecular masses between 25 and 50 kDa (Figure 1). In electrophoretic profiles of glutelins, the tested landrace genotypes showed bands ranging from 25 to 150 kDa, with higher band intensity in molecular masses between 37 and 75 kDa (Figure 2). Globulins and glutelins showed differences in band intensities in protein profile analysis using one-dimensional SDS-PAGE (Figures 3 and 4); this difference may be attributed to the inhibition or overexpression of specific proteins.

### Cotyledons protein identification


Proteins from landrace genotypes of common bean were extracted and separated via one-dimensional SDS-PAGE (Figure 1 and 2), followed by identification using MALDI-ToF MS.

The conversion of proteins identified by MS into a binary matrix and subsequent hierarchical clustering allowed the grouping of genotypes according to the similarities in proteins in the pool profile formed at different hydration times (0, 9, 12, 15, 18, and 21 hours).


For the globulin group (Figure 5), high-vigor genotypes (55 and 81) showed a greater similarity in the protein pool present in three excised bands, whereas low-vigor genotypes (23 and 50) did not show any similarity in the detected proteins, specifically between bands 1 and 3. This technique of band characterization achieved a greater separation and better detection sensitivity of the expressed proteins in each genotype during germination. However, due to certain technical limitations, some proteins were detected in multiple excised bands. PHAE, V5YN37, and IOJ8I4 were detected in band 3 alone in high-vigor genotypes (55 and 81) and in band 1 alone in a low-vigor genotype (50; Figure 5). Other proteins including Q8RVH3, Q8RVY0, and V7C654 were detected in band 1 alone in a low-vigor genotype (50) and in band 3 alone in high-vigor genotypes (55 and 81).




**Figure 1.** One-dimensional SDS-PAGE of cotyledons globulins in landrace genotypes of common bean during germination. Excised bands are shown (#1-3). M: molecular range marker.



**Figure 2.** One-dimensional SDS-PAGE of cotyledons glutelins in landrace genotypes of common bean during germination. Excised bands are shown (#1-5). M: molecular range marker.



**Figure 3.** Differential intensity of cotyledons globulins between low- (23 and 50) and high-vigor (55 and 81) genotypes using GelAnalyzer. Excised bands are shown (#1-3).



**Figure 4.** Differential intensity of cotyledons glutelins between low- (23 and 50) and high-vigor (55 and 81) genotypes using GelAnalyzer. Excised bands are shown (#1-5).

Based on the clusters formed within the glutelin group, high- and low-vigor genotypes did not correspond to one another in the five excised bands (Figure 6). Although the proteins corresponding to bands 2, 4, and 5 did not allow for the separation of genotypes based on vigor, band 1 corresponded to a protein detected only in high-vigor genotypes (55 and 81), lipoxygenase (LOX; Table 2).

### Functional classification of the identified cotyledons proteins


The results generated lists of abundant proteins per extract, and numerous proteins were identified per band excised from different proteins extracts. The proteins identified by one-dimensional SDS-PAGE and MALDI-ToF MS analyses were grouped into categories according to their possible biological functions (molecular or biological) using the UniProtKB and SwissProt databases.

In electrophoretic profiles of globulins (Figure 1 and 3), genotype 81 showed different intensity of the excised bands compared with the other genotypes. Some identified proteins, such as alpha-phaseolin and phaseolin, serve the molecular function of nutrient reserve activity. However, they may show differences in


amino acid number or diverse post-translational modifications. Phaseolin (alpha-type) is 436-amino acid long, while phaseolin (beta-type) is 421-amino acid long, and both proteins are glycosylated. The excised band 2 showed a higher intensity, which was consistent with phaseolin. Phaseolin exhibited a high conversion efficiency of nutrient reserves in all genotypes, because some cotyledons proteins were not completely hydrolyzed and mobilized to the embryonic axis due to different hydration times.

In electrophoretic profiles of glutelins (Figures 2 and 4), high-vigor genotypes (55 and 81) showed different intensities of the five excised bands compared with low-vigor genotypes. For both high-vigor genotypes, band 1 corresponded to the enzyme LOX. This enzyme is 860-amino acid long, and it may be involved in diverse physiological processes in plants.

LOX was described in 1932 as a fatty acid-oxidizing enzyme; however, studies on the molecular and functional characteristics of this enzyme started only in the mid-90s when the role of jasmonic acid, which is one of the basic products of the LOX pathway, was discovered and extensively studied (Babenko, Shcherbatiuk, Skaterna, & Kosakivska, 2017). Among various described LOX-derived plant lipid mediators, jasmonic acid has received the most attention due to its important roles in defense against pathogens and insect pests via the activation of specific genes in plants (Bannenberg, Martínez, Hamberg, & Castresana, 2009). In general, lipid oxidation is a common metabolic reaction in all biological systems, and it occurs as a developmentally regulated process as well as in response to abiotic and biotic stresses (Andreou & Feussner, 2009).



**Figure 5.** Hierarchical clustering of cotyledons globulins between low- (23 and 50) and high-vigor (55 and 81) genotypes from the matrix plot generated with PAST using the Jaccard similarity coefficient. Excised bands are shown (#1-3). Arrows indicate proteins detected in multiple bands.



**Figure 6.** Hierarchical clustering of cotyledons glutelins between low- (23 and 50) and high-vigor (55 and 81) genotypes from the matrix plot generated with PAST using the Jaccard similarity coefficient. Excised bands are shown (#1-5).

Biochemically, LOXs catalyze polyunsaturated fatty acid oxygenation, the first step in the biosynthesis of a large group of biologically active fatty acid metabolites, collectively named oxylipins (Bannenberg et al., 2009). Phyto-oxylipins likely play pivotal roles in immune defense, because they act as signal molecules, protective compounds (antibacterial and wound-healing agents), or cutin constituents (Blée, 2002). According to Yang, Jiang, and Yu (2012), LOXs may be involved in stress responses through the formation of growth regulators, signal molecules, and volatile compounds in cucumber (*Cucumis sativus* L.). In soybean vegetative tissues and seeds, LOXs were considered to be storage proteins and might be associated with immune response (Porta & Rocha-Sosa, 2002). In *Phaseolus vulgaris* seedlings, LOX accumulation in response to phytohormone, cold, and desiccation stress was detected (Porta & Rocha-Sosa, 2002). During seed germination, LOXs are essential enzymes involved in the mobilization of carbon reserves (Aanangi et al., 2016). Another possibility is the effect of genetic diversity among the genotypes tested in this study. Differences in seed vigor may be observed regardless of similar germination indices (Michels et al., 2014).

In the process of germination, LOXs removed reactive oxygen species (ROS) during reserve mobilization in soybean seeds. These enzymes use molecular oxygen during their function in lipids degradation and may therefore help relieve oxidative stress; this likely explains the negligible abundance of proteins involved in redox regulation in soybean seeds (Han, Yin, He, & Yang, 2013). Similarly, Naguib (2019) reported that LOX activity was increased during germination in cotton seeds. Although the results of this study generated an extensive list of proteins identified per band excised from different proteins extracts, as described earlier,

LOX was detected in the glutelin profiles of genotypes 55 and 81 alone. As demonstrated in soybean and cotton studies, LOX may help remove ROS during reserve mobilization in common bean seed germination. Lack of LOX detection in the excised bands of low-vigor genotypes does not rule out the possibility that this enzyme plays a key role in antioxidant mechanisms of these genotypes; it may be present at a trace amount and may thus remain undetected.

**Table 2.** Cotyledons glutelins of landrace genotypes of common bean identified by mass spectrometry.

| Band | Protein                                    | UniProtKB Accession | M <sub>r</sub> (Da) | 23 | 50 | 55 | 81 |
|------|--|---------------------|---------------------|----|----|----|----|
| 1    | Aspartate aminotransferase                 | Q8HQQ0_PHAVU        | 50560               | +  | -  | -  | -  |
|      | Lipoxygenase                               | V7BX14_PHAVU        | 97527               | -  | -  | +  | +  |
| 2    | Uncharacterized protein                    | V7B9V4_PHAVU        | 94545               | -  | +  | -  | -  |
|      | Uncharacterized protein                    | V7CFI9_PHAVU        | 85178               | +  | +  | +  | +  |
|      | Formate dehydrogenase (FDH), mitochondrial | D2DWA5_PHAVU        | 41524               | -  | +  | -  | -  |
|      | Formate dehydrogenase (FDH), mitochondrial | V7C179_PHAVU        | 42578               | -  | +  | -  | -  |
| 3    | Formate dehydrogenase (FDH), mitochondrial | V7C2C9_PHAVU        | 44388               | -  | +  | -  | -  |
|      | Purple acid phosphatase                    | PPAF_PHAVU          | 52880               | -  | -  | +  | -  |
|      | Purple acid phosphatase                    | V7AEU9_PHAVU        | 45278               | -  | -  | +  | -  |
|      | Uncharacterized protein                    | V7AWN2_PHAVU        | 59095               | +  | -  | -  | +  |
|      | Uncharacterized protein                    | V7BFL3_PHAVU        | 57523               | -  | -  | -  | +  |
|      | Alpha-phaseolin                            | Q41115_PHAVU        | 48533               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5CHV7_PHAVU        | 48511               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5CHV9_PHAVU        | 48686               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5CHW3_PHAVU        | 48531               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5CHW7_PHAVU        | 48714               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5CHX2_PHAVU        | 48501               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5CN36_PHAVU        | 48501               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5CN52_PHAVU        | 48487               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5CQ77_PHAVU        | 48577               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5CQ79_PHAVU        | 48570               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5CQ83_PHAVU        | 48543               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5CQ86_PHAVU        | 48514               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5CQ89_PHAVU        | 48541               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5CQ94_PHAVU        | 48547               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5D3I5_PHAVU        | 48516               | +  | +  | +  | +  |
| 4    | Alpha-phaseolin                            | X5D3J1_PHAVU        | 48550               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5D3J3_PHAVU        | 48591               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5D3J8_PHAVU        | 48502               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5D5C2_PHAVU        | 48586               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5D5C4_PHAVU        | 48759               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5D5D0_PHAVU        | 48684               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5D5D4_PHAVU        | 48516               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5D5D7_PHAVU        | 48517               | +  | +  | +  | +  |
|      | Alpha-phaseolin                            | X5D5E0_PHAVU        | 48633               | +  | +  | +  | +  |
|      | Phaseolin                                  | A0A0A0P203_PHAVU    | 47449               | +  | +  | +  | +  |
|      | Phaseolin                                  | Q43632_PHAVU        | 47525               | +  | +  | +  | +  |
|      | Phaseolin                                  | Q43633_PHAVU        | 48448               | +  | +  | +  | +  |
|      | Phaseolin, alpha-type                      | PHSA_PHAVU          | 49241               | +  | +  | +  | +  |
|      | Phaseolin, beta-type                       | PHSB_PHAVU          | 47536               | +  | +  | +  | +  |
|      | Uncharacterized protein                    | V7BFL2_PHAVU        | 41272               | +  | +  | +  | +  |
|      | Uncharacterized protein                    | V7BFL4_PHAVU        | 41678               | +  | +  | +  | +  |
| 5    | Uncharacterized protein                    | V7BN55_PHAVU        | 33677               | +  | +  | +  | +  |

'+' indicates the presence of a protein and '-' the absence of a protein in a band.

Future studies are warranted to investigate the association between antioxidant enzyme activity and seed vigor in landrace genotypes of common bean under various stresses. In future studies, the biochemical and physiological responses of these genotypes to oxidative stress through the antioxidant defense system as well as the mechanisms underlying ROS removal and oxidative damage relief in the presence of specific biotic or abiotic stresses will be evaluated.

## Conclusion

The quantification of soluble protein fractions during bean seed germination associated with mass spectrometry (MS) technique proved to be an efficient tool in these genotypes characterization. Although

globulins are the most abundant fraction, the high-vigor genotypes differed from those of low-vigor in not abundant fractions, the glutelins. In the tested landrace genotypes of common bean, LOX was present at a concentration within the detection limit only in high-vigor genotypes. Thus, further studies with additional common bean genotypes with contrasting vigor characteristics are recommended. Assessment of additional genotypes will allow for the verification any association between seed storage protein composition and seed vigor, and also, if LOX may be a possible biochemical marker candidate protein for high-vigor genotypes of common bean.

### Acknowledgements

The authors acknowledge the *Coordenação de Aperfeiçoamento de Pessoal de Nível Superior* (Capes), *Conselho Nacional de Desenvolvimento Científico e Tecnológico* (CNPq), *Fundação de Amparo à Pesquisa e Inovação de Santa Catarina* (Fapesc) and *Programa de Apoio à Pesquisa* (PAP) of the *Universidade do Estado de Santa Catarina* (Udesc) together Fapesc (PAP/Udesc/Fapesc) for providing financial support for developing this study and corresponding author (Coelho, C. M. M.) thanks CNPq for productivity scholarship. The authors also acknowledge *Centro de Biologia Molecular Estrutural* of *Universidade Federal de Santa Catarina* (Cebime-UFSC) for Mass Spectrometry facilities.

### References

- Aanangi, R., Kotapati, K. V., Palaka, B. K., Kedam, T., Kanika, N. D., & Ampasala, D. R. (2016). Purification and characterization of lipoxygenase from mung bean (*Vigna radiata* L.) germinating seedlings. *3 Biotech*, 6(1), 113. DOI: <https://doi.org/10.1007/s13205-016-0427-5>
- Andreou, A., & Feussner, I. (2009). Lipoxygenases – structure and reaction mechanism. *Phytochemistry*, 70(13-14), 1504-1510. DOI: <https://doi.org/10.1016/j.phytochem.2009.05.008>
- Anzala, F., Paven, M.-C. M.-L., Fournier, S., Rondeau, D., & Limami, A. M. (2006). Physiological and molecular aspects of aspartate-derived amino acid metabolism during germination and post-germination growth in two maize genotypes differing in germination efficiency. *Journal of Experimental Botany*, 57(3), 645-653. DOI: <https://doi.org/10.1093/jxb/erj054>
- Babenko, L. M., Shcherbatiuk, M. M., Skaterna, T. D., & Kosakivska, I. V. (2017). Lipoxygenases and their metabolites in formation of plant stress tolerance. *Ukrainian Biochemical Journal*, 89(1), 5-21. DOI: <https://doi.org/10.15407/ubj89.01.005>
- Bannenberg, G., Martínez, M., Hamberg, M., & Castresana, C. (2009). Diversity of the enzymatic activity in the lipoxygenase gene family of *Arabidopsis thaliana*. *Lipids*, 44(2), 85-95. DOI: <https://doi.org/10.1007/s11745-008-3245-7>
- Blée, E. (2002). Impact of phyto-oxylipins in plant defense. *Trends in Plant Science*, 7(7), 315-322. DOI: [https://doi.org/10.1016/s1360-1385\(02\)02290-2](https://doi.org/10.1016/s1360-1385(02)02290-2)
- Bojórquez-Velázquez, E., Barrera-Pacheco, A., Espitia-Rangel, E., Herrera-Estrella, A., & Rosa, A. P. B. (2019). Protein analysis reveals differential accumulation of late embryogenesis abundant and storage proteins in seeds of wild and cultivated amaranth species. *BMC Plant Biology*, 19(59), 1-17. DOI: <https://doi.org/10.1186/s12870-019-1656-7>
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72(1-2), 248-254. DOI: [https://doi.org/10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3)
- Carović-Stanko, K., Liber, Z., Vidak, M., Barešić, A., Grdiša, M., Lazarević, B., & Šatović, Z. (2017). Genetic diversity of Croatian common bean landraces. *Frontiers in Plant Science*, 8(604), 1-8. DOI: <https://doi.org/10.3389/fpls.2017.00604>
- Chéreau, D., Videcoq, P., Ruffieux, C., Pichon, L., Motte, J.-C., Belaid, S., ... Lopez, M. (2016). Combination of existing and alternative technologies to promote oilseeds and pulses proteins in food applications. *Oilseeds & Fats Crops and Lipids*, 23(4), D406. DOI: <https://doi.org/10.1051/ocl/2016020>
- D'Hooghe, P., Picot, D., Brunel-Muguet, S., Kopriva, S., Avice, J.-C., & Trouverie, J. (2019). Germinative and post-germinative behaviours of *Brassica napus* seeds are impacted by the severity of s limitation applied to the parent plants. *Plants*, 8(1), 12. DOI: <https://doi.org/10.3390/plants8010012>

- Ehrhardt-Brocardo, N. C. M., & Coelho, C. M. M. (2016). Hydration patterns and physiologic quality of common bean seeds. *Semina: Ciências Agrárias*, 37(4), 1791-1800. DOI: <http://dx.doi.org/10.5433/1679-0359.2016v37n4p1791>
- Gu, J., Hou, D., Li, Y., Chao, H., Zhang, K., Wang, H., ... Li, M. (2019). Integration of proteomic and genomic approaches to dissect seed germination vigor in *Brassica napus* seeds differing in oil content. *BMC Plant Biology*, 19(1), 21. DOI: <https://doi.org/10.1186/s12870-018-1624-7>
- Hammer, Ø., Harper, D. A. T., & Ryan, P. D. (2001). Past: paleontological statistics software package for education and data analysis. *Palaeontologia Electronica*, 4(1), 1-9.
- Han, C., Yin, X., He, D., & Yang, P. (2013). Analysis of proteome profile in germinating soybean seed, and its comparison with rice showing the styles of reserves mobilization in different crops. *PLoS ONE*, 8(2), e56947. DOI: <https://doi.org/10.1371/journal.pone.0056947>
- Landry, J., Delhay, S., & Damerval, C. (2000). Improved method for isolating and quantitating  $\alpha$ -amino nitrogen as nonprotein, true protein, salt-soluble proteins, zeins, and true glutelins in maize endosperm. *Cereal Chemistry*, 77(5), 620-626. DOI: <https://doi.org/10.1094/CCHEM.2000.77.5.620>
- Ma, Y., & Bliss, F. A. (1978). Seed proteins of common bean. *Crop Science*, 18(3), 431-437. DOI: <https://doi.org/10.2135/cropsci1978.0011183X001800030018x>
- Martínez, M. F., López-Pedrouso, M., Alonso, J., Ferradás, M. S., Pedreira, A. M. R., Álvarez, G., & Zapata, C. (2012). In-depth characterization of the phaseolin protein diversity of common bean (*Phaseolus vulgaris* L.) based on two-dimensional electrophoresis and mass spectrometry. *Food Technology and Biotechnology*, 50(3), 315-325. DOI: <https://doi.org/10.2661/97442>
- Michels, A. F., Souza, C. A., Coelho, C. M. M., & Zilio, M. (2014). Qualidade fisiológica de sementes de feijão crioulo produzidas no oeste e planalto catarinense. *Revista Ciência Agronômica*, 45(3), 620-632. DOI: <https://doi.org/10.1590/S1806-66902014000300025>
- Naguib, D. M. (2019). Metabolic profiling during germination of hydro primed cotton seeds. *Biocatalysis and Agricultural Biotechnology*, 17(2), 422-426. DOI: <https://doi.org/10.1016/j.bcab.2018.12.025>
- Porta, H., & Rocha-Sosa, M. (2002). Plant lipoxygenases. Physiological and molecular features. *Plant Physiology*, 130(1), 15-21. DOI: <https://doi.org/10.1104/pp.010787>
- Rajjou, L., Duval, M., Gallardo, K., Catusse, J., Bally, J., Job, C., & Job, D. (2012). Seed germination and vigor. *Annual Review of Plant Biology*, 63, 507-533. DOI: <https://doi.org/10.1146/annurev-arplant-042811-105550>
- Rossi, G. B., Valentim-Neto, P. A., Blank, M., Faria, J. C., & Arisi, A. C. M. (2017). Comparison of grain proteome profiles of four Brazilian common bean (*Phaseolus vulgaris* L.) cultivars. *Journal of Agricultural and Food Chemistry*, 65(34), 7588-7597. DOI: <https://doi.org/10.1021/acs.jafc.7b03220>
- Shevchenko, A., Tomas, H., Havli, J., Olsen, J. V., & Mann, M. (2006). In-gel digestion for mass spectrometric characterization of proteins and proteomes. *Nature Protocols*, 1(6), 2856-2860. DOI: <https://doi.org/10.1038/nprot.2006.468>
- Silva, F. A. S., & Azevedo, C. A. V. (2016). The Assistat software version 7.7 and its use in the analysis of experimental data. *African Journal of Agricultural Research*, 11(39), 3733-3740. DOI: <https://doi.org/10.5897/AJAR2016.11522>
- UniProt Consortium. (2017). UniProt: the universal protein knowledgebase. *Nucleic Acids Research*, 45(D1), D158-D169. DOI: <https://doi.org/10.1093/nar/gkw1099>
- Yang, X.-Y., Jiang, W.-J., & Yu, H.-J. (2012). The expression profiling of the lipoxygenase (LOX) family genes during fruit development, abiotic stress and hormonal treatments in cucumber (*Cucumis sativus* L.). *International Journal of Molecular Sciences*, 13(2), 2481-2500. DOI: <https://doi.org/10.3390/ijms13022481>