



Revista Colombiana de Química  
ISSN: 0120-2804  
ISSN: 2357-3791  
rcolquim\_fcbog@unal.edu.co  
Universidad Nacional de Colombia  
Colombia

## Synthesis, characterization, X-ray crystal structure and DFT calculations of 4-([2,2':6',2''-terpyridin]- 4'-yl)phenol

**Castro Agudelo, Brian; Ochoa-Puentes, Cristian; Rodriguez-Córdoba, William; Reiber, Andreas; Sierr, César A.**

Synthesis, characterization, X-ray crystal structure and DFT calculations of 4-([2,2':6',2''-terpyridin]- 4'-yl)phenol  
Revista Colombiana de Química, vol. 47, no. 1, 2018

Universidad Nacional de Colombia, Colombia

**Available in:** <https://www.redalyc.org/articulo.oa?id=309055254010>

**DOI:** <https://doi.org/10.15446/rev.colomb.quim.v47n1.66281>

Los contenidos se distribuyen con la licencia Creative Commons 4.0, que permite a los usuarios copiar, distribuir, adaptar y comunicar públicamente, con cualquier propósito siempre y cuando se reconozcan los créditos de la obra.



This work is licensed under Creative Commons Attribution 4.0 International.

# Synthesis, characterization, X-ray crystal structure and DFT calculations of 4-([2,2':6',2''-terpyridin]-4'-yl)phenol

Síntesis, caracterización, estructura cristalina y cálculos DFT para el 4-([2,2':6',2''-terpiridin]-4'-il)fenol

Síntese, caracterização, estrutura de cristal e cálculos DFT para o 4-([2,2':6',2''-terpiridina]-4'-il)fenol

Brian Castro Agudelo

Universidad Nacional de Colombia, Colombia

DOI: [https://doi.org/10.15446/](https://doi.org/10.15446/rev.colomb.quim.v47n1.66281)

rev.colomb.quim.v47n1.66281

Redalyc: [https://www.redalyc.org/articulo.oa?](https://www.redalyc.org/articulo.oa?id=309055254010)

id=309055254010

Cristian Ochoa-Puentes

Universidad Nacional de Colombia, Colombia

William Rodriguez-Córdoba

Universidad Nacional de Colombia, Colombia

Andreas Reiber

Universidad de los Andes, Colombia

César A. Sierr

Universidad Nacional de Colombia, Colombia

casierraa@unal.edu.co

Received: 17 June 2017

Accepted: 25 September 2017

## ABSTRACT:

The synthesis of new terpyridine (Tpy) derivatives has been subject of extensive research due to its potential as functional materials for solar energy conversion, among other applications. In this contribution, the 4-([2,2':6',2''-terpyridin]-4'-yl)phenol (TpyOH) has been synthesized, characterized and studied through several methods, including X-ray crystallography and computational approaches. Single crystal X-ray structure analysis shows that TpyOH is essentially planar, with dihedral angles of about 5.03° between the central pyridinyl and the phenolic ring, and also 6.05 and 12.2° in the terpyridine moiety. In the crystal, molecules are linked by intermolecular hydrogen bonds and through  $\pi$ - $\pi$  stacking interactions. Using a time-dependent density functional theory approach and taking into account bulk solvent effects, the absorption and fluorescence spectra of TpyOH were investigated and compared. The TD-DFT  $S_0 \rightarrow S_n$  and  $S_1 \rightarrow S_0$  transition energies are in good agreement with experimental results. The frontier molecular orbitals analysis showed that the low-energy absorption band has an intraligand charge transfer character (ICT), while the high-energy band is a common feature of p-p\* transitions of the Tpy moiety. The  $S_1 \rightarrow S_0$  emission transition also has an ICT character, with a 90% contribution from the HOMO  $\rightarrow$  LUMO transitions.

**KEYWORDS:** Terpyridine, Kröhnke reaction, Crystal structure, TD-DFT.

## RESUMEN:

La síntesis de derivados terpiridinicos (Tpy) se ha investigado ampliamente debido a su potencial para la conversión de energía solar. En este artículo se sintetizó y caracterizó el 4-([2,2':6',2''-terpiridin]-4'-il)fenol (TpyOH), a través de varias metodologías como la cristalografía de rayos X y herramientas computacionales. El análisis de rayos X de monocristal mostró que el TpyOH es plano, con ángulos diedros de 5,03° entre el piridinilo central y el anillo fenólico, con presencia de ángulos de 6,05 y 12,2° en la porción terpiridinica. En el cristal, las moléculas están unidas por enlaces de hidrógeno intermoleculares y mediante interacciones de apilamiento  $\pi$ - $\pi$ . Utilizando cálculos DFT dependientes del tiempo (TD-DFT) y teniendo en cuenta el efecto de los disolventes, se investigaron y compararon los espectros de absorción y fluorescencia de TpyOH. Las energías de transición TD-DFT de  $S_0 \rightarrow S_n$  y  $S_1 \rightarrow S_0$  concuerdan con los resultados experimentales. El análisis de orbitales moleculares de frontera mostró que la banda de absorción de baja energía corresponde a transferencia de carga intraligando (ICT); mientras que la banda de alta energía es común en las transiciones p-p\* del resto Tpy. La emisión debido a la transición  $S_1 \rightarrow S_0$  corresponde a ICT, con una contribución del 90% proveniente de transiciones HOMO  $\rightarrow$  LUMO.

**PALABRAS CLAVE:** Terpiridina, reacción de Kröhnke, estructura cristalina, TD-DFT.

**RESUMO:**

A síntese de derivados de terpiridina (Tpy) tem sido estudada devido ao seu potencial para a conversão de energia solar. Nesta contribuição, o 4-([2,2':6',2''-terpiridina]-4'-il) fenol (TpyOH) foi sintetizado, caracterizado e estudado por vários métodos. A análise de estrutura de raios X de cristal único mostra que o TpyOH é plano, com ângulos diedros de 5,03 ° entre o piridinilo central e o anel fenólico, e também 6,05 e 12,2 ° na porção de terpiridina. No cristal, as moléculas são ligadas por ligações intermoleculares de hidrogênio e através de interações de empilhamento  $\pi$ - $\pi$ . Usando uma abordagem da teoria funcional da densidade dependente do tempo e levando em consideração os efeitos do solvente em massa, foram investigados e comparados os espectros de absorção e fluorescência do TpyOH. As energias de transição TD-DFT  $S_0 \rightarrow S_n$  e  $S_1 \rightarrow S_0$  estão de acordo com os resultados experimentais. A análise de orbitais moleculares de fronteira mostrou que a banda de absorção de baixa energia possui um caráter de transferência de carga intraligando (TIC), enquanto a banda de alta energia é uma característica comum das transições  $p \rightarrow p^*$  da fração Tpy. A transição de emissão  $S_1 \rightarrow S_0$  também tem um caráter TIC, com uma contribuição de 90% das transições HOMO<sup>\*</sup>LUMO.

**PALAVRAS-CHAVE:** Terpiridine, reação de Kröhnke, estrutura de cristal, TD-DFT.

**INTRODUCTION**

Currently, terpyridines (Tpys) have a very important presence in the scientific literature due principally to their optoelectronic properties and coordination capability (1, 2, 3). The different characteristics of Tpys are provided by the three neighbouring pyridine rings and their spatial disposition, which form a very particular and changeable electronic space. As a result, for example, Tpys are capable of forming coordination compounds with various transition metals. This makes Tpys very promissory moieties in the development of new materials with high technological potentials like photovoltaics (4), LEDs (5), solar cells (6, 7, 8, 9) and chemo-sensor probes (10).

This interest in Tpys has represented a considerable effort on the synthetic field. Many methodologies have been widely studied for the synthesis of Tpys, like the Potts, Jameson, Sasaki, Stille cross-coupling, and the Kröhnke reaction (11), the latter being the most useful methodology to synthesize 4-aryl substituted Tpys using very simple reagents under feasible conditions. The Kröhnke reaction involves the use of the appropriate aryl-aldehydes and 2-acetylpyridine to give the corresponding 4-substituted Tpy in moderate yields. This is a very important advantage over the aforementioned methodologies which require very expensive reagents non-commercially available. However, the Kröhnke methodology has a very important limitation regarding the synthesis of Tpys, since this synthetic route can only be used for the synthesis of 4-aryl substituted Tpys.

The spatial disposition of the nitrogen atoms over the pyridine rings of Tpys is very important for the physical and chemical properties. As a result of the free rotation presented by the sigma bonds that connect the aromatic rings, Tpys have different structural conformations, which can present high or low  $\pi$ -electronic delocalization, properties that can affect the optoelectronic properties drastically. Interestingly, the conformation in Tpys can be altered or modified by specific guest, such as metal ions, through coordination bonds. Therefore, correlation of spatial conformations, crystallographic structure, and optoelectronic properties of Tpys are very important for technological applications such as chemical sensors. Herein, the synthesis, characterization, and the crystal structure of 4-([2,2':6',2''-terpyridin]-4'-yl)phenol (TpyOH) are reported, a versatile tridentate ligand which may be used as building-block for the synthesis of multicomponent systems with interesting photochemical and photophysical properties.

## MATERIALS AND METHODS

### Synthesis

4-Hydroxybenzaldehyde and 2-acetylpyridine were purchased from Sigma-Aldrich (St. Louis - MO, USA) and were used without further purification. Sodium hydroxide was obtained from Alfa-Aesar (Haverhill - MA, USA). Following the Kröhnke reaction (Figure 1), TpyOH was obtained in moderate yield.




Figure 1. Synthetic route for 4-aryl substituted TpyOH

FIGURE 1  
Synthetic route for 4-aryl substituted TpyOH

### General procedure

#### *Synthesis of 4-([2,2':6',2''-terpyridin]-4'-yl)phenol (TpyOH)*

2.41 mL of 2-acetylpyridine (20 mmol) were added dropwise to a solution of 1.3 g of 4-hydroxybenzaldehyde (10 mmol) and 2.1 g of sodium hydroxide (50 mmol) in 25 mL of ethanol at 0 °C. Two hours later having reached the room temperature, 10 mL of concentrated ammonium hydroxide were added and the solution was refluxed overnight. After this, filtration at room temperature gave a yellowish solid that was recrystallized from ethanol-water (1:1) to yield 2.03 g (57%) of light yellow crystals. M.p. > 300 °C (lit. (11) 302-306°C). FT-IR (cm<sup>-1</sup>): 3441, 3055-3039, 1597, 1527 and 1095. <sup>1</sup>H-RMN (400 MHz, CDCl<sub>3</sub>) δ (ppm): 8,79 (d, 2H), 8,75 (d, 2H), 8,70 (s, 2H), 7,72–7,78 (m, 2H), 7,38–7,41 (m, 6H). <sup>13</sup>C-RMN (100 MHz, CDCl<sub>3</sub>) δ (ppm): 116.52, 117.53, 121.39, 125.02, 128.31, 128.58, 139.12, 149.71, 155.59, 160.13 MW: Calculated: 325.1201 Found: 325.171(MALDI-TOF).

### Steady-state spectroscopy

All measurements were performed at room temperature (20 ± 1°C) in HPLC quality ethanol. The stationary absorption spectra were measured using a Shimadzu UV-2401 PC double channel spectrophotometer (Kyoto, Japan) in a 1 cm quartz cell. The steady-state fluorescence emission and excitation spectra were recorded using a PTI Quantamaster™ 4000 spectrofluorometer (New Jersey, USA).

### Computational details

All calculations were performed using the Gaussian09 quantum chemical program package (12). The molecular structures of TpyOH in the electronic ground state were fully optimized with density functional

theory (DFT) method using the parameter-free PBE0 hybrid functional and the 6-31+G(d,p) basis sets. To include the solvent effects, we used the Polarized Continuum Model (PCM) with the integral equation formalism (IEFPCM) developed by Tomasi et al (13, 14).


To simulate the absorption and emission spectra, vertical excitation energies in the singlet and triplet manifold were calculated using the PCM/TD-PBE0/6-31+G(d,p) level of theory. This approach has proven to predict accurate descriptions of excited states for several organic (15) and metal-organic complexes (16, 17, 18). Geometry optimizations for the S0 and S1 states were performed without symmetry constraints and were confirmed to be stationary points through vibrational frequency analysis.

## RESULTS AND DISCUSSION

### Synthesis and crystal structure

TpyOH was synthesized under mild conditions following the Kröhnke reaction with a yield similar to the previously reported (11). After purification, TpyOH was crystallized in an orthorhombic crystal system with  $P2_12_12_1$  symmetry. Full information of the structure was deposited at the CSD Database under the deposition number 1454127 (Table 1). The crystal-packing diagram (Figure 2) shows that the asymmetric unit of TpyOH contains one crystallographically independent molecule as part of four molecules per unit cell with similar bond distance and dihedral angles. The geometry of TpyOH corresponds to a nearly coplanar structure, where the central pyridinyl ring (Py2) makes a dihedral angle of about  $5.03^\circ$  with the phenolic ring (PhOH), and  $6.05$  and  $12.2^\circ$  with the pyridinyl substituent rings Py1 and Py3, respectively.

The differences in the dihedral angles observed on the terpyridyl moiety can be attributed to the intermolecular hydrogen bond  $O25-H26 \cdots N18$  formed between two adjacent molecules. As shown in Figure 1, the molecules are column-like stapled along the a-axis of the unit cell with a packing dominated mainly by intermolecular hydrogen bonds ( $1.85(3)$  Å), between the pyridinyl nitrogen of Py3 and the -OH over the phenol-group at the neighbour molecule and by  $\pi$ - $\pi$  interactions ( $3.52(3)$  Å) between molecules of the same staple. Although the crystal structure of the target compound was published earlier (19) and the crystallographic data are quite similar, the cell volume here reported is a little bit smaller (about  $68$  Å<sup>3</sup>), showing the possibility of polymorph formation for this compound (20).



**Figure 2.** X-Ray molecular structure of TpyOH. 1, 2, and 3 are the dihedral angles defined by atoms  $C_{12}-C_{11}-C_{19}-C_{24}$ ,  $C_1-C_2-C_7-C_{12}$ , and  $C_{18}-C_{13}-C_9-C_{10}$ , respectively (left). And Cell packing of the four molecules in the unit cell along the a-axes, showing hydrogen bonding as red lines. The molecules are column-like stapled along the a-axis (right).

**FIGURE 2**

X-Ray molecular structure of TpyOH. f1, f2, and f3 are the dihedral angles defined by atoms  $C_{12}-C_{11}-C_{19}-C_{24}$ ,  $C_1-C_2-C_7-C_{12}$ , and  $C_{18}-C_{13}-C_9-C_{10}$ , respectively (left). And Cell packing of the four molecules in the unit cell along the a-axes, showing hydrogen bonding as red lines. The molecules are column-like stapled along the a-axis (right)

TABLE 1

Crystal data and structure refinement for TpyOH

**Table 1.** Crystal data and structure refinement for TpyOH.

Crystal data	
Chemical formula	C <sub>21</sub> H <sub>13</sub> N <sub>3</sub> O
<i>M<sub>r</sub></i>	325.36
Crystal system, space group	Orthorhombic, <i>P</i> 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>
Temperature (K)	100
<i>a</i> , <i>b</i> , <i>c</i> (Å)	4.5788 (6), 16.204 (2), 20.798 (3)
<i>V</i> (Å <sup>3</sup> )	1543.1 (3)
<i>Z</i>	4
Radiation type	Mo <i>K</i> α
<i>μ</i> (mm <sup>-1</sup> )	0.09
Crystal size (mm)	0.18 × 0.16 × 0.13
Data collection	
Diffractometer	SuperNova, Dual, Cu at zero, Atlas diffractometer
Absorption correction	Multi-scan <i>CrysAlis PRO</i> , Agilent Technologies, Version 1.171.36.28 (release 01-02-2013 <i>CrysAlis171.NET</i> ) (compiled Feb 1 2013, 16:14:44) Empirical absorption correction using spherical harmonics, implemented in <i>SCALE3 ABSPACK</i> scaling algorithm
<i>T<sub>min</sub></i> , <i>T<sub>max</sub></i>	0.984, 0.989
No. of measured, independent and observed [ <i>I</i> > 2σ( <i>I</i> )] reflections	9237, 2803, 2597
<i>R<sub>int</sub></i>	0.036
(sin θ/λ) <sub>max</sub> (Å <sup>-1</sup> )	0.600
Refinement	
<i>R</i> [ <i>F</i> <sup>2</sup> > 2σ( <i>F</i> <sup>2</sup> )], <i>wR</i> [ <i>F</i> <sup>2</sup> ], <i>S</i>	0.040, 0.100, 1.10
No. of reflections	2803
No. of parameters	230
No. of restraints	0
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
Δρ <sub>max</sub> , Δρ <sub>min</sub> (e Å <sup>-3</sup> )	0.13, -0.18


## Molecular Hirshfeld surface

The Hirshfeld surface is a useful tool for describing the surface characteristics of molecules. The molecular Hirshfeld surface of TpyOH was generated by the CrystalExplorer 17.5 program (21), using a standard high surface resolution with the 3D dnorm surface (Figure 3) mapped over a fixed color scale of -0.66 (red) to 1.2 Å (blue). The Hirshfeld surface is shown to be transparent to permit visualization of the molecular moiety, and the 3D dnorm surface was used to identify close intermolecular interactions. The values of dnorm are negative or positive either the intermolecular contacts are shorter or respectively longer than the Van der Waals radii. The dnorm values were pictured onto the Hirshfeld surface using a red-blue-white color scheme, where red regions represent a closer contact with a negative dnorm value, a blue region represents longer



contacts with a positive  $d_{\text{norm}}$  value, and a white region represents exactly the distances by van der Waals radii with a  $d_{\text{norm}}$  value of zero.

Intermolecular contact atom- atom	Surface Area %
C...C	6,9
C...H	30,6
C...N	5,3



**Figure 3.** 3D  $d_{\text{norm}}$  Hirshfeld-surface of the TPyOH molecule. Red spots indicate short distances between neighbour molecules. The two shortest distances between molecules are formed by hydrogen-bonds between OH-groups and nitrogen-atoms of neighbour molecules

### FIGURE 3

3D  $d_{\text{norm}}$  Hirshfeld-surface of the TPyOH molecule. Red spots indicate short distances between neighbour molecules. The two shortest distances between molecules are formed by hydrogen-bonds between OH-groups and nitrogen-atoms of neighbour molecules

The 2D fingerprint plots (Figure 4) can be interpreted as particular atom pair contacts between neighbour molecules. It enables the separation of different interaction types that overlap in the full fingerprint (Figure 4a). The 2D fingerprints are symmetrical along the diagonal. The de- and di-axes represent the distances from the molecule surface to the nearest nucleus external to the surface (de) and the distance from the surface to the nearest nucleus internal to the surface (di). These values provide an immediate picture of the nature of intermolecular contacts in the crystal.




FIGURE 4

2D-Fingerprints including reciprocal contacts between atoms of neighbour molecules: a) overall; b) N-H and the surface area of a TpyOH molecule which has the nearest contact to surrounding molecules discriminated by intermolecular atom-atom distances (inserted table)



Intermolecular contact atom- atom	Surface Area %
C...C	6,9
C...H	30,6
C...N	5,3
H...H	42
N...H	7,8
N...N	0,9
O...H	6,5
O...O	0
C...O	0
N...O	0

FIGURE 4 (CONT.)

2D-Fingerprints including reciprocal contacts between atoms of neighbour molecules: a) overall; b) N-H and the surface area of a TpyOH molecule which has the nearest contact to surrounding molecules discriminated by intermolecular atom-atom distances (inserted table)

In Figure 4b the spikes represent the nearest contact between N- and H-atoms ( $d_i=1.008\text{Å}$ ,  $d_e=0.730\text{Å}$ ) but in this type of interaction only 7.8% of the Hirshfeld surface is involved, the main contact area between the surrounding molecules is formed between H--H (42% of the surface) and between C--H atoms (30.6% of the surface).

## Steady-state spectroscopy

The absorption and emission spectra of TpyOH in ethanol solution are shown in Figure 5. As can be observed, the absorption spectrum shows an intense band centred at  $\lambda = 286\text{ nm}$  and a weak long-wavelength absorption in the 330–410 nm range. The low-energy band is assigned to the intraligand charge transfer transition (ICT), while the high-energy band is a common feature of p-p\* transitions of the Tpy moiety (22, 23, 24, 25). From the comparison between the UV-visible spectrum of TpyOH and the corresponding terpyridine systems bearing electron-donating or electron-withdrawing groups, is clear that the intensity

of the ICT band is mainly governed by the electron donating strength of the substituent. Specifically, terpyridines carrying an  $\text{-NR}_2$  electron donor substituent show a strong absorption band around 360-399 nm.

On the contrary, for the case of  $\text{NO}_2$  substitution, the spectrum shows only one absorption band at higher energy, as expected for a  $\text{p-p}^*$  transition centred on the ligand (22, 23, 24, 25). Furthermore, the emission spectrum of TpyOH shows a broad emission band centred at  $\lambda = 384$  nm, with a very large Stokes shift of almost 100 nm, indicating that TpyOH undergoes with difficulty self-absorption processes. This suggests that TpyOH could be a good candidate for applications such as chemo-sensors. This emission band corresponds to energy relaxation from a nearly coplanar  $\text{S}_1$  state formed after the photo-absorption process (as it is shown in the following Figure 5).




FIGURE 5

Absorption (blue line) and emission (red line) spectra of TpyOH in ethanol. The fluorescence spectrum was taken with an excitation wavelength of 285 nm. The vertical sticks correspond to the vertical absorption and emission transitions calculated at the PCM/TD-PBE0/6-31+G(d,p) level of theory

## DFT ground state properties and conformational analysis

DFT geometry optimization of TpyOH in the ground state was performed taking into account three different isomers, generally named as transoid, cisoid, and a mixture between them (from now on labelled as TpyOH 1, TpyOH 2, and TpyOH 3), which differ in the spatial orientation of the terpyridyl nitrogen atoms. Optimized ground state structures of these Tpy isomers in ethanol are depicted in Figure 6, while selected structural parameters are included in Table 2. For the TpyOH 1 isomer, the phenol substituent forms a dihedral angle of approximately  $34^\circ$  relative to central terpyridine ring, while the three terpyridine aromatic rings are nearly planar, with  $\text{f}_2$  and  $\text{f}_3$  of about  $-1.8^\circ$  and  $-1.9^\circ$ , respectively. For the TpyOH 2 isomer, the substituted phenol ring and the terminal terpyridine rings are twisted for about  $38^\circ$  and  $-139^\circ$ , respectively, while for the TpyOH 3 corresponding dihedral angles (see Figure 2 for angles designations) are  $\text{f}_1 = 36^\circ$ ,  $\text{f}_2 = -144^\circ$  and  $\text{f}_3 = -3.7^\circ$ .




FIGURE 6

Optimized molecular structures of TpyOH 1, TpyOH 2 and TpyOH 3 conformers in ethanol, calculated at the PCM/TD-PBE0/6-31+G(d,p) level of theory

The results showed that TpyOH 1 is the most stable conformer, with the TpyOH 2 and TpyOH 3 conformers lying higher in energy at 13.0 and 5.51 kcal/mol in the gas phase, and 6.52 and 3.12 kcal/mol in ethanol. Comparing Figures 2 and 6, it is clear that TpyOH 1 corresponds to the crystal structure, for this reason, in the following section, we will only limit the analysis to this conformer.

TABLE 2

Selected structural parameters of TpyOH by X-ray, and theoretical calculations for isomers 1, 2, and 3

**Table 2.** Selected structural parameters of TpyOH by X-ray, and theoretical calculations for isomers 1, 2, and 3

		(PCM)/TD-PBE0/6-31+G(d,p)					
	Experimental	TpyOH 1		TpyOH 2		TpyOH 3	
Bond lengths (Å)	X-ray	Gas Phase	EtOH	Gas Phase	EtOH	Gas Phase	EtOH
N <sub>1</sub> -C <sub>2</sub>	1.34498	1.33981	1.34142	1.33750	1.34050	1.33843	1.39998
C <sub>2</sub> -C <sub>7</sub>	1.48502	1.48801	1.48932	1.49029	1.49077	1.48994	1.49068
C <sub>7</sub> -N <sub>8</sub>	1.33936	1.33667	1.33744	1.33426	1.33705	1.33383	1.33640
N <sub>8</sub> -C <sub>9</sub>	1.34203	1.33668	1.33746	1.33419	1.33704	1.33564	1.33805
C <sub>9</sub> -C <sub>13</sub>	1.49163	1.48800	1.48933	1.49030	1.49076	1.48745	1.48942
C <sub>13</sub> -N <sub>18</sub>	1.33723	1.33988	1.34144	1.33750	1.34052	1.33965	1.34145
C <sub>11</sub> -C <sub>19</sub>	1.48729	1.47714	1.47726	1.47739	1.47687	1.47743	1.47710
C <sub>22</sub> -O <sub>25</sub>	1.36008	1.35871	1.35736	1.35827	1.35694	1.35858	1.35719
O <sub>25</sub> -H <sub>26</sub>	0.90706	0.96283	0.96483	0.96284	0.96486	0.96285	0.96486
Bond angles (°)							
N <sub>1</sub> -C <sub>2</sub> -C <sub>7</sub>	116.672	116.882	116.952	116.937	117.037	116.913	120.706
C <sub>2</sub> -C <sub>7</sub> -N <sub>8</sub>	115.859	116.949	116.791	116.797	116.928	116.794	116.971
N <sub>8</sub> -C <sub>9</sub> -C <sub>13</sub>	114.966	116.944	116.792	116.796	116.928	116.455	116.697
C <sub>9</sub> -C <sub>13</sub> -N <sub>18</sub>	118.047	116.901	116.955	116.947	117.037	117.039	116.976
C <sub>12</sub> -C <sub>11</sub> -C <sub>19</sub>	121.225	121.361	121.382	121.545	121.557	121.389	121.486
C <sub>23</sub> -C <sub>22</sub> -O <sub>25</sub>	118.259	117.496	117.505	117.466	117.505	117.483	117.516
C <sub>22</sub> -C <sub>25</sub> -H <sub>26</sub>	110.134	109.800	110.189	109.887	110.205	109.828	110.190
Dihedral angles (°)							
C <sub>12</sub> -C <sub>11</sub> -C <sub>19</sub> -C <sub>24</sub>	5.034	33.979	33.285	38.194	34.229	35.958	33.667
N <sub>1</sub> -C <sub>2</sub> -C <sub>7</sub> -C <sub>12</sub>	6.054	-1.780	-1.752	-138.787	-142.652	-143.706	-144.672
N <sub>18</sub> -C <sub>13</sub> -C <sub>9</sub> -C <sub>10</sub>	12.203	-1.943	-2.027	-138.469	-142.489	-3.737	-3.590
C <sub>21</sub> -C <sub>22</sub> -O <sub>25</sub> -H <sub>26</sub>	-7.140	-0.374	-0.314	-0.388	-0.282	-0.716	-0.510

## S<sub>0</sub> --> S<sub>n</sub> excitation energies calculations. Absorption spectrum

TD-DFT computational studies were performed to elucidate the electronic structures of the S<sub>0</sub> state of TpyOH 1. The calculated excitation energies with their contribution, oscillator strengths, and coefficients are reviewed in Table 3. The best agreement with the experimental data was obtained using the PCM/TD-PBE0/6-31+G(d,p) level of theory. Specifically, the calculated S<sub>0</sub> --> S<sub>1</sub> ICT transition is 3.98 eV (311 nm), while the S<sub>0</sub> --> S<sub>2</sub> transition is 4.15 eV (299 nm). This table also shows that the lowest-energy singlet --> singlet transition at 311 nm has its main contribution from the highest occupied molecular orbital HOMO (DFT orbital 85) to the lowest unoccupied molecular orbital LUMO (DFT orbital 86) and is assigned as an ICT transition.

TABLE 3

Excitation energies (eV/nm), largest excitation coefficients, electronic transition configurations and oscillator strengths for TpyOH 1 in ethanol computed at the PCM/TD-PBE0/6-31+G(d,p) level of theory.

**Table 3.** Excitation energies (eV/nm), largest excitation coefficients, electronic transition configurations and oscillator strengths for TpyOH 1 in ethanol computed at the PCM/TD-PBE0/6-31+G(d,p) level of theory.

Electronic transitions	Composition	Coefficient	E(eV)/(nm)	Oscillator
$S_0 \rightarrow S_1$	(85) H $\rightarrow$ (86) L (67%)	0.57893	3.9832/311.27	0.0317
	(84) H-1 $\rightarrow$ (87) L+1 (28%)	0.37622		
$S_0 \rightarrow S_2$	(85) H $\rightarrow$ (87) L+1 (96%), (85) H $\rightarrow$ (86) L (2%)	0.69121	4.1458/299.06	0.4651
$S_0 \rightarrow S_3$	(84) H-1 $\rightarrow$ (87) L+1 (66%), (85) H $\rightarrow$ (86) L (29%)	0.57394 0.37919	4.2764/289.93	0.3646
$S_0 \rightarrow S_4$	(83) H-2 $\rightarrow$ (87) L+1 (74%)	0.60879	4.3729/283.53	0.0099
	(80) H-5 $\rightarrow$ (86) L (11%)	0.23649		
$S_0 \rightarrow S_5$	(83) H-2 $\rightarrow$ (86) L (41%)	0.45506	4.3898/282.44	0.1411
	(84) H-1 $\rightarrow$ (86) L (43%)	0.46312		

The frontier molecular orbitals and its composition are shown in Figure 7 and Table 4, respectively. From Table 4, on the one hand, it can be observed that the HOMO is composed of 34.0% p (Tpy) and 66.0% p (PhOH), whereas the LUMO orbital is localized on the terpyridine moiety with a 99.0%  $\pi^*$  (Tpy). On the other hand, the next high-energy transitions at 299 nm ( $f = 0.4651$ , Table 3) and 289 nm ( $f = 0.3646$ , Table 3) dominate the energy absorption band. Table 3 also shows that the S2 state has its main contribution from the H  $\rightarrow$  L+1 transitions (96%), while the S3 state has contributions from the H  $\rightarrow$  L ICT transitions (29%) mixed with the H-1  $\rightarrow$  L+1 (66%) transition. Both states have a p-p\* character due to the p-p\* transitions of the terpyridine and the phenol ring.




FIGURE 7

Frontier molecular orbitals of TpyOH 1 in ethanol calculated at the PCM/TD-PBE0/6-31+G(d,p) level of theory (isosurface value: 0.02)

TABLE 4

Frontier molecular orbital compositions (%) in the ground state for TpyOH 1 at the PCM/TD-PBE0/6-31+G(d,p) level of theory

**Table 4.** Frontier molecular orbital compositions (%) in the ground state for TpyOH 1 at the PCM/TD-PBE0/6-31+G(d,p) level of theory

Orbital	Energy (eV)	Contribution (%)		Main bond type
		Tpy	PhOH	
(88) LUMO+2	-0.9421779214848	96.0	4.00	$\pi^*$ (Tpy) + $\pi^*$ (PhOH)
(87) LUMO+1	-1.76680044386092	88.0	12.0	$\pi^*$ (Tpy) + $\pi^*$ (PhOH)
(86) LUMO	-1.76837348931256	99.0	1.00	$\pi^*$ (Tpy)
(85) HOMO	-6.5419507154456	34.0	66.0	$\pi$ (Tpy) + $\pi$ (PhOH)
(84) HOMO-1	-6.7969372037268	100	0.00	$\pi$ (Tpy)

## Optimization of the S1 excited state. Fluorescence emission spectrum

Optimization of the S1 Franck–Condon state of TpyOH 1 in both gas phase and ethanol solution resulted in a more coplanar structure (Figure 8). Specifically, the dihedral angles f1, f2 and f3 decrease 15.1, 1.19, and 1.41° in the gas phase, and 12.5, 1.68, and 1.21° in ethanol, respectively. According to these results, we can conclude that the resonance interaction is more efficient in the first excited state than in the ground state. Table 5 lists the calculated emission energy, electronic transition configuration, and frontier molecular orbital compositions (%) in the S1 excited state of TpyOH 1. As can be observed, the calculated S1 -->



S<sub>0</sub> transition at 355 nm in ethanol has its main contribution from the HOMO and LUMO orbitals (90% contribution) and oscillator strength of 0.2383. This result shows the ICT character of the emission band.




FIGURE 8

S<sub>1</sub> optimized molecular structures of TpyOH 1 in a) gas phase and b) ethanol solution calculated at the (PCM)/TD-PBE0/6-31+G(d,p) level of theory

In the S<sub>1</sub> excited state, the HOMO is composed of 35.0% p (Tpy) and 65.0% p (PhOH), whereas the LUMO orbital is mainly localized on the terpyridine moiety with a 94.0%  $\pi^*$  (Tpy). The frontier molecular orbitals in the S<sub>1</sub> excited state of TPyOH 1 in ethanol are depicted in Figure 9.

TABLE 5

Excitation energy (eV/nm), largest excitation coefficients, electronic transition configuration, oscillator strength *f*, and frontier molecular orbital compositions (%) in the S<sub>1</sub> excited state for TpyOH 1 calculated at the PCM/TD-PBE0/6-31+G(d,p) level of theory

**Table 5.** Excitation energy (eV/nm), largest excitation coefficients, electronic transition configuration, oscillator strength *f*, and frontier molecular orbital compositions (%) in the S<sub>1</sub> excited state for TpyOH 1 calculated at the PCM/TD-PBE0/6-31+G(d,p) level of theory.

Electronic transitions	Composition	Coefficient	E(eV)/(nm)	Oscillator
S <sub>1</sub> → S <sub>0</sub>	(85) H → (86) L (90%)	0.67100	3.49/355	0.2383
	(84) H-1 → (86) L (5%)	-0.16353		
Molecular orbital compositions				
		Contribution (%)		
Orbital	Energy (eV)	Tpy	PhOH	Main bond type
(86) LUMO	-2.05	94.0	6.00	π* (Tpy) + π* (PhOH)
(85) HOMO	-6.26	35.0	65.0	π (Tpy) + π (PhOH)
(84) HOMO-1	-6.70	100	0.00	π (Tpy)




FIGURE 9

Frontier molecular orbitals in the S1 excited state of TPyOH 1 in ethanol calculated at the PCM/TD-PBE0/6-31+G(d,p) level of theory (isosurface value: 0.02).

## CONCLUSIONS

In this contribution, the 4-([2,2':6',2''-terpyridin]-4'-yl)phenol compound was synthesized in a moderate yield following the Kröhnke conditions and its structure was unambiguously determined by X-ray single-crystal diffraction,  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR, and UV-vis techniques. Crystallographic and DFT results showed that the transoid conformation (TPyOH 1 isomer) corresponds to the most stable and predominant structure for the model compound. Comparison between the geometrical parameters of the optimized ground state and the crystal data shows that the differences are related to the intermolecular hydrogen bond formed between two adjacent molecules in the crystal packing.

The  $S_0 \rightarrow S_n$  and  $S_1 \rightarrow S_0$  transition energies calculated with the PCM/TD-PBE0/631+G(d,p) level of theory are in good agreement with experimental results. TD-PBE0 results show that both the absorption band observed in the 330–410 nm range and the emission band have their main contribution from the HOMO to the LUMO orbitals and are assigned as intraligand charge transfer (ICT) transitions.

## ACKNOWLEDGMENTS

For financial support, we are thankful to Colciencias (Grants 534-2011 and 111865843739) and to Direccion Nacional de Investigaciones-UNAL (Grant 18296).

## REFERENCES

1. Sun, Y.; El Ojaimi, M.; Hammitt, R.; Thummel, R. P.; Turro, C. Effect of Ligands with Extended  $\pi$ -System on the Photophysical Properties of Ru(II) Complexes. *Phys. Chem. B.* 2010, 114, 14664-14670. DOI: <http://dx.doi.org/10.1021/jp102613n>.
2. Schubert, U. S.; Winter, A.; Newkome, G. R. Terpyridine-based Materials: For Catalytic, Optoelectronic and Life Science Applications. Wiley-VCH. Weinheim, Alemania, 2011; pp 260-280. DOI: <http://dx.doi.org/10.1002/9783527639625>
3. Schubert, U. S.; Hofmeier, H.; Newkome, G. R. Modern Terpyridine Chemistry. Wiley-VCH. Weinheim, Alemania, 2006; pp 26-41. DOI: <http://dx.doi.org/10.1002/3527608486>.
4. Duprez, V.; Biancardo, M.; Spanggaard, H.; Synthesis of Conjugated Polymers Containing Terpyridine–Ruthenium Complexes: Photovoltaic Applications. *Macromolecules.* 2005, 38, 10436-10448. DOI: <http://dx.doi.org/10.1021/ma051274f>.

5. Holder, E.; Marin, V.; Meier, M.; Schubert, U.; A Novel Light-Emitting Mixed-Ligand Iridium(III) Complex with a Terpyridine-Poly(ethylene glycol) Macroligand. *Macromolecular Rapid Comm.* 2004, 25, 1491-1496. DOI: <http://dx.doi.org/10.1002/marc.200400206>.
6. Knoll, J. D.; Albani, B. A.; Turro, C.; New Ru(II) Complexes for Dual Photoreactivity: Ligand Exchange and  $1O_2$  Generation. *Acc. Chem. Res.* 2015, 48, 2280-2287. DOI: <http://dx.doi.org/10.1021/acs.accounts.5b00227>.
7. Yang, S. H.; Wu, K. L.; Chi, Y.; Cheng, Y. M.; Chou, P. T.; Tris(thiocyanate) ruthenium(II) sensitizers with functionalized dicarboxyterpyridine for dye-sensitized solar cells. *Angew Chem. Int. Ed. Engl.* 2011, 50, 8270-8274. DOI: <http://dx.doi.org/10.1002/anie.201103515>.
8. Li, F.; Jiang, Y.; Zhang, B.; Huang, F.; Gao, Y.; Sun, L.; Towards A Solar Fuel Device: Light-Driven Water Oxidation Catalyzed by a Supramolecular Assembly. *Angew Chem. Int. Ed.* 2012, 51, 2417-2420. DOI: <http://dx.doi.org/10.1002/anie.201108051>.
9. Lin, R. G.; A new zinc(II) coordination compound of meta-aminobenzoate and 4, 4' -bipyridine with broad-band photoluminescence emission. *Inorg. Chim. Acta.* 2015, 432, 46-49. DOI: <http://dx.doi.org/10.1016/j.ica.2015.03.03>.
10. Thornleya, P.; Starkeya, J.; Zibasereshta, R.; Polsona, M.; Wikairaa, M. and Hartshorna, R. 4' -(o-Toluy1)-2, 2' :6' , 2' ' -terpyridine: synthesis, bromination, complexation, and X-ray crystallographic characterization. *J. Coord. Chem.* 2011, 64, 145-158. DOI: <http://dx.doi.org/10.1080/00958972.2010.546397>.
11. Anthonysamy, A.; Balasubramanian, S.; Shanmugaiah, V.; Mathivanan, N. Synthesis, characterization and electrochemistry of 4' -functionalized 2, 2' :6' , 2'' -terpyridine ruthenium(II) complexes and their biological activity. *Dalton Transactions*, 2008, 16, 2136-2143. DOI: <http://dx.doi.org/10.1039/b716011a>.
12. Frisch, M.; Trucks, G.; Schlegel, G.B.; Scuseria, G.; Robb, M.; Cheeseman, J.; Scalmani, et al. Gaussian 09 package: Gaussian 09, Revision E.01. Gaussian, Inc., Wallingford CT. 2009.
13. Cancès, E.; Mennucci, B.; Tomasi, J.; A new integral equation formalism for the polarizable continuum model: Theoretical background and applications to isotropic and anisotropic dielectrics. *J. Chem. Phys.* 1997, 107, 3032-3041. DOI: <http://dx.doi.org/10.1063/1.474659>.
14. Mennucci, B.; Cancès, E.; Tomasi, J. Evaluation of Solvent Effects in Isotropic and Anisotropic Dielectrics and in Ionic Solutions with a Unified Integral Equation Method: Theoretical Bases, Computational Implementation, and Numerical Applications. *J. Phys. Chem. B.* 1997, 101, 10506-10517. DOI: <http://dx.doi.org/10.1021/jp971959k>.
15. Plaza-Medina, E.; Rodríguez-Córdoba, W.; Morales-Cueto, R.; Peon, J. Primary Photochemistry of Nitrated Aromatic Compounds: Excited-State Dynamics and NO $\cdot$  Dissociation from 9-Nitroanthracene. *J. Phys. Chem. A.* 2011, 115, 577-585. DOI: <http://dx.doi.org/10.1021/jp109041y>.
16. Velmurugana, G.; Venuvanalingam, P. Luminescent Re(I) terpyridine complexes for OLEDs: what does the DFT/TD-DFT probe reveal? *Dalton Trans.* 2015, 44, 8529-8542. DOI: <http://dx.doi.org/10.1039/c4dt02917h>.
17. Zhang, T.; Jia, J.; Ren, Y.; Wu, H. Ligand Effects on Structures and Spectroscopic Properties of Pyridine-2-aldoxime Complexes of Re(CO) $_3$ +: DFT/TDDFT Theoretical Studies. *J. Phys. Chem. A.* 2011, 115, 3174-3181. DOI: <http://dx.doi.org/10.1021/jp200872b>.
18. Martinez Saavedra, H.; Ragone, F.; Ruiz, G.; Gara, P.; Wolcan, E. Solvent Dependent Switching of 3MLLCT and 1IL Luminescent States in [ClRe(CO) $_3$ (bathocuproinedisulfonate)] $_2$  - Spectroscopic and Computational Study. *J. Phys. Chem. A.* 2014, 118, 9661-9674. DOI: <http://dx.doi.org/10.1021/jp506890r>.
19. Darabi, F.; Hadadzadeh, H.; Simpson, J.; Shahpiric, A. A water-soluble Pd(II) complex with a terpyridine ligand: experimental and molecular modeling studies of the interaction with DNA and BSA; and in vitro cytotoxicity investigations against five human cancer cell lines. *New J. Chem.* 2016, 40, 9081-9097. DOI: <http://dx.doi.org/10.1039/C6NJ01880G>.
20. Gavezzotti, A.; Fillippini, G. Polymorphic forms of organic crystals at room conditions: Thermodynamic and structural implications. *J. Am. Chem. Soc.* 1995, 117, 12299-12305. DOI: <http://dx.doi.org/10.1021/ja00154a032>.

21. M. J. Turner, J. J. McKinnon, S. K. Wolff, D. J. Grimwood, P. R. Spackman, D. Jayatilaka and M. A. Spackman, CrystalExplorer17. University of Western Australia. <http://hirshfeldsurface.net>. 2017.
22. Tessore, F.; Roberto, D.; Ugo, R.; Pizzotti, M. Terpyridine Zn(II), Ru(III), and Ir(III) Complexes: The Relevant Role of the Nature of the Metal Ion and of the Ancillary Ligands on the Second-Order Nonlinear Response of Terpyridines Carrying Electron Donor or Electron Acceptor Groups. *Inorg. Chem.* 2005, 44, 8967-8978. DOI: <http://dx.doi.org/10.1021/ic050975q>.
23. Kanis, D.; Lacroix, P.; Ratner, M.; Marks, T. Electronic Structure and Quadratic Hyperpolarizabilities in Organotransition-Metal Chromophores Having Weakly Coupled  $\pi$ -Networks. Unusual Mechanisms for Second-Order Response. *J. Am. Chem. Soc.* 1994, 116, 10089-10102. DOI: <http://dx.doi.org/10.1021/ja00101a030>.
24. Roberto, D.; Ugo, R.; Bruni, S.; Cariati, E.; Cariati, F.; Fantucci, P.; Invernizzi, I. Quadratic Hyperpolarizability Enhancement of para-Substituted Pyridines upon Coordination to Organometallic Moieties: The Ambivalent Donor or Acceptor Role of the Metal. *Organometallics* 2000, 19, 1775-1788. DOI: <http://dx.doi.org/10.1021/om990865p>.
25. Bruni, S.; Cariati, E.; Cariati, F.; Porta, F. A.; Quici, S.; Roberto, D. Determination of the quadratic hyperpolarizability of trans-4-[4-(dimethylamino)styryl]pyridine and 5-dimethylamino-1,10-phenanthroline from solvatochromism of absorption and fluorescence spectra: a comparison with the electric-field-induced second-harmonic generation technique. *Spectrochim. Acta Part A.* 2001, 57, 1417-1426. DOI: [http://dx.doi.org/10.1016/S1386-1425\(00\)00483-2](http://dx.doi.org/10.1016/S1386-1425(00)00483-2).

#### ALTERNATIVE LINK

<https://revistas.unal.edu.co/index.php/rcolquim/article/view/66281> (html)