

Nutrición Hospitalaria

Nutrición Hospitalaria

ISSN: 0212-1611

ISSN: 1699-5198

Grupo Arán

López Gómez, Juan José; Izaola Jáuregui, Olatz; Torres Torres, Beatriz; Gómez Hoyos, Emilia; Castro Lozano, María Ángeles; Ortolá Buigues, Ana; Martín Ferrero, Miguel Ángel; de Luis Román, Daniel A.

Influence of a meal-replacement diet on quality of life in women with obesity and knee osteoarthritis before orthopedic surgery

Nutrición Hospitalaria, vol. 35, no. 1, 2018, January-February, pp. 71-77
Grupo Arán

DOI: <https://doi.org/10.20960/nh.1148>

Available in: <https://www.redalyc.org/articulo.oa?id=309258226013>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative



Trabajo Original

Obesidad y síndrome metabólico

Influence of a meal-replacement diet on quality of life in women with obesity and knee osteoarthritis before orthopedic surgery

Influencia de una dieta modificada en la calidad de vida en mujeres con obesidad y artrosis de rodilla antes de la cirugía ortopédica

Juan José López Gómez^{1,3}, Olatz Izaola Jáuregui^{1,3}, Beatriz Torres Torres^{1,3}, Emilia Gómez Hoyos^{1,3}, María Ángeles Castro Lozano^{1,3}, Ana Ortolá Buigues^{1,3}, Miguel Ángel Martín Ferrero² and Daniel A. de Luis Román^{1,3}

Departments of ¹Endocrinology and Nutrition and ²Traumatology. Hospital Clínico Universitario de Valladolid. Valladolid, Spain. ³Investigation Institute of Endocrinology and Nutrition. University of Valladolid. Valladolid, Spain

Abstract

Background: Knee osteoarthritis is a disease with a high prevalence in our environment, especially in women. Weight loss can improve the quality of life of these patients before surgery.

Objectives: To evaluate the effect of a meal-replacement diet on weight loss, body composition, and the improvement of the quality of life in obese women with knee osteoarthritis pending surgery.

Methods: One branch intervention study was performed over three months on 81 women with a body mass index greater than 30 kg/m² with knee osteoarthritis before surgery. Patients received a hyperproteic meal-replacement diet with two bottles of an oral nutrition supplement in lunch and dinner (1,035 kcal). Anthropometric parameters, and body composition were measured. The quality of life was assessed by WOMAC and SF-36 test.

Results: The mean age of the patients was 62.23 (8.50) years. The percentage of weight loss was 8.23% (4.04). An improvement in the SF-36 total score was observed (basal: 49.35 [20.41], three months: 58.71 [17.07], $p < 0.01$). There was an improvement in WOMAC test (basal: 49.24% [25.53], three months: 40.59% [21.76], $p < 0.01$). It was observed that a 10% improvement in the SF-36 test was independently related to weight loss (OR: 1.2 [1.03-1.36], $p < 0.02$) adjusted by age and changes in body composition.

Conclusions: In women with osteoarthritis of the knee treated with a meal-replacement diet, there is a significant decrease in weight and fat mass with a relative increase of the latter. There is an improvement in the quality of life according to SF-36 and WOMAC. There is an independent relationship between weight loss and SF-36 improvement.

Key words:

Knee osteoarthritis.
Quality of life. Diet.
Weight loss.

Resumen

Introducción: la artrosis de rodilla es una enfermedad con alta prevalencia en nuestro medio, especialmente en mujeres. La pérdida de peso puede mejorar la calidad de vida de estas pacientes antes de la cirugía.

Objetivos: evaluar el efecto de una dieta de sustitución de comidas en la pérdida de peso, la composición corporal y la mejora de la calidad de vida en mujeres obesas con artrosis de rodilla pendientes de cirugía ortopédica.

Métodos: se realizó un estudio de intervención de una rama durante tres meses en 81 mujeres con un índice de masa corporal superior a 30 kg/m² con artrosis de rodilla antes de la cirugía. Las pacientes recibieron una dieta de sustitución de comidas hiperproteica con dos botellas de un suplemento nutricional oral en almuerzo y cena (1.035 kcal). Se midieron los parámetros antropométricos y la composición corporal. La calidad de vida fue evaluada por WOMAC y la prueba SF-36.

Resultados: la edad media de las pacientes fue de 62,23 (8,50) años. El porcentaje de pérdida de peso fue de 8,23% (4,04). Se observó una mejora en la puntuación total del SF-36 (basal: 49,35% [20,41], tres meses: 58,71% [17,07], $p < 0,01$). Hubo una mejora en la prueba WOMAC (basal: 49,24% [25,53], tres meses: 40,59% [21,76], $p < 0,01$). Se observó que una mejora del 10% en la prueba SF-36 que se relacionó independientemente con la pérdida de peso (OR: 1,2 [1,03-1,36], $p < 0,02$) ajustada por edad y cambios en la composición corporal.

Conclusiones: en mujeres con osteoartritis de la rodilla tratada con una dieta de sustitución de comidas hay una disminución significativa en el peso y la masa grasa, con un aumento relativo de esta última. Se observa una mejora en la calidad de vida según SF-36 y WOMAC y existe una relación independiente entre la pérdida de peso y la mejora del SF-36.

Palabras clave:

Artrosis de rodilla.
Calidad de vida.
Dietas. Pérdida de peso.

Received: 25/03/2017 • Accepted: 14/05/2017

López Gómez JJ, Izaola Jáuregui O, Torres Torres B, Gómez Hoyos E, Castro Lozano MÁ, Ortolá Buigues A, Martín Ferrero MÁ, de Luis Román DA. Influence of a meal-replacement diet on quality of life in women with obesity and knee osteoarthritis before orthopedic surgery. Nutr Hosp 2018;35:71-77
DOI: <http://dx.doi.org/10.20960/nh.1148>

Correspondence:

Juan José López Gómez. Department of Endocrinology and Nutrition. Hospital Clínico Universitario Valladolid.
Av. Ramón y Cajal, 3. 47003 Valladolid, Spain
e-mail: jjlopez161282@hotmail.com

INTRODUCTION

Osteoarthritis is a joint disease that affects 12-15% of the population aged 25 to 74 years old. The prevalence of this disease increases significantly with age (1).

The most common site of osteoarthritis is the knee. In the United States, this entity represents 37% of the population over 60 years of age; 12% are symptomatic and have functional repercussions. Prevalence is higher in people with older age and higher body mass index (BMI). There is a predominance of females in both the prevalence of osteoarthritis of the knee (2) and the replacement surgery (3). In Spain, the prevalence of osteoarthritis in relation to overweight in 2006 was 32.16% (34.9% in men and 20.93% in women) (4). The prevalence of knee osteoarthritis in the general population in the 60-69 age group was 28.1% (18.1% in men and 37.2% in women) (5).

Overweight and obesity have been shown to be a predictive factor for the development of osteoarthritis. A BMI greater than 25 kg/m² over 40 years increases the risk of symptomatic knee osteoarthritis (6). The mechanisms that link osteoarthritis with obesity are related to the chronic joint load (7), the altered body composition (8), and the situation of subacute inflammation (8).

The basis of any treatment of obesity is the modification of the hygienic-dietary patterns. It has been estimated that if the prevalence of obesity were reverted to levels ten years ago, 111,206 complete knee arthroplasties would be prevented (9). Several studies have estimated that up to half of cases of knee osteoarthritis can be avoided if obesity is eliminated as a risk factor (10).

The improvement of pain and functional disability is the main objective of the patient when considering a surgical treatment of arthropathy. The weight loss through conventional diet therapy has a positive effect on knee arthropathy (11). However, due to physical limitation this weight loss is discrete and can impair muscle mass (12). An alternative to conventional dietary treatment in the surgical patient is the meal-replacement diets. The effectiveness of these diets in the short term (three months) is contrasted. A meta-analysis in 2003 showed an average weight loss of between 6.19-6.50 kg (7% of total weight) compared to the control group, where a loss of 3.23-3.99 kg (4% of initial weight) was observed (13). Even so, there is little evidence of the effect of meal-replacement diets on quality of life in patients with arthropathy.

For this reason, we carried out a study with a meal-replacement diet on women with obesity and knee arthropathy candidates for orthopedic surgery. The main objective was to investigate the weight loss and the modification of body composition secondary to this type of diet, to evaluate the modification of the quality of life after the dietary intervention and to value its relationship with the change of anthropometric parameters and body composition.

METHODS

STUDY DESIGN

An intervention study of one branch with a meal-replacement diet with an oral nutritional supplement (Vegestart®) for three

months was carried out in women with obesity (BMI greater than 30 kg/m²) and knee osteoarthritis pending orthopedic surgery. Patients with active oncologic disease and previous history of alcohol or toxic abuse were excluded.

The study was performed from January 2014 until July 2016. The study was carried out in patients belonging to the health area of Valladolid Este in Spain. These patients were referred from the Department of Traumatology to the Department of Endocrinology and Nutrition of the Hospital Clínico Universitario de Valladolid for weight loss prior to orthopedic knee surgery.

PROCEDURES

All participants provided informed consent to a protocol approved by the local ethical review boards. This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving patients were approved by the Hospital Clínico Universitario de Valladolid (HCUVA) ethics committee on 31-1-2014 with the code PI 14-151 CINV 13-60. All patients signed the informed consent and were included in the study. Patients received nutritional education and a low-fat hyperproteic hypocaloric diet (Table I). The diet was structured in six meals (breakfast, mid-morning, lunch, snack, dinner and late night snack). The lunch and dinner was replaced by an oral nutritional supplement called VEGEStart Complete®, whose nutritional characteristics are described in table I.

The adherence of these diets was assessed each seven days with a phone call by a dietitian to improve compliance of the calorie restriction and macronutrient distribution. The diet compliance was verified with a 24-hour telephone dietary questionnaire every seven days and a 48-hour dietary survey of in face-to-face visits. All parameters were measured at baseline and these variables were repeated after three months.

STUDY VARIABLES

Anthropometry

The anthropometric evaluation of the subjects was performed by determination of weight, height and body mass index (BMI).

The weight was measured without clothing with an accuracy of ± 0.1 kg using a hand scale to the nearest 0.1 kg (Seca, Birmingham, UK). The height was measured with the patient standing up to the nearest centimeter using a stadiometer (Seca, Birmingham, UK). BMI was calculated using the formula: weight (kg)/height x height (m²).

The percentage of weight loss (% WL) was used to assess the relative weight difference.

Bioelectrical impedance measurement

A bioelectrical impedance analysis (BIA) was performed in all subjects. These measurements were performed before the start of the dietary intervention and three months after the intervention.

The BIA was performed on all subjects after a fast of at least five hours. It may be influenced by the degree of hydration, so subjects were warned that they could not exercise or drink alcohol within 48 hours prior to the test.

It was determined by a four-point single-decubitus device. An alternating current of 0.8 mA at 50 kHz produced by a calibrated signal generator (EFG, Akern It) was used and applied to the skin by adhesive electrodes placed on the back of the hand and right foot. Body composition was estimated with the Bodygraff® software.

The parameters analyzed with the BIA were: fat free mass and fat mass. All of them are represented as weight (kg) and percentage with respect to the total weight.

Quality of life test

Two different tests were used to evaluate the impact of the dietary intervention on health status and quality of life. The Short Form Health Survey (SF-36) test was used to evaluate the patient's quality of life before and after treatment. It is an instrument developed from an extensive battery of questionnaires used in the Medical Outcome Study (MOS) (14). In our design, we used the Spanish-translated version (15). It consists of 36 themes, which explore eight dimensions of health status: physical functioning, role physical, bodily pain, general health, vitality, social functioning, role emotional and mental health. The score is directly proportional to the state of health (score ranges from 0 to 100).

The Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) (16) was used to assess the functional limitation and influence of diet therapy in patients with osteoarthritis of the knee and/or hip. We used the validated Spanish version (17). The test was performed face-to-face in front of the patient. This test is used to assess the functionality of the patient from three subscales with different scores: pain, stiffness, physical function, total (results from the sum of the three scores). The interpretation of the test is that the higher score means a worsening in any of the three areas (pain, stiffness and physical function) or in the total. To homogenize the data,

Table I. Macronutrient composition of diet and oral nutritional supplement Vegestart®

Macronutrients	Complete diet	Oral nutrition supplement (200 ml)
Caloric value (kcal)	1035	200
Proteins (g [%TCV])	64.4 (25%)	15.4 (31%)
Lipids (g [%TCV])	19.1 (17%)	5.2 (23%)
<i>Cholesterol (mg)</i>	21	0
Carbohydrates (g [%TCV])	151.6 (59%)	21(42%)
Fiber (g)	15.9	4.2

%TCV: total caloric value percentage.

the scores have been standardized from 0 to 100, where 0 means no alteration and 100 is the situation with the greatest alteration.

STATISTICAL ANALYSIS

The data were analyzed using the SPSS statistical package (SPSS for Windows version 15.0, 2008 SPSS INC, Chicago, USA). An analysis of patients by intention to treat was performed. Sample size was calculated to detect a percentage of weight loss over 5%. The level of significance was conventionally set at $p \leq 0.05$.

RESULTS

DEVELOPMENT OF THE STUDY

Eighty-one women with obesity and knee osteoarthritis pending orthopedic surgery were recruited between January 2014 and July 2016.

After three months of intervention, 75 (92.6%) patients maintained complete therapeutic adherence while six (7.40%) patients started the diet but did not complete it for different reasons (Fig. 1).

INITIAL EVALUATION

The mean age of the individuals was 62.23 (8.50) years.

Anthropometry and body composition

The initial weight was 99.33 (14.61) kg, while BMI was 40.80 (4.40) kg/m².




Figure 1.

Therapeutic intervention scheme.

In the analysis of body composition, fat mass and fat free mass were evaluated. A percentage of fat mass of 47.67% (5.19) with respect to the total weight and a percentage of fat free mass of 52.25% (5.24) were observed.

Quality of life

Test SF-36

To evaluate the patients' quality of life, the structured SF-36 test was performed in eight dimensions:

- Decreased values were observed in general health, physical functioning, role-physical, bodily pain and mental health. Decreased vitality values were also observed (Table II).
- In the areas of role emotional and social functioning, no decrease was observed (Table II).

WOMAC

This test was used for the assessment of pain and its implication in the functionality of the patient. All patients had altered values at all levels (pain: 52.94% [26.08]; stiffness: 50% [25-75]; functional capacity: 49.19% [27.01]); as well as in total index: 49.24% [25.53]).

EFFECT OF DIETARY TREATMENT

Anthropometry and body composition

A weight loss rate of 8.23% (4.04) was observed after a three-month meal-replacement hypocaloric diet. There was a decrease in all body components. When assessing the change in body composition at three months, there was a relative decrease in the percentage of fat mass 2.66% (0.53-3.92) and a relative increase in the percentage of fat-free mass 2.30% (3.72) (Table III).

Quality of life

SF-36 test

An improvement was observed in total score (basal time: 49.35 [20.41], at three months: 58.71 [17.07], $p < 0.01$). We obtained an improvement rate of 15.97% (28.08).

In the analysis of the different spheres (Table II) there was a significant improvement in all spheres except for social functioning (Fig. 2).

WOMAC

The total score of WOMAC showed a significant improvement (basal time: 49.24% [25.53], at three months: 40.59% [21.76], $p < 0.01$). The decrease proportion of this test after intervention

Table II. Modification in quality of life test: SF-36 and WOMAC before and after three months of treatment

		Basal	3 months	p
SF-36	General health	41.49 (16.53)	48.79 (13.63)	< 0.01
	Physical functioning	25 (10-45)	35 (20-50)	< 0.01
	Role physical	25 (0-100)	75 (12.5-100)	< 0.01
	Role emotional	69.13 (42.74)	81.85 (36.59)	< 0.01
	Social functioning	75 (50-100)	87.5 (50-100)	0.10
	Bodily pain	43.95 (23.68)	54.23 (27.76)	< 0.01
	Vitality	44.22 (23.68)	57.71 (54.34)	0.03
	Mental health	59.80 (27.40)	68.49 (22.98)	< 0.01
WOMAC test	Pain	52.94 (26.08)	45.25 (23.57)	< 0.01
	Stiffness	50 (25-75)	25 (12.5-50)	0.02
	Functional capacity	49.19 (27.01)	40.16 (22.06-54.41)	< 0.01

Table III. Modification of parameters of anthropometry and body composition before and after three months of treatment

	Basal	3 months	p
Weight (kg)	99.33 (14.61)	91.02 (13.01)	< 0.01
BMI (kg/m ²)	40.80 (4.40)	37.02 (5.00)	< 0.01
Fat mass (kg)	46.60 (10.20)	40.19 (9.65)	< 0.01
Fat mass (%)	47.67 (5.19)	44.93 (6.43)	< 0.01
Fat free mass (kg)	55.49 (12.03)	52.87 (10.77)	< 0.01
Fat free mass (%)	52.25 (5.24)	54.51 (5.96)	< 0.01

BMI: body mass index.

was 13.64% (0-35.68). There was significant improvement in the three spheres (Table II and Fig. 2).

Relationship of anthropometry changes with improvement of the quality of life

The percentage of weight loss showed a positive correlation with the percentage of variation of SF-36 ($r = 0.23$; $p = 0.04$), with the WOMAC pain domain ($r = 0.34$; $p < 0.01$), with the functional capacity domain ($r = 0.31$; $p < 0.01$), and the percentage change in WOMAC total score ($r = 0.35$; $p < 0.01$).

A multivariate analysis was performed to determine the influence of anthropometric changes on the modification of the quality of life scores. Improvement in quality of life was observed to be related to weight loss regardless of changes in body composition and age (Table IV). There was no relationship between the WOMAC test, weight loss and body composition (Table IV).




Figure 2. Variation of the different areas of the SF-36 and WOMAC tests before the start of the intervention and three months later. *Significant difference ($p < 0.05$).

Table IV. Relation between improvement quality of life test (SF-36 and WOMAC) with anthropometry and body composition

Improvement 10% SF-36 test (Total) (Yes/No)			
	OR	CI (95%)	p
Age	0.96	(0.90-1.02)	0.16
Weight loss	1.20	(1.03-1.36)	0.02
Fat mass loss	1.03	(0.94-1.13)	0.49
Fat free mass gain	0.98	(0.89-1.09)	0.73
Improvement 10% WOMAC test (Total) (Yes/No)			
Age	0.98	(0.93-1.04)	0.61
Weight loss	1.01	(0.88-1.15)	0.92
Fat mass loss	1.05	(0.97-1.14)	0.23
Fat free mass gain	1.02	(0.92-1.12)	0.75

DISCUSSION

The effect of dietary treatment of obesity has shown several benefits on the complications associated with knee osteoarthritis. Even so, there are few studies that analyze the effect on quality of life in these patients when using a meal-replacement diet.

The SF-36 is a very useful test to evaluate the quality of life in a global way and in terms of different aspects affecting it. When comparing the score of the SF-36 test with the values of the test for the general Spanish population, a decrease in all values is observed except in the areas of role emotional, social functioning and mental health (18). This could be justified by habituation to chronic symptoms, or because of the difficulty of the test to assess these types of symptoms.

When comparing our data to those of similar populations (obese with degenerative arthropathy), quite a few similarities in the alteration of the different spheres of the SF-36 test are observed. When analyzing the excess weight, a previous study compared the obese *versus* non-obese population; in the age group between 55 and 64 years worse scores in the population with obesity in spheres of physical functioning, role physical, pain, general health and vitality were observed, while in other spheres the score was similar (19). Similar levels were also observed in populations with the same characteristics as ours in which different comorbidities are combined. In this way, it has been shown that the SF-36 physical spheres score worsens with the increase in the number of comorbidities, and particularly with the presence of obesity (20,21).

Given the special characteristics of these patients, a more specific test for arthropathy, such as WOMAC, was performed. When analyzing the WOMAC, a score of pain, stiffness, functional capacity and the total of the sum close to half were observed. The levels of pain and altered functional capacity in the WOMAC test are variable according to the study ranged from lower scores around 20% (22) till higher scores around 40-50% in populations with obesity (23). This data means that higher values

of BMI produce worse values of WOMAC (24). This would justify that the values obtained in our study were in a range of greater pain and disability. The pain and physical situation scores would resemble those of the other test performed (SF-36). This situation has been evaluated in other studies, with similar findings and improvement after arthroplasty replacement (25). The effect of dietary treatment on these parameters does not have such clear evidence in the literature.

The results on weight loss in our study are similar to those performed with the same intervention. In a study by De Luis et al. in a population of patients with arthropathy, a greater weight loss was observed in the treatment group with a commercial diet compared to dietary advice (26), with a mean weight loss of 7.7 kg compared to 3.92 kg in the control group. In another meta-analysis by Anderson, it was found that there was a greater percentage of weight loss, up to 9.3% in men and 8.6% in women, with an initial BMI ranged from 28 to 35 kg/m². These data were similar to what was done in our study (27).

When the change in body composition was analyzed, a decrease in the percentage of fat mass that was associated with a relative increase of fat-free mass was observed. This data shows that maintaining an adequate protein intake can help to reduce excess muscle loss, despite caloric restriction. This topic has been observed in different studies with hyperproteic meal-replacement diets of one or more meals (28,29).

The main symptoms that are usually evaluated in these patients are the improvement of pain and stiffness. In the WOMAC test, an improvement in the parameters of stiffness and physical function was observed. This change in functional capacity score is seen in multiple studies associated with weight loss in patients with arthropathy. In 2010, a meta-analysis on long-term weight loss in obese elderly patients showed that functional impairment and overall quality of life improved with weight loss in obese elderly patients (11).

We observed an improvement in all of the items of the SF-36 test after intervention except for social functioning. This situation could be due to two reasons: the baseline scores were close to the average of the general Spanish population, and the measurement of this sphere through a single question with three degrees makes it difficult to show differences.

In the SF-36 test, although there were no differences in social functioning, there was an improvement in mental health. This situation related to the decrease of pain and the improvement of the functional capacity in the evaluated tests relates the influence of these two characteristics to the psychic situation of the patients.

The reduction of weight in the context of the treatment of osteoarthritis of different locations, especially those of load (hip and knee), has a positive effect on pain. The effect starts to be effective with weight losses above 5% (30). This relationship has been analyzed in many studies of osteoarthritis of the knee using the WOMAC (31) with greater effect in those patients with greater weight loss. This effect is enhanced if there is a long-term maintenance of weight loss (32).

In our study, there were isolated correlations between the greater weight loss and the improvement of the WOMAC in its total

score, in the field of pain and functional capacity. This approach has recently been observed in a study where the percentage of weight change was associated with the WOMAC score, both gain and weight loss (33). In the case of the analysis of the improvement of the articular functionality after the loss of weight the data are variable in the literature. In several cases, a functional improvement was observed in relation to weight loss in diet-only management (34), with intensive diet (35), or a combination of diet and exercise combined (36).

An independent relationship was observed in the weight loss with the intensity of the SF-36 test improvement. This relationship did not occur with the WOMAC test. The differences found between both tests in obese patients with arthropathy in relation to weight loss can be related to a partial improvement in pain and functional capacity, and with the positive effect that the weight loss had on the patient's well-being. It has been observed that this score improves more in obese patients than in non-obese patients after moderate weight control (37). Therefore, these results may be due to the influence of weight loss on quality of life in other aspects than mechanical ones, such as psychological and social factors.

The main limitation of our study was the absence of a control group with a different dietary intervention to be able to categorize the impact of the diet itself on the quality of life. Second, the inclusion of only women limits the extrapolation of study data, but this decision was made due to the higher prevalence of osteoarthritis of the knee in women. Finally, it would be interesting to carry out a long-term assessment of the maintenance of weight loss and the change of the quality of life of the patient.

CONCLUSIONS

A short term dietary treatment through a meal replacement diet in obese women with knee osteoarthritis pending surgery showed: a weight decrease between 5-10% with a relative decrease of fat mass and relative increase of fat-free mass; an improvement in the quality of life measured by the SF-36 test in all its spheres except for social functioning; and improvement in all dimensions of the WOMAC test. Adjusting for age and body composition weight loss showed an independent relationship with SF-36 improvement. Further studies are needed to evaluate this dietary intervention in patients with other arthropathy and during long term interventions.

TRANSPARENCY DECLARATION

The lead author affirms that this manuscript is an honest, accurate, and transparent account of the study being reported. The lead author affirms that no important aspects of the study have been omitted and that any discrepancies from the study as planned have been explained. This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving patients were approved by the Hospital

Clínico Universitario de Valladolid (HCUVA) ethics committee on the 31-1-2014 with the code PI 14-151 CINV 13-60.

REFERENCES

- Guilak F. Biomechanical factors in osteoarthritis. *Best Pract Res Clin Rheumatol* 2011;25(6):815-23.
- Núñez M, Núñez E, Sastre S, Del-Val JL, Segur JM, Macule F. Prevalence of knee osteoarthritis and analysis of pain, rigidity and functional incapacity. *Orthopedics* 2008;31:753.
- Kremers HM, Larson DR, Crowson CS, Kremers WK, Washington RE, Steiner CA, et al. Prevalence of total hip and knee replacement in the United States. *J Bone Joint Surg Am* 2015;97(17):1386-97.
- Martín Ramiro JJ, Álvarez-Martín E, Gil-Prieto R. Discapacidad atribuible al exceso de peso en España. *Med Clin (Barc)* 2014;143(4):150-6.
- Fernández-López JC, Laffon A, Blanco FJ, Carmona L, EPISER Study Group. Prevalence, risk factors, and impact of knee pain suggesting osteoarthritis in Spain. *Clin Exp Rheumatol* 2008;26(2):324-32.
- Sridhar MS, Jarrett CD, Xerogeanes JW, Labib SA. Obesity and symptomatic osteoarthritis of the knee. *J Bone Joint Surg Br* 2012;94:433-40.
- Felson DT, Goggins J, Niu K, Zhang Y, Hunter DJ. The effect of body weight on progression of knee osteoarthritis is dependent on alignment. *Arthritis Rheum* 2004;50:3904-09.
- Wluka AE, Lombard CB, Cicuttini FM. Tackling obesity in knee osteoarthritis. *Nat Rev Rheumatol* 2013;9(4):225-35.
- Messier SP. Obesity and osteoarthritis: Disease genesis and nonpharmacologic weight management. *Rheum Dis Clin North Am* 2008; 34(3):713-29.
- Koonce RC, Bravman JT. Obesity and osteoarthritis: More than just wear and tear. *J Am Acad Orthop Surg* 2013;21:161-9.
- Witham MD, Avenell A. Interventions to achieve long-term weight loss in obese older people. A systematic review and metaanalysis. *Age Ageing* 2010;39:176-84.
- Henriksen M, Christensen R, Danneskiold-Samsøe B, Bliddal H. Changes in lower extremity muscle mass and muscle strength after weight loss in obese patients with knee osteoarthritis: A prospective cohort study. *Arthritis Rheum* 2012;64(2):438-42.
- Heysfield SB, Van Mierlo CA, Van der Knaap HC, Heo M, Frier HI. Weight management using a meal replacement strategy: Meta and pooling analysis from six studies. *Int J Obes* 2003;27:537-54.
- Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36) (I). Conceptual framework and item selection. *Med Care* 1992;30:473-83.
- Alonso J, Prieto L, Antó JM. The Spanish version of the SF-36 Health Survey (the SF-36 health questionnaire): An instrument for measuring clinical results. *Med Clin (Barc)* 1995;104(20):771-6.
- Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of WOMAC: A health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol* 1988;15(12):1833-40.
- Escobar A, Quintana JM, Bilbao A, Azkárte J, Güenaga JI. Validation of the Spanish version of the WOMAC questionnaire for patients with hip or knee osteoarthritis. *Western Ontario and McMaster Universities Osteoarthritis Index. Clin Rheumatol* 2002;21(6):466-71.
- Alonso J, Regidor E, Barrio G, Prieto L, Rodríguez C, De la Fuente L. Population reference values of the Spanish version of the Health Questionnaire SF-36. *Med Clin (Barc)* 1998;111(11):410-6.
- Anandacoomarasamy A, Caterson ID, Leibman, Smith GS, Sambrook PN, Fransen M. Influence of BMI on health-related quality of life: Comparison between an obese adult cohort and age-matched populations norms. *Obesity (Silver Spring)* 2009;17(11):2114-8.
- Wesseling J, Welsing PM, Bierma-Zeinstra MA, Dekker J, Gorter KJ, Kloppenburg M, et al. Impact of self-reported comorbidity on physical and mental health status in early symptomatic osteoarthritis: The CHECK (Cohort Hip and Cohort knee) study. *Rheumatology* 2013;52:180-8.
- Van Dijk GM, Veenhof C, Schellevis F, Hulsmans H, Bakker JP, Arwert H, et al. Comorbidity, limitations in activities and pain in patients with osteoarthritis of the hip or knee. *BMC Musculoskelet Disord* 2008;9:95.
- Collins JE, Katz J, Dervan E, Losina E. Trajectories and risk profiles of pain in persons with radiographic, symptomatic knee osteoarthritis: Data from the osteoarthritis initiative. *Osteoarthritis Cartilage* 2014;22(5):622-30.
- Garver MJ, Focht BC, Dials J, Rose M, Lucas AR, Devor ST, et al. Weight status and differences in mobility performance, pain symptoms, and physical activity in older, knee osteoarthritis patients. *Arthritis* 2014;375909.
- Sanghi D, Srivastava RN, Singh A, Kumari R, Mishra R, Mishra A. The association of anthropometric measures and osteoarthritis knee in no-obese subjects: A cross sectional study. *Clinics (Sao Paulo)* 2011;66(2):275-9.
- Núñez M, Lozano L, Núñez E, Segur JM, Sastre S, Maculé F, et al. Total knee replacement and health-related quality of life: Factors influencing long-term outcomes. *Arthritis Rheum* 2009;61(8):1062-9.
- De Luis DA, Izaola O, García Alonso M, Aller R, Cabezas G, De la Fuente B. Randomized clinical trial between nutritional counseling and commercial hypocaloric diet in weight loss in obese patients with chronic arthropathy. *Med Clin (Barc)* 2009;132(19):735-9.
- Anderson JW, Luan J, Hoie LH. Structured weight-loss programs: Meta-analysis of weight loss at 24 weeks and assessment of effects of intervention intensity. *Adv Ther* 2004;21:61-75.
- Magrans-Courtney T, Wilborn C, Rasmussen C, Ferreira M, Greenwood L, Campbell B, et al. Effects of diet type and supplementation of glucosamine, chondroitin, and MSM on body composition, functional status, and markers of health in women with knee osteoarthritis initiating a resistance-based exercise and weight loss program. *J Int Soc Sports Nutr* 2011;8(1):8.
- Treyzon L, Chen S, Hong K, Yan E, Carpenter CL, Thames G, et al. A controlled trial of protein enrichment of meal replacements for weight reduction with retention of lean body mass. *Nutr J* 2008;7:23.
- Bliddal H, Leeds AR, Christensen R. Osteoarthritis, obesity and weight loss: Evidence, hypotheses and horizons - A scoping review. *Obes Rev* 2014;15:578-86.
- Weiss E. Knee osteoarthritis, body mass index and pain: Data from the osteoarthritis initiative. *Rheumatology (Oxford)* 2014;53(11):2095-9.
- Bliddal H, Leeds AR, Stigsgaard L, Astrup A, Christensen R. Weight loss as treatment for knee osteoarthritis symptoms in obese patients: 1-year results from a randomized controlled trial. *Ann Rheum Dis* 2011;70(10):1798-803.
- Teichtahl AJ, Wluka AE, Tanamas SK, Wang Y, Strauss BJ, Proietto J, et al. Weight and change in tibial cartilage volumens and symptoms in obese adults. *Ann Rheum Dis* 2015;74(6):1024-9.
- Riddle DL, Stratford PW. Body weight changes and corresponding changes in pain and function in persons with symptomatic knee osteoarthritis. A cohort study. *Arthritis Care Res (Hoboken)* 2013;65(1):15-22.
- Foy CG, Lewis CE, Hairston KG, Miller GD, Lang W, Jakicic JM, et al. Intensive lifestyle intervention improves physical function among obese adults with knee pain: Findings from the look AHEAD trial. *Obesity (Silver Spring)* 2011;19(1):83-93.
- Messier P, Mihalko SL, Legault C, Miller GD, Nicklas BJ, De Vita P, et al. Effects of intensive diet and exercise on knee joint loads, inflammation, and clinical outcomes among overweight and obese adults with knee osteoarthritis. *JAMA* 2013;310(12):1263-73.
- Dowsey MM, Liew D, Stoney JD, Choong PF. The impact of obesity on weight change and outcomes at 12 months in patients undergoing total hip arthroplasty. *Med J Aust* 2010;193(1):17-21.