

Nutrición Hospitalaria

Nutrición Hospitalaria

ISSN: 0212-1611

ISSN: 1699-5198

Grupo Arán

Ajejas Bazán, M.^a Julia; Jiménez Trujillo, María Isabel; Wärnberg, Julia;
Domínguez Fernández, Silvia; López de Andrés, Ana; Pérez Farinós, Napoleón
Prevalence of childhood overweight/obesity in Spain 1993-2011 and associated risk factors in 2011
Nutrición Hospitalaria, vol. 35, no. 1, 2018, January-February, pp. 84-89
Grupo Arán

DOI: <https://doi.org/10.20960/nh.1355>

Available in: <https://www.redalyc.org/articulo.oa?id=309258226015>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative



Trabajo Original

Obesidad y síndrome metabólico

Prevalence of childhood overweight/obesity in Spain 1993-2011 and associated risk factors in 2011

Prevalencia de sobrepeso y obesidad infantil en España (1993-2011) y factores asociados en 2011

M.^a Julia Ajejas Bazán^{1,2}, María Isabel Jiménez-Trujillo³, Julia Wärnberg⁴, Silvia Domínguez Fernández¹, Ana López-de-Andrés³ and Napoleón Pérez-Farinós⁵

¹Department of Nursing. Nursing, Physiotherapy and Podology Faculty. Universidad Complutense de Madrid. Madrid, Spain. ²Institute of Defense Preventive Medicine. Ministerio de Defensa. Madrid, Spain. ³Preventive Medicine and Public Health Teaching and Research Unit. Health Sciences Faculty. Universidad Rey Juan Carlos. Alcorcón, Madrid, Spain.

⁴Department of Nursing. Health Sciences Faculty. Málaga, Spain. ⁵Department of Public Health and Psychiatry. Medicine Faculty. Universidad de Málaga. Málaga, Spain

Abstract

Introduction: Childhood obesity is a recognized public health problem. The present work reports the changing prevalence of childhood overweight/obesity in Spanish boys and girls over the period 1993-2011, and examines the risk factors apparent in 2011.

Methods: Children with a body mass index (BMI) of ≥ 25 were deemed overweight, and those with a BMI of ≥ 30 were deemed obese. Overweight and obesity was consistently more common among boys than among girls.

Results: The prevalence of overweight and obesity in Spain increased over the study period.

Conclusions: According to the 2011 data, children who undertook no physical activity, or whose parents/guardians had a low level of education, showed the highest prevalence of obesity.

Key words:

Overweight. Pediatric obesity. Prevalence. Risk factors.

Resumen

Introducción: la obesidad infantil es un problema de salud pública a nivel mundial. El objetivo de este trabajo es describir la evolución de la prevalencia de sobrepeso/obesidad infantil en niños y niñas durante el periodo 1993-2011 y analizar cuáles son los factores de riesgo asociados, utilizando la Encuesta Nacional de Salud (ENS) de 2011.

Métodos: los niños con índice de masa corporal (IMC) ≥ 25 se consideraron con sobrepeso y aquellos con IMC ≥ 30 , como obesos. Se hizo uso de la ENS desde 1993 a 2011.

Resultados: la prevalencia de sobrepeso y obesidad en España aumentó durante el periodo de estudio. La prevalencia de sobrepeso y obesidad fue más elevada en niños que en niñas.

Conclusiones: según los datos de 2011, los niños que no realizaban ningún tipo de actividad física o cuyos padres tenían un nivel de educación bajo mostraron la mayor prevalencia de obesidad.

Palabras clave:

Obesidad. Sobrepeso. Pediatría. Prevalencia. Factores de riesgo.

Received: 14/06/2017 • Accepted: 13/08/2017

Ajejas Bazán MJ, Jiménez-Trujillo MI, Wärnberg J, Domínguez Fernández S, López-de-Andrés A, Pérez-Farinós N. Prevalence of childhood overweight/obesity in Spain 1993-2011 and associated risk factors in 2011. Nutr Hosp 2018;35:84-89

DOI: <http://dx.doi.org/10.20960/nh.1355>

Correspondence:

María Julia Ajejas Bazán. Department of Nursing. Nursing, Physiotherapy and Podology Faculty. Universidad Complutense de Madrid. Plaza de Ramón y Cajal, s/n. 28040 Madrid, Spain
e-mail: majejas@ucm.es

INTRODUCTION

The prevalence of childhood obesity has increased across Europe (1) and it is now a recognized public health problem. According to the World Health Organization (WHO), obesity is a chronic disease that has reached epidemic proportions; in some parts of the world, 17.6% of children under five years of age are obese (2). Recent studies in different countries have shown 10% of all school-age children to have excess body fat, increasing their risk of developing chronic diseases (3). Indeed, overweight/obesity has become one of the most common nutritional problems among children of developed countries, though it is by no means limited to them (3,4).

Obesity is increasingly affecting all age groups and is associated with increased morbidity and mortality (5,6). The complications of obesity most commonly appear during adulthood, although nowadays they are also being detected in childhood. Many factors are associated with its appearance; genetic, sociodemographic (including sex and level of parental education) and lifestyle factors are all involved, including the duration of breastfeeding, the daily taking of breakfast, getting enough sleep, time spent in front of monitors (7) and the amount of physical activity undertaken (6,8). Tackling obesity is commonly one of the best set out objectives in healthcare strategies (9), but understanding the factors that encourage its appearance is vital if it is to be prevented (10,11).

The aim of the present work was to describe the prevalence of overweight and obesity in Spanish children (boys and girls) using data provided by the National Health Surveys (*Encuestas Nacionales de Salud*) of 1993-2011, and to determine the main socioeconomic and lifestyle factors influencing the appearance of these problems in 2011.

METHODS

DESIGN

This descriptive, cross-sectional study examined data provided by the 1993-2011 Spanish National Health Surveys (Ministerio de Sanidad, 1993; 1995; 1997; 2001; 2003; 2006; 2011). These surveys question members of the public regarding their health status and provide information on perceived morbidity, risk factors, the use of health services, and the following of preventive practices.

SUBJECTS

The data examined were those of children aged 2-15 years, chosen by stratified multi-stage sampling. Entries for children of this age were stratified by region, then by province (as conglomerates), then by "habitat" (urban or rural area). Schools were then chosen at random within these "habitats", then classes by age group between 2 and 15 years, followed by random sampling of

the children in these age groups. The final sample size (which included data from only fully completed survey questionnaires) was $n = 23,237$ (11,663 girls and 11,574 boys). The children's heights and weights (as recorded by their parents/guardians) were used to calculate their body mass indices (BMI). Children with a $BMI \geq 25$ were deemed overweight, and those with a $BMI \geq 30$ were deemed obese.

STATISTICS

The prevalence (with 95% confidence intervals) of overweight and obesity was analyzed for each survey year separately.

The association of overweight and obesity with each independent variable as measured in 2011, i.e., sex, age at the time of the survey, recommended number of hours of sleep (3-4 years 12 h, 4-5 years 11 h, 5-11 years 10 h and 11+ years 9 h) (10,13), physical activity (measured as yes/no), hours spent watching TV or using other monitors (< 2 or ≥ 2 h per day) (14), having been breast-fed for the first three or six months (yes/no for each), taking breakfast daily (yes/no), level of education of head of family (no education or obligatory schooling, school baccalaureate or technical award, higher education) (15,16), was examined using the Chi-squared test. Significance was set at $p \leq 0.05$. Adjusted odds ratios (OR) were obtained by bivariate and consequent multivariate regression analysis. All calculations were performed using SPSS v.21.0 for Windows.

ETHICS

Since all data used were rendered anonymous and were in the public domain (www.ine.es), no ethical approval was required.

RESULTS

Table I shows the prevalence of childhood overweight and obesity for girls and boys over the study period. In 1993, 7.0% of boys were overweight, but in 2011 this figure had risen to 13.2%; for girls, the figures were 6.0% and 10.2% respectively. In 1993, 13.8% of boys and 8.7% of girls were obese, while in 2011, 13.6% of boys and 9.9% of girls were obese. Overweight and obesity was consistently more common among boys than in girls.

Table II reveals that, in 2011 (note that 2011 health survey data were used to determine all risk factor associations), those children who belonged to a family whose head had no or only low-level studies showed a higher prevalence of overweight (boys 13.2%, girls 7.4%) compared to those who had higher education (boys 19.1%, girls 9.0%) ($p < 0.05$ in all cases). Similar results were seen for obesity (boys 16.3% and girls 5.5% compared to 10.7 and 8.1%; $p < 0.05$ for the comparison between boys).

Those children who did not sleep the recommended number of hours were more commonly obese (boys 16.3%, girls 12.6%)

Table I. Prevalence of overweight/obesity in boys and girls, as recorded in different National Health Surveys (1993-2011)

		Both sexes			Boys			Girls		
		n	%	(95% CI)	n	%	(95% CI)	n	%	(95% CI)
Normal-weight	1993	3,926	82.3	(81.1-83.5)	1,937	79.3	(77.5-81.1)	1,989	85.4	(83.8-87)
	1995	1,517	83.9	(82.1-85.7)	764	82.3	(79.6-85)	753	85.5	(83.0-88.0)
	1997	1,452	81.7	(79.7-83.7)	718	79.1	(76.1-82.1)	734	84.5	(81.9-87.1)
	2001	3,663	80.6	(79.3-81.9)	1,826	78.2	(76.3-80.1)	1,837	83	(81.3-84.7)
	2003	4,258	78.9	(77.7-80.1)	2,125	76.6	(74.8-78.4)	2,133	81.3	(79.6-83.0)
	2006	4,640	77.4	(76.2-78.6)	2,286	74.2	(72.4-76.0)	2,354	80.7	(79.1-82.3)
	2011	2,864	76.3	(74.7-77.9)	1,470	73.2	(70.9-75.5)	1,394	79.9	(77.8-82.0)
Overweight	1993	309	6.5	(5.9-7.0)	170	7	(3.2-10.8)	139	6	(2.1-9.9)
	1995	131	7.2	(2.8-11.6)	65	7	(0.8-13.2)	66	7.5	(1.1-13.9)
	1997	138	7.8	(3.3-12.3)	73	8	(1.4-14.2)	65	7.5	(1.1-13.9)
	2001	362	8	(5.2-10.8)	203	8.7	(4.8-12.6)	159	7.2	(3.2-11.2)
	2003	598	11.1	(8.6-13.6)	309	11.1	(7.6-14.6)	289	11	(7.4-14.6)
	2006	677	11.3	(8.9-13.7)	366	11.9	(8.6-15.2)	311	10.7	(7.3-14.1)
	2011	443	11.8	(8.8-14.8)	265	13.2	(9.1-17.3)	178	10.2	(5.8-14.6)
Obese	1993	538	11.3	(8.6-14)	336	13.8	(10.1-17.5)	202	8.7	(4.8-12.6)
	1995	161	8.9	(4.5-13.3)	99	10.7	(4.6-16.8)	62	7	(0.6-1.4)
	1997	187	10.5	(6.1-14.9)	117	12.9	(6.8-19)	70	8.1	(1.7-14.5)
	2001	522	11.5	(8.8-14.2)	305	13.1	(9.3-16.9)	217	9.8	(5.8-13.8)
	2003	544	10.1	(7.6-12.6)	341	12.3	(8.8-15.8)	203	7.7	(4.0-11.4)
	2006	680	11.3	(8.9-13.7)	427	13.9	(10.6-17.2)	253	8.7	(5.2-12.2)
	2011	445	11.9	(8.9-14.9)	272	13.6	(9.5-17.7)	173	9.9	(5.4-14.4)

n: sample size; CI: confidence intervals.

Table II. Distribution of normal-weight, overweight and obesity with respect to sex and lifestyle variables according to the 2011 National Health Survey results

		Boys n (%)			Girls n (%)		
		Normal-weight	Overweight	Obese	Normal-weight	Overweight	Obese
Education level of family head	No studies or just obligatory schooling	670 (70.5)	125 (13.2)*	155 (16.3)*	458 (87.1)	39 (7.4)*	29 (5.5)
	Baccalaureate or technical training	521 (74.7)	90 (13.0)*	86 (12.3)*	197 (83.1)	17 (7.2)	23 (9.7)
	Higher education	275 (77.2)	50 (12.1)*	31 (10.7)*	92 (82.9)	10 (9.0)*	9 (8.1)
Hours of sleep	Recommended no. of hours not slept	842 (70.9)	152 (12.8)*	193 (16.3)*	778 (78.0)	94 (9.4)*	126 (12.6)*
	Recommended no. of hours slept	628 (76.6)	113 (13.8)	79 (9.6)*	616 (82.5)	84 (11.2)	47 (6.3)*
Physical activity	No	170 (62.0)	38 (13.9)	66 (24.1)	276 (74.4)	43 (11.6)	52 (14.0)
	Yes	1,300 (75.1)	227 (13.1)*	205 (11.8)*	1,116 (81.3)	135 (9.8)*	121 (8.8)
Time spent at a monitor (TV, computer, video-gaming)	≤ 2 h	101 (76.5)	15 (11.4)	16 (12.1)	83 (80.6)	11 (10.7)	9 (8.7)
	> 2 h	832 (71.5)	165 (14.2)	166 (14.3)	729 (77.6)	112 (11.9)	99 (10.5)
Breast-fed until 3 months old	No	1,278 (73.9)	241 (13.9)*	210 (12.1)*	1,221 (81.1)	157 (10.4)*	127 (8.4)
	Yes	192 (69.1)	24 (8.6)*	62 (22.3)	173 (72.1)	21 (8.8)*	46 (19.2)
Breast-fed until 6 months old	No	1,332 (73.6)	248 (13.7)	230 (12.7)	1,273 (80.7)	165 (10.5)	140 (8.9)*
	Yes	138 (70.1)	17 (8.6)	42 (21.3)	121 (72.5)	13 (7.8)	33 (19.8)
Took daily breakfast	No	7 (36.8)	5 (26.3)	7 (36.8)	14 (82.4)	2 (11.8)	1 (5.9)
	Yes	1,463 (73.6)	260 (13.1)	265 (13.3)*	1,380 (79.9)	176 (10.2)	172 (10.0)

n: Sample size. *p < 0.05.

than those who slept the recommended amount (boys 9.6%, girls 6.3%) ($p < 0.05$ both sexes). Those children who undertook some kind of physical activity were less commonly overweight (boys 13.1%, girls 9.8%) than those who undertook none (boys 13.9%, girls 11.6%) ($p < 0.05$ for both sexes); they were also less commonly obese (boys 11.8%, girls 8.8%) than those who undertook no physical activity (boys 24.1%, girls 14%) ($p < 0.05$ for the comparison between boys). No significant differences were seen between children who used monitors (TV, playing videogames, etc.) more than two hours per day compared to those who used them for less time, either in terms of overweight (boys 14.2%, girls 11.9% vs boys 11.4%, girls 10.7%) or obesity (boys 14.3%, girls 10.5% vs boys 12.1%, girls 8.7%).

Those children exclusively breast-fed for their first three months of life were less commonly overweight (boys 8.6%, girls 8.8%) than those who were not so fed (boys 13.9%, girls 10.4%) ($p < 0.05$ for boys and girls). No significant differences were seen between children who were exclusively breast-fed for the first six months of life and those who were not in terms of overweight (boys 8.6%, girls 7.8% vs boys 13.7%, girls 10.5%). No comparisons were made for obesity given the small sample size available, neither was any significant difference seen in terms of overweight between children who ate or did not eat breakfast (boys 26.3%, girls 11.8% vs boys 13.1% girls 10.2%).

Table III shows the results of the multivariate analysis. Age, male sex, being breast-fed until three months, not undertaking physical activity, and the level of education of the head of the family were independently and significantly associated with being overweight. The same variables were independently and significantly associated with being obese.

DISCUSSION

The present results show that the prevalence of overweight and obesity increased over the study period in both boys and girls. In addition, overweight and obesity was consistently more common among boys.

The present prevalence data for 1997 were compared to those of the enKid study (16) which was performed in 1997-1998. The prevalence of overweight was higher in this latter study (boys 14.3%, girls 15.6%) than in the present work (boys 8.0%, girls 7.5%). In addition, the present prevalence data for 2011 were compared to those reported by the Aladino 2011 (17). This latter study reported a higher prevalence of overweight (boys 22.9%, girls 17.4% vs the present boys 13.2%, girls 10.2%), but a lower prevalence of obesity (boys 7.7%, girls 4.7% vs the present boys 13.6%, girls 9.9%). These discrepancies might be due to differences in the methodologies employed.

According to the 2011 health survey results (used for all associations with lifestyle factors), the prevalence of overweight and obesity was higher among children of families whose heads were the least well educated. The enKid study reported similar results for obesity (15.6% vs 10.9% in families with the head in the highest education bracket) (16). Similarly, the Aladino 2011 study reported that, among normal-weight children, 58.5% of their mothers and 59.4% of their fathers had university studies, while among obese children only 14.6% of their mothers and 14.7% of their fathers had university studies. This might reflect a limitation of economic resources faced by families with low education levels, or their more limited knowledge of what makes a healthy diet, or perhaps a different aesthetic outlook (17).

Table III. Multivariate analysis of factors associated with overweight/obesity according to the data collected by the 2011 National Health Survey

		Overweight		Obesity	
		OR	(CI 95%)	OR	(CI 95%)
Sex	Boys	1.22***	(1.15-1.30)	1.4***	(1.29-1.51)
	Girls	1		1	
Age	Years	1.03***	(1.02-1.04)	0.88***	(0.87-0.89)
Breast-fed until 3 months old	Yes	1		1	
	No	1.24**	(1.07-1.43)	1.24**	(1.07-1.42)
Physical activity	Yes	1		1	
	No	1.03**	(1.01-1.06)	1.22***	(1.11-1.35)
Education level of family head	No studies or just obligatory schooling	1.21***	(1.17-1.26)	1.36***	(1.21-1.52)
	Baccalaureate or technical training	1.10***	(1.06-1.13)	1.22**	(1.09-1.39)
	Higher education	1		1	

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Those children who slept the recommended number of hours for their age were less commonly obese than those who did not get enough sleep. Very few studies performed in Spain have examined the link between the number of hours spent sleeping and overweight/obesity. The Aladino 2011 study did report, however, results very similar to those of the present work: 10.7% of those who slept sufficient hours were obese, rising to 15.2% among those who did not, whereas in the present study these figures were boys 9.6% vs 16.3%, and girls 6.3% vs 12.6% (17).

No differences were seen in overweight or obesity between those children who spent under or over two hours per day using monitors. This is in contrast to that reported in other studies. For example, in Sweden, children who had a television in their bedrooms, or who watched television for more than two hours per day, were more likely to be overweight/obese (OR 1.26 and 1.55 respectively) (7). This discrepancy might be due to the small numbers of children in the present study who actually spent < 2 h per day in this activity.

The prevalence of obesity was lowest among those children who undertook physical activity (boys 11.8%, girls 8.8%), and highest among those who undertook none (boys 24.1%, girls 14.0%). Another paper reported similar results, with obesity more prevalent among those who undertook no physical activity (boys 21.7%, girls 9.8% vs boys 9.7%, girls 4.3% in those who did) (9). Other authors also report children who played sport to be less commonly obese than those who did not (10.3% vs 22.7%) (16). In the present work, children who spent more time in physical activity spent less time using monitors.

Overweight was less common among children who had been breast-fed until three months of age than in those not so fed. Similar results have been reported from Korea (18), and in an earlier systematic review of 300 articles from around the world (19). No significant differences were found for children who were/were not breast-fed until six months of age, perhaps due to the small sample size available for comparison.

Overweight was no less prevalent among children who ate breakfast (no result is available for obesity given the small numbers involved). In contrast, the AVENA study (20) reported that the habit of not taking breakfast was more common among children who were overweight or obese (normal weight children 5.9%, overweight children 11.6%, obese children 13.3%, $p < 0.05$). The Aladino 2011 study also reported similar results, with 3.1% of normal weight children and 4.2% of obese children not eating breakfast. Not taking breakfast might reflect a wider range of poor food habits, or might leave children hungry later in the day, leading to their eating with poorer control. The discrepancy with the present results is probably due to the low sample size available for those who took no breakfast (17).

CONCLUSIONS

The prevalence of obesity increased in Spain between 1993 and 2011, and was more common among boys than in girls over

this study period. Child overweight/obesity was found to be more prevalent in families in which the head of the family had a low level of education. Breastfeeding for three months appears to reduce the prevalence of overweight, while undertaking physical activity reduces the prevalence of obesity.

REFERENCES

1. Kipping RR, Jago R LD. Obesity in children. Part 1: Epidemiology, measurement, risk factors, and screening. *BMJ* 2008;15:33-1824.
2. Organization WH. Health Impact Assessment Methods and Strategies. Copenhagen: World Health Organization Regional Office for Europe. Cited 2016 Apr 1. Available from: <http://www.euro.who.int/healthimpact>
3. Kosti RI, Panagiotakos DB, Kosti RI, Panagiotakos DB. The epidemic of obesity in children and adolescents in the world. *Cent Eur J Public Health* 2015;14:151-9.
4. Tadesse Y, Derso T, Alene KA, Wassie MM. Prevalence and factors associated with overweight and obesity among private kindergarten school children in Bahirdar Town, Northwest Ethiopia: Cross-sectional study. *BMC Res Notes* 2017;10(1):22.
5. Jelastopulu E, Kallianezos P, Merikoulias G, Alexopoulos EC, Sapountzi-Krepia D. Prevalence and risk factors of excess weight in schoolchildren in West Greece. *Nurs Heal Sci* 2012;14:372-80.
6. Wu H, Zong X, Li H. The prevalence of overweight, obesity and stunting in school children aged 6-19 years in Beijing, China. *Ann Hum Biol* 2016;43(6).
7. Garby P, Clausson EK, Nyberg P, Jakobsson U. Overweight and television and computer habits in Swedish school-age children and adolescents: A cross-sectional study. *Nurs Heal Sci* 2014;16(2):143-8.
8. Thasanasuwan W SW. Low sleeping time, high TV viewing time, and physical inactivity in school are risk factors for obesity in pre-adolescent Thai children. *J Med Assoc Thai* 2016;99(3):314-21.
9. Bartrina J, Rodrigo C. Epidemiología y factores determinantes de la obesidad infantil y juvenil en España. *Rev Pediatr* 2005;VII:13-47.
10. Vázquez IA, Zapico RB, Díez JH, Rodríguez CF. Actividad física, ocio sedentario, falta de sueño y sobrepeso infantil. *Psicothema* 2008;20(4):516-20.
11. Carriere C, Cabaussel C, Bader C, Barberger-Gateau P, Barat P TH. Multi-disciplinary care management has a positive effect on paediatric obesity and social and individual factors are associated with better outcomes. *Acta Paediatr* 2016;105(11).
12. Ministerio de Sanidad Servicios Sociales e Igualdad. Encuesta Nacional de Salud 2011. Available from: <http://pestadistico.inteligenciadegestion.msssi.es/publicoSNS/comun/ArbolNodos.aspx>
13. Anton-Paduraru D, Teslariu O, Mocanu V. Influence of sleep on obesity in children. *Rev Med Chir Soc Med Nat Iasi* 2016;120(2):239-43.
14. Zapico RB, Vázquez IA, Díez JH, Rodríguez CF. La relación entre la falta de sueño, el ocio sedentario y el sobrepeso infantil. *Análisis Modif Conduct* 2006;32:391-401.
15. Vázquez IA, Zapico RB. La obesidad infantil como resultado de un estilo de vida obesogénico. *Endocrinol Nutr* 2007;54(10):530-4.
16. Serra L, Ribas L, Aranceta J. Obesidad infantil y juvenil en España. Resultados del estudio EnKid (1998-2000). *Med Clin* 2003;121(19):725-32.
17. Ortega RM, López AM, Perea JM, González L. Estudio Aladino, estudio de vigilancia del crecimiento, alimentación, actividad física, desarrollo infantil y obesidad en España. 2011. Available from: http://www.naos.aesan.msssi.gob.es/en/naos/ficheros/investigacion/Articulo_ALADINO.pdf
18. Park J, Chu HS, Chu SH, Jekal YS, Lee JY. The effect of predominant breast-feeding on the risk of obesity in Korean preschool children. *Nurs Heal Sci* 2015;17:77-83.
19. Lefebvre CM, John RM. The effect of breastfeeding on childhood overweight and obesity: A systematic review of the literature. *J Am Assoc Nurse Pract* 2014;26(7):386-401.
20. Wärnberg J, Ruiz JR, Ortega FB, Romeo J, González Gross M, Moreno LA, et al. Grupo AVENA. Estudio AVENA (Alimentación y valoración del estado nutricional en adolescentes). Available from: http://www.estudioavena.es/downloads/Publicaciones/Warnberg_J_2006PediatrInt.pdf