

## Nutrición Hospitalaria

Nutrición Hospitalaria

ISSN: 0212-1611

ISSN: 1699-5198

Grupo Arán

Guimarães da Silva, Danielle Cristina; Aparecida da Cunha, Kelly;  
Segheto, Wellington; Guimarães Reis, Vanessa; Araújo Coelho, France;  
O. Morais, Sílvia Helena; Pessoa, Milene Cristine; Zarbato Longo, Giana  
Behavioral patterns that increase or decrease risk of abdominal adiposity in adults  
Nutrición Hospitalaria, vol. 35, no. 1, 2018, January-February, pp. 90-97  
Grupo Arán

DOI: <https://doi.org/10.20960/nh.1228>

Available in: <https://www.redalyc.org/articulo.oa?id=309258226016>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative



## Trabajo Original

## Obesidad y síndrome metabólico

### Behavioral patterns that increase or decrease risk of abdominal adiposity in adults

### *Patrones de comportamiento que aumentan o disminuyen el riesgo de adiposidad abdominal en adultos*

Danielle Cristina Guimarães da Silva<sup>1</sup>, Kelly Aparecida da Cunha<sup>1</sup>, Wellington Segheto<sup>1</sup>, Vanessa Guimarães Reis<sup>1</sup>, France Araújo Coelho<sup>1</sup>, Sílvia Helena O. Morais<sup>1</sup>, Milene Cristine Pessoa<sup>2</sup> and Giana Zerbato Longo<sup>1</sup>

<sup>1</sup>Department of Nutrition and Health. Federal University of Viçosa. Viçosa, Minas Gerais, Brazil. <sup>2</sup>Department of Nursing. Federal University of Minas Gerais, Belo Horizonte, Minas Gerais, Brazil

### Abstract

**Introduction:** The identification of risk or protective behavioral patterns associated with abdominal adiposity may aid in prevention and health promotion measures.

**Objective:** To identify and to associate behavioral patterns of risk and protection to abdominal adiposity in adults in a Brazilian city.

**Material and methods:** A population-based cross-sectional study was carried out in Viçosa, Brazil, with 1,226 adults of both sexes. Information on social-demographic characteristics, food intake, level of physical activity, alcohol consumption and smoking were collected by using a questionnaire. The anthropometric measurement of waist circumference and anthropometric indices waist/hip ratio and waist/height ratio were indicators of abdominal adiposity. To identify behavioral patterns, exploratory factor analysis was applied for the variables considered as risk or protective factors. The association of the identified patterns with abdominal adiposity was estimated by multiple linear regression, adjusted for gender, age and social economical class.

**Results:** Two patterns were obtained, "healthy" and "risk". The "healthy" pattern, comprised of the clustering of the variables food consumption, fruits, fresh fruit juices, raw and cooked vegetables and the appropriate level of physical activity, was negatively associated with abdominal adiposity identified by waist circumference ( $p = 0.048$ ), waist/hip ( $p = 0.013$ ) and waist/height ( $p = 0.018$ ) indices. The "risk" pattern, composed of smoking, alcohol beverage abuse and habit of consuming visible fat in fat-rich red meat or poultry skin, was positively associated with abdominal adiposity identified by waist circumference ( $p = 0.002$ ) and waist/hip ( $p = 0.007$ ) and waist/height indices ( $p = 0.006$ ).

**Conclusions:** Two behavioral patterns were identified, a risk pattern and a protective pattern for abdominal adiposity in the assessed population. The study shows the importance of conducting clustering of multiple risk and protective factors to better explain the health conditions of a group.

#### Key words:

Abdominal obesity.  
Factorial analysis.  
Protective factors.  
Risk factors.

### Resumen

**Introducción:** la identificación de los riesgos o los patrones de comportamiento de protección asociados con la adiposidad abdominal puede ayudar en las medidas de prevención y promoción de la salud.

**Objetivo:** identificar y establecer la asociación entre los patrones de comportamiento de riesgo y de protección y la adiposidad abdominal en adultos en una ciudad brasileña.

**Material y métodos:** se llevó a cabo un estudio transversal basado en la población en Viçosa, Brasil, con 1.226 adultos de ambos sexos. Se recogió información sobre las características sociodemográficas, la ingesta de alimentos, el nivel de actividad física, el consumo de bebidas alcohólicas y el hábito tabáquico mediante un cuestionario. La medición antropométrica de la circunferencia de la cintura y de los índices antropométricos cintura/cadera y cintura/altura fueron los indicadores de adiposidad abdominal. Para identificar los patrones de comportamiento, se aplicó un análisis factorial exploratorio de las variables de riesgo o factores de protección considerados. La asociación de los patrones identificados con la adiposidad abdominal se estimó por regresión lineal múltiple, ajustada por género, edad y nivel socioeconómico.

**Resultados:** se establecieron dos patrones, "sano" y "riesgo". El patrón "sano", compuesto por la agrupación de las variables consumo de alimentos, frutas, zumos de fruta fresca, verdura cruda y cocida y el nivel apropiado de actividad física, se asoció negativamente con la adiposidad abdominal identificada por la circunferencia de la cintura ( $p = 0,048$ ) y los índices cintura/cadera ( $p = 0,013$ ) y cintura/altura ( $p = 0,018$ ). El patrón de "riesgo", compuesto por hábito tabáquico, abuso de alcohol y consumo de grasa visible en carnes rojas ricas en grasa o piel de las aves, se asoció positivamente con la adiposidad abdominal identificada por la circunferencia de la cintura ( $p = 0,002$ ) y las ratios cintura/cadera ( $p = 0,007$ ) y cintura/altura ( $p = 0,006$ ).

**Conclusiones:** fueron identificados dos patrones de comportamiento, el patrón de riesgo y el patrón de protección, relacionados con la adiposidad abdominal en la población estudiada. El estudio muestra la importancia de agrupar múltiples factores de riesgo y de protección para explicar mejor las condiciones de salud de un grupo.

#### Palabras clave:

Obesidad abdominal.  
Análisis factorial.  
Factores de protección. Factores de riesgo.

Received: 26/04/2017 • Accepted: 17/08/2017

Silva DCG, Cunha KA, Segheto W, Reis VG, Coelho FA, Morais SHO, Pessoa MC, Longo GZ. Behavioral patterns that increase or decrease risk of abdominal adiposity in adults. Nutr Hosp 2018;35:90-97

DOI: <http://dx.doi.org/10.20960/nh.1228>

#### Correspondence:

Danielle Cristina Guimarães da Silva. Department of Nutrition and Health. Federal University of Viçosa. Av. PH Rolfs, s/n. 36571-000 Viçosa, Minas Gerais, Brazil  
e-mail: [daniellenut@hotmail.com](mailto:daniellenut@hotmail.com)

## INTRODUCTION

Obesity is characterized by excessive accumulation of body fat (1) and represents a serious public health issue due to its trend of increasing prevalence and the impact it has on society (2). Body mass index (BMI) has been the most frequently used method to define excess body weight; however, it is unable to differentiate between lean mass and fat mass (3) and so central obesity measures have been recently recommended due to a strong association with morbidity and mortality (4-6).

Abdominal adiposity is defined as the accumulation of intra-abdominal fat and it has been considered as an important risk factor for major chronic non-communicable diseases (NCDs) (7). Although there are sophisticated methods for evaluating abdominal fat, from an epidemiological point of view, the anthropometric indicators are considered for being used in population studies due to their low cost and ease of use (8). Waist circumference, waist-to-hip ratio and waist-to-height ratio are alternative proposals for the detection of abdominal obesity and have been strongly associated with cardiometabolic risk (5). However, even though there is consensus about the importance of measuring such anthropometric measures, epidemiological studies have not been able to demonstrate the measure or index that shows a better association with unhealthy behavioral patterns (7,9,10), making it necessary to carry out further studies that compare these indices to verify which one is more associated with behavioral patterns.

Studies in some countries have shown that abdominal adiposity in adults has increased over the past few years (11-13). This increase is attributed to changes in eating patterns and physical activity, increased stress and endocrine disruptors (14), and socioeconomic factors such as education level, income and occupation (8). Adequate dietary intake of fruits and vegetables and level of leisure physical activity, no smoking and no alcohol consumption has been associated to a lower risk of abdominal obesity (15-17).

The exploratory factor analysis has been increasingly used since this technique allows to identify the clustering of variables, showing the interrelations among them (18,19).

However, in nutritional epidemiology, most of the published papers that used this technique have focused on the clustering of foods/nutrients in the definition of dietary patterns (20,21), rather than on identifying abdominal adiposity risk or protection patterns by clustering behavioral and dietary variables. In addition, some unhealthy behaviors may interact, producing an even greater risk than the individual ones (22).

However, studies that make associations between abdominal adiposity risk or protection factors are rare and inconclusive (23,24). Therefore, it is of great importance to identify behavioral risk and protection patterns in predicting abdominal obesity so that intervention strategies to prevent and to control this type of worsening may be better targeted, with actions that promote health.

## MATERIAL AND METHODS

This work is an epidemiological cross-sectional designed study conducted in the urban area of the city of Viçosa, Brazil, from September 2012 to April 2014. It was carried out by the health

and nutrition study group of Viçosa (ESA/Viçosa). Data collection involved two steps: visits to the subjects' residences with the application of a structured questionnaire and anthropometric measurements on the university facilities. The fieldwork flow and all instruments used in the data collection were previously calibrated and tested in a pilot study (25,26).

The study sample was made up of adults of both sexes; 50.8% were women, from 20 to 59 years old. For sample calculation, the formula for prevalence estimates was used, considering the total number of individuals between 20 and 59 years old living in the urban area of Viçosa, totaling 43,431 people (27), an estimated prevalence of 50% (since this study is part of a thematic research project where other outcomes were analyzed), confidence level of 95%, sampling error of 4.1 percentage points and estimated *deff* (design effect sampling per cluster) of 1.55. An increase of 20% occurred for losses and refusals, and 10% for control of confusion factors. By using Epi-Info, version 3.5.2® (28), calculations evidenced a minimal sample size (n) of 1,137 participants.

The following were considered as exclusion criteria: pregnant women, postpartum women, bedridden individuals or those unable to be measured, and individuals with cognitive/intellectual difficulty or who were not able to answer the questionnaire. Individuals who reported having a previous diagnosis of diabetes or cardiovascular disease were not excluded from this study because the prevalence of these outcomes was considered to be low.

Two-stage cluster sampling was carried out. First, census sectors were randomly selected. After that, a block and a corner were selected, from which fieldwork was started in a clockwise direction. Thirty census sectors were randomly selected from 99 census sectors existing in Viçosa by simple random sampling with no replacement.

The information collected from the questionnaire consisted of social-demographic characteristics, food intake, physical activity level, alcohol consumption and smoking. The questions used in this survey were based on the Surveillance System of Risk and Protective Factors for Non Communicable Diseases through Telephone Interviews (*Vigilância de Fatores de Risco e Proteção para Doenças Crônicas por Inquérito Telefônico*, VIGITEL) (26), held annually in Brazil. The anthropometric measurements used in this study were performed by measuring the height and the circumferences of waist and hip (29).

The social demographic variables were as follows: age (completed years), education degree in completed years of study (defined as 0-3, 4-7, and  $\geq 8$  years of schooling) and socioeconomic status, determined by tools of the Brazilian Association of Research Companies (30) and classified into A and B, C and D and E.

The risk behavior indicators for abdominal adiposity included in this study were: smoking, abusive consumption of alcoholic beverages, excessive screen time, consumption of sugar-sweetened beverages at least five times a week, considered regular, and habit of consuming saturated fat source food, such as whole milk, red meat with visible fat and poultry skin.

Smoking was categorized into smokers, former smokers and non-smokers, regardless of the frequency and intensity of using tobacco (31). The abusive consumption of alcohol was considered

as present when the intake of at least five shots in a single sitting occurred, based on the reference of the past 30 days for men and at least four for women. The standard was considered as the consumption of half a bottle or a can of beer, a glass of wine or a shot of distilled drink (32). The screen time was evaluated by summing the time the individual spent watching television or using the computer on weekdays and weekends. The cut-off point, indicator of a sedentary behavior, was the sum of the time longer than or equal to four hours a day watching TV or using the computer (33).

Among the behavioral indicators related to protective factors for abdominal adiposity, the following were included: the appropriate level of physical activity in leisure time and regular consumption of fruits, fresh fruit juices, raw salads, cooked vegetables and beans.

The level of physical activity during leisure time was assessed by the specific section of the long version of the International Physical Activity Questionnaire. The cut-off for physical activity practice time longer than or equal to 150 minutes per week to classify individuals as physically active (34) was used. Dietary intake associated with protection for abdominal adiposity was regarded from the report of intake of fruits, fresh fruit juices, vegetables and beans for at least five days a week, which was considered as regular consumption.

Abdominal adiposity was evaluated by means of anthropometric parameters of waist circumference (WC), waist-to-hip ratio (WHR) and waist-to-height ratio (WHER). Waist circumference was measured by using an nonelastic 2-m long measuring tape (Sanny®, São Paulo, SP, Brazil), the measurement being made at the mid-point between the iliac crest and the last rib. Hip circumference was measured at the most protuberant area of the buttocks, and height was measured by a fixed rod stadiometer (Welmy®, in wall, Santa Bárbara D'Oeste, SP, Brazil), with a length of 2.5 m and resolution of 0.1 cm.

The cut-offs used were those proposed by Grundy et al. (35) to determine the prevalence of abdominal adiposity based on the evaluation of waist circumference, which considers that men with a WC  $\geq 90$  cm and women with a WC  $\geq 80$  cm have central obesity. WHR was determined by the following equation:  $[WHR = WC / \text{hip circumference (cm)}]$ , and the cut-offs used were men with a ratio  $\geq 1$  and women  $\geq 0.85$  were considered as having central obesity (36). The cut-offs proposed by Ashwell and Hsieh (37), which consider that values  $\geq 0.5$  for men and women indicate abdominal adiposity, were used to determine adiposity prevalence from the evaluation of the waist-to-height ratio. Measurements were performed in triplicate by a single evaluator, using the average of the measurements in the analysis.

For the identification of behavioral patterns, the methodology of exploratory factor analysis was applied to the answers obtained from the structured questionnaire. Prior to the calculation of the principal component analysis (PCA), the coefficient of Kaiser-Meyer-Olkin (KMO) and Bartlett's sphericity test were estimated to assess the quality of the correlations among the variables. After that, the factors were extracted by the principal components analysis method, rotated by an orthogonal transformation (varimax), retaining those with eigenvalues  $> 1.3$ , defined according to the scree plot graph of the variance for the number of components.

The exploratory factor structure was obtained from the indicators that presented factor loadings larger than 0.3, a score was determined for each behavioral pattern using principal components factor analysis, and the patterns were named according to the retained indicators.

The association among the patterns and abdominal obesity anthropometric indicators was analyzed using linear regression models. The WC, WHR and WHER variables presented a symmetric distribution ( $p > 0.05$ ; Shapiro-Wilk test). The main independent variable was the score of the patterns, those which had a  $p < 0.20$  in the linear regression analysis. The models were adjusted by confounding variables such as gender, age and social economical class. A significance level of 5% was considered in the study.

Quality control of this study was performed by applying, at random, questions of the questionnaire to 10% of the sample by phone calls and double data input. After checking data consistency, analyses were performed in the STATA statistical package, version 13.1, by taking into account the effect of sample design by the "svy" command group, which considered the complex sample design (sampling by clustering in two steps and prior stratification through census sectors). Sample weights, considering gender, age and education, were calculated in order to equalize differences in the social-demographic composition of the sample in relation to the composition of the adult population of the city, according to the 2010 census distribution (27).

The project was submitted and approved by the Ethics Committee, under protocol number 008/12. Participants of the study were requested to sign the terms of free consent, which was explained to them before data collection.

## RESULTS

In this study 1,226 interviews were conducted, of which 50.8% were answered by women, 32.7% of the interviewees were 20 to 29 years old, 71.7% reported to have been at school for more than eight years and 64.6% were in middle C social economical class.

The frequencies of variables used as of risk or protection indicators for abdominal adiposity in the principal component analysis are presented in table I.

For protection indicators, regular consumption of raw salads (53.5%) and regular consumption of beans (86.2%) were reported by the interviewees.

In relation to risk indicators, the habit of consuming whole milk and red meat with visible fat and/or poultry skin was reported by 59.0% and 55.3% of the respondents, respectively.

The values presented in the tests for evaluation of the correlation between the indicators of protection or risk of abdominal adiposity and the adjustment for using factor analysis to identify behavioral patterns were satisfactory for the PCA (KMO = 0.60 and Bartlett's sphericity = 0.000). By means of the exploratory factor analysis, five components were found, two of which had eigenvalues larger than 1.3, explaining 23.78% of the components variation. After orthogonal rotation, two components remained in the correlation matrix, representing a behavioral pattern of risk and one of pro-

tection. Of the 12 variables regarded as risk or protection factors for the tested abdominal obesity, those considered to be valid to remain in each component were the indicators with saturation higher than 0.3 (indicated in italics), shown in table II.

The means of waist circumference, waist-to-hip ratio and waist-to-height ratio among men were 87.4 cm (95% CI 85.3-89.5), 0.91 (95% CI 0.89-0.92) and 0.51 (95% CI 0.50-0.53), respectively. Abdominal adiposity evaluated by waist circumference was present in 12.8% (95% CI 9.4-17.1) of the male participants, by means of the waist-to-hip ratio in 25.3% (95% CI 21.0-30.2) and waist-to-height ratio in 44.7% (95% CI 39.4-50.1).

For women, the average of waist circumference was 80.6 cm (95% CI 78.5-82.6), the mean of waist-to-hip ratio was 0.82 (95% CI 0.85-0.88) and the average of waist-to-height ratio was 0.55 (95% CI 0.35-0.56). Abdominal adiposity evaluated by measurement of waist circumference was present in 23.9% of women (95% CI 19.6-28.7), by waist-to-hip ratio in 55.3% (95% CI 49.8-60.5) and by waist-to-height ratio in 29.1% (95% CI 24.6-34.0).

The first behavioral pattern found in the study was termed "Healthy Pattern". It explained 12.58% of the variance and

included the regular consumption of fruits, fresh fruit juices, raw and cooked vegetables and the appropriate level of physical activity.

The second behavioral pattern was characterized by the habit of smoking, and by the consumption of alcohol and fat-rich meats (red meat with visible fat and/or poultry skin). This pattern was identified as "Risk Pattern" and presented a variance of 11.20%.

**Table II. Rotated factor loadings for the first two factors of principal component analysis of the population, Viçosa, Brazil, 2012-2014**

Indicators	"Pattern 1" Healthy Pattern	"Pattern 2" Risk Pattern	Communality
Regular consumption of fruits <sup>†</sup>	<i>0.39</i>	-0.27	0.42
Regular consumption of fresh fruit juice <sup>†</sup>	<i>0.35</i>	-0.15	0.48
Regular consumption of raw salads <sup>†</sup>	<i>0.71</i>	-0.01	0.55
Regular consumption of cooked vegetables <sup>†</sup>	<i>0.64</i>	-0.10	0.46
Regular consumption of beans <sup>†</sup>	0.23	0.02	0.41
Regular consumption of sugar-sweetened beverages <sup>†</sup>	0.01	0.21	0.51
Habit of consuming whole milk	-0.01	-0.06	0.61
Habit of consuming fat-rich meat <sup>‡</sup>	-0.16	<i>0.57</i>	0.45
Smoking habit	-0.11	<i>0.33</i>	0.60
Over consumption of alcohol drinks <sup>§</sup>	0.01	<i>0.77</i>	0.63
Physically active <sup>  </sup>	<i>0.43</i>	0.28	0.49
Excessive screen time <sup>¶</sup>	-0.02	-0.03	0.77
Eigenvalues	1.51	1.34	
Percentage of explained variance	12.58	11.20	
Percentage of explained cumulative variance	12.58	23.78	

Observation: indicators with factor loading larger than or equal to 0.3 are in italics. <sup>†</sup>Consumption of fruit, fresh fruit juice, raw salads and cooked vegetables, beans and sugar-sweetened beverages five or more days a week. <sup>‡</sup>Consumption of red meat without removing visible fat or poultry skin from the food. <sup>§</sup>Consumption of > 5 shots (man) and > 4 shots (woman) in ≥ 1 day in the last 30 days. <sup>||</sup>Time of physical activity practice longer than or equal to 150 minutes per week. <sup>¶</sup>Time spent in front of TV or computer longer than or equal to four hours a day.

**Table I. Frequency of risk and protective factor for abdominal adiposity in the population, Viçosa, Brazil, 2012-2014**

Indicator	Proportion* (%)	Confidence interval (95% CI)
Regular consumption of fruits <sup>†</sup>	34.8	(30.3-39.5)
Regular consumption of fresh fruit juice <sup>†</sup>	16.6	(13.1 -20.8)
Regular consumption of raw salads <sup>†</sup>	52.5	(44.1-60.7)
Regular consumption of cooked vegetables <sup>†</sup>	30.0	(26.7-33.5)
Regular consumption of beans <sup>†</sup>	86.7	(83.9-89.5)
Regular consumption of sugar-sweetened beverages <sup>†</sup>	27.6	(23.0-32.6)
Habit of consuming whole milk	59.3	(54.5-63.9)
Habit of consuming fat-rich meat <sup>‡</sup>	57.4	(52.2-62.5)
Smoking habit	17.9	(14.1-22.4)
Over consumption of alcohol <sup>§</sup>	42.6	(37.6-47.7)
Physically active <sup>  </sup>	20.9	(15.4-27.8)
Excessive screen time <sup>¶</sup>	17.8	(13.8-22.5)

\*Proportion weighed by gender, age and schooling. <sup>†</sup>Consumption of fruit, fresh fruit juice, raw salads and cooked vegetables, beans and sugar-sweetened beverages five or more days a week. <sup>‡</sup>Consumption of red meat without removing visible fat or poultry skin from the food. <sup>§</sup>Consumption of > 5 shots (man) and > 4 shots (woman) in ≥ 1 day in the last 30 days. <sup>||</sup>Time of physical activity practice longer than or equal to 150 minutes per week. <sup>¶</sup>Time spent in front of TV or computer longer than or equal to four hours a day.



Table III shows the association between the two behavioral patterns found in abdominal adiposity, determined by waist circumference and by the indices of waist-to-hip ratio and waist-to-height ratio.

Negative associations among the “healthy” protection pattern and waist circumference, waist-to-hip ratio and waist-to-height ratio were found, showing that the increment in the consumption of fruits, fresh fruit juices, raw or cooked vegetables and the appropriate level of physical activity were associated with the reduction in abdominal adiposity in this population. Contrary to that, the pattern termed “risk” was positively associated with abdominal adiposity. In addition, the multiple linear regression analysis showed that the association between risk and protection patterns and the parameters of abdominal adiposity were better explained by the waist-to-height ratio and waist circumference since they presented 36.5% and 32.9% of the variability explained for the “healthy” pattern and 36.5% and 36.9% of the variability explained for the “risk” pattern.

## DISCUSSION

This study investigated the clustering of risk factors and the association between risk and protection behavioral patterns with abdominal adiposity indicators among adult individuals of Viçosa, using the method of exploratory factor analysis. Two patterns were established, one of which is considered as a protection factor for abdominal adiposity, where variables of healthy food consumption, such as raw salads or cooked vegetables, fruits, fresh fruit juices and physical activity, were included. The second pattern identified was considered as risky because it grouped variables considered to be risk factors such as smoking, abusive alcohol consumption and the habit of consuming fat-rich meats.

Physical inactivity, smoking, alcohol drinking and inappropriate diet, regarded as the main risk factors, have been analyzed each in several studies. However, this way of analyzing may be inefficient, since, according to Hofstetter et al. (6), risk behaviors do not occur in isolation. Furthermore, populations simultaneously presenting risk behavior factors are more likely to present major non-communicable chronic diseases than individuals with only one risk factor (38). Therefore, using statistical techniques to reduce variables into clusters with similar profiles is recom-

mended to assist in obtaining knowledge of possible associations of risk behaviors or health protection among them.

Once the analyses were adjusted, the “healthy” pattern was found to be negatively associated with abdominal adiposity. Similar to what was found in this study, McNaughton et al. (39), when using the exploratory factor analysis, found that the dietary pattern characterized by high consumption of fruits and vegetables was inversely associated with body mass index, blood pressure and waist circumference. Neumann et al. (40), when using the same statistical technique, found that the termed “modern” dietary pattern, made up of fat-free dairy products, seafood, fresh fruit juices and fruit was inversely associated with risk factors for cardiovascular diseases, and the waist-to-hip ratio measure was among these evaluated variables. Ribeiro et al. (41), when describing the frequency of fruit and vegetable consumption of adults living in the city of São Paulo, São Paulo state, found a positive correlation between the consumption of these foods and physical activity during leisure time.

Our findings can be explained by the positive and healthy actions of eating habits and lifestyle. The effects of fruits and vegetables on general and central obesity are believed to relate to their soluble fiber content, which is associated with increased satiety, delayed gastric emptying and enhanced insulin sensitivity (42). In relation to the practice of physical activity, the energy expenditure independent of intensity reduces abdominal fat (43).

When evaluating the aggregation of behavioral patterns of British university students, identified by means of cluster analysis, Dodd et al. (44) identified three subgroups, characterized by high prevalence of physical inactivity (70%), inadequacy in the regular consumption of fruits and vegetables (66%) and abusive alcohol consumption (56%). By using the same technique, Costa et al. (22) found the simultaneity of health protective behaviors, such as the occurrence of a proper consumption of fruit and the practice of physical activity during leisure time in 35.5% of men and 33.3% of adult women living in the city of Florianópolis, state of Santa Catarina, Brazil.

The data previously shown illustrate the limitation regarding the comparison of this study with other studies due to variations in the methodology used for determining a standard or a variable clustering, since the identification of standards involving behavioral or anthropometric variables has been frequently carried out by the statistical technique of clustering, which, according to Hofstetter et al. (6), place individuals with similar interrelationships in the same clusters.

**Table III.** Associations of risk and protection behavioral patterns with abdominal adiposity, Viçosa, Brazil, 2012-2014

Indicators	Waist circumference (cm)			Waist/hip ratio			Waist/height ratio		
	Regression coefficient $\beta$	Value $p^*$	R <sup>2</sup>	Regression coefficient $\beta$	Value $p^*$	R <sup>2</sup>	Regression coefficient $\beta$	Value $p^*$	R <sup>2</sup>
Pattern 1 “Healthy”	-0.991	0.048	32.9	-0.003	0.013	25.0	-0.007	0.018	36.5
Pattern 2 “Risk”	1.219	0.002	36.9	0.007	0.003	25.6	0.007	0.006	36.5

\* $p \leq 0.05$ . Models adjusted according to gender, age and social economical class.

The second pattern found in the study, termed "risk", was positively associated with abdominal adiposity. Schuit et al. (45), in a cross-sectional study with European adult subjects, showed that the associations between the consumption of alcohol and smoking are strong especially among individuals with low education degree and income. Furthermore, the influence of smoking and consumption of alcoholic drinks on abdominal fat has been shown in other studies (46,47) by measurement of the waist circumference as well as waist-to-hip ratio.

A transversal study carried out by Faria et al. (47) with men living in Cuiabá, state of Mato Grosso do Sul, Brazil, found that the means of waist-to-hip ratio indices were higher among smokers and the means of waist circumference and waist-to-height ratio were higher among those who reported higher consumption of alcoholic drinks. Studies have shown a higher prevalence of abdominal obesity in former smokers, a result from the increment in the caloric intake and a reduction in the basal metabolic rate after quitting the addiction (48). Despite being evident, the literature has not been able to determine a relationship between cause and effect in relation to the association between consumption of alcohol drinks and abdominal fat due to the large methodological variation in the frequency analysis and the amount consumed (8).

Smoking, abusive alcohol consumption and the habit of consuming fatty meats (red meat with visible fat and/or chicken with skin) are considered as important risk factors for chronic non-communicable diseases and have been associated with abdominal adiposity. The relation between cigarette smoking and body fat distribution is complex and incompletely understood. Biological mechanisms involving higher levels of cortisol increase lipogenesis, differentiation of adipocytes and deposition of abdominal fat (47). As for the consumption of alcohol, its metabolism may trigger endocrine changes, such as increased cortisol, predisposing to changes in fat distribution. The habit of consuming fat-rich meats is associated with abdominal adiposity because this food group provides excess calories to individuals.

The combination of risk factors such as smoking, alcoholism and inadequate food intake has been investigated in studies in Brazil (22,31) and worldwide (44,45). The prevalence of those three risk factors, evaluated by the cluster statistical technique in a Brazilian study, showed that 4.9% of men and 1.9% of women had the occurrence of those factors at the same time (20). The same methodological work proposal in European countries presented prevalence data higher than these, which were 15.3% for men and 14.4% for women.

The identification of risk or protection behavioral patterns for non-communicable chronic diseases was carried out by using the PCA technique in a sample with 108,706 Brazilian adults, participants of the VIGITEL. That investigation showed results similar to those of this present study, where it was found that the behavioral pattern considered risk was defined by smoking, excessive alcohol consumption and regular intake of meat with visible fat (31). Therefore, strategies to control the increase in the prevalence of such risk factors in the population may be directed to these three factors, which have a tendency to aggregate.

The population-based transversal study with 2,732 adults, conducted in Pelotas, state of Rio Grande do Sul, Brazil, to eval-

uate the prevalence of the combination of behavioral risk factors, showed that the combination of physical inactivity factors and regular consumption of visible meat fat was present in 18.2% of the men and 17.0% of the women in the study. Vilela et al. (21), when evaluating the dietary pattern of adults in Cuiabá, state of Mato Grosso do Sul, Brazil, found a positive association between "Western" eating pattern, characterized by high intake of pasta, fatty meats, sugar-sweetened beverages and sweets, and the increase in the waist circumference and waist-to-hip ratio among women.

The parameters of abdominal adiposity, waist-to-height ratio and waist circumference presented a higher capacity for explaining the behavioral patterns than did waist-to-hip ratio, this is observed since of different found regression coefficients, that can show the superiority tool for discriminating the behavioral patterns. Waist-to-height ratio has a direct regulation with growth and waist circumference, so it minimizes erroneous assessment of health risk in individuals with different heights, which is the greatest advantage of this measure. In addition, this indicator has been considered as a good marker of abdominal obesity related to cardiovascular risk factors. Waist circumference is already recognized as an important and simple indicator of central adiposity and risk for chronic diseases (35).

This is the first population-based study conducted in Viçosa and one of the few works that identifies and explores risk and protection factors by using the statistical technique of principal components and makes associations with abdominal adiposity. Lam et al. (3) mentioned that the measures of central adiposity are more closely associated with cardiovascular risk factors and metabolic diseases than BMI. Furthermore, the body mass index does not consider body fat distribution, which is a limitation since there are suggestions that the metabolic complications of obesity are more closely related to visceral adiposity than to overall adiposity. However, some limitations should be considered. It is noteworthy that this is a cross-sectional study, which prevents the establishment of temporality of associations. Self-reports of risk behaviors tend to be underestimated because of being socially undesirable. In addition, menopausal women were not excluded because they presented lower sample representativity.

---

## CONCLUSION

This study may have important implications for public policies because it was found that by clustering risk or protection factors, health conditions of a population can be better explained. In addition, multiple interventions may be more effective and efficient in promoting health.

---

## AUTHOR'S CONTRIBUTION

Danielle Cristina Guimarães da Silva, Kelly Aparecida da Cunha, Wellington Segheto, Vanessa Guimarães Reis, France Araújo Coelho and Sílvia Helena O. Morais, post-graduate students in Science

of Nutrition, contributed to the data collection and analyses of this study, wrote the initial draft of the manuscript and assembled the final version.

Milene Cristine Pessoa, professor, contributed to the study design, data analyses and participated in the approval of the final version of the manuscript.

Giana Zarbato Longo, professor, supervised this study, contributed to the study design, and developed analyses and data analyses.

## ACKNOWLEDGEMENTS

The authors thank all the volunteers who contributed to the study and the National Council for Scientific and Technological Development (CNPq process no. 481418/2011-3) for financially funding the project, and also the Foundations for Supporting Research in the states of Minas Gerais (FAPEMIG process no. APQ-00296-12).

## FINANCIAL SUPPORT

National Council for Scientific and Technological Development (CNPq process no. 481418/2011-3) and Foundations for Supporting Research in the states of Minas Gerais (FAPEMIG process no. APQ-00296-12).

## REFERENCES

- Friedman JM. Obesity: Causes and control of excess body fat. *Nature* 2009;459(7245):340-2.
- World Health Organization. Obesity and overweight. Fact Sheet Number 311. Geneva (CH): WHO (2012). Available from: <http://www.who.int/mediacentre/factsheets/fs311/en/index.html>
- Lam BCC, Koh GCH, Chen C, Wong MTK, Fallows SJ. Comparison of body mass index (BMI), body adiposity index (BAI), waist circumference (WC), waist-to-hip ratio (WHR) and waist-to-height ratio (WHtR) as predictors of cardiovascular disease risk factors in an adult population in Singapore. *PLoS ONE* 2015;10(4):e0122985.
- Lanas F, Serón P, Muñoz S, Margozzini P, Puig T. Central obesity measurements better identified risk factors for coronary heart disease risk in the Chilean National Health Survey (2009-2010). *J Clin Epidemiol* 2017;86:111-6.
- Lim LL, Seubsman S, Sleight A, Bain C. Validity of self-reported abdominal obesity in Thai adults: A comparison of waist circumference, waist-to-hip ratio and waist-to-stature ratio. *Nutr Metab Cardiovasc Dis* 2012;22(1):42-9.
- Hofstetter H, Dusseldorp E, Van Empelen P, Paulussen TW. A primer on the use of cluster analysis or factor analysis to assess co-occurrence of risk behaviors. *Prev Med* 2014;67:141-6.
- Vasques ACJ, Priori SE, Rosado LEFPL, Franceschini SCC. The use of anthropometric measures to assess visceral fat accumulation. *Rev Nutr* 2010;23(1):107-18.
- Pinho CPS, Diniz AS, Arruda IKG, Batista Filho M, Coelho PC, Sequeira LAS, et al. Prevalence of abdominal obesity and associated factors among individuals 25 to 59 years of age in Pernambuco State, Brazil. *Cad. Saúde Pública* 2013;29(2):313-24.
- Park SH, Choi SJ, Lee KS, Park HY. Waist circumference and waist-to-height ratio as predictors of cardiovascular disease risk in Korean adults. *Circ J* 2009;73(9):1643-50.
- Bevilacqua MR, Gimeno SGA. Abdominal obesity in Japanese-Brazilians: Which measure is best for predicting all-cause and cardiovascular mortality? *Cad. Saúde Pública* 2011;27(10):1986-96.
- Ford ES, Maynard LM, Li C. Trends in mean waist circumference and abdominal obesity among US adults, 1999-2012. *JAMA* 2014;312(11):1151-3.
- Walls HL, Stevenson CE, Mannan HR, Abdullah A, Reid CM, McNeil JJ, et al. Comparing trends in BMI and waist circumference. *Obesity* 2011;19(1):216-9.
- Linhares RS, Horta BL, Gigante DP, Dias-da-Costa JS, Olinto MT. Distribution of general and abdominal obesity in adults in a city in southern Brazil. *Cad. Saúde Pública* 2012;28(3):438-47.
- Friedman DS, Ford ES. Are the recent secular increases in the waist circumference of adults independent of changes in BMI? *Am J Clin Nutr* 2015;101(3):425-31.
- Castanho GKF, Marsola FC, Mclellan KCP, Nicola M, Moreto F, Burini RC. Consumption of fruit and vegetables associated with the metabolic syndrome and its components in an adult population sample. *Cien Saúde Colet* 2013;18(2):385-92.
- Chau JY, Chau JY, Grunseit AC, Stamatakis E, Brown WJ, Matthews CE, et al. Daily sitting time and all-cause mortality: A meta-analysis. *PLoS One* 2013;13(8):e80000.
- Mielke GI, Hallal PC, Malta DC, Lee IM. Time trends of physical activity and television viewing time in Brazil: 2006-2012. *Int J Behav Nutr Phys Act* 2014;15(11):101.
- Marchioni DML, Latorre MRDO, Eluf-Neto, Wünsch-Filho V, Fisberg RM. Identification of dietary patterns using factor analysis in an epidemiological study in São Paulo. *São Paulo Med J* 2005;123(3):124-7.
- Sato RC, Zouain DM. Factor analysis for the adoption of nuclear technology in diagnosis and treatment of chronic diseases. *Einstein* 2012;10(1):62-6.
- Olinto MTA, Willett WC, Gigante DP, Victora CG. Sociodemographic and lifestyle characteristic in relation to dietary patterns among young Brazilian adults. *Public Health Nutr* 2011;14(1):150-9.
- Vilela AAF, Sichieri R, Pereira RA, Cunha DB, Rodrigues PRM, Gonçalves-Silva RMV, et al. Dietary patterns associated with anthropometric indicators of abdominal fat in adults. *Cad. Saúde Pública* 2014;30(3):502-10.
- Costa FF, Benedet J, Leal DB, Altenburg de Assis MA. Clustering of risk factors for non-communicable diseases in adults from Florianópolis, SC. *Rev Bras Epidemiol* 2013;16(2):398-408.
- Romaguera D, Ångquist L, Du H, Jakobsen MU, Forouhi NG, Halkjær J, et al. Food composition of the diet in relation to changes in waist circumference adjusted for body mass index. *PLoS One* 2011;6(8):e23384.
- Funtikova AN, Subirana I, Gómez SF, Fitó M, Elosua R, Benítez-Arciniegua AA, et al. Soft drink consumption is positively associated with increased waist circumference and 10-year incidence of abdominal obesity in Spanish adults. *J Nutr* 2015;145(2):328-34.
- Segheto W, Silva DCG, Coelho FA, Reis VG, Morais SHO, Marins JCB, et al. Body adiposity index and associated factors in adults: Method and logistics of a population-based study. *Nutr Hosp* 2015;32(1):101-9.
- Brazil Ministry of Health. VIGITEL Brasil 201. Vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília: Ministério da Saúde; 2012.
- Brazilian Institute of Geography and Statistics. Censo Demográfico 2010. Rio de Janeiro: Brazilian Institute of Geography and Statistics; 2010.
- Dean AG, Dean JA, Colombari, Brendel KA, Smith DC, Burton AH, et al. Epi Info, version 6: A word processing, database, and statistics program for epidemiology on microcomputers. Atlanta, Georgia: Centers for Disease Control and Prevention; 1994.
- Stewart A, Marfell-Jones M, Olds T, Carter L. International standards for anthropometric assessment. New Zealand: Lower Hutt; 2011.
- Brazilian Association of Research Companies (ABEP). Critério de classificação econômica Brasil, 2011. Available from: [www.abep.com.br](http://www.abep.com.br). Accessed 25 February 2015.
- Steele EM, Claro RM, Monteiro C. Behavioural patterns of protective and risk factors for non-communicable diseases in Brazil. *Public Health Nutr* 2014;17(2):369-75.
- Furlan-Viebig R, Pastor-Valero M. Development of a food frequency questionnaire to study diet and non-communicable diseases in adult population. *Rev Saúde Públ* 2004;38(4):581-4.
- Viner RM, Cole TJ. Who changes body mass between adolescence and adulthood? Factors predicting change in BMI between 16 year and 30 years in the 1970 British Birth Cohort. *Int J Obes* 2006;30:1368-74.
- Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical activity and public health: Updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 2007;39:1423-34.
- Grundey SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. American Heart Association; National Heart, Lung, and Blood Institute.



- Diagnosis and management of the metabolic syndrome: An American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. Published corrections appear in *Circulation* 2005;112:e297 and *Circulation* 2005;112:e298. *Circ J* 2005;112(e297):2735-52.
36. World Health Organization. Obesity: Preventing and managing the global epidemic - Report of a WHO Consultation (TRS894). Geneva: World Health Organization; 2000.
  37. Ashwell M, Hsieh SD. Six reasons why the waist-to-height ratio is a rapid and effective global indicator for health risks of obesity and how its use could simplify the international public health message on obesity. *Int J Food Sci Nutr* 2005;56(5):303-7.
  38. Lippke S, Nigg CR, Maddock JE. Health-promoting and health-risk behaviors: Theory-driven analyses of multiple health behavior change in three international samples. *Int J Behav Med* 2012;19(1):1-13.
  39. McNaughton SA, Mishra GD, Stephen AM, Wadsworth ME. Dietary patterns throughout adult life are associated with body mass index, waist circumference, blood pressure and red cell folate. *J Nutr* 2007;137(1):99-105.
  40. Neumann AICP, Martins IS, Marcopito LF, Araujo EA. Dietary patterns associated with risk factors for cardiovascular disease in a Brazilian city. *Rev Panam Salud Public* 2007;22(5):329-39.
  41. Ribeiro AG, Cotta RMM, Ribeiro SMR. The promotion of health and integrated prevention of risk factors for cardiovascular diseases. *Cien Saúde Colet* 2012;17(1):7-17.
  42. Rolls BJ, Ello-Martin JA, Tohill BC. What can intervention studies tell us about the relationship between fruit and vegetable consumption and weight management? *Nutr Rev* 2004;62(1):1-17.
  43. Kay SJ, Fiatarone Singh MA. The influence of physical activity on abdominal fat: A systematic review of the literature. *Obes Rev* 2006;7(2):183-200.
  44. Dodd LJ, Al-Nakeeb Y, Nevill A, Forshaw MJ. Lifestyle risk factors of students: A cluster analytical approach. *Prev Med* 2010;51(1):73-7.
  45. Schuit AJ, Van Loon AJ, Tijhuis M, Ocké M. Clustering of lifestyle risk factors in a general adult population. *Prev Med* 2002;35(3):219-24.
  46. Rouillier P, Bertrais S, Daudin JJ, Bacro JN, Hercberg S, Boutron-Ruault MC. Drinking patterns are associated with variations in atherosclerotic risk factors in French men. *Eur J Nutr* 2006;45(2):79-87.
  47. Faria CS, Botelho C, Silva RMVG, Ferreira MG. Smoking and abdominal fat in blood donors. *J Bras Pneumol* 2012;38(3):356-63.
  48. Chatkin R, Chatkin JM. Smoking and changes in body weight: Can physiopathology and genetics explain this association? *J Bras Pneumol* 2007;33(6):712-19.