

Nutrición Hospitalaria

Nutrición Hospitalaria

ISSN: 0212-1611

ISSN: 1699-5198

Grupo Arán

Neri-Sánchez, Marisol; Martínez-Carrillo, Beatriz-Elina; Valdés-Ramos, Roxana; Soto-Piña, Alexandra-Estela; Vargas-Hernández, Joel-Alberto; Donají-Benítez-Arciniega, Alejandra
Dietary patterns, central obesity and serum lipids concentration in Mexican adults
Nutrición Hospitalaria, vol. 36, no. 1, 2019, January-February, pp. 109-117
Grupo Arán

DOI: <https://doi.org/10.20960/nh.2002>

Available in: <https://www.redalyc.org/articulo.oa?id=309260651018>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UAEH 

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative



Trabajo Original

Epidemiología y dietética

Dietary patterns, central obesity and serum lipids concentration in Mexican adults *Patrones de dieta, obesidad central y concentración de lípidos séricos en adultos mexicanos*

Marisol Neri-Sánchez¹, Beatriz Elina Martínez-Carrillo², Roxana Valdés-Ramos², Alexandra Estela Soto-Piña²,
Joel Alberto Vargas-Hernández² and Alejandra Donaji Benítez-Arciniega²

Faculties of ¹Nursing and Obstetrics and ²Medicine. Autonomous University of the State of Mexico. Toluca, Mexico

Abstract

Introduction: dietary patterns (DP) analyse the relationship between consumption of foods or nutrients and disease or health outcomes. High prevalence of obesity in adults in Mexico is associated with factors such as high consumption of certain food groups and nutrients. However, few studies have been conducted to explore associations between dietary patterns and obesity in apparently healthy adults.

Objective: to identify major dietary patterns (DP) associated with central-obesity (CO) and lipid concentrations in healthy adults.

Methods: longitudinal study including usual dietary intakes derived from multiple 24-hour-recalls. Waist-circumference (WC) and biochemical measurements were obtained by standardized procedures and DP by principal component analysis. Adjusted-logistic regression was used to examine associations between DP, CO and serum-lipid concentrations.

Results: three DP were identified: healthy-DP, risky-DP and empty-DP. Participants in the healthy-DP were more likely to have lower risk for central-obesity according to WC criteria (OR = 0.31, CI = 0.12, 0.82), $p = 0.017$, but also had the highest risk for elevated LDL-cholesterol (OR = 2.98, CI = 1.16, 7.66), $p = 0.030$. There was no significant association between risky and empty DP with obesity or overweight by body mass index (BMI), central-obesity by WC or serum lipid abnormalities.

Conclusions: the healthy-DP is associated with lower risk for CO, with higher risk for elevated LDL-cholesterol. It is necessary to develop longitudinal studies of foods and nutritional analysis of the diet to clarify these associations, to promote the reduction of modifiable risk factors.

Key words:

Dietary-patterns.
Central-obesity.
Serum lipids.

Resumen

Introducción: los patrones de dieta (PD) analizan la relación entre el consumo de alimentos o nutrientes con la salud y el desarrollo de enfermedades en poblaciones. La elevada prevalencia de obesidad en adultos mexicanos se asocia con factores como el elevado consumo de ciertos grupos de alimentos y nutrientes. Sin embargo, pocos estudios han explorado la asociación entre los patrones dietéticos y la obesidad en adultos aparentemente sanos.

Objetivo: identificar los patrones dietéticos (PD) asociados con la obesidad central (OC) y las concentraciones séricas de lípidos en adultos.

Métodos: estudio longitudinal del consumo dietético obtenido de múltiples recordatorios de consumo de 24 horas. La circunferencia de cintura (CC) y las mediciones bioquímicas se obtuvieron mediante procedimientos estandarizados; los PD, por análisis del componente principal. Mediante regresión logística se identificaron las asociaciones entre PD, OC y las concentraciones séricas de lípidos.

Resultados: se identificaron tres PD: PD saludable, PD de riesgo y PD vacío. Los participantes del PD saludable presentaron menor riesgo de OC de acuerdo con los criterios de la CC (OR = 0.31, CI = 0.12, 0.82), $p = 0.017$, pero también fueron los que presentaron mayor riesgo de cifras elevadas de colesterol-LDL (OR = 2.98, CI = 1.16, 7.66), $p = 0.030$. No hubo asociación estadísticamente significativa entre el PD de riesgo y el PD vacío con obesidad o sobrepeso por IMC, OC por CC o con la presencia de dislipidemias.

Conclusiones: el PD saludable se asocia con un menor riesgo para OC pero con mayor riesgo de elevación del colesterol-LDL. Se necesitan estudios longitudinales para esclarecer estas asociaciones para promover la reducción de factores de riesgo modificables.

Palabras clave:

Patrones de dieta.
Obesidad central.
Lípidos séricos.

Received: 20/04/2018 • Accepted: 30/10/2018

Neri-Sánchez M, Martínez-Carrillo BE, Valdés-Ramos R, Soto-Piña AE, Vargas-Hernández JA, Benítez-Arciniega AD. Dietary patterns, central obesity and serum lipids concentration in Mexican adults. Nutr Hosp 2019;36(1):109-117

DOI: <http://dx.doi.org/10.20960/nh.2002>

Correspondence:

Alejandra Donaji Benítez-Arciniega. Faculty of Medicine. Autonomous University of the State of Mexico. Paseo Tollocan y Jesús Carranza, s/n. Col. Moderna de la Cruz. 50080 Toluca, México
e-mail: abeniteza@uaemex.mx

INTRODUCTION

Diet related diseases are on the rise worldwide. Poor diet increases risk for chronic and metabolic diseases such as obesity and dyslipidemias. Dietary patterns analysis (DP) has emerged as a useful method for the description of relationships between diet and health. The principal component analysis method (PCA) extracts correlation coefficients between foods or dietary elements to classify subjects in groups according to their consumption (1,2).

Evidence based on prospective studies indicates that prudent DP (high intakes of fruits, vegetables, whole grains and fish, and low intakes of saturated fats) were associated with decreased cardiovascular disease in the general population, while those DP with high intake of red or processed meats, sugars or alcoholic beverages were associated with an increased risk (3,4). The evidence is still inconsistent, especially in young and healthy people; only a few epidemiological studies have demonstrated association between DP and serum lipid concentrations (5,6). Therefore, low consumption of fruits, vegetables and fiber, and high consumption of energy, sugars, saturated fats and animal protein increase the risk of chronic non-communicable diseases such as obesity, considered as a public health problem worldwide. Since 2009, a cross-sectional study revealed that DP including cereals, small fish, crackers, fruits-vegetables, and olive oil were related to a beneficial cardiovascular disease risk profile, in contrast with those DP with sweets, red meat, margarine, nuts with salt and cheese, which were related to an unfavorable risk profile in adults (7).

A recent meta-analysis determined that prudent DP have positive effects on biomarkers associated with adiposity, insulin resistance and especially with inflammation in adults (8). Also, in a multi-city cohort study, authors reported that a prudent DP was associated with reduced plasma concentrations of apo B, total cholesterol and LDL-cholesterol, but a western DP, characterized by higher intake of eggs, pastry-cakes, pizza, snacks, refined grains, red meat, vegetable oils and poultry was not (9).

Mexican adults have reported a rapid increase in hypercholesterolemia, hypertriacylglyceridemia and low HDL-cholesterol trends since past decades (10), being the most common lipid abnormality low HDL-cholesterol, a condition that promotes cardiovascular risk factors for morbidity and mortality in addition to the burden of chronic diseases in Mexico (11).

According to the last National Health and Nutrition Survey carried out in 2016 (ENSANUT 2016), the prevalence of abdominal obesity in Mexican adults was 65.4% for men and 87.7% for women. It is known that obesity increases the risk for diabetes mellitus (12,13). Particularly in young healthy adults in Mexico, researchers have found that abdominal obesity is the most prevalent metabolic syndrome component (14).

With respect to diet, the first study of adherence to the World Health Organization dietary recommendations in Mexico reported the presence of a double burden of malnutrition in some segments of the population. A very low percentage of Mexican adults (< 35%) consume the recommended intake of fruits and vegetables, making Mexico one of the countries with the lowest

consumption of fruits and vegetables worldwide (12). This dietary situation seems to remain currently, since ENSANUT 2016 reported that one-half of the adults in Mexico eat fruits while only 42.3% eat vegetables. In contrast, the food group with the highest consumption was sweetened beverages (15). These conditions make diet poor in quality; therefore, the study of DP has special interest.

For this study, our principal aim was to identify major dietary patterns associated with central-obesity and serum lipid concentrations in healthy Mexican adults.

MATERIAL AND METHODS

PARTICIPANTS

The data used in this study were obtained from 102 healthy adults. During seven months follow-up, a sample of 102 from a total of 134 adult school-workers (76% participation rate) completed anthropometric and dietary data. LDL-, HDL- serum total cholesterol and triacylglycerides were determined in all subjects for analysis.

All participants denominated "administrative staff" carried out activities such as laboratory technicians, clerks, cleaning or maintenance in the School of Medicine of the Autonomous University of the State of Mexico, located in the central region of México.

Participants eligible for inclusion were men or women, aged 20-70 years old, with body mass index (BMI) ≥ 20 kg/m² according to the World Health Organization (WHO) criteria, who agreed to provide a serum-sample as well as to participate in dietary and anthropometric evaluations. Health or previous disease conditions in each participant was registered, but it was not considered as an exclusion criteria. The study was approved by the Ethics Committee of the School of Medicine (CONBIOETICA-15CEI03420140313). Written informed consents were obtained from all participants prior to their voluntary participation.

Gender, age, physical activity, smoking habits, concomitant medication and level of education were obtained from a self-administered questionnaire. To compare variables between sexes, Student's *t* test was used for age, BMI and WC; Chi-square Fisher's exact test was used to compare physical activity, smoking, concomitant medication and level of education. No statistically significant differences were observed for age, physical activity, concomitant medication or level of education by sex. BMI and WC were considered for obesity diagnosis.

ANTHROPOMETRIC MEASUREMENTS

All measurements were performed by trained-standardized personnel using standard procedures. Body weight was recorded to the nearest 0.1 kg using a digital electronic scale (Tanita® model 1631); height was recorded to the nearest 0.1 cm using a wall stadiometer (Seca® model 1013522). BMI was computed as a ratio of weight (kg): height squared (m²). BMI cut-off points defined by the WHO according to height and sex were categorized

as < 18.5: underweight; 18.5-24.9: normal; 25-29.9: overweight; and ≥ 30.0 : obese (16,17). Waist circumference was expressed in centimeters, and recorded to the nearest 0.1 cm using a flexible vinyl Gülick tape at the level of midpoint between the iliac crest and the lower border of tenth rib, at the end of normal expiration in supine position. Central-obesity was defined by waist circumference ≥ 80 cm for females, and ≥ 90 cm for males according to the International Diabetes Federation criteria (IDFC) (18). The percentage of fat mass was determined by bioimpedance with the same calibrated electronic scale used for weight measurement. High body fat proportion was defined as $> 25\%$ for men and $> 32\%$ for women (19).

BIOCHEMISTRY DATA

Participants came to the Nutrition Research Laboratory facilities, after an overnight fasting for 10-12 hours. Two venous blood samples were obtained in suitable Vacutainer® tubes and centrifuged. Total-cholesterol, LDL-cholesterol, VLDL-cholesterol, HDL-cholesterol and triacylglycerides (mg/dl) were measured in a Selectra® autoanalyzer equipment by automatized enzymatic-colorimetric methods. LDL-cholesterol was calculated by the Friedewald equation. Altered serum lipid concentrations were classified according to these criteria: total-cholesterol: ≥ 200 mg/dl; triacylglycerides: ≥ 150 mg/dl; LDL-cholesterol: ≥ 150 mg/dl; and low HDL-cholesterol: ≤ 50 mg/dl for women and ≤ 40 mg/dl for men (20,21).

DIETARY ASSESSMENT

Multiple 24-hour recalls (24-h recall)

According to the information reported by each participant, frequency, type and form of food consumption were used to assess habitual dietary intakes; 24-hour recalls interviews were carried out over a seven month period, including weekdays and weekends. Household measures registered were converted into units of volume or weight according to the Mexican equivalents food system. Nutrimind® software 2015 was used to obtain nutrients from 144 different registered foods. Sixteen food-groups were considered depending on the foods items derived from participant's recalls: peanuts-nuts-almonds, vegetables, fruits, no fat cereals, fish and sea food, legumes, lean meat, high fat meat, alcoholic beverages, dairy products, oils and fats, saturated fats, sugars, high fat cereals and fast food (ultra-processed food).

Dietary patterns (DP)

After categorizing 144 food-items into 16 food-groups, factor analysis by principal component analysis (PCA) procedures with Varimax orthogonal rotation were used to identify DP, based on the eating frequency of each food group. Factor loadings refer

the coefficients that define the components in each DP, named according to the food-groups with the highest factor loadings. The number of factors was based on the components with an eigenvalue of 1 and the interpretability of the factors. The eigenvalues signify the amount of variance explained by each of the factors. If a food item had an absolute correlation ≥ 0.5 , it was considered to load on a factor. Factorability of data was confirmed with the Kaiser-Meyer-Olkin (KMO) measure of sampling adequacy and Bartlett's test of sphericity achieving correct estimation of statistical significance. Each factor corresponds to each dietary pattern, therefore, three dietary patterns were derived: healthy-pattern, risk-pattern and empty-pattern, based on the above.

The healthy-pattern included highest consumption of nuts, almonds, peanuts, vegetables, fruits, no fat cereals and fish or sea food, although the risk pattern presented the highest consumption of cereals with fat, high fat meat, fast food and oils and fats. The empty dietary pattern included the highest intakes of alcoholic beverages and sugars. We named dietary patterns considering general dietetic recommendations for healthy adults. All dietary data were adjusted by energy intake.

Statistical analysis

Participants' characteristics were compared by gender, presented as means, standard deviations, and percentages for categorical variables. Student's t test, Chi-square test or ANOVA were used to compare means; logistic regression models were used to explore the association between DP and central-obesity (present/absent) and serum lipid concentrations (desirable/elevated) by calculating odds ratio (OR) with 95% confidence intervals (95% CI). The reference condition (OR = 1.0) represented the chances of being included in the healthy pattern, according to the different anthropometric or lipid categories. Statistical significance was considered at $p < 0.05$. Statistical analyses were performed with SPSS® 21.0 software (UAEM).

RESULTS

CHARACTERISTICS OF PARTICIPANTS

The characteristics of participants are summarized in table I. The mean BMI and WC were 28.3 kg/m² and 97.1 cm for men and 26.2 kg/m² and 85.2 cm for women, respectively. BMI and WC as continuous variables were higher in men ($p < 0.05$). As expected, mean of body fat mass percentage was higher among women; however, the prevalence of excessive body fat mass was not dependent on sex by classifying body fat mass percentage. Also, there were no significant differences between women and men on age nor physical activity, concomitant medication and educational level ($p > 0.05$). Smoking habit was more prevalent in males.

DIETARY PATTERNS

Table II shows factor loadings for the three DP between participants. We extracted three distinct DP with factor loadings > 0.50: the first one was characterized by a high intake of fruits,

vegetables, non-fat cereals, fish and shellfish, legumes and fats with protein, named *healthy pattern*; the second one, named *risk pattern*, was highly correlated with high-fat food, high-saturated fats, fast-foods, high-fat meals, sausages, and sweet-beverages; and the third one, named *empty pattern*, was highly correlated

Table I. Characteristics of participants

	Men	Women	All	<i>p</i> [†]
n (%)	24 (23.5)	78 (76.5)	102	
Age (years)*	38.5 ± 8.9	38.14 ± 11.0	38.23 ± 10.6	0.872
BMI (kg/m ²)*	28.3 ± 3.5	26.2 ± 4.4	26.68 ± 4.3	0.036
WC (cm)*	97.1 ± 10.5	85.2 ± 10.8	88.01 ± 11.8	0.001
Body fat mass percentage (%)*	26.7 ± 5.6	33.8 ± 7.5	32.15 ± 7.7	0.001
Prevalence of excessive body fat mass percentage, n (%)	14 (13.7)	50 (49.0)	64 (62.7)	0.390
Physical activity, n (%)	14 (58.3)	34 (43.6)	48 (47.1)	0.246
Smokers, n (%)	12 (50.0)	14 (17.9)	26 (25.5)	0.003
Concomitant medication, n (%)	6 (25.0)	14 (17.9)	20 (19.6)	0.312
Basic level of education, n (%)	18 (75.0)	48 (61.5)	66 (64.7)	0.168

BMI: body mass index; WC: waist circumference. *Age, body mass index (BMI), waist circumference (WC) and body fat mass percentage are presented as mean ± standard deviation (SD). Categorical variables are presented as a percentage. †Student's *t* test was used for continuous variables and Chi-square, Fisher's test, for categorical variables.

Table II. Factor loadings for three dietary patterns among adults derived by principal component analysis*

Healthy pattern n = 40		Risk pattern n = 35		Empty pattern n = 27	
Variance explained (%)	23.7		17.4		10.2
Nuts, almonds, peanuts	0.879	Cereals with fat	0.815	Alcoholic beverages	0.742
Vegetables	0.780	High fat meat	0.786	Sugars	0.529
Fruits	0.670	Fast food	0.735	Cereals (no fat)	0.429
Cereals (no fat)	0.636	Oils and fats	0.547	Fast food	0.147
Fish/sea food	0.539	Sugars	0.444	Meat (high fat)	0.051
Legumes	0.461	Meat (lean)	0.396	Nuts, almonds, peanuts	0.048
Meat (lean)	0.279	Fats (saturated)	0.353	Fats (saturated)	0.029
Alcoholic beverages	0.240	Cereals (no fat)	0.248	Legumes	0.007
Dairy	0.226	Legumes	0.215	Cereals (with fat)	0.006
Oils and fats	0.218	Alcoholic beverages	0.140	Fish and sea food	-0.084
Fats (saturated)	0.089	Dairy	0.136	Vegetables	-0.087
Meat (high fat)	0.000	Fruits	0.079	Fruits	-0.277
Sugars	-0.048	Fish and sea food	0.048	Oils and fats	-0.277
Cereals (with fat)	-0.063	Vegetables	-0.088	Meat (lean)	-0.406
Fast food	-0.136	Nuts, almonds, peanuts	-0.120	Dairy	-0.523

Correlation values ≥ 0.50 are presented in bold style, for easy identification of the major food group intakes. Variance explained is expressed as a percentage.

*Adjusted for energy intake.

with alcohol and soft-drinks. These three DP explained 23.7%, 17.4% and 10.2% of the variance of food intake, respectively.

Average consumption of food groups, energy and nutrients is presented in table III. Participants in the healthy DP were those who consumed fewer cereals with fat, dairy products and alcoholic beverages, while those in the risk-DP had greater amounts of high fat meat. Adults in the empty-DP group had the highest intake of dairy products and alcohol ($p < 0.05$). Adults in the healthy-DP group had the lowest cholesterol consumption ($p = 0.007$) and the highest monounsaturated fats consumption, with a p-value close to statistical significance ($p = 0.056$). For the other components, while there were no significant differences between DP, although it was observed that adults in the healthy-DP presented lower energy consumption, those in the risk-DP ate more proteins and fats, and those in the empty-DP

consumed higher amounts of carbohydrates, polyunsaturated, and saturated fats than adults in the other DP. In addition, it is notorious that with respect to the recommendations for healthy adults, average fiber intake was low in the three DP (< 30 g/day).

DIETARY PATTERNS, CENTRAL-OBESITY AND SERUM LIPID CONCENTRATIONS

Table IV shows the data on obesity and overweight, central-obesity, body fat-mass percentage, total cholesterol, LDL-cholesterol, HDL-cholesterol, triacylglycerides, and atherogenic index according to DP. Since there were more women in the healthy pattern, it was expected that the highest body fat-mass percentage was

Table III. Intake of dominant food-groups, energy and nutrients for three dietary patterns derived by principal component analysis among healthy adults*

	Dietary pattern			<i>p</i> [†]
	Healthy pattern n = 40	Risk pattern n = 35	Empty pattern n = 27	
Food groups				
Vegetables (g/day)	188.0 ±147.9	182.7 ± 71.5	129.6 ± 76.5	0.078
Fruits (g/day)	229.8 ± 232.7	237.0 ± 128.0	219.6 ± 157.9	0.933
Cereals without fats (g/day)	208.7 ± 102.2	216.5 ± 74.9	193.4 ± 136.5	0.688
<i>Cereals with fats (g/day)</i>	<i>38.9 ± 21.7</i>	<i>58.2 ± 38.3</i>	<i>41.1 ± 22.0</i>	<i>0.010</i>
Legumes (g/day)	52.4 ± 46.2	44.9 ± 42.1	29.3 ± 23.8	0.241
Meat (lean) (g/day)	47.4 ± 28.4	58.0 ± 28.6	57.7 ± 34.2	0.230
Fish and sea food (g/day)	12.4 ± 14.6	16.8 ± 12.8	13.2 ± 10.7	0.318
<i>High fat meat (g/day)</i>	<i>10.0 ± 8.8</i>	<i>15.9 ± 15.8</i>	<i>9.8 ± 6.8</i>	<i>0.046</i>
<i>Dairy products (g/day)</i>	<i>103.4 ± 86.7</i>	<i>144.6 ± 92.0</i>	<i>172.3 ± 170.9</i>	<i>0.045</i>
Oils and fats (g/day)	17.2 ± 14.7	22.5 ± 17.3	19.8 ± 14.8	0.355
High saturated oils/fats (g/day)	3.5 ± 1.9	4.9 ± 3.2	4.0 ± 2.9	0.108
Oils and fats with protein (g/day)	14.6 ± 14.9	9.9 ± 6.2	11.2 ± 9.7	0.169
Sugars (g/day)	93.7 ± 98.9	129.7 ± 121.6	115.4 ± 113.8	0.372
Fast food (g/day)	8.2 ± 7.4	12.7 ± 14.9	7.8 ± 5.9	0.104
<i>Alcohol in beverages (g/day)</i>	<i>11.2 ± 13.2</i>	<i>15.9 ± 14.1</i>	<i>28.9 ± 43.4</i>	<i>0.019</i>
Energy and nutrients				
Energy (kcal/day)	1,774.5 ± 376.8	1,879.2 ± 477.7	1,853.4 ± 424.4	0.544
Carbohydrates (g/day)	225.9 ± 54.9	238.8 ± 62.8	247.2 ± 68.1	0.361
Proteins (g/day)	73.5 ± 16.4	77.6 ± 16.1	75.2 ± 14.4	0.526
Lipids (g/day)	61.9 ± 19.1	68.1 ± 21.7	62.7 ± 16.9	0.344
Fiber (g/day)	20.5 ± 5.4	19.6 ± 5.5	20.5 ± 5.5	0.767
<i>Cholesterol (mg/day)</i>	<i>149.5 ± 46.5</i>	<i>187.4 ± 59.6</i>	<i>181.1 ± 56.7</i>	<i>0.007</i>
<i>Monounsaturated fats (g/day)</i>	<i>8.54 ± 2.4</i>	<i>5.9 ± 2.5</i>	<i>6.4 ± 2.6</i>	<i>0.056</i>
Polyunsaturated fats (g/day)	5.6 ± 2.8	5.4 ±2.9	6.5 ± 3.3	0.335
Saturated fats (g/day)	7.7 ± 2.9	7.6 ± 3.3	8.9 ± 3.1	0.599

*All values are mean $\pm$ standard deviation, adjusted for energy intake except for the energy variable.

† p-values calculated with ANOVA test adjusted by energy intake.

Table IV. Association between obesity, central-obesity, body fat mass percentage, serum lipid concentrations and dietary patterns in healthy adults*

	Dietary pattern			<i>p</i> -value [‡]
	Healthy n = 40	Risk n = 35	Empty n = 27	
Male, n (%)	4 (16.7)	8 (33.3)	12 (50.0)	0.135
Female, n (%)	36 (46.2)	27 (34.6)	15 (19.2)	0.014
BMI (kg/m ²)	27.2 ± 3.7	25.9 ± 4.2	26.9 ± 4.6	0.687
Prevalence of obesity/overweight by BMI [†] (%)	28.4	19.6	16.6	0.372
Prevalence of obesity by BMI (%)	8.8	4.9	5.9	0.619
WC (cm)	88.4 ± 9.6	86.5 ± 12.4	90.5 ± 12.5	0.605
Prevalence of central-obesity by WC (%)	80.0	62.9	66.7	0.206
Body fat mass percentage (%)	34.6 ± 6.7	30.8 ± 7.3	30.3 ± 8.2	0.033
Body fat mass percentage in women (%)	34.9 ± 7.2	32.3 ± 8.8	33.8 ± 5.3	0.001
Body fat mass percentage in men (%)	31.6 ± 4.0	25.6 ± 6.1	25.7 ± 5.2	0.001
Prevalence of excessive body fat mass, n (%)	28 (42.4)	20 (30.3)	18 (27.3)	0.595
Prevalence of excessive body fat mass in women, n (%)	25 (69.4)	13 (65.0)	8 (44.4)	0.208
Prevalence of excessive body fat mass in men, n (%)	3 (10.7)	7 (35.0)	10 (55.6)	0.861
Total cholesterol (mg/dl)	213.9 ± 52.4	209.0 ± 54.5	213.7 ± 53.5	0.998
LDL-cholesterol (mg/dl)	110.4 ± 43.3	96.1 ± 42.3	95.5 ± 38.7	0.236
HDL-cholesterol (mg/dl)	36.5 ± 8.9	34.2 ± 8.7	34.4 ± 9.4	0.712
Triacylglycerides (mg/dl)	146.7 ± 85.4	160.2 ± 75.2	183.7 ± 108.4	0.250
Atherogenic index	6.4 ± 1.9	6.7 ± 2.1	6.5 ± 2.1	0.888

BMI: body mass index; WC: waist circumference. *All values are mean ± standard deviation. [†]According to BMI overweight and obesity in the same category. [‡]*p*-value calculated with ANOVA test, adjusted by energy intake for continuous variables; Chi-square test for categorical variables. [§]≥ 32% for females, ≥ 25% for males.

observed in this dietary pattern ($p < 0.05$). Nonetheless, it should be noted that body fat mass percentage was the highest in women and men in the healthy pattern ($p < 0.05$). Also, it should be noted that the mean of body fat mass percentage was above the healthy limits among adults in the three DP. In addition, for a more complete analysis, a subsample of 24 women was extracted to analyze their distribution in the three dietary patterns based on the consumption of the characteristic food groups of each of them. As no statistically significant differences were found, we assume that sex does not determine the classification of participants in dietary patterns.

Even when differences were not statistically significant ($p > 0.05$), it is interesting to observe that according to BMI and WC, obesity/overweight, obesity, central obesity, the highest average HDL-cholesterol and lowest atherogenic index were more prevalent in adults in the healthy-DP group, but all participants had elevated averages of total cholesterol, triacylglycerides and low HDL-cholesterol concentrations.

Finally, ordinal logistic regression is shown in table V. Participants in the healthy-DP group were more likely to have lower risk for central obesity according to WC criteria (OR = 0.31, CI = 0.12, 0.82), $p = 0.017$. Interestingly, participants in the healthy-DP group also had the highest risk for elevated LDL-cholesterol (OR = 2.98, CI = 1.16, 7.66), $p = 0.030$. No significant associa-

tion was found between risky and empty DP with obesity or overweight by BMI, central-obesity by WC or serum lipid abnormalities.

DISCUSSION

In this study in Mexican healthy adults, three major DP were identified and their association with central obesity and serum lipid concentrations was investigated. With the use of PCA, we applied a complex epidemiological method recently used in Mexico. Clearly these three DP characterize the diet of young urban adults, describing a large variety of fresh, industrialized and traditional foods as habitual diet. It is interesting that with no differences between the three DP, consumption of dairy products is an important feature in this population, which may indicate that it is a habit derived from decades of dairy consumption, possibly due to a high availability of milk in childhood.

The first dietary pattern, which we considered as healthier because it included consumption of nuts, vegetables, fruits, cereals without fats, fish and the lowest intake of high fat meat and dairy, was associated with a low intake of cholesterol and a high intake of mono-unsaturated fats. The risk-DP and the empty-DP were considered as unhealthy because they had high amounts of non-recommended foods like high-fat cereals and meats, fast-foods, alcohol and sugars.

Adults from the healthy-DP group were more likely to eat less

Table V. Odds ratios (OR) and 95% confidence intervals (95% CI) of the association between dietary patterns, obesity, abdominal obesity and elevated serum lipid concentrations in 102 healthy adults

	Dietary pattern		
	Healthy	Risky	Empty
BMI (kg/m²)			
Normal (18.5-24.9)	1		
Obesity/overweight (≥ 25.0)	1.78 (0.75, 4.21)	0.61 (0.26, 1.42)	0.90 (0.36, 2.25)
<i>p</i> -value	0.20	0.28	0.81
Waist circumference (cm)			
Normal	1		
Abdominal obesity (≥ 80 for women; ≥ 90 for men)	0.31 (0.12, 0.82)	1.82 (0.76, 4.31)	1.77 (0.71, 4.43)
<i>p</i> -value	0.01	0.18	0.23
Serum lipid concentrations (mg/dl)			
TC desirable	1		
TC elevated (≥ 200)	1.02 (0.45, 2.32)	0.75 (0.33, 1.73)	1.36 (0.55, 3.38)
<i>p</i> -value	1.0	0.52	0.64
LDL-C desirable	1		
LDL-C elevated (≥ 150)	2.98 (1.16, 7.66)	0.42 (0.14, 1.24)	0.63 (0.21, 1.91)
<i>p</i> -value	0.03	0.14	0.59
HDL-C desirable	1		
HDL-C low (≤ 50 for female; ≤ 40 for male)	0.73 (0.29, 1.84)	1.15 (0.44, 3.03)	1.26 (0.44, 3.58)
<i>p</i> -value	0.63	0.81	0.79
Triacylglycerides desirable	1		
Triacylglycerides elevated (≥ 150)	0.55 (0.24, 1.26)	1.31 (0.57, 3.01)	1.47 (0.61, 3.59)
<i>p</i> -value	0.21	0.53	0.49

BMI: body mass index; TC: total cholesterol; LDL: low density lipoproteins; VLDL: very low density lipoproteins; HDL: high density lipoproteins; TG: triglycerides.

fats and to drink less alcoholic beverages, possibly in concordance with a healthier behavior. Furthermore, this healthy DP was also associated with a lower risk for central obesity in adults, although adults in this healthy-DP presented higher body fat mass. In our study, adults with higher body fat mass percentage were more likely to eat healthier, while no significant association was found regarding risk or empty DP and prevalence of obesity. This probably could be because when adults perceive they have overweight or obesity, they improve their diet; in other words, selection of healthier foods and, as a consequence, a healthier diet is perhaps the product of overweight or obesity.

Analyzing the overall diet components and combined nutrients resulted in a better way to examine dietary effects on health outcomes and a good interpretation of possible risk factors in the future. It is very important to remember that participants in this study were healthy young adults, and the importance of the dietary analysis is related to the promotion of modification of risk factors for metabolic conditions (22).

Another important finding in our results was the low consumption of fiber in all participants; even in participants of the healthy-DP group, who presented higher consumption of vegetables and

fruits, the intake of dietary fiber does not cover recommendations for a healthy population (23). However, we agree with the evidence that promotes the study of all the components of the diet, rather than one of its components, particularly with respect to fruits and vegetables consumption (24). In this study, it must be noted that PCA is one of the best techniques for the extraction of dietary patterns in nutritional epidemiology (25).

With respect to the transcendence of chronic diseases related to diet, some studies mention that the nutritional epidemiological transition leads to chronic diseases as the growing epidemic obesity in Latin America (26,27). From the perspective of public health, obesity represents the most important challenge. In Mexico, the total cost of overweight and obesity and their indirect costs are estimated to reach 150.860 billion pesos (8.122 billion dollars) for 2017 (28).

The imbalance in the diet is crucial to the development of obesity, especially the high consumption of fats and sugars, comprised in the risky-DP group in our study, which is similar to previous findings (29-32).

Likewise, associations between high consumption of risky or empty foods with metabolic disorders such as hypercholester-

olemia and hypertriglyceridemia had been described previously (33,34), and similarly with our results, despite not having found significant associations, abnormal concentrations of total cholesterol and triacylglycerides were observed in apparently healthy subjects. In this regard, McNaughton et al. have reported that a high intake of white bread, chips, sugar in hot-beverages, hamburgers and soft drinks and a low intake of vegetables are associated with an increase in serum total-cholesterol, triacylglycerides and low concentration of HDL-cholesterol (35). These are conditions that, unless they are modified, could lead to the development of obesity as well (34,36,37).

On the other hand, we found in this study a healthy-DP integrated with food groups such as fruits and vegetables considered a prudent or healthy-DP as in other studies, which have been linked to the prevention of chronic diseases such as obesity, diabetes, hypertension and cancer (34,38-40). These could be used for the assessment of diet quality in future studies, or in interventions to modify risk factors such as diet.

CONCLUSIONS

In this sample of Mexican adults, three DP were identified: a healthy-DP, a risk-DP and an empty-DP. The healthy-DP was associated with a low risk of central obesity. Our results confirm that higher intakes of vegetables, fruit, lean protein, nuts and lower intakes of dairy and alcohol are likely beneficial for the prevention of central obesity, but not for desirable lipid concentrations.

REFERENCES

- Slattery ML, Boucher KM, Caan BJ, Potter JD, Ma KN. Eating patterns and risk of colon cancer. *Am J Epidemiol* 1998;148:4-16.
- Hu FB, Rimm EB, Stampfer MJ, Ascherio A, Spiegelman D, Willett WC. Prospective study of major dietary pattern and risk of coronary heart disease in men. *Am J Clin Nutr* 2000;72:912-21. DOI: 10.1093/ajcn/72.4.912
- Rodríguez-Monforte M, Flores-Mateo G, Sánchez E. Dietary patterns and CVD: A systematic review and meta-analysis of observational studies. *Br J Nutr* 2015;114:1341-59. DOI: 10.1017/S0007114515003177.
- Kimokoti RW, Gona P, Zhu L, Newby P, Millen BE, Brown LS, et al. Dietary patterns of women are associated with incident abdominal obesity but not metabolic syndrome. *J Nutr* 2012;142:1720-7. DOI: 10.3945/jn.112.162479
- Schulze MB, Hoffmann K, Kroke A, Boeing H. Risk of hypertension among women in the EPIC-Potsdam study: comparison of relative risk estimates for exploratory and hypothesis-oriented dietary patterns. *Am J Epidemiol* 2003;158:365-73. DOI: 10.1093/aje/kwg156
- United States Department of Agriculture. A series of systematic reviews on the relationship between dietary patterns and health outcomes; 2014. Accessed on Sept 8th 2016. Available from: <http://www.nel.gov/vault/2440/web/files/DietaryPatterns/DPRptFullFinal.pdf>
- Panagiotakos D, Pitsavos C, Chrysoshoou C, Palliou K, Lentzas I, Skoumas I, et al. Dietary patterns and 5-year incidence of cardiovascular disease: a multivariate analysis of the ATTICA study. *Nutr Metab Cardiovasc Dis* 2009;19:253-63. DOI: 10.1016/j.numecd.2008.06.005
- Neale EP, Batterham MJ, Tapsell LC. Consumption of a healthy dietary pattern results in significant reductions in C-reactive protein levels in adults: a meta-analysis. *Nutr Res* 2016;36(5):391-401. DOI: 10.1016/j.nutres.2016.02.009
- Poggio R, Elorriaga N, Gutiérrez L, Irazola V, Rubinstein A, Danaei G. Associations between dietary patterns and serum lipids, apo and C-reactive protein in an adult population: evidence from a multi-city cohort in South America. *Br J Nutr* 2017;117(4):548-55. DOI: 10.1017/S0007114517000514
- Villalpando S, Shamah-Levy T, Rojas R, Aguilar-Salinas CA. Trends for type 2 diabetes and other cardiovascular risk factors in Mexico from 1993-2006. *Salud Publica Mex* 2010;52(1):S72-9.
- Escobedo-de-la-Peña J, De Jesús Pérez R, Scharngrodsy H, Champagne B. Prevalencia de dislipidemias en la ciudad de México y su asociación con otros factores de riesgo cardiovascular. Resultados del estudio CARMELA. *Gac Med Mex* 2014;150:128-36.
- Secretaría de Salud. Encuesta Nacional de Salud y Nutrición 2016 Medio Camino. ENSANUT 2016. Informe final de resultados. Instituto Nacional de Salud Pública; 2016. Available from: http://promocion.salud.gob.mx/dgps/descargas1/doctos_2016/ensanut_mc_2016-31Oct.pdf
- Alegre-Díaz J, Herrington W, López-Cervantes M, Gnatiuc L, Ramírez R, Hill M, et al. Diabetes and cause-specific mortality in Mexico City. *N Engl J Med* 2016;375(20):1961-71. DOI: 10.1056/NEJMoa1605368.
- Urquidez-Romero R, Murguía-Romero M, Esparza-Romero J, Díaz-Torres BA, Rodríguez-Tadeo A, Medrano-Donlucas G, et al. Abdominal obesity is strongly associated to blood pressure in young Mexicans. *Nutr Hosp* 2017;34:357-62. DOI: 10.20960/nh.171
- Ponce X, Rodríguez-Ramírez S, Mundo-Rosas V, Shamah T, Barquera S, De Cossio TG. Dietary quality indices vary with sociodemographic variables and anthropometric status among Mexican adults: a cross-sectional study. Results from the 2006 National Health and Nutrition Survey. *Public Health Nutr* 2014;17(8):1717-28. DOI: 10.1017/S1368980013002462
- World Health Organization (WHO). 2008. Global strategy on diet, physical activity, and health. Available from: <http://www.who.int/dietphysicalactivity/publications/facts/obesity/en>. World Health Organization (2000) Obesity: Preventing and Managing the Global Epidemic Report on a WHO Consultation. WHO Technical Report Series no. 894. Geneva: WHO.
- Secretaría de Gobernación. Norma oficial mexicana NOM-174-SSA1-98, para el manejo integral de la obesidad. México; 1998.
- International Diabetes Federation (IDF). The IDF consensus worldwide definition of the metabolic syndrome. IDF; 2006.
- Deurenberg P, Yap M, Staveren WA. Body mass index and percent body fat: a meta-analysis among different ethnic groups. *Int J Obes Relat Metab Disord* 1998;22:1164-71.
- Goncalves JP, Oliveira A, Severo M, Santos AC, López C. Cross-sectional and longitudinal associations between serum uric acid and metabolic syndrome. *Endocrine* 2012;41(3):450-7. DOI: 10.1007/s12020-012-9629-8
- Secretaría de Gobernación. Norma oficial mexicana NOM-037-SSA2-2002 para la prevención, tratamiento y control de las dislipidemias. México; 2002.
- Slattery ML. Analysis of dietary patterns in epidemiological research. *Appl Physiol Nutr Metab* 2010;35:207-10. DOI: 10.1139/H10-006
- Kim Y, Je Y. Dietary fibre intake and mortality from cardiovascular disease and all cancers: a meta-analysis of prospective cohort studies. *Arch Cardiovasc Dis* 2016;109(1):39-54. DOI: 10.1016/j.acvd.2015.09.005
- Alissa EM, Ferns GA. Dietary fruits and vegetables and cardiovascular diseases risk. *Crit Rev Food Sci Nutr* 2017;57(9):1950-62. DOI: 10.1080/10408398.2015.1040487
- Sugawara N, Yasui-Furukori N, Sato Y, Saito M, Furukori H, Nakagami T, et al. Dietary patterns are associated with obesity in Japanese patients with schizophrenia. *BMC Psychiatry* 2014;14:184. DOI: 10.1186/1471-224X-14-184
- Aballay LR, Eynard AR, Díaz MP, Navarro A, Muñoz SE. Overweight and obesity: a review of their relationship to metabolic syndrome, cardiovascular disease, and cancer in South America. *Nutr Rev* 2013;71:168-79. DOI: 10.1111/j.1753-4887.2012.00533.x
- Chatenoud L, Bertuccio P, Bosetti C, Malvezzi M, Levi F, Negri E, et al. Trends in mortality from major cancers in the Americas: 1980-2010. *Ann Oncol* 2014;25:1843-53. DOI: 10.1093/annonc/mdl206
- Rtveladze K, Marsh T, Barquera S, Sánchez Romero LM, Levy D, Meléndez G, et al. Obesity prevalence in Mexico: impact on health and economic burden. *Public Health Nutr* 2014;17:233-9. DOI: 10.1017/S1368980013000086
- Gutiérrez-Pliego LE, Camarillo-Romero ES, Montenegro-Morales LP, Garduño-García JJ. Dietary patterns associated with body mass index (BMI) and lifestyle in Mexican adolescents. *BMC Public Health* 2016;16(1):850. DOI: 10.1186/s12889-016-3527-6
- Pou SA, Del Pilar Díaz M, De La Quintana AG, Forte CA, Aballay LR. Identification of dietary patterns in urban population of Argentina: study on diet-obesity relation in population-based prevalence study. *Nutr Res Pract* 2016;10(6):616-22. DOI: 10.4162/nrp.2016.10.6.616
- Cunha DB, De Almeida RM, Sichieri R, Pereira RA. Association of dietary patterns with BMI and waist circumference in a low-income neighborhood in Brazil. *Br J Nutr* 2010;104:908-13. DOI: 10.1017/S0007114510001479

32. Shang X, Li Y, Liu A, Zhang Q, Hu X, Du S, et al. Dietary pattern and its association with the prevalence of obesity and related cardiometabolic risk factors among Chinese Children. *PLoS ONE* 2012;7(8):e43183. DOI: 10.1371/journal.pone.0043183
33. Sadakane A, Tsutsumi A, Gotoh T, Ishikawa S, Ojima T, Kario K, et al. Dietary patterns and levels of blood pressure and serum lipids in a Japanese population. *J Epidemiol* 2008;18(2):58-67.
34. Fung TT, Rimm EB, Spiegelman D, Rifai N, Tofler GH, Willett WC, et al. Association between dietary patterns and plasma biomarkers of obesity and cardiovascular disease risk. *Am J Clin Nutr* 2001;73:61-7. DOI: 10.1093/ajcn/73.1.61
35. McNaughton SA, Mishra GD, Brunner EJ. Food patterns associated with blood lipids are predictive of coronary heart disease: the Whitehall II study. *Br J Nutr* 2009;102(4):619-24. DOI: 10.1017/S0007114509243030
36. Limosin F, Gasquet I, Leguay D, Azorin JM, Rouillon F. Body mass index and prevalence of obesity in a French cohort of patients with schizophrenia. *Acta Psychiatr Scand* 2008;118:19-25. DOI: 10.1111/j.1600-0447.2008.01208.x
37. Togo P, Osler M, Sørensen TI, Heitmann BL. Food intake patterns and body mass index in observational studies. *Int J Obes Relat Metab Disord* 2001;25:1741-51. DOI: 10.1038/sj.ijo.0801819
38. Newby PK, Muller D, Hallfrisch J, Qiao N, Andres R, Tucker KL. Dietary patterns and changes in body mass index and waist circumference in adults. *Am J Clin Nutr* 2003;77:1417-25. DOI: 10.1093/ajcn/77.6.1417
39. Kaiser KA, Brown AW, Bohan Brown MM, Shikany JM, Mattes RD, Allison DB. Increased fruit and vegetable intake has no discernible effect on weight loss: a systematic review and meta-analysis. *Am J Clin Nutr* 2014;100:567-76. DOI: 10.3945/ajcn.114.090548
40. Rezazadeh A, Rashidkhani B. The association of general and central obesity with major dietary patterns of adult women living in Tehran, Iran. *J Nutr Sci Vitaminol (Tokyo)* 2010;56(2):132-8. DOI: 10.3177/jnsv.56.132