

Endosonographic Characteristics of Subepithelial Lesions of the Upper Digestive Tract: Experience of a Referral Center in Colombia

Ileana Rocío Bautista-Parada,^{1*} Ángel Rojas-Espinosa,² Lázaro Antonio Arango-Molano,³ Andrés Sánchez-Gil,³ Claudia Díaz-Tobar.³

OPEN ACCESS

Citation:

Bautista-Parada IR, Rojas-Espinosa A, Arango-Molano LA, Sánchez-Gil A, Díaz-Tobar C. Endosonographic Characteristics of Subepithelial Lesions of the Upper Digestive Tract: Experience of a Referral Center in Colombia. *Revista colomb. Gastroenterol.* 2023;38(3):264-270. <https://doi.org/10.22516/25007440.1014>

¹ General surgeon, clinical-surgical gastroenterology specialist, Fundación FOSUNAB. Bucaramanga, Colombia.

² General surgeon, clinical-surgical gastroenterology specialist. Gastroenterologist, Fundación FOSUNAB. Bucaramanga, Colombia.

³ General surgeon, clinical-surgical gastroenterology specialist. Professor at the Universidad de Caldas. Unión de Cirujanos. Manizales, Colombia.

*Correspondence: Ileana Rocío Bautista-Parada. ibautista4@hotmail.com

Received: 29/01/2023

Accepted: 13/04/2023



Abstract

Introduction: Subepithelial lesions (SELs), described as bulges or masses covered by healthy-looking mucosa, are usually found incidentally during endoscopic studies. They are typically asymptomatic and are estimated to be identified in 1% of esophago-gastroduodenoscopies performed. **Materials and methods:** A descriptive study was conducted with retrospective data collection. We included all patients treated at the Unión de Cirujanos, a referral gastroenterology unit of the Coffee Region in Manizales, between January 2020 and January 2022, who underwent endoscopic ultrasonography to study subepithelial-looking lesions located in the esophagus, stomach, and duodenum. **Results:** 152 endoscopic ultrasounds were performed, finding 108 SELs; 66.6% of the patients were women, and the average age was 58. Most SELs were located in the stomach (78.7%), the antrum being the most frequent location. The average diameter of the gastric SELs was 14.6 mm, and 47% of the lesions depended on the fourth echolayer; the most frequent presumptive diagnoses were gastrointestinal stromal tumor (GIST; 65.8%) and lipoma (11.7%). **Conclusions:** SELs of the GI tract originate in the muscularis mucosae, submucosa, or muscularis propria. They are most frequently located in the stomach, and their characterization usually requires endoscopic ultrasonography and histopathology. Treatment of these lesions remains controversial due to their low frequency, histological variety, and low malignant potential.

Keywords

Endoscopic ultrasonography, subepithelial lesion, GIST, follow-up.

INTRODUCTION

Subepithelial lesions (SELs), described as lumps or masses covered by healthy-appearing mucosa, are usually found incidentally during endoscopic studies; they are typically asymptomatic and are estimated to be identified in 1% of esophagogastroduodenoscopies (EGDs) performed^(1,2). Recently, their incidence has increased; however, this phenomenon may be due to the greater availability of endoscopic studies in our setting. The present research intends to characterize the endosonographic lesions in the upper GI tract

in patients treated at a gastroenterology referral center in the Coffee Region between January 2020 and January 2022.

MATERIALS AND METHODS

A descriptive study was conducted with a retrospective collection of information. It included all patients treated at Unión de Cirujanos (gastroenterology referral center for the Coffee Region located in Manizales, Colombia) from February 2020 to January 2022 who underwent endoscopic ultrasound as part of the study of lesions with a sube-

pithelial appearance located in the esophagus, stomach, or duodenum that were found incidentally in previous endoscopic studies.

A descriptive analysis of the recorded data was performed, and the median was calculated for the numerical variables, while the qualitative variables were described with frequencies. Endoscopic ultrasound (EUS) was performed with a high-resolution Fujinon radial scanning endoscopic ultrasound machine, SU-1 processor, with variable frequencies of 7.5, 12, and 20 MHz, by three experienced endosonographers, and all procedures were performed under intravenous sedation. Endosonographic characteristics were prospectively recorded for all lesions: location, maximum diameter, growth pattern, echolayer of origin, and echogenicity.

RESULTS

One hundred fifty-two EUSs were performed; mucosal lesions were found in 44 cases, extrinsic compressions in nine, and a diverticular formation in one. The remaining 108 cases indeed corresponded to SELs (**Figure 1**).




Figure 1. Studies performed, patients excluded, and the number of studies by anatomical location. SELs: subepithelial lesions; EUS: endoscopic ultrasound. Image owned by the authors.

Of the 108 patients with SEL, 72 (66.6%) were female, and the average age was 58. Most of the SELs evaluated were located in the stomach (78.7%); of these, the most frequent location was the antrum. The most common presumptive diagnosis based on endosonographic findings was gastrointestinal stromal tumor (GIST; 60.1%), followed by ectopic pancreas (14.8%) and lipoma (12%).

The average age, sex, echolayer of origin, and presumptive diagnoses of the lesions according to their location are described in **Tables 1, 2, and 3**.

DISCUSSION

Most SELs are detected incidentally during procedures requested for another indication. They are found with the same frequency in men and women, generally after the fifth decade of life; these data are consistent with those in this work. It has been reported that gastric SEL is found in up to 0.3% of middle-aged adults⁽³⁾.

SEL can be classified as non-neoplastic, which includes inflammatory changes, cysts, ectopic pancreatic tissue, and varicose veins, and neoplastic, with a low malignant potential such as lipomas, leiomyomas or with a high malignant potential such as GIST, neuroendocrine tumors or lymphomas⁽⁴⁾. The majority are asymptomatic since they usually do not involve the mucosa; however, in some cases, they can manifest with digestive bleeding, iron deficiency anemia, abdominal pain, or signs of intestinal obstruction depending on the location and size of the lesion, mainly when they are close to the pylorus, cardia, ileocecal valve, or rectum⁽⁵⁾.

Diagnosis is usually challenging since the endoscopic appearance is usually insufficient, and the diagnostic yield of conventional biopsies is usually meager⁽⁶⁾. Most are identified in the endoscopic evaluation as small lumps (less than 20 mm) covered with normal-looking mucosa, data consistent with those in our paper, although some have endoscopic characteristics that may suggest their etiology, such as lipomas. They are usually identified as yellowish in color and soft in consistency when pressed with forceps; this sign is known as the *pillow sign* and has a reported sensitivity of 98% for this diagnosis^(2,7). On the other hand, ectopic pancreatic tissue is usually identified as a nodular lesion with central umbilication in the antrum (**Figure 2**)⁽⁸⁾.

In order of frequency, they are located in the stomach, esophagus, duodenum, and colon, data consistent with the results described in our population⁽²⁾. Leiomyomas are the subepithelial lesions most frequently found in the distal two-thirds of the esophagus, and GISTs are the most frequent in the stomach (**Table 4**)⁽⁵⁾.

EUS is the gold standard in the evaluation of SELs of the gastrointestinal tract⁽⁹⁾, as it allows differentiating extrinsic compressions from intramural lesions, establishing the echolayer of origin, size, echogenicity, and margins, as well as identifying regional lymphadenopathy and obtain tissue samples if necessary⁽¹⁰⁾. There are pathognomonic endosonographic features for lipomas and varicose veins; however, for other types of lesions, its diagnostic accuracy is reported between 43% and 67%⁽¹¹⁾.

We recommend taking a biopsy of SEL suggestive of GIST, those with high-risk endosonographic characteristics (irregular edges, cystic degeneration, ulceration, echogenic foci, heterogeneity), or those greater than 20 mm⁽¹²⁾.

Table 1. Endosonographic characteristics and presumptive diagnoses of gastric subepithelial lesions

Gastric subepithelial lesions			Presumptive diagnosis					
			Echolayer	GIST	EP	LEI	LIP	NET
Average age	59.3							
Sex	Male	29						
	Female	56						
Average diameter	14.6 mm		2	16		1		17
Location	Antrum	38	2 and 3	1	2			3
	Body	35	3		11		10	25
	Fundus	11	4	39	1			4
	Cardia	1	Total	56	14	1	10	4
								85

GIST: gastrointestinal stromal tumors; LEI: leiomyoma; LIP: lipoma; EP: ectopic pancreas; NET: neuroendocrine tumor. Table prepared by the authors.

Table 2. Endosonographic characteristics and presumptive diagnoses of esophageal subepithelial lesions

Esophageal subepithelial lesions			Presumptive diagnosis				Total
			Echolayer	GIST	LEI	CYST	
Average age	53.9						
Sex	Male	3	1 and 2			1	1
	Female	8	2	1	4	1	6
Average diameter	18.1 mm		3			1	1
Location	Proximal	0	3 and 4		1		1
	Middle	3	4	2			2
	Distal	8	Total	3	5	3	11

GIST: gastrointestinal stromal tumors; LEI: leiomyoma; CYST: intestinal duplication cyst. Table prepared by the authors.

Table 3. Endosonographic characteristics and presumptive diagnoses of duodenal subepithelial lesions

Duodenal subepithelial lesions			Presumptive diagnosis					Total
			Echolayer	GIST	EP	LIP	CYST	
Average age	56.4							
Sex	Male	4	2	2				2
	Female	8	3		2	3	1	6
Average diameter	13 mm		3 and 4	1				1
Location	Bulb	11	4	3				3
	D2	1	Total	6	2	3	1	12

GIST: gastrointestinal stromal tumors; LEI: leiomyoma; CYST: intestinal duplication cyst; Ep: ectopic pancreas. Table prepared by the authors.

Due to the low yield of conventional biopsies, multiple strategies have been designed to perform them, such as biopsy-on-biopsy and using jumbo forceps or loops. It can also be performed under endosonographic guidance or with a mucosal incision that allows the lesion to be exposed and the biopsy to be taken⁽¹³⁾. This last technique is recommended for lesions smaller than 20 mm, and its general diagnostic yield is 89%, given the technical difficulty in taking needle samples in lesions of this size⁽¹⁴⁾. When com-

paring the diagnostic performance of fine needle aspiration (FNA) with fine needle biopsy (FNB), there is evidence in favor of the use of FNB without a recommendation of a specific type of needle^(15,16). The diagnostic accuracy with FNB has been reported between 83% and 100%⁽¹⁷⁾.

On the one hand, asymptomatic lesions with endoscopic and endosonographic characteristics compatible with varicose veins, ectopic pancreatic tissue, or lipomas do not require resection or additional follow-up. On the other




Figure 2. Endoscopic view of SEL of the antrum suggestive of an ectopic pancreas. **A.** Endosonographic appearance. **B.** Lesion involving the third echolayer (arrow); endoscopic view of the distal gastric body SEL. **C.** Endosonographic appearance. **D.** Third echolayer lesion consistent with a lipoma (arrow); endoscopic view of the gastric body SEL. **E.** Endosonographic appearance. **F.** Fourth echolayer lesion suggestive of GIST (dotted lines); endoscopic view of an elevation of a subepithelial appearance of the gastric antrum. **G.** Endosonographic image. **H.** Endosonographic image with extrinsic compression of the gallbladder (arrow). Image owned by the authors.

Table 4. Endoscopic and endosonographic characteristics of the most common subepithelial lesions

Subepithelial lesion	Endoscopic appearance	Endosonographic appearance	Compromised echolayer	Most frequent location
Lipoma	Yellowish, positive pillow sign	Hyperechogenic, homogeneous	3. rd	Any location of the GIT
Leiomyoma	No specific features	Hypoechoic, well-defined edges	2. nd , 3. rd , 4. th	Esophagus, stomach
Duplication cyst	Smooth, regular surface, compressible	Anechoic, negative Doppler signal	3. rd	Any location of the GIT
Ectopic pancreas	Central umbilication	Hypoechoic, heterogeneous	3. rd , 4. th	Antrum, gastric body, duodenum
Varicose veins	Bluish hue, tortuous	Anechoic, positive Doppler signal	3. rd	Any location of the GIT
Low-risk GIST	Subepithelial elevation without mucosal ulceration or bleeding, <30 mm	Hypoechoic, heterogeneous, hypervascular	2. nd or 4. th	Esophagus, stomach, small intestine, rectum
High-risk GIST	Areas of ulceration or bleeding, >30 mm	Hypoechoic, heterogeneous areas of cystic degeneration or hyperechoic foci	2. nd or 4. th	Esophagus, stomach, small intestine, rectum
Schwannoma	No specific features	Hypoechoic, homogeneous	4. th	Gastric body
Lymphoma	No specific features	Hypoechoic	2. nd , 3. rd , 4. th	Stomach, small intestine
Neuroendocrine tumors	Rounded, yellowish or reddish hue	Hypoechoic or hyperechoic	2. nd , 3. rd	Stomach, small intestine, rectum

GIT: gastrointestinal tract. Table prepared by the authors.

hand, lesions diagnosed histologically as benign, such as leiomyomas, lipomas, ectopic pancreas, granular cell tumors, schwannomas, and glomus tumors, among others, do not require any follow-up or additional treatment since the risk of malignancy is very high low. Lesions with malignant confirmation require individualized treatment⁽⁵⁾. In the case of neuroendocrine tumors, type 1 tumors smaller than 10 mm could be candidates for annual endoscopic follow-up since the risk of malignancy is low⁽¹⁸⁾.

Managing GISTs smaller than 20 mm without suspicious endosonographic characteristics is controversial since the lesions' follow-up and resection are accepted in the literature, given that they have a low risk of malignancy^(12,19). If follow-up is chosen, the method of choice is EUS, and even though there is no consensus on the best follow-up strategy, EUS has been recommended every 1-2 years for lesions between 10 and 20 mm and every 2-3 years for lesions smaller than 10 mm. Surgical resection of GISTs diagnosed in organs other than the stomach is advisable⁽²⁰⁾.

When a histological diagnosis is not available, as is frequently the case in our environment, the symptoms, the location of the lesion, and the endosonographic characteristics must be taken into account. Lesions smaller than 20 mm in an asymptomatic gastric or esophageal area have a low risk of malignancy; therefore, their follow-up could

be considered the best option. Although the follow-up interval and method are not standardized, esophagogastroduodenoscopy (EGD) and EUS have been suggested at 3-6 months to verify the stability of the lesion and, subsequently, EGD or EUS every 1-3 years⁽²¹⁾.

Given the retrospective nature of the study and that long-term follow-up of the lesions was not carried out, it is impossible to establish the incidence of malignancy or changes in lesions that would allow suggesting follow-up intervals in the study population.

Subepithelial lesions of the colon and rectum always require histological evaluation to establish the appropriate treatment since there is no evidence to support their follow-up as an adequate management strategy⁽⁵⁾.

Endoscopic resection aims to obliterate the lesions; the indication exists for lesions with the potential for malignancy that cause symptoms or patients who are candidates for bariatric surgery. The type of resection depends on the involved echolayer and its anatomical location.

Mucosal resection (MR) or endoscopic submucosal dissection (ESD) are proposed strategies for esophageal lesions. In the former, the use of bands has reported a technical success of up to 100% in lesions smaller than 20 mm⁽²²⁾; for more extensive lesions up to 40 mm, endoscopic full-thickness resection (EFTR) could be considered⁽²³⁾.

In gastric SEL with an indication for resection, the endoscopic route using MRI, ESD, or EFTR for lesions smaller than 40 mm or laparoscopic wedge resection is an option⁽⁵⁾.

CONCLUSION

SELs of the gastrointestinal tract originate in the muscularis mucosa, submucosa, or muscularis propria. They are most frequently located in the stomach, and their characterization usually requires endoscopic ultrasound and, in some cases, histopathological study.

The results are relevant since it is the first study in our environment that characterizes this type of lesion. However, given that the treatment described in the literature remains controversial (due to its low frequency, histological variety, and low malignant potential), it would be beneficial to conduct prospective studies in which long-term endosonographic follow-up is performed to establish the incidence of malignant transformation in our population and, thus, propose follow-up intervals or the need for additional interventions such as biopsies or resection of the lesions.

REFERENCES

1. Attila T, Aydin Ö. Lesion size determines diagnostic yield of EUS-FNA with onsite cytopathologic evaluation for upper gastrointestinal subepithelial lesions. *Turkish J Gastroenterol.* 2018;29(4):436-41. <https://doi.org/10.5152/tjg.2018.17876>
2. Menon L, Buscaglia JM. Endoscopic approach to subepithelial lesions. *Therap Adv Gastroenterol.* 2014;7(3):123-30. <https://doi.org/10.1177/1756283X13513538>
3. Nishida T, Kawai N, Yamaguchi S, Nishida Y. Submucosal tumors: Comprehensive guide for the diagnosis and therapy of gastrointestinal submucosal tumors. *Dig Endosc.* 2013;25(5):479-89. <https://doi.org/10.1111/den.12149>
4. Cho JW. Current guidelines in the management of upper gastrointestinal subepithelial tumors. *Clin Endosc.* 2016;49(3):235-40. <https://doi.org/10.5946/ce.2015.096>
5. Deprez PH, Moons L, O'toole D, Gincul R, Seisean A, Pimentel-Nunes P, et al. Endoscopic management of subepithelial lesions including neuroendocrine neoplasms: European Society of Gastro- intestinal Endoscopy (ESGE) Guideline. *Endoscopy.* 2022;54(4):412-429. <https://doi.org/10.1055/a-1751-5742>
6. Franco MC. Opinion: How to manage subepithelial lesions of the upper gastrointestinal tract? *World J Gastrointest Endosc.* 2015;7(18):1262. <https://doi.org/10.4253/wjge.v7.i18.1262>
7. Eckardt AJ, Jenssen C. Current endoscopic ultrasound-guided approach to incidental subepithelial lesions: Optimal or optional? *Ann Gastroenterol.* 2015;28(2):160-72.
8. Arango L, Sánchez A, Bautista I, Díaz C. Endosonographic Findings of Gastroduodenal Ectopic Pancreas: Experience of A Reference Center in Colombia and Literature Review. *J Gastroenterol Hepatol.* 2021;21(5):1-7. <https://doi.org/04.2021/1.1010>
9. Landi B, Palazzo L. The role of endosonography in submucosal tumours. *Best Pract Res Clin Gastroenterol.* 2009;23(5):679-701. <https://doi.org/10.1016/j.bpg.2009.05.009>
10. Hu ML, Wu KL, Changchien CS, Chuah SK, Chiu YC. Endosonographic surveillance of 1-3 cm gastric submucosal tumors originating from muscularis propria. *World J Gastroenterol.* 2017;23(12):2194-200. <https://doi.org/10.3748/wjg.v23.i12.2194>
11. Park EY, Kim GH. Diagnosis of gastric subepithelial tumors using endoscopic ultrasonography or abdominopelvic computed tomography: Which is better? *Clin Endosc.* 2019;52(6):519-20. <https://doi.org/10.5946/ce.2019.188>
12. Hohenberger P, Raut CP, Rutkowski P. Gastrointestinal stromal tumors. *Visc Med.* 2018;34(5):332-3. <https://doi.org/10.1159/000494077>
13. Sanaei O, Fernández-Esparrach G, De La Serna-Higuera C, Carrara S, Kumbhari V, El Zein MH, et al. EUS-guided 22-gauge fine needle biopsy versus single-incision with needle knife for the diagnosis of upper gastrointestinal subepithelial lesions: a randomized controlled trial. *Endosc Int Open.* 2020;08(03):E266-73. <https://doi.org/10.1055/a-1075-1900>
14. Dhaliwal A, Kolli S, Dhindsa BS, Devani K, Ramai D, Sayles H, et al. Clinical efficacy and safety of mucosal incision-assisted biopsy for the diagnosis of upper gastrointestinal subepithelial tumors: A systematic review and meta-analysis. *Ann Gastroenterol.* 2020;33(2):155-61. <https://doi.org/10.20524/aog.2020.0460>
15. de Moura DTH, McCarty TR, Jirapinyo P, Ribeiro IB, Flumignan VK, Najdawai F, et al. EUS-guided fine-needle biopsy sampling versus FNA in the diagnosis of subepithelial lesions: a large multicenter study. *Gastrointest Endosc.* 2020;92(1):108-119.e3. <https://doi.org/10.1016/j.gie.2020.02.021>
16. Trindade AJ, Benias PC, Alshelleh M, Bazarbashi AN, Tharian B, Inamdar S, et al. Fine-needle biopsy is superior to fine-needle aspiration of suspected gastrointestinal stromal tumors: a large multicenter study. *Endosc Int Open.*

- 2019;07(07):E931-6.
<https://doi.org/10.1055/a-0953-1640>
17. Kim GH, Cho YK, Kim EY, Kim HK, Cho JW, Lee TH, et al. Comparison of 22-gauge aspiration needle with 22-gauge biopsy needle in endoscopic ultrasonography-guided subepithelial tumor sampling. *Scand J Gastroenterol.* 2014;49(3):347-54.
<https://doi.org/10.3109/00365521.2013.867361>
 18. Oberg K, Couvelard A, Delle Fave G, Gross D, Grossman A, Jensen RT, et al. ENETS Consensus Guidelines for the Standards of Care in Neuroendocrine Tumors: Biochemical Markers. *Neuroendocrinology.* 2017;105(3):201-11.
<https://doi.org/10.1159/000472254>
 19. Esmo T, Sarcoma E, Working N. Gastrointestinal stromal tumours: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2014;25(Suppl 3):iii21-6.
<https://doi.org/10.1093/annonc/mdu255>
 20. Landi B, Blay JY, Bonvalot S, Brasseur M, Coindre JM, Emile JF, et al. Gastrointestinal stromal tumours (GISTs): French Intergroup Clinical Practice Guidelines for diagnosis, treatments and follow-up (SNFGE, FFCD, GERCOR, UNICANCER, SFCD, SFED, SFRO). *Dig Liver Dis.* 2019;51(9):1223-31.
<https://doi.org/10.1016/j.dld.2019.07.006>
 21. Dumonceau JM, Deprez PH, Jenssen C, Iglesias-Garcia J, Larghi A, Vanbiervliet G, et al. Indications, results, and clinical impact of endoscopic ultrasound (EUS)-guided sampling in gastroenterology: European Society of Gastrointestinal Endoscopy (ESGE) Clinical Guideline - Updated January 2017. *Endoscopy.* 2017;49(7):695-714.
<https://doi.org/10.1055/s-0043-109021>
 22. Kumar S, Chandrasekhara V, Kochman ML, Ahmad N, Attalla S, Ho IK, et al. Ligation-assisted endoscopic mucosal resection for esophageal granular cell tumors is safe and effective. *Dis Esophagus.* 2020;33(8):1-5.
<https://doi.org/10.1093/dote/doaa027>
 23. Peng W, Tan S, Huang S, Ren Y, Li H, Peng Y, et al. Efficacy and safety of submucosal tunneling endoscopic resection for upper gastrointestinal submucosal tumors with more than 1-year' follow-up: a systematic review and meta-analysis. *Scand J Gastroenterol.* 2019;54(4):397-406.
<https://doi.org/10.1080/00365521.2019.1591500>



Available in:

<https://www.redalyc.org/articulo.oa?id=337782275003>

How to cite

Complete issue

More information about this article

Journal's webpage in redalyc.org

Scientific Information System Redalyc
Diamond Open Access scientific journal network
Non-commercial open infrastructure owned by academia

Ileana Rocío Bautista-Parada, Ángel Rojas-Espinosa,
Lázaro Antonio Arango-Molano, Andrés Sánchez-Gil,
Claudia Díaz-Tobar

**Endosonographic Characteristics of Subepithelial Lesions
of the Upper Digestive Tract: Experience of a Referral
Center in Colombia**

**Características endosonográficas de las lesiones
subepiteliales del tracto digestivo superior: experiencia
de un centro de referencia en Colombia**

Revista colombiana de Gastroenterología
vol. 38, no. 3, p. 264 - 270, 2023
Asociación Colombiana de Gastroenterología,
ISSN: 0120-9957
ISSN-E: 2500-7440

DOI: <https://doi.org/10.22516/25007440.1014>