


Microbiological identification of pathogens in water from educational centers in Norte de Santander

Research Article

 Open access

Identificación microbiológica de patógenos en aguas de centros educativos de Norte de Santander

Identificação microbiológica de patógenos em água de centros educacionais do Norte de Santander



How to cite this article:

Soto Javier Andrés, Martínez-Marciales Karen Piedad. Microbiological identification of pathogens in water from educational centers in Norte de Santander. *Revista Cuidarte*. 2025;16(1):e4052. <https://doi.org/10.15649/cuidarte.4052>

Highlights

- Water is food, which is why it is necessary to consume it daily, as it allows us to stay hydrated for developing vital functions.
- Water contamination is caused by microbiological, physical, and chemical characteristics.
- For proper water care, it is necessary to verify the storage conditions.
- It is always necessary to know the quality and safety of the water; this is done through periodic microbiological and physicochemical analyses that allow the detection of any non-compliance with the criteria of suitability for human consumption.


Revista Cuidarte

Rev Cuid. 2025; 16(1): e4052

<https://doi.org/10.15649/cuidarte.4052>



E-ISSN: 2346-3414

 Javier Andrés Soto¹

 Karen Piedad Martinez-Marciales²

1. Universidad de Santander, Faculty of Medical and Health Sciences, Masira Research Institute, Cúcuta, Colombia. E-mail: jav.soto@mail.udes.edu.co
2. Universidad de Santander, Faculty of Medical and Health Sciences, Masira Research Institute, Cúcuta, Colombia. E-mail: ka.martinez@mail.udes.edu.co

Abstract

Introduction: Water is an essential resource for survival, and therefore, its quality and safety must be a priority, especially for susceptible population groups. **Objectives:** To determine the presence of mesophilic aerobes, coliforms, and *Pseudomonas* in drinking water in schools from three municipalities of Norte de Santander. **Materials and Methods:** Maintenance personnel were inquired about water storage. Samples from different sources were collected and processed using the membrane filtration method to identify aerobic mesophilic bacteria, *E. coli*, coliforms, *Pseudomonas* spp., and *Salmonella* spp. following the technical standards established for each microorganism. **Results:** Mesophilic bacteria growth was identified in 77.50% of the samples, total coliforms in 84.00%, fecal coliforms in 72.00%, and *Escherichia coli* in 21%. *Pseudomonas* spp. was also identified in 73.00% of the samples and *Salmonella* spp. in 10.50%. **Discussion:** These findings reflect non-compliance with current regulations due to the presence of indicator organisms such as mesophiles and the indicator par excellence in water quality: coliforms, a fact that is ratified by the presence of *Pseudomonas* spp. and *Salmonella* spp. **Conclusions:** The presence of these microorganisms is associated with failures in the water purification process, which allows us to expose the need for corrective actions to guarantee the microbiological quality of water and ensure health.

Keywords: Coliforms; Food Safety; Water Monitoring.

Received: May 27th, 2024

Accepted: December 9th, 2024

Published: April 30th 2025

 *Correspondence

Karen Piedad Martinez-Marciales

E-mail: ka.martinez@mail.udes.edu.co

Identificación microbiológica de patógenos en aguas de centros educativos de Norte de Santander

Resumen

Introducción: El agua es un recurso esencial para la supervivencia y por ello su calidad e inocuidad debe ser una prioridad, especialmente para grupos poblacionales susceptibles. **Objetivo:** Determinar la presencia de aerobios mesófilos, coliformes y *Pseudomonas* en agua de consumo en colegios de 3 municipios de Norte de Santander. **Materiales y Métodos:** Se realizó indagación con el personal de mantenimiento sobre el almacenamiento del agua. Se recolectaron muestras de diferentes fuentes, las cuales se procesaron por el método de filtración por membrana para identificar bacterias Aerobias Mesófilas, *E. coli*, coliformes, *Pseudomonas* spp y *Salmonella* spp. mediante las normas técnicas establecidas para cada microorganismo. **Resultados:** Se identificó el crecimiento de bacterias Mesófilas en un 77,50%, coliformes totales y fecales en un 84,00% y 72,00% respectivamente y *Escherichia coli* en un 21%. También se identificó *Pseudomonas* spp. en un 73,00% y *Salmonella* spp. con 10,50%. **Discusión:** Estos hallazgos reflejan el incumplimiento de normativa vigente debido a la presencia de bacterias indicadoras como los mesófilos, y el indicador por excelencia en calidad de aguas: los coliformes; hecho que se ratifica con *Pseudomonas* spp. y *Salmonella* spp. **Conclusiones:** La presencia de estos microorganismos está asociada a fallas en el proceso de potabilización, permitiendo exponer la necesidad de acciones correctivas que permitan garantizar la calidad microbiológica del agua y la salud.

Palabras Clave: Coliformes; Inocuidad de los Alimentos; Monitoreo del Agua.

Identificação microbiológica de patógenos em água de centros educacionais do Norte de Santander

Resumo

Introdução: A água é um recurso essencial para a sobrevivência e, portanto, sua qualidade e segurança devem ser prioridade, especialmente para grupos populacionais suscetíveis. **Objetivo:** Determinar a presença de aeróbios mesófilos, coliformes e *Pseudomonas* na água potável em escolas de 3 municípios do Norte de Santander. **Materiais e Métodos:** A equipe de manutenção foi questionada sobre o armazenamento de água. Amostras foram coletadas de diferentes fontes, as quais foram processadas pelo método de filtração por membrana para identificar bactérias aeróbicas mesófilas, *E. coli*, coliformes, *Pseudomonas* spp. e *Salmonella* spp. através das normas técnicas estabelecidas para cada microrganismo. **Resultados:** O crescimento de bactérias mesófilas foi identificado em 77,5%, coliformes totais e fecais em 84% e 72% respectivamente, e *Escherichia coli* em 21%. *Pseudomonas* spp. também foi identificada. em 73% e *Salmonella* spp. com 10,5%. **Discussão:** Estes achados refletem o não cumprimento das normas vigentes devido à presença de bactérias indicadoras como os mesófilos, e o indicador por excelência da qualidade da água: os coliformes; fato que é confirmado por *Pseudomonas* spp. e *Salmonella* spp. **Conclusões:** A presença desses microrganismos está associada a falhas no processo de purificação, permitindo expor a necessidade de ações corretivas para garantir a qualidade microbiológica da água e a saúde.

Palavras-Chave: Coliformes; Inocuidade dos Alimentos; Monitoramento da Aguas.

Introduction

Water is an essential element for survival and is a need that directly influences health. Poor-quality water for human consumption is associated with diseases that arise from the lack of access to water sources and inadequate sanitation conditions. Schools located in rural areas often serve communities with a high prevalence of diseases related to inadequate water, sanitation, and hygiene conditions¹. According to the United Nations Educational, Scientific and Cultural Organization (UNESCO), more than 800,000 people die each year from diarrhea caused by unsafe drinking water, poor sanitation, and inadequate hand hygiene, about 36% of whom are children under five years of age².

Drinking water is a basic service in any part of the world³. Ensuring its distribution and expanding access to potable water have been priorities monitored over the last few years⁴. It is necessary to validate water quality through in situ analysis of characteristics such as pH and chlorine levels, along with microbiological testing to identify any risk of contamination promptly⁵.

In Mexico, in their study on the microbiological quality of drinking water in public schools, Iñiguez et al. found that 59% of the water samples analyzed exceeded permissible limits. While some samples met the required level of chlorine, others did not. Similarly, pathogens such as *Escherichia coli* were detected in 13.6% of the samples, and strains of *Citrobacter freundii*, *Klebsiella* spp., and *Pseudomonas* spp.⁶ were also isolated. In Venezuela, a study conducted in a context comparable to that of Iñiguez et al. identified contamination in nearly all the samples analyzed. The authors concluded that the indicator microorganisms found posed a health risk⁷. Continuing with evidence from Latin America, a study in Argentina on the microbiological quality of school drinking water revealed that 73% of the samples were unfit for consumption⁸.

These studies from Central and South America reveal coincidences regarding the lack of adequate water management and storage and the growing need for integrated water management in educational institutions. This is why research on water quality and safety is being conducted, particularly in Latin America, such as the research conducted by Mucinhato in 2022⁹.

In Colombia, supplying potable water in rural areas is a challenge¹⁰. Enteric pathogens present in water pose a public health risk^{11,12}. Global regulations infer the presence of these pathogens primarily through the analysis of fecal indicator organisms and recommend targeted studies of specific enteric genera only in high-risk areas. Water monitoring is critical, whether the source is a treated drinking water system or untreated water since water characteristics depend on external factors that exert selective pressure on the natural microbial population¹³, contributing to the proliferation of pathogenic microorganisms among vulnerable populations¹⁴. In our country, the Ministry of Health and Social Protection, the Ministry of Housing, City and Territory, the Superintendency of Household Public Utilities, and the National Health Institute publish the National Water Quality Report annually. This report serves as a reference for authorities and stakeholders involved in decision-making and implementing control measures¹⁵.

The city of San José de Cúcuta reported a Water Quality Risk Index of 4.0 in 2020, placing it in the 'no risk' category¹⁶. However, the greatest microbiological risk associated with drinking water—and the primary focus of monitoring efforts—stems from a group of microorganisms known as coliforms. This group includes the genera *Escherichia*, *Edwardsiella*, *Enterobacter*, *Klebsiella*, *Serratia*, and *Citrobacter*. These microorganisms can be found in large quantities in surface waters, vegetation, and soils. Notably, spaces of *Enterobacter* and *Klebsiella* can colonize the inner surfaces of water pipes and storage tanks, where they form biofilms under conditions such as the presence of nutrients, warm temperatures, low disinfectant concentrations, and prolonged storage¹⁷. Biofilm formation in drinking water distribution systems is a widely studied topic¹⁸. Over time, bacterial communities inevitably establish themselves on the internal surfaces of pipelines, whether under conditions of high or low water flow or varying shear stress. Opportunistic pathogens can survive longer and exhibit greater resistance to chlorine when embedded within biofilms³. Coliform microorganisms and *Escherichia coli* are the best indicators of fecal

contamination. *Escherichia coli* is traditionally used to assess the microbiological quality of water and continues to serve as a key parameter in monitoring and surveillance efforts¹⁹.

Pseudomonas spp., due to its high resistance to chlorine, is considered an indicator of water disinfection efficiency²⁰. The presence of *Salmonella* spp. is particularly concerning because of its association with fecal contamination, high viability, and pathogenic potential over time²⁰. In this context, the objective of the present study was to detect the presence of *Escherichia coli*, *Salmonella* spp., *Pseudomonas* spp., and mesophilic bacteria in drinking water from 20 schools located in three municipalities of Norte de Santander, with the aim of learning about the quality of the water in these key educational settings.

Materials and Methods

Specimen collection and sampling period

This research followed a descriptive, field-based study design. A non-probabilistic sampling method was used to collect a total of 80 water samples over four sampling rounds conducted across 6 months. Educational institutions were selected primarily based on their geographic location, with strategically geo-positioned schools in the municipalities of Cúcuta, Pamplona, and Los Patios included to ensure representative coverage of both urban and rural areas within the city of Cúcuta. Another selection criterion was that the sample should represent the different water supply sources found in these institutions since not all have the same infrastructure and storage capacity, which are critical variables in the project's outcomes. The participating institutions were organized sequentially to build a database containing the results obtained from each site.

Samples were manually collected directly from the outlet valves of water dispensers and faucets after disinfection, using Whirl-Pak[®] bags, and transported to the laboratory under a cold chain at temperatures between 4 and 6 °C. Sample collection forms were duly completed.

Another methodological approach used in the study to gather information on the condition of drinking water supply sources involved surveying personnel to learn about the condition of the storage tanks. Additionally, *in situ* variables such as temperature and pH were measured using pH testers (HANNA instruments). The process began by adding 10 ml of the water sample to a container. The device was then turned on and immersed in the sample for measurement. It is important to note that the equipment was calibrated before each use.

A DPD checker disc (HANNA Instruments) was used to determine residual chlorine. Two cuvettes—one for the control and one for the sample—were each filled with 10 mL of water. A powder pillow of DPD free chlorine reagent (HI93701-0) was then added to the sample cuvette. Both cuvettes were capped and gently shaken for a few seconds before reading the result.

Microbiological analysis

Detection of *Escherichia coli* and mesophilic aerobes. The membrane filtration method was used following the Colombian Technical Standard NTC 4772²¹, employing nitrocellulose membranes with a 0.45- μ m pore size. A volume of 100 mL from each sample was filtered, and the membrane was then placed on Plate Count Agar (Merck, Frankfurt, Germany) for mesophilic aerobes growth and on Chromocult[®] Coliform Agar (Merck) for the detection of coliforms and *Escherichia coli*. Plates were incubated for 24 hours at 37 °C.

Detection of *Pseudomonas* spp. In accordance with NTC 4940²², tubes with asparagine broth (Merck, Frankfurt, Germany) were inoculated and incubated at 37 °C for 48 hours. Tubes exhibiting turbidity and fluorescence were recorded as positive. For confirmation, 0.1 mL from each positive tube was streaked

onto Cetrimide Agar (Oxoid, Hampshire, UK) and incubated at 37 °C for 24 hours. Oxidase tests were subsequently performed.

Detection of *Salmonella* spp. The method outlined in the NTC 4574²³ was used. A volume of 25 mL of the sample was added to 225 mL of peptone water (Acumedia, Madrid, Spain) and incubated at 37 °C for 24 hours. Subsequently, 0.1 mL of this pre-enrichment culture was transferred to Rappaport broth (Merck, Frankfurt, Germany) and incubated at 42 °C for 24 hours. Differential isolation was performed by streaking onto Hektoen Enteric Agar and SS Agar (Merck, Frankfurt, Germany), followed by incubation at 37 °C for 24 hours. The process concluded with biochemical confirmation.

Statistical analysis

Descriptive measures were calculated for the physicochemical and microbiological variables, and the results were summarized using contingency tables to facilitate comparisons between municipalities. The data generated in this study were stored in the Mendeley Data repository, a platform developed by Elsevier for the storage of general data²⁴.

Ethical considerations

The research project was approved by the Vice-Rector's Office for Research at the Universidad de Santander, under initiation record No. 065-16 as part of the Internal Focused Call for Research and Technological Development Projects 2016-2017, identified by code PICF0216540231765EJ. Sample collection was authorized by the legal representatives of the educational institutions.

Results

According to the results of the physical and chemical tests of water condition regarding residual chlorine levels, no residual chlorine was found in 70.00% of the schools, indicating non-compliance with the established standards for this parameter. For the pH parameter, 87.50% of the water samples from the schools complied with the established ranges in 4 of the 4 sampling rounds, while 12.50% failed to meet the standards in 4 of the 4 sampling rounds. Surface water temperature averaged 18.8°C in Pamplona, while in Cúcuta and Los Patios, the average was 26°C, with variations depending on location, altitude, and atmospheric conditions (Table 1).

Hydrosanitary characteristics of the analyzed schools

The information collected from school personnel and field observations made it possible to learn about water storage and treatment conditions within the educational institutions. From the data obtained, it was inferred that most schools are over 20 years old (95.00%). Only 40.00% of them reported information regarding improvements of water pipes, a common situation across the three municipalities. In 60.00% of the institutions, water tanks are installed above ground, mainly in the municipality of Pamplona, where all schools have this type of infrastructure. In contrast, underground tanks are more common in Los Patios (80.00%), while in the institutions located in Cúcuta, the proportion of overhead and underground tanks is similar. The predominant tank material is a combination of plastic and cement. This type of tank was observed more frequently in schools of Los Patios and Pamplona (80.00% and 60.00%, respectively). An important fact is that most of the water storage tanks in the institutions are covered with lids. Shortcomings were also identified in the frequency of maintenance, as most institutions reported performing it only once a year (55.00%), which poses a risk that is more frequent among schools located in Cúcuta. Regarding the cleaning and disinfection of the tanks, the most frequently used combination is chlorine and soap (35.00%). Most of the school personnel surveyed were able to recognize and identify the areas where water purification takes place. Only 20.00% of the schools reported conducting microbiological analysis of the water, and only 5.00% reported performing physical-chemical analyses (Table 2).

Table 1. Physicochemical characteristics of drinking water from samples collected *in situ*

Code	Physicochemical analysis <i>in situ</i>											
	Sampling 1			Sampling 2			Sampling 3			Sampling 4		
	T °C	pH	Chlorine	T °C	pH	Chlorine	T °C	pH	Chlorine	T °C	pH	Chlorine
1	17.3	7.3	0	16.7	7.3	0	21	8.6	0	21.3	9.5	0
2	16.5	7.4	1.5	17.8	7.2	0.9	17.2	8.8	0.2	18.2	8.6	1.5
3	18.6	7.1	0	19.1	7.3	0	16.9	9.3	0	18.9	8.8	0
4	16.5	6.5	0	17	6.8	0	16.2	9.7	0	17.6	9.4	0
5	16.8	7	1	17.5	7.5	1.2	17.4	8.4	0	19.2	8.4	0
6	26.9	7.2	0	29.5	8.5	0	27.2	8.6	0	25.9	8.6	0
7	10.1	8.8	0	28.3	9	0	10.7	9	0	26.6	8.7	0
8	27.2	8.8	0	28.1	8.9	0	25.7	8.4	0	25.8	9.2	0
9	27.5	8.1	2	29.1	8.2	2	25.4	7.8	2	28	8.4	2
10	27.9	8.6	0	29	8.5	0	27.5	8.5	0	26.5	8.8	0
11	26.6	7.8	0.9	27.1	9.1	0.7	25.3	8.7	1.5	27.2	7	2
12	26.3	8.2	1.5	27.1	8.6	0.5	25.7	8.6	1	26.9	7.3	1.2
13	27.2	7.1	1	26.9	8.6	1.5	26.5	8.5	1.2	27.3	8	1.2
14	26.8	8.9	0.1	27.7	9.1	0.2	26.4	8.5	0	27.8	8	0
15	25.2	5.7	0	11.9	8.5	0	27.1	8.6	0	21.9	8.2	0
16	23.6	8.5	0	24.6	9.2	0	17.1	8.8	0	21	8.9	0
17	29.9	8.8	0	28.4	9.2	0	27.1	8.7	0.5	27.1	8.6	0
18	26.5	7.7	0.3	27.1	8.6	1.2	26.1	8.5	1.7	26	8.3	0
19	21.5	8.6	0	22.9	8.4	0	25.7	8.4	0	23	8.3	0
20	27	9	0	23.4	8	0	26	9	0	25	8	0

Permissible limits (Resolution 2115 of 2007): Residual chlorine 0.3 to 2.0 mg/L pH between 6.5 to 9.0

Table 2. Water storage and treatment conditions of the schools studied

Variable	Municipality			
	Pamplona	Los Patios	Cúcuta	Total
Water piping improvements				
Yes (%)	40	40	40	40
No (%)	60	60	60	60
Water tank location				
Overhead (%)	100	20	60	60
Underground (%)	0	80	40	40
Storage tank materials				
Plastic (%)	40	20	30	30
Cement (%)	20	20	40	30
Plastic and cement (%)	60	80	30	50
Tanks are covered with lids				
Yes (%)	60	100	90	85
No (%)	40	0	100	60
Frequency of tank maintenance				
Every 4 months (%)	0	0	20	10
Every 6 months (%)	40	80	10	35
Every 12 months (%)	60	20	70	55

Variable	Municipality			
	Pamplona	Los Patios	Cúcuta	Total
Products used for cleaning/disinfection				
Soap (%)	0	20	20	15
Chlorine (%)	20	40	40	35
Soap/chlorine (%)	40	40	30	35
Other (%)	40	0	10	15
Complaints from students				
Yes (%)	20	0	10	10
No (%)	80	100	90	90
Knowing where the water is made potable				
Yes (%)	60	100	90	85
No (%)	40	0	10	15
Knowing about laboratory analysis performed				
Yes (%)	40	0	20	20
No (%)	60	100	80	80
Performing temperature, pH, and residual chlorine analysis				
Yes (%)	0	0	10	5
No (%)	100	100	90	95

Table 3. Drinking water analysis for mesophilic aerobes, total coliforms, fecal coliforms, and *Escherichia coli*

School code	Aerobes maximum 100 CFU/100ml				Total coliforms 0 CFU /100 ml				Fecal coliforms 0 CFU /100 ml				<i>E. coli</i> 0 CFU /100 ml			
	Samplings				Samplings				Samplings				Samplings			
	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4
1	10	0	44	50	0	0	32	25	0	0	1	0	0	0	0	0
2	49	116	152	120	0	0	0	0	0	0	0	0	0	0	0	0
3	16	100	16	20	0	0	0	0	0	0	0	0	0	0	0	0
4	12	160	112	105	0	24	20	18	0	12	1	0	0	0	0	0
5	*100	100	72	20	80	72	0	0	0	8	0	0	0	0	0	0
6	86	152	*100	*100	20	20	*100	8	20	12	0	0	0	0	0	0
7	*100	*100	0	*100	1200	84	0	240	0	3	0	40	0	0	0	0
8	*100	*100	*100	*100	488	336	0	48	0	0	0	4	0	0	0	0
9	37	51	*100	*100	0	0	*100	12	0	0	0	0	0	0	0	0
10	*100	*100	*100	*100	31	36	*100	80	0	2	16	0	0	0	0	0
11	*100	*100	*100	*100	30	1120	264	52	30	0	12	0	0	0	0	0
12	*100	992	*100	*100	2	280	0	0	1	0	0	0	0	0	0	0
13	*100	1920	*100	*100	12	112	16	0	4	0	0	0	0	0	0	0
14	98	*100	*100	*100	14	180	0	40	5	0	0	4	0	0	0	0
15	200	*100	*100	*100	90	100	2100	40	0	28	0	0	0	0	0	0
16	106	210	240	*100	4	10	8	*100	4	2	8	0	0	0	8	0
17	*100	18	392	*100	2	16	14	12	2	1	14	0	2	0	14	0
18	104	254	300	*100	29	0	12	8	29	10	12	0	29	0	12	0
19	86	301	1000	*100	86	5	40	35	0	5	40	35	0	0	40	0
20	*100	*100	*100	*100	*100	*100	*100	*100	20	140	182	150	0	0	0	0

* = Too numerous to count, and estimates were considered due to colony grouping

Microbiological analysis

Regarding the detection of microorganisms, this study identified atypical values across measurements, particularly in educational institutions located in Cúcuta. Based on the data, only 22.5% of the total samples analyzed met the permissible limit for microorganisms as established by current regulations (<100 colony-forming units [CFU]/100 mL)²⁵. Only 25.00% of the schools complied with the established values at least once, and only one of the 20 institutions (a school located in Pamplona) met the standard in all four sampling rounds. In contrast, 45% of the schools failed to meet the standard in any of the four rounds (Table 3).

Mesophilic aerobes

As shown in Table 3, the highest mesophilic aerobic counts—1920, 1000, and 992 CFU/100 mL—were recorded in water samples from schools 12, 13, and 19, all located in Cúcuta. These results indicate a higher level of contamination by mesophilic aerobes.

Total and fecal coliforms

Regarding the presence of coliforms, the situation is far from that observed with *E. coli*. Only two schools met the standard for total coliforms (i.e., absence in all four sampling rounds), representing just 10.00% of the schools analyzed. Considered differently, only 27.50% of the samples fell within the limits established by current regulations.

As expected, the findings regarding the presence of fecal coliforms in the drinking water of the schools studied were similar to those for total coliforms; in relation to the number of schools that showed no presence of these microorganisms in any of the 4 sampling rounds, only 15.00% of the institutions met this standard. However, from an overall perspective, 60.00% of the total samples complied with the regulations. Notably, all samples collected from school #20 exhibited particularly high CFU counts for these bacteria (Table 3).

Escherichia coli

In the cases where *E. coli* growth was observed, the samples always came from schools in the municipality of Cúcuta. The highest recorded value was 40 CFU/100 ml (Table 3).

Salmonella spp. and Pseudomonas spp

Salmonella spp. is a microorganism strongly associated with the contamination of food or water sources. It was detected in only 10% of the schools studied, and only on a single occasion. Institution #20 showed the presence of *Pseudomonas* spp. in all samples analyzed (Table 4).

Resolution 2115 of 2007²⁵ does not mandate testing for *Salmonella* spp. and *Pseudomonas* spp. However, these analyses were conducted, and these microorganisms were detected in at least one school in each municipality during at least one of the four sampling rounds. Interestingly, mirroring the analysis performed to detect fecal coliforms, institution #20 showed the presence of *Pseudomonas* spp. in all samples (Table 4). The lowest average CFU counts for this bacterium were observed in schools located in the municipality of Pamplona.

Table 4. Drinking water analysis for *Salmonella* spp. and *Pseudomonas* spp

School codes	<i>Salmonella</i> spp. Absence or presence in 25 ml				<i>Pseudomonas aeruginosa</i> bact/ml (MPN)*			
	Sampling 1	Sampling 2	Sampling 3	Sampling 4	Sampling 1	Sampling 2	Sampling 3	Sampling 4
1	A	A	A	A	0	0	9	9
2	A	A	A	A	0	0	43	43
3	A	A	A	A	0	7	43	43
4	A	A	A	A	0	0	0	0
5	A	A	A	A	0	0	0	0
6	A	A	A	A	0	0	0	0
7	A	A	A	A	0	0	21	21
8	A	A	A	A	0	0	23	0
9	A	A	A	A	0	0	23	0
10	A	A	A	A	0	0	11	0
11	A	A	A	A	0	0	21	0
12	A	A	A	A	0	0	21	0
13	A	A	A	A	0	0	93	0
14	P	A	A	A	0	0	0	21
15	P	A	A	A	7	7	0	0
16	A	A	A	A	0	2400	0	0
17	A	A	A	A	7	0	0	0
18	A	A	A	A	0	0	0	0
19	A	A	A	A	7	2400	0	0
20	A	A	A	A	7	7	7	7

* MPN: Most Probable Number. A: Absence. P: Presence.

Discussion

The World Health Organization (WHO) Guidelines for Drinking Water Quality²⁶, the European Union Council Directive 98/83/EC²⁷, sanitary regulations from each country, and in Colombia, Resolution 2115 of 2007 issued by the Ministry of Health and Social Protection²⁵, all establish that water intended for human consumption must meet specific physical, chemical, and microbiological standards. Water is fit for consumption if it is free from pathogenic microorganisms²⁸.

This study confirmed the presence of mesophiles at levels higher than those accepted, with a frequency of 77.5%. In other studies, mesophilic aerobes exceeded acceptable levels in 56% of the analyses performed by Pacheco Marza in 2023²⁹. In comparison, Obando et al. affirmed that assessing water quality involves the evaluation of its chemical, physical, and biological characteristics in relation to its natural quality, anthropogenic impacts, and uses³⁰. They also asserted that the detection of waterborne pathogens remains a global problem. Several countries in sub-Saharan Africa have adopted the WHO Guidelines for Drinking Water Quality (GDWQ) to guide the design of their sampling programs³¹.

According to Omar et al. (2017), the World Health Organization (WHO) and the International Water Association (IWA) promote a preventive risk management approach to ensure the provision of safe drinking water through the implementation of "Water Safety Plans" (WSPs)³². These plans address all the stages of the water supply³². A study by Setty et al. in 2017 assessed the outcomes of WSP implementation in drinking water systems in France and Spain by analyzing quality and compliance indicators between 2003 and 2015³³. Quality indicators included *E. coli*, fecal streptococci, total coliforms, heterotrophic plate counts, and disinfectants (residual and total chlorine). Implementing WSPs improved water quality and compliance with regulatory thresholds at most sites. Reported adverse effects were minimal, further supporting evidence that WSPs offer operational performance benefits³³. Future research should focus on identifying the factors contributing to the successful implementation of WSPs and on determining best practices. Diduch et al.³⁴ in 2016 noted that a group of microorganisms in groundwater are heterotrophic bacteria. Low concentrations of organic available matter in the environment (oligotrophic conditions) are conducive to these waterborne heterotrophic microorganisms.

In Colombia, water quality requirements are imposed to ensure control and promote efforts to meet the conditions that make water safe for consumption. Suescún in 2021 mentioned that aerobic mesophilic bacteria present low counts, which reduces the likelihood of adverse effects on both consumers and water quality. This control on microorganisms is possible thanks to the concentration of chlorine³⁵.

In this study, coliforms and *Escherichia coli* were detected at levels above the limits established by the standard (0 CFU/100 mL) in at least one of the samples taken from every school, with the lowest average levels observed in Pamplona. Research conducted by Cueva in 2021, which aimed to remove total coliforms from stored water, found that the water consumed by residents of El Rosal de San Diego contained total coliforms and was unfit for human consumption according to the Maximum Permissible Limits (MPL) established in the Water Quality Regulation for Human Consumption (DS N°031-2010- SA36)³⁶. The same findings applied to the presence of *E. coli*³⁷. According to the study conducted by Estupiñan et al. 2020, which assessed bacteriological and physical characteristics of drinking water in the municipality of Une, Cundinamarca, *Escherichia coli* counts ranged between 32 and 65 CFU/100 ml in 10 samples (76.9%), indicating non-compliance with Colombian water quality regulations. The authors argue that the presence of fecal coliforms or *E. coli* in a water system is a sign of recent contamination by human or animal feces from sewers, septic systems, or animal enclosures³⁸. Lucas et al., in Ecuador, reported that from the microbiological perspective, the presence of total and fecal coliforms posed a problem in all the localities studied, with values exceeding the permissible limits for water intended for human consumption³⁹. Similarly, in a study conducted in Ecuador, observed contamination with *Escherichia coli* serotype O157:H7 in 80% of the water samples (40/50).

There are environmental factors that cause water pollution, many of which are linked to human activity, such as discharges or waste from industrial processes, imperfect water treatment plants, leaking oil pipelines, petroleum derivatives, and improper disposal or dumping of waste, especially plastics⁴⁰. This evidence confirms the need for a systematic and continuous review of the entire water supply network system, including the source, purification processes, distribution, adequate storage, along with control and monitoring.

Among the enterobacteria capable of forming biofilms in drinking water distribution networks are the genera *Shigella* and *Salmonella*. These bacteria are responsible for diseases such as

bacillary dysentery (in the case of *Shigella*), gastroenteritis (caused by *Salmonella enterica serotype Typhimurium*)¹⁷, and typhoid fever (caused by *Salmonella enterica serotype Typhi*). One of the objectives of this study was to detect *Salmonella* spp, given their important role in posing a threat to public health. Studies conducted in 2020 indicate that bacteria within biofilms live grouped in three-dimensional cellular clusters of aerobic and anaerobic bacterial cells. While many factors influence biofilm formation, the material used to construct pipes is one of the most important, as it affects not only the number of microorganisms but also the structure of the microbial community within the pipe. In addition, it is essential to understand each of these factors and their effect as a whole, in order to find solutions to biofilm-related problems in water distribution systems^{41,42}. In 2017, Kovačić et al. successfully isolated *Salmonella enterica* subsp. *enterica* serovar *Enteritidis* from stool samples of patients who had consumed untreated water and groundwater⁴³. Similarly, Douterlo et al. in 2018 inferred that biofilms offer several advantages to microorganisms, including the sharing of nutrients and metabolic byproducts, as well as enhanced resistance to environmental stressors⁴⁴. Biofilms can directly impact the infrastructure of water distribution systems by promoting the biocorrosion of metal pipes. Additionally, they can alter the general characteristics of water, air, and soil.

A high presence of *Pseudomonas* was observed in this study. This bacterium thrives at temperature between 30°C and 37°C, enabling it to survive in a wide range of environments and making it very easy to find in aquatic and terrestrial environments. These findings highlight the need for proper maintenance in water distribution systems¹². The genus *Pseudomonas* is more resistant to chlorine than many other microorganisms and can inhibit coliforms, the most commonly used indicator of contamination. As a result, the probability of consuming water with a zero coliform count but inhibited by *Pseudomonas* increases, facilitating enteric colonization. A local source of contamination can become a regional health problem. Among the water samples analyzed, 68.75% had pH levels within the ranges established by Resolution 2115 of 2007²⁵, and the temperature varied according to the municipality. Chlorine levels were found to be low in 50% of the institutions evaluated, and 10 of the 20 institutions failed to maintain the required residual chlorine level. This situation creates an ideal environment for bacterial growth and biofilm formation and correlates directly with the increased presence of mesophilic aerobes, total coliforms, fecal coliforms, and *Pseudomonas aeruginosa*.

Although Colombian regulations do not require testing for *Pseudomonas* spp. and *Salmonella* spp. in drinking water, their inclusion in this study provides a more complete view of the microbiological quality of water in educational institutions. The aim is to set a precedent and encourage the replication of this approach in other broader scenarios. This comprehensive perspective is crucial for identifying multiple risks and designing effective mitigation strategies.

Conclusion

The study on the identification of pathogenic bacteria in drinking water in schools in Norte de Santander highlights the relevance of public health in protecting the most vulnerable populations. Implementing corrective measures based on these findings will not only support processes to improve water quality but will also enhance the health and well-being of students, thereby contributing to a safer and healthier school environment.

Water storage conditions at each school were characterized using surveys and compliance profiles aligned with Colombian regulatory requirements. The physicochemical characteristics of the water

were identified in situ by measuring residual chlorine levels, pH, and water temperature. Different data and instances of non-compliance of water quality were found that could lead to disease from the consumption of unsafe water.

Bacterial pathogens were identified using appropriate techniques, revealing failures in water reception and storage processes. The study also identified the need for greater monitoring of the filters installed for students to drink directly from these devices, as the findings of coliforms, *Escherichia coli*, *Pseudomonas* spp., and *Salmonella* spp., suggest possible cases of diseases not only among students but also among school staff.

The findings of this study serve as input for implementing public policies and health programs aimed at improving water treatment infrastructure in schools. This includes training for maintenance personnel and the adoption of more advanced water treatment technologies.

The identification of microbial contaminants underscores the importance of meeting water quality regulations. Shortcomings in the water purification process indicate the need to improve water treatment and storage practices.

Conflict of Interest: The authors declare no conflict of interest regarding any aspect of this article's publication.

Financing: This research was funded by Universidad de Santander through its internal research call in 2018.

Acknowledgment: The authors would like to thank Universidad de Santander, the Crisálida Research Group, the Orugas Research Group, and each school in Cúcuta, Los Patios, and Pamplona that participated in the study.

References

1. **Adams J, Bartram J, Chartier Y, Sims J, and W.H.** Organization. Normas sobre agua, saneamiento e higiene para escuelas en contextos de escasos recursos. 2010. Consulta: Mayo 20, 2024. Disponible en: https://iris.who.int/bitstream/handle/10665/44349/9789243547794_spa.pdf?sequence=1&isAllowed=y
2. **Koncagül E, Tran M, Connor R.** The United Nations world water development report 2021: valuing water; facts and figures.[Internet] 2021. [Cited: May 20 2024]. Available from: <https://unesdoc.unesco.org/ark:/48223/pf0000375751>
3. **Alba R. JD, Ortega S. JL, Álvarez H. G, Cervantes F. M, Ruiz B. E, Urtiz E. N, et al.** Riesgos microbiológicos en agua de bebida: una revisión clínica. *Química Viva*, 2013;12(3):215-233. <https://www.redalyc.org/pdf/863/86329278004.pdf>
4. **Orobio AOA, Osorio VOV, Rodríguez NRN, Ramírez NRN, León LLL, Hernández NHN, et al.** Problemas y desafíos que afronta Colombia respecto a la salud ambiental, un enfoque basado en el plan decenal de Salud. *Biociencias*, 2017;1(1) Disponible en: <https://hemeroteca.unad.edu.co/index.php/Biociencias/article/view/2220/2380>
5. **Martinez K, Caballero R, Díaz E, Pérez K, Suarez E.** Evaluación de la calidad del agua en restaurantes de la ciudad de San José de Cúcuta, de diferentes estratos, para contribuir con la seguridad alimentaria. *@limentech, Ciencia y Tecnología Alimentaria*, 2015;13(1):66-71 <http://dx.doi.org/10.24054/16927125.v1.n1.2015.1651>

6. **Iñiguez-Muñoz LE, Anaya-Esparza LM, Castañeda-Villanueva AA, Martínez-Esquivias F, Carvajal-Hernández M, Mendez-Robles MD.** Calidad microbiológica del agua potable utilizada en escuelas públicas de la ciudad de Tepatitlán, Jalisco. *Boletín de Ciencias Agropecuarias del ICAP*, 2022;8(15):33-39 Disponible en: <https://repository.uaeh.edu.mx/revistas/index.php/icap/article/view/7958/8718>
7. **Ravelo CU.** Estimación de la calidad microbiológica (indicadores bacteriológicos) del agua de consumo en puntos de suministro en escuelas ubicadas en municipios del estado Bolívar, Venezuela. *Guayana Moderna*, 2019.8(8):7-17 <https://revistasenlinea.saber.ucab.edu.ve/index.php/guayanamoderna/article/download/5368/4551/18173>
8. **Nicora B, Barranquero RS, Etcheverría SG, Tabera A, Quiroga M, Landa R, et al.** Evaluación integral de la gestión del agua subterránea en escuelas rurales en Tandil, Argentina. *Revista de Ciencias Ambientales*, 2021;55(1):294-316 <http://dx.doi.org/10.15359/rca.55-1.14>
9. **Mucinhato RMD, Zanin LM, Carnut L, Quintero-Flórez A, Stedefeldt E.** Inocuidad y calidad del agua y alimentación escolar: enfoques en América Latina y el Caribe. *Revista Panamericana de Salud Pública*, 202;46:e28 <https://doi.org/10.26633/RPSP.2022.28>
10. **Rodríguez Rueda V, Monroy Vargas ER, Bedoya Ríos DF, Bohórquez Joya MA.** Implementación de un sistema de tratamiento individual para mejorar la calidad del agua de la vereda Sabaneta Alta, San Francisco, Colombia. *INVENTUM*. 2022;17(32):78-85 <https://doi.org/10.26620/uniminuto.inventum.17.32.2022.78-85>
11. **Guzmán BL, Nava G, Díaz Bevilacqua P.** La calidad del agua para consumo humano y su asociación con la morbilidad en Colombia, 2008-2012. *Biomédica*. 2015;35(2):177-190. <http://dx.doi.org/10.7705/biomedica.v35i0.2511>
12. **Rodríguez SC, Asmundis CL, Ayala MT, Arzú OR.** Presencia de indicadores microbiológicos en agua para consumo humano en San Cosme (Corrientes, Argentina). *Revista veterinaria*. 2018;29(1):9-12 <http://dx.doi.org/10.30972/vet.2912779>
13. **Liu G, Zhang Y, Knibbe WJ, Feng C, Liu W, Medema G, et al.** Potential impacts of changing supply-water quality on drinking water distribution: A review. *Water research*. 2017;116:135-148 <https://doi.org/10.1016/j.watres.2017.03.031>
14. **Acevedo Osorio GO, Duran Ospina P, Betancourt CL.** Calidad microbiológica del agua en dos instituciones de salud del eje cafetero, Colombia 2015. *Archivos de Medicina (Manizales)*. 2016;16(2):246-256 <https://www.redalyc.org/pdf/2738/273849945004.pdf>
15. **Ministerio de la Protección Social.** Decreto 1575. Por el cual se establece el Sistema para la Protección y Control de la Calidad del Agua para Consumo Humano. Mayo 9 2007. Consulta: Febrero 11, 2025. Disponible en: <https://www.funcionpublica.gov.co/eva/gestornormativo/norma.php?i=30007#:~:text=El%20objeto%20del%20presente%20decreto,consumo%2C%20exceptuando%20el%20agua%20envasada>
16. **Instituto Nacional de Salud.** Boletín de Vigilancia del Agua para consumo Humano. 2020. Consulta: Mayo 20, 2024. Disponible en: <https://www.ins.gov.co/BibliotecaDigital/boletin-vigilancia-calidad-del-agua-octubre-2020.pdf>
17. **Ríos-Tobón S, Agudelo-Cadavid RM, Gutiérrez-Builes LA.** Patógenos e indicadores microbiológicos de calidad del agua para consumo humano. *Revista Facultad Nacional de Salud Pública*. 2017;35(2):236-247 <http://dx.doi.org/10.17533/udea.rfnsp.v35n2a08>
18. **Rueda Ariza JC, Martínez AJ, Calvo DC.** Efecto del desprendimiento de las biopelículas formadas en una red de acueducto sobre la calidad del agua. *Revista de Ingeniería*, 2013;(39):6-11 http://www.scielo.org.co/scielo.php?script=sci_arttext&pid=S0121-49932013000200002#:~:text=El%20desprendimiento%20de%20las%20biopel%C3%ADculas,Jorand%20%26%20Block%2C%202003
19. **Gurgel RS, da Silva LS, Silva LA.** Investigaçãõ de coliformes totais e Escherichia coli em água de consumo da comunidade Lago do limão, Município de Iranduba-AM. *Brazilian Applied Science Review*. 2020;4(4):2512-2529 <https://doi.org/10.34115/basrv4n4-028>
20. **Hernández Urzúa MÁ.** Microbiología de los alimentos: fundamentos y aplicaciones en ciencias de la salud. 2016: Editorial Médica Panamericana.
21. **ICONTEC.** Norma Técnica NTC Colombiana 4772. Calidad del agua. Detección y re-cuento de Escherichia coli y coliformes: Parte 1: Metodo de filtracion por membrana. Consulta: Febrero 11, 2025. Disponible en: <https://tienda.icontec.org/gp-calidad-del-agua-deteccion-y-recuento-de-escherichia-coli-y-de-bacterias-coliformes-parte-1-metodo-de-filtracion-por-membrana-ntc4772-2008.html>

22. **ICONTEC.** Norma Técnica NTC Colombiana 4940. Calidad del agua. Enumeración de pseudomona aeruginosa. Técnica del número más probable, NMP. 2001. Consulta: Mayo 20, 2024. Disponible en: <https://docplayer.es/159691682-Norma-tecnica-colombiana-4940.html>
23. **ICONTEC.** Norma Técnica NTC Colombiana 4574. Método horizontal para la detección de Salmonella spp. 2007. Consulta: Mayo 20, 2024. Disponible en: <https://archive.org/details/manualzilla-id-6211278>
24. **Martínez-Marciales KP, Soto JA.** Identificación microbiológica de patógenos en aguas de centros educativos de Norte de Santander. *Mendeley Data V1* 2024. <https://doi.org/10.17632/j9f93ffsxj.2>
25. **Ministerio de Ambiente, Vivienda y Desarrollo Territorial y Ministerio de Protección social.** Resolución 2115 de 2007 (junio 22). 2007. Consulta: Mayo 20, 2024. Disponible en: <https://minvivienda.gov.co/sites/default/files/normativa/2115%20-%202007.pdf>
26. **Organización Mundial de la Salud.** Guías para la calidad del agua potable. 2011. Consulta: Mayo 20, 2024. Disponible en: <https://www.who.int/es/publications/i/item/9789241549950>
27. **Consejo de la Unión Europea.** Directiva 98/83/CE del Consejo, de 3 de noviembre de 1998, relativa a la calidad de las aguas destinadas al consumo humano. *DOCE L.* 1998; 330:05-12. <https://www.boe.es/doue/1998/330/L00032-00054.pdf>
28. **Franco PA, López LA, Orozco ME.** Calidad microbiológica del agua destinada para consumo humano en siete municipios de la región Caribe Colombiana. *Ciencia Actual*; 2011;1(2):84-93 https://www.researchgate.net/publication/305193670_Microbiological_quality_of_water_intended_for_human_consumption_in_seven_municipalities_in_the_Colombian_Caribbean_region
29. **Pacheco Marza OM, Martínez Sánchez O.** Calidad microbiológica de aguas de mesa en los mercados los pozos y la ramada de la ciudad Santa Cruz junio 2022. *Ciencia Latina Revista Científica Multidisciplinar.* 2023;7(1):5491-5508. https://doi.org/10.37811/cl_rcm.v7i1.4842
30. **Obando JA, Murillo DF, Hernández CA, Torres DM, Cardenas D.** La Gobernanza del agua y su calidad en tres acueductos de Villavicencio (Colombia). *Revista espacios.* 2019;40(30):10. <https://www.revistaespacios.com/a19v40n30/a19v40n30p10.pdf>
31. **Taylor DDJ, Khush R, Peletz R and Kumpel E.** Efficacy of microbial sampling recommendations and practices in sub-Saharan Africa. *Water research.* 2018;134:115-125. <https://doi.org/10.1016/j.watres.2018.01.054>
32. **Omar YY, Parker A, Smith JA, Pollard SJ.** Risk management for drinking water safety in low and middle income countries-cultural influences on water safety plan (WSP) implementation in urban water utilities. *Science of the Total Environment.* 2017; 576:895-906. <http://dx.doi.org/10.1016/j.scitotenv.2016.10.131>
33. **Setty KE, Kayser GL, Bowling M, Enault J, Loret JF, Serra CP, et al.** Water, quality, compliance, and health outcomes among utilities implementing Water Safety Plans in France and Spain. *International Journal of Hygiene and Environmental Health.* 2017;220(3):513-530. <https://doi.org/10.1016/j.ijheh.2017.02.004>
34. **Diduch M, Polkowska Z, Namieśnik J.** The role of heterotrophic plate count bacteria in bottled water quality assessment. *Food Control,* 2016;61:188-195. <https://doi.org/10.1016/j.foodcont.2015.09.024>
35. **Suescun Gamboa KC.** Análisis microbiológico y fisicoquímico del azúcar, agua potable y agua envasada [Trabajo de Grado Pregrado] Universidad de Pamplona: Repositorio Hulago Universidad de Pamplona;2021. <http://repositoriodspace.unipamplona.edu.co/jspui/handle/20.500.12744/3612>
36. **Arriaza AE, Waight SE, Contreras CE, Ruano AB, López A, Ortiz D.** Determinación bacteriológica de la calidad del agua para consumo humano obtenida de filtros ubicados dentro del campus central de la Universidad de San Carlos de Guatemala. *Revista Científica de la Facultad de Ciencias Químicas y Farmacia.* 2015; 25(2):21-29. <https://doi.org/10.54495/Rev.Cientifica.v25i2.88>
37. **Cueva Díaz A.** Remoción de coliformes totales en agua almacenada en tanques elevados domésticos mediante nanoburbujas de aire [Tesis de Grado Pregrado Ingeniería Ambiental] Perú: Universidad Privada del Norte; 2021. <https://repositorio.upn.edu.pe/handle/11537/28062>
38. **Estupiñán-Torres SM, Ávila de Navia SL, Barrera Aguirre D, Baquero Torres R, Díaz Ibañez DA, Rodríguez Ramírez HA.** Características bacteriológicas, físicas y pH del agua de consumo humano del municipio de Une-Cundinamarca. *Nova.* 2020;18(33):101-112. <https://doi.org/10.22490/24629448.3702>

39. **Lucas L, Carreño A.** Calidad de agua de consumo humano en las comunidades Balsa en Medio, Julián y Severino de la microcuenca Carrizal, Ecuador. *Revista IIGEO*. 2018;21(42). <https://doi.org/10.15381/iigeo.v21i42.15785>
40. **Ramos Mancheno ADJ.** Efectos del consumo de agua contaminada en la calidad de vida de las personas. *Polo del Conocimiento: Revista científico-profesional*. 2024;9(1):614-632. <https://doi.org/10.23857/pc.v9i1.6396>
41. **Kilb B, Lange B, Schaule G, Flemming HC, Wingender J.** Contamination of drinking water by coliforms from biofilms grown on rubber-coated valves. *International journal of hygiene and environmental health*. 2003;206(6):563-573. <https://doi.org/10.1078/1438-4639-00258>
42. **Wingender J, Flemming HC.** Contamination potential of drinking water distribution network biofilms. *Water Science and Technology*, 2004;49(11-12):277-286 <https://doi.org/10.2166/wst.2004.0861>
43. **Kovačić A, Huljev Ž, Sušić E.** Ground water as the source of an outbreak of Salmonella Enteritidis. *Journal of epidemiology and global health*. 2017;7(3):181-184 <https://doi.org/10.1016/j.jegh.2017.05.001>
44. **Douterelo I, Fish KE, Boxall JB.** Succession of bacterial and fungal communities within biofilms of a chlorinated drinking water distribution system. *Water Research*. 2018;141:74-85. <https://doi.org/10.1016/j.watres.2018.04.058>



Available in:

<https://www.redalyc.org/articulo.oa?id=359582314006>

How to cite

Complete issue

More information about this article

Journal's webpage in redalyc.org

Scientific Information System Redalyc
Diamond Open Access scientific journal network
Non-commercial open infrastructure owned by academia

Javier Andrés Soto, Karen Piedad Martínez-Marciales
Microbiological identification of pathogens in water from educational centers in Norte de Santander
Identificación microbiológica de patógenos en aguas de centros educativos de Norte de Santander
Identificação microbiológica de patógenos em água de centros educacionais do Norte de Santander

Revista Cuidarte

vol. 16, no. 1, e4052, 2025

Programa de Enfermería, Facultad de Ciencias de la Salud,
Universidad de Santander UDES,

ISSN: 2216-0973

ISSN-E: 2346-3414

DOI: <https://doi.org/10.15649/cuidarte.4052>