

COVID-19: Factores asociados con los síntomas de ansiedad y depresión en UCI

COVID-19: Fatores associados a sintomas de ansiedade e depressão na UTI



How to cite this article:

Díaz-Mayorga Laura Andrea, Vivas-López Harol Giovanni, Torres Contreras Claudia Consuelo, Rojas Lyda Z, Serrano Norma C, Serrano-García Angie Yarlady, Parra Ortiz Margarita Rosa, Quintero-Lesmes Doris Cristina. COVID-19: Factors associated with symptoms of anxiety and depression in the ICU. *Revista Cuidarte*. 2025;16(1):e3998. <https://doi.org/10.15649/cuidarte.3998>

Highlights

- The COVID-19 pandemic posed a challenge to healthcare institutions, especially ICU personnel, exposing them to physical and mental health risks.
- Healthcare professionals, including nurses and doctors, faced anxiety and depression due to increased workloads and fear of infection.
- COVID-19 symptoms and working in both general ICUs and COVID-19 ICUs were identified as significant factors contributing to anxiety and depression among healthcare staff.
- Women in ICU healthcare roles show a slightly higher susceptibility to anxiety and depression symptoms compared to men.

Revista Cuidarte

Rev Cuid. 2025; 16(1): e3998

<https://doi.org/10.15649/cuidarte.3998>

E-ISSN: 2346-3414

- Laura Andrea Díaz-Mayorga¹
- Harol Giovanni Vivas-López²
- Claudia Consuelo Torres Contreras³
- Lyda Z. Rojas⁴
- Norma C. Serrano⁵
- Angie Yarlady Serrano-García⁶
- Margarita Rosa Parra Ortiz⁷
- Doris Cristina Quintero-Lesmes⁸

1. Universidad de Santander. Facultad de Ciencias Médicas y de la Salud. Instituto de Investigación Masira, Bucaramanga, Santander, Colombia. E-mail: lauraandima2021@gmail.com
2. Universidad de Santander. Facultad de Ciencias Médicas y de la Salud. Instituto de Investigación Masira, Bucaramanga, Santander, Colombia. E-mail: haroldgiovanni06@gmail.com
3. Universidad de Santander. Facultad de Ciencias Médicas y de la Salud. Instituto de Investigación Masira, Bucaramanga, Santander, Colombia. E-mail: clau.torres@mail.udes.edu.co
4. Fundación Cardiovascular de Colombia, Floridablanca, Santander, Colombia. E-mail: lydarojas@fcv.org
5. Fundación Cardiovascular de Colombia, Floridablanca, Santander, Colombia. E-mail: normaserrano@fcv.org
6. Fundación Cardiovascular de Colombia, Floridablanca, Santander, Colombia. E-mail: asgarcia96@gmail.com ; angieserranogarcia@fcv.org
7. Universidad Militar Nueva Granada, Bogotá DC, Colombia. E-mail: est.margarita.parra@unimilitar.edu.co
8. Fundación Cardiovascular de Colombia, Floridablanca, Santander, Colombia. E-mail: dorisquintero@fcv.org

Abstract

Introduction: The COVID-19 pandemic led to a high prevalence of anxiety and depression among healthcare personnel. **Objective:** To assess the prevalence and independent risk factors associated with anxiety and depression symptoms among healthcare staff working in Intensive Care Units (ICUs) during the COVID-19 pandemic in Bucaramanga and its metropolitan area. **Materials and Methods:** This was an analytical cross-sectional study. Anxiety and depression were measured using the Hopkins Symptom Checklist-25 (HSCL-25). Bivariate and multivariate analyses were conducted using linear regressions to investigate associated factors. **Results:** A total of 288 people were included in the study. The prevalence of anxiety and depression symptoms was 8.34% (95% CI: 5.41-12.14%). In the bivariate analysis, six factors were associated with depression and anxiety symptoms; however, only three remained in the multivariate analysis: female sex ($\beta=0.085$, 95% CI: 0.019 - 0.151), experiencing COVID-19 symptoms in the past 14 days ($\beta=0.115$, 95% CI: 0.024 - 0.205), and having worked in general ICUs and COVID-19 ICUs ($\beta=0.009$, 95% CI: 0.025 - 0.173). **Discussion:** The prevalence of anxiety and depression symptoms was considerably lower than reported in the scientific literature. **Conclusions:** In the studied population, although the prevalence of depression and anxiety symptoms was low, three independent factors were found to be statistically associated with the presence of these mental symptoms.

Keywords: Anxiety; Depression; Healthcare Personnel; Intensive Care Units; COVID-19.

Received: May 14th 2024

Accepted: November 17th 2024

Published: March 26th 2025

*Correspondence

Doris Cristina Quintero-Lesmes

E-mail: dorisquintero@fcv.org

COVID-19: Factores asociados con los síntomas de ansiedad y depresión en UCI

Resumen

Introducción: La pandemia por COVID-19 provocó una elevada prevalencia de ansiedad y depresión en el personal de la salud. **Objetivo:** Evaluar la prevalencia y los factores de riesgo asociados con los síntomas de ansiedad y depresión en personal de salud que trabaja en las unidades de cuidado intensivo (UCI) durante la pandemia de COVID-19 en Bucaramanga y su área metropolitana. **Materiales y Métodos:** Se realizó un estudio transversal analítico. La ansiedad y la depresión se midieron usando la Hopkins Symptom Checklist-25 (HSCL-25). Se realizaron análisis bivariados y multivariados con regresiones lineales para identificar los factores asociados. **Resultados:** Se incluyó a un total de 288 personas. La prevalencia de síntomas de ansiedad y depresión fue de 8.34% (IC 95% 5.41-12.14%). En el análisis bivariado, seis factores fueron asociados con los síntomas de depresión y ansiedad, sin embargo, solo tres de ellos se mantuvieron en el análisis multivariado: ser mujer ($\beta=0.085$, IC 95% 0.019 - 0.151), síntomas de COVID-19 en los últimos 14 días ($\beta= 0.115$, IC 95% 0.024 - 0.205) y haber laborado en UCI general y UCI-COVID-19 ($\beta =0.009$, IC 95% 0.025 - 0.173). **Discusión:** La prevalencia de síntomas de ansiedad y depresión fue considerablemente menor a la reportada en la literatura científica. **Conclusiones:** En la población estudiada, aunque la prevalencia de síntomas de depresión y ansiedad fue baja, se encontraron tres factores independientes asociados estadísticamente con la presencia de estos síntomas mentales.

Palabras Clave: Ansiedad; Depresión; Personal de Salud; Unidades de Cuidados Intensivos; COVID-19.

COVID-19: Fatores associados a sintomas de ansiedade e depressão na UTI

Resumo

Introdução: A pandemia de COVID-19 levou a uma alta prevalência de ansiedade e depressão entre os profissionais de saúde. **Objetivo:** Avaliar a prevalência e os fatores de risco independentes associados a sintomas de ansiedade e depressão em profissionais de saúde que trabalham em Unidades de Terapia Intensiva (UTIs) durante a pandemia de COVID-19 em Bucaramanga e sua área metropolitana. **Materiais e métodos:** Este foi um estudo transversal analítico. Ansiedade e depressão foram medidas usando a Escala Hopkins Symptom Checklist-25 (HSCL-25). Análises bivariadas e multivariadas usando regressões lineares foram conduzidas para investigar fatores associados. **Resultados:** Um total de 288 pessoas foram incluídas. A prevalência de sintomas de ansiedade e depressão foi de 8,34% (IC 95% 5,41-12,14%). Na análise bivariada, seis fatores foram associados a sintomas de depressão e ansiedade; entretanto, apenas três deles se mantiveram na análise multivariada: ser mulher ($\beta=0,085$, IC 95% 0,019 - 0,151), apresentar sintomas de COVID-19 nos últimos 14 dias ($\beta= 0,115$, IC 95% 0,024 - 0,205) e ter trabalhado em áreas combinadas de UTI e UTI-COVID-19 ($\beta =0,009$, IC 95% 0,025 - 0,173). **Discussão:** A prevalência de sintomas de ansiedade e depressão foi consideravelmente menor do que a relatada na literatura científica. **Conclusões:** Na população estudada, embora a prevalência de sintomas de depressão e ansiedade tenha sido baixa, foram encontrados três fatores independentes estatisticamente associados à presença desses sintomas mentais.

Palavras-Chave: Ansiedade; Depressão; Pessoal de Saúde; Unidades de Terapia Intensiva; COVID-19.

Introduction

Pandemics caused by infectious diseases, mainly viral infections, have experienced a resurgence in recent years. Healthcare personnel have faced highly contagious diseases such as SARS, influenza A (H1N1), avian flu, swine flu (H1N1), MERS, and, more recently, COVID-19¹. The coronavirus disease (COVID-19), caused by the SARS-CoV-2 virus, originated in Wuhan, China, in December 2019 and rapidly spread worldwide, being declared a pandemic by the WHO in March 2020². This viral disease causes severe respiratory problems, requiring hospitalization in many cases, emergency care, or even admission to an Intensive Care Unit (ICU)³. In addition to facing the risk of severe illness, healthcare personnel are also at high risk of developing mental health issues, such as anxiety and depression³⁻⁵.

Multiple factors negatively impact the mental health of ICU healthcare personnel: fear of contagion, long working hours, demanding protocols, uneven patient distribution, feelings of powerlessness, disease complexity, difficult ethical decisions, social isolation, and drastic changes in work methods^{4,6,7}. The consequences of anxiety and depression in this group include reduced quality of life, medical leave, absenteeism, and increased demand for health services⁸.

Different studies reveal a high prevalence of anxiety and depression among healthcare personnel, especially those caring for COVID-19 patients⁸. It has been observed that one in five healthcare professionals has reported depression and anxiety symptoms, which are more prevalent in women and nursing staff⁹. This has led to the urgent need to implement measures to prevent or reduce these effects³. This study is relevant and novel in its analysis of factors associated with anxiety and depression symptoms among ICU healthcare personnel in Bucaramanga, allowing for support strategies tailored to specific local needs. Theoretically, it contributes to understanding how factors such as gender, recent COVID-19 exposure, and work in both general ICUs and COVID-19 ICUs affect mental health. Methodologically, the use of the HSCL-25 ensures the reliability of results. Practically, the findings highlight the need for adapted psychological support policies, improving the healthcare workers' well-being and the responsiveness of the local healthcare system.

Institutions must implement strategies and spaces to assess the mental health status of healthcare personnel and provide psychological support. The pandemic has highlighted the importance of protecting the well-being of healthcare personnel, who are fundamental pillars in patient care. Therefore, the objective of this study was to evaluate the prevalence and independent risk factors associated with anxiety and depression symptoms among healthcare personnel working in Intensive Care Units (ICUs) during the COVID-19 pandemic in Bucaramanga and its metropolitan area.

Materials and Methods

Design and Population

An analytical cross-sectional study was conducted on frontline healthcare workers caring for patients diagnosed with COVID-19 in healthcare institutions in Bucaramanga and its metropolitan area between March and April 2021.

Eligibility Criteria

This study is a sub-analysis of a previously published research Project, *"Fortalecimiento de Capacidades en Ciencia y Tecnología del Laboratorio de Biología Molecular de la Fundación Cardiovascular de*

Colombia para atender problemáticas asociadas con agentes biológicos de alto riesgo para la salud humana en Bucaramanga/Santander¹⁰. For this study, all records of participants from the healthcare area working in the ICUs were included. At the same time, administrative or ancillary staff with shifts of less than 12 consecutive hours were excluded. No sample size calculation was performed; instead, all records of individuals meeting the inclusion criteria and not meeting the exclusion criteria were analyzed, resulting in a total of 288 participants (Figure 1). The initial study sample was selected based on convenience sampling.




Figure 1. Flowchart of participant selection

Measurements

The dependent variable was the presence of anxiety and depression symptoms, measured using the Hopkins Symptom Checklist-25 (HSCL-25), a shortened version of the Hopkins Symptom Checklist-58 (HSCL-58) was applied. This freely available scale does not require publication permission. It was developed by Derogatis et al.¹¹ and later translated and validated into Spanish by Claveria et al.¹². This checklist consists of two dimensions: the first assesses anxiety symptoms (Items 1-10), while the second focuses on depression symptoms (Items 11-25). Participants respond using a four-point Likert scale ranging from 1 to 4, where 1 represents "Not at all," 2 "A little," 3 "Quite a bit," and 4 "A lot." Originally developed as a self-report symptom inventory, the scale's total score is obtained by summing item scores and dividing by 25 (the total number of items). A total score ≥ 1.75 raises suspicion for a diagnosis of depression or anxiety. This same cut-off point is applied separately to assess anxiety and depression, using the first 10 and remaining 15 items, respectively. Regarding psychometric properties, the HSCL-25 shows good internal consistency, with a Cronbach's alpha coefficient of 0.92 ($\alpha=0.88$ for depression and $\alpha=0.83$ for anxiety). Confirmatory factor analysis revealed the presence of a global factor and two highly correlated factors (0.84). When compared with the gold standard (Composite International Diagnostic Interview-CIDI), the area under the curve (AUC) was 0.89 (95% CI: 0.86-0.93%). For a cut-off point of 1.75, the sensitivity was 88.1% (95% CI: 77.1-95.1%), and the specificity was 76.7% (95% CI: 73.3-79.8%). Furthermore, the scale demonstrated good test-retest reliability, with an intra-class correlation coefficient of 0.92 (95% CI: 0.87-0.95)¹³.

Sociodemographic characteristics were considered independent variables, which included sex, age, socioeconomic status, municipality of residence, level of education, marital status, household

size, mode of transportation to the workplace, and whether any household member had tested positive for COVID-19. Also, personal history was inquired, such as comorbidities, presence of COVID-19 symptoms in the last 14 days, COVID-19 vaccination status, number of vaccine doses received, and presence of COVID-19 antibodies. Finally, work-related aspects were studied, such as the type of healthcare institution, occupation, work area, whether a co-worker had been diagnosed with COVID-19, sharing a break room with co-workers without masks at a distance less than 1 meter for more than 15 minutes, and sharing a dining area with co-workers at a distance less than 1 meter.

Information collection procedure

First, the various healthcare institutions providing services to COVID-19 patients were identified. Subsequently, these institutions were invited to participate in the study. Upon receiving a positive response, they were asked to provide a list of frontline staff or those most exposed to the virus, along with their contact details. The selected personnel were invited to participate in the study by email. In addition, we had the support of the coordination offices of each institution to disseminate the invitation among its staff. The email invitation included a link to an informed consent form and a data collection form developed on the RedCap platform. Once participants completed the form, they obtained a signed copy of the informed consent form and a unique code for identification.

Data Set

The validated information was stored in GitLab¹⁴. GitLab is an online repository platform based on free software that follows an open-core model, offering a free version that does not require publication permissions.

Analysis of data

A descriptive analysis of the studied population's sociodemographic characteristics and personal and work history was conducted. Categorical variables were presented as absolute and relative values (percentages). In contrast, continuous variables, which did not follow a normal distribution according to the Shapiro-Wilk test, were reported as medians with first and third quartiles (Q1-Q3). Anxiety and depression symptoms were described using absolute and relative values, and the overall prevalence was calculated along with its 95% confidence interval. The dependent variable was treated as a continuous variable; therefore, bivariate and multivariate analyses were conducted using linear regression models. Initially, independent variables were selected based on the background in the literature and biological plausibility for those with an impact on the mental health of healthcare workers in the context of the pandemic. Subsequently, variables with p values ≤ 0.20 were included in multivariate analysis using a backward modeling approach. Finally, variables with p values < 0.05 were considered statistically significant, and the assumptions of the linear model were evaluated. All analyses were performed using Stata version 15 statistical software.

Ethical Considerations

The study was conducted following the guidelines of the Declaration of Helsinki and Resolution 8430 of 1993 in Colombia. It was approved by the Research Ethics Committee of the Fundación Cardiovascular de Colombia (protocol code CEI-2020-01858, December 15, 2020). All participants provided written informed consent after being fully informed about the study's objectives, potential risks, and benefits, as well as their right to withdraw at any time without repercussions. To ensure confidentiality, each participant was assigned a unique anonymized code, keeping personal identifiers separate from research data. This information was securely stored in a restricted-access database, accessible only to authorized research team members.

Results

Sociodemographic characteristics, personal and work history

A total of 288 health personnel participants who worked in the ICUs of eight health institutions in Bucaramanga and its metropolitan area were included in the study. Women accounted for 69.69% of the sample, with a median age of 34 years (Q1=27; Q3=40). It was observed that the majority of participants belonged to a middle socioeconomic status (60.85%) and mainly resided in Floridablanca (43.40%) and Bucaramanga (34.72%). Regarding education, 34.73% had technical training, while 31.60% held university degrees.

The most common mode of transportation to work was a personal vehicle (33.33%). Regarding personal history, the most frequent comorbidities were obesity (8.33%), COPD or asthma (6.60%), hypothyroidism (4.51%), arterial hypertension (3.13%), diabetes mellitus (2.08%), and dyslipidemia (1.74%). Additionally, 48.06% of participants had completed the vaccination schedule. Regarding job characteristics, 92.36% worked in private healthcare institutions, 37.15% were professional nurses, and 31.38% were nursing assistants. In addition, 21.18% worked in both the general ICU and the COVID-19 ICU. Regarding workplace exposure, 90.97% had a colleague diagnosed with COVID-19, while 83.68% reported not having shared a break room without masks with colleagues (Table 1).

Table 1. Sociodemographic characteristics, personal and work history (n=288)

Characteristics	n (%)
Sociodemographic	
Sex	
Female	200 (69.69)
Male	87 (30.31)
Age (years)*	34 (Q1=27; Q3=40)
Socioeconomic status	
Low	67 (23.84)
Middle	171 (60.85)
High	43 (15.30)
Municipality of residence	
Bucaramanga	100 (34.72)
Metropolitan area**	188 (65.28)
Education level	
Technical training	100 (34.72)
College degree	91 (31.60)
Completed postgraduate education	79 (27.43)
Incomplete postgraduate education	18 (6.25)
Marital status	
Single	139 (48.26)
Married or cohabiting	137 (47.57)
Separated, divorced, or widowed	12 (4.17)

Characteristics	n (%)
Household size*	3 (Q1=2; Q3=4)
Has anyone in your household been diagnosed with COVID-19? (Yes)	63 (21.88)
Personal history	
COVID-19 symptoms in the past 14 days (Yes)	37 (12.85)
Received COVID-19 vaccine (Yes)	206 (71.53)
COVID-19 vaccine doses received	
First dose	107 (51.94)
Second dose	99 (48.06)
Antibodies against COVID-19	
IgM (+) / IgG (+)	10 (3.47)
IgM (+) / IgG (-)	56 (19.44)
IgM (-) / IgG (+)	44 (15.28)
IgM (-) / IgG (-)	178 (61.81)
Job Characteristics	
Type of healthcare institution	
Private	266 (92.36)
Public	22 (7.64)
Occupation	
Registered nurse	107 (37.15)
Nursing assistant	99 (34.38)
Medical specialist	38 (13.19)
General practitioner	28 (9.72)
Respiratory therapist	8 (2.78)
Physiotherapist	6 (2.08)
Psychologist	2 (0.69)
Work area	
ICU	227 (78.82)
ICU and COVID-19 ICU	61 (21.18)
Has any co-worker been diagnosed with COVID-19?	
Yes	262 (90.97)
No/don't know	26 (9.03)
Have you shared a break room with co-workers without masks and at a distance of less than 1 meter for more than 15 minutes? (Yes)	47 (16.32)
Have you shared a dining area with co-workers at a distance of less than 1 meter? (Yes)	99 (34.38)

*Median (quartile 1 [Q1] and quartile 3 [Q3]). ** *Floridablanca, Piedecuesta, Girón, and Lebrija*

Prevalence of anxiety and depression

The combined prevalence of anxiety and depression symptoms was 8.34 % (95% CI: 3.41-12.14%).

Independent risk factors associated with anxiety and depression symptoms

In the bivariate analysis, six independent variables were associated with anxiety and depression: sex, as women had a higher prevalence compared to men ($p=0.025$); household size ($p=0.046$); experiencing COVID-19 symptoms in the past 14 days ($p=0.012$); not being vaccinated against COVID-19 ($p=0.033$); occupation, as registered nurses had a higher prevalence compared to medical specialists ($p=0.030$), and working in the ICU or COVID-19 ICU ($p=0.017$) (Table 2).

Table 2. Bivariate analysis of independent risk factors associated with anxiety and depression symptoms (n=288)

Characteristics	Coefficient (β)	CI 95%	p-value*
Sex			
Male	Reference		
Female	0.076	0.009 - 0.143	0.025
Age (years)*	-0.002	-0.005 - 0.002	0.333
Socioeconomic status			
Low	Reference		
Middle	-0.009	-0.083 - 0.063	0.788
High	-0.023	-0.122 - 0.076	0.650
Education level			
Incomplete/completed postgraduate	Reference		
Technical training	-0.006	-0.080 - 0.068	0.882
Academic	0.027	-0.049 - 0.103	0.489
Marital status			
Single	Reference		
Married or cohabiting	-0.033	-0.096 - 0.030	0.307
Separated, divorced or widowed	-0.080	-0.237 - 0.077	0.316
Household size	0.019	0.0003 - 0.037	0.046
Has anyone in your household been diagnosed with COVID-19?			
No	Reference		
Yes	0.027	-0.047 - 0.102	0.475
Presence of comorbidities			
No	Reference		
Yes	0.069	-0.004 - 0.143	0.064
COVID-19 symptoms in the past 14 days			
No	Reference		
Yes	0.117	0.026 - 0.208	0.012
Vaccine against COVID-19			
Yes	Reference		
No	0.074	0.006 - 0.141	0.033
Type of healthcare institution			
Private	Reference		
Public	-0.021	-0.137 - 0.095	0.720

Characteristics	Coefficient (β)	CI 95%	p-value*
Occupation			
Medical specialist	Reference		
Nursing assistant	0.044	-0.056 - 0.143	0.387
Registered nurse	0.109	0.010 - 0.207	0.030
General practitioner	0.097	-0.032 - 0.227	0.141
Respiratory therapist	0.122	-0.080 - 0.325	0.235
Physiotherapist and psychologist	0.017	-0.185 - 0.219	0.866
Work area			
ICU	Reference		
ICU and COVID-19 ICU	0.091	0.016 - 0.166	0.017
Has any co-worker been diagnosed with COVID-19?			
No/don't know	Reference		
Yes	0.032	-0.075 - 0.139	0.559
Have you shared a break room with co-workers without masks and at a distance of less than 1 meter for more than 15 minutes?			
No	Reference		
Yes	0.081	-0.002 - 0.164	0.055
Have you shared a dining area with co-workers at a distance of less than 1 meter?			
No	Reference		
Yes	0.057	-0.007 - 0.122	0.082

*p-value from a linear regression; CI: confidence interval.

Finally, in the multivariate analysis, three variables were associated with anxiety and depression symptoms: female sex (female vs. male; $\beta=0.085$; 95% CI: 0.019 - 0.151), presence of COVID-19 symptoms in the past 14 days (Yes vs. No; $\beta=0.115$; 95% CI: 0.024 - 0.205), and work area (ICU vs. ICU and COVID-19 ICU; $\beta=0.099$; 95% CI: 0.025 - 0.173) (Table 3). These variables explained 5.27% of the variability in anxiety and depression symptoms (adjusted R-squared). The model did not meet the assumptions of normality or homoscedasticity, but no collinearity was evident (variance inflation factor < 10 for all variables).

Table 3. Multivariate analysis of risk factors associated with anxiety and depression symptoms

Characteristics	Coefficient (β)	CI 95%	p-value*
Sex			
Male	Reference		
Female	0.085	0.019 - 0.151	0.011
COVID-19 symptoms in the past 14 days			
No	Reference		
Yes	0.115	0.024 - 0.205	0.014
Work area			
ICU	Reference		
ICU and COVID-19 ICU	0.099	0.025 - 0.173	0.009

*p value from a linear regression; CI: confidence interval.

Discussion

In this cross-sectional study of ICU healthcare professionals during the COVID-19 pandemic, a low prevalence of anxiety and depression symptoms was observed. Three factors were associated with these symptoms: Female sex, presence of COVID-19 symptoms in the past 14 days, and work area.

The prevalence of anxiety and depression symptoms in this study was considerably lower than that reported in other studies. A systematic review⁹ evaluating 24 studies showed a prevalence of anxiety from 14.5% to 44.6% and depression from 8.9% to 50.4%. Another systematic review and meta-analysis conducted by Salari et al.⁸ also presented prevalence rates similar to those previously reported, with a prevalence of depression of 24.3% and a prevalence of anxiety of 25.8%. Other cross-sectional studies, such as that by Peng et al.¹⁵, showed that up to 65.9% and 58.7% of ICU workers presented symptoms of depression and anxiety, respectively, with no significant difference between first-line and second-line workers in contact with COVID-19 patients. The lower prevalence we observed may be due to sample size, the instruments used to measure anxiety and depression, the sociodemographic characteristics of the study population, and the timing of the research (second wave of the COVID-19 pandemic).

Regarding the factors associated with the development of anxiety and depression symptoms, our study found that being a woman was associated with a higher prevalence rate, similar to what has been reported in the literature^{9,15}. This association is likely since the nursing staff, where women constitute the majority, had the greatest exposure to COVID-19-infected patients¹⁶. On the other hand, our study showed a significant relationship between experiencing COVID-19 symptoms in the past 14 days and the presence of anxiety and depression symptoms. This association was also observed in other studies. Motahedi et al.¹⁷ found higher anxiety levels among healthcare workers with prior COVID-19 infection. Chen et al.¹⁸ showed the presence of respiratory symptoms as an independent risk factor for suffering from anxiety and depression. The above findings could be influenced by health professionals' fear of transmitting the virus to their family members⁹.

The work area was also identified as an associated risk factor, with a higher prevalence of mental symptoms among participants who worked shifts in both the ICU and the COVID-19 ICU. This association was predictable, given the high workload experienced by these health professionals, as demonstrated by Peng et al.¹⁵, who found that longer working hours in the ICU were related to a deterioration in mental health. This situation was significantly exacerbated in referral hospitals, where patient volume was higher¹⁹. This study included institutions that faced this reality while meeting local and national healthcare demands.

Among the strengths of this study, we highlight that it included and evaluated a sample of health workers from various institutions, especially those at higher risk of presenting any mental health symptoms related to the COVID-19 pandemic, such as ICU staff, without making distinctions based on healthcare professions. However, a significant limitation is that the sample size was selected in a non-probabilistic and limited way, which could introduce selection bias and limit the ability to find associations with some factors due to insufficient statistical power. Therefore, the results of the study are not generalizable to the entire population of healthcare workers. Additionally, the model had limited explanatory power of the outcome of interest, and some assumptions were not met. This may be explained by the fact that it was a secondary data analysis, and it is possible that relevant variables were not included in the initial study; however, it is worth noting that the associated factors found have biological plausibility and are supported by findings from previous research.

Conclusion

This study identifies female sex, experiencing COVID-19 symptoms in the past 14 days, and working in specific work areas as factors associated with the risk of presenting anxiety and depression symptoms. However, in our study population, the prevalence of these symptoms was low. The higher prevalence of mental health symptoms among nursing professionals is notable and underlines the need to pay them special attention, especially to their workload. Further long-term research is needed to evaluate the impact of the COVID-19 pandemic on the mental health of healthcare workers.

Conflicts of Interest: The authors declare no conflicts of interest.

Funding: This research received funding from the Sistema General de Regalías of the Department of Santander, Colombia (BPIN code: 2020000100082), under the supervision of the Ministerio de Ciencia, Tecnología e Innovación (Minciencias).

Acknowledgments: The authors are grateful to the participants of the study.

References

1. **Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, et al.** A new coronavirus associated with human respiratory disease in China. *Nature*. 2020;579(7798):265–9. <http://dx.doi.org/10.1038/s41586-020-2008-3>
2. **World Health Organization.** WHO Director-General's opening remarks at the media briefing on COVID-19 [Internet] 2020 [Cited: March 07, 2024]. Available from: <https://www.who.int/director-general/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>
3. **Danet Danet A.** Impacto psicológico de la COVID-19 en profesionales sanitarios de primera línea en el ámbito occidental. Una revisión sistemática. *Med Clínica*. 2021;156(9):449–58. <http://dx.doi.org/10.1016/j.medcli.2020.11.009>
4. **Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al.** Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Netw Open*. 2020;3(3):e203976. <http://dx.doi.org/10.1001/jamanetworkopen.2020.3976>
5. **Wang C, Pan R, Wan X, Tan Y, Xu L, McIntyre RS, et al.** A longitudinal study on the mental health of general population during the COVID-19 epidemic in China. *Brain Behav Immun*. 2020;87:40–8. <http://dx.doi.org/10.1016/j.bbi.2020.04.028>
6. **Shanafelt T, Ripp J, Trockel M.** Understanding and Addressing Sources of Anxiety Among Health Care Professionals During the COVID-19 Pandemic. *JAMA*. 2020;323(21):2133. <http://dx.doi.org/10.1001/jama.2020.5893>
7. **Xiang YT, Yang Y, Li W, Zhang L, Zhang Q, Cheung T, et al.** Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. *Lancet Psychiatry*. 2020;7(3):228–9. [https://doi.org/10.1016/S2215-0366\(20\)30046-8](https://doi.org/10.1016/S2215-0366(20)30046-8)
8. **Salari N, Khazaie H, Hosseini-Far A, Khaledi-Paveh B, Kazemini M, Mohammadi M, et al.** The prevalence of stress, anxiety and depression within front-line healthcare workers caring for COVID-19 patients: a systematic review and meta-regression. *Hum Resour Health*. 2020;18(1):100. <http://dx.doi.org/10.1186/s12960-020-00544-1>
9. **Pappa S, Ntella V, Giannakas T, Giannakoulis VG, Papoutsis E, Katsaounou P.** Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and meta-analysis. *Brain Behav Immun*. 2020;88:901–7. <http://dx.doi.org/10.1016/j.bbi.2020.05.026>
10. **Colmenares-Mejía C, Quintero-Lesmes DC, Acosta IS, Suárez DP, Meneses L, Sopó Rincón OL, et al.** Seroprevalence of SARS-CoV-2 Infection among Vaccinated Health Workers and Hospital Staff. *Arch Clin Biomed Res*. 2022;06(02):290–295. <https://www.fortunejournals.com/articles/seroprevalence-of-sarscov2-infection-among-vaccinated-health-workers-and-hospital-staff.html>

11. **Derogatis LR, Lipman RS, Rickels K, Uhlenhuth EH, Covi L.** The Hopkins Symptom Checklist (HSCL): A self-report symptom inventory. *Behav Sci.* 1974;19(1):1–15. <http://dx.doi.org/10.1002/bs.3830190102>
12. **Clavería A, Rodríguez-Barragán M, Fernández-San-Martín MI, Nabbe P, Le Reste JY, Miguéns-Blanco I, et al.** Traducción y adaptación transcultural al español, catalán y gallego de la escala Hopkins Symptom Checklist-25 para la detección de depresión en Atención Primaria. *Aten Primaria.* 2020;52(8):539–47. <http://dx.doi.org/10.1016/j.aprim.2020.05.017>
13. **Rodríguez-Barragán M, Fernández-San-Martín MI, Clavería-Fontán A, Aldecoa-Landesa S, Casajuana-Closas M, Llobera J, et al.** Validation and Psychometric Properties of the Spanish Version of the Hopkins Symptom Checklist-25 Scale for Depression Detection in Primary Care. *Int J Environ Res Public Health.* 2021;18(15):7843. <http://dx.doi.org/10.3390/ijerph18157843>
14. **Díaz-Mayorga LA, Vivas-López HG, Torres Contreras CC, Rojas LZ, Serrano NC, Serrano-García AY, et al.** Base Datos COVID-19 - COVID-19: Factors associated with symptoms of anxiety and depression in the ICU. GitLab [Internet] 2024 [Cited: March 06, 2024]. Available from: https://gitlab.com/investigaciones1/Covid-19_UCI
15. **Peng X, Meng X, Li L, Hu C, Liu W, Liu Z, et al.** Depressive and Anxiety Symptoms of Healthcare Workers in Intensive Care Unit Under the COVID-19 Epidemic: An Online Cross-Sectional Study in China. *Front Public Health.* 2021;9:603273. <http://dx.doi.org/10.3389/fpubh.2021.603273>
16. **Liu Z, Wu J, Shi X, Ma Y, Ma X, Teng Z, et al.** Mental Health Status of Healthcare Workers in China for COVID-19 Epidemic. *Ann Glob Health.* 2020;86(1):128. <http://dx.doi.org/10.5334/aogh.3005>
17. **Motahedi S, Aghdam NF, Khajeh M, Baha R, Aliyari R, Bagheri H, et al.** Anxiety and depression among healthcare workers during COVID-19 pandemic: A cross-sectional study. *Heliyon.* 2021;7(12):e08570. <http://dx.doi.org/10.1016/j.heliyon.2021.e08570>
18. **Chen J, Liu X, Wang D, Jin Y, He M, Ma Y, et al.** Risk factors for depression and anxiety in healthcare workers deployed during the COVID-19 outbreak in China. *Soc Psychiatry Psychiatr Epidemiol.* 2021;56(1):47–55. <http://dx.doi.org/10.1007/s00127-020-01954-1>
19. **Lasalvia A, Amaddeo F, Porru S, Carta A, Tardivo S, Bovo C, et al.** Levels of burn-out among healthcare workers during the COVID-19 pandemic and their associated factors: a cross-sectional study in a tertiary hospital of a highly burdened area of north-east Italy. *BMJ Open.* 2021;11(1):e045127. <http://dx.doi.org/10.1136/bmjopen-2020-045127>



Available in:

<https://www.redalyc.org/articulo.oa?id=359582314010>

How to cite

Complete issue

More information about this article

Journal's webpage in redalyc.org

Scientific Information System Redalyc
Diamond Open Access scientific journal network
Non-commercial open infrastructure owned by academia

Laura Andrea Díaz-Mayorga, Harol Giovanni Vivas-López,
Claudia Consuelo Torres Contreras, Lyda Z. Rojas,
Norma C. Serrano, Angie Yarlady Serrano-García,
Margarita Rosa Parra Ortiz, Doris Cristina Quintero-Lesmes
**COVID-19: Factors associated with symptoms of anxiety
and depression in the ICU**
**COVID-19: Factores asociados con los síntomas de
ansiedad y depresión en UCI**
**COVID-19: Fatores associados a sintomas de ansiedade e
depressão na UTI**

Revista Cuidarte
vol. 16, no. 1, e3998, 2025
Programa de Enfermería, Facultad de Ciencias de la Salud,
Universidad de Santander UDES,
ISSN: 2216-0973
ISSN-E: 2346-3414

DOI: <https://doi.org/10.15649/cuidarte.3998>